




Comhairle Contae Mhaigh Eo
Mayo County Council



Draft

**Mayo County
Development Plan**

2021 - 2027

**Natura
Impact Report**

JBA Project Manager

Ross Bryant
Unit 24 Grove Island
Corbally,
Limerick,
Ireland

Revision history

Revision Ref/Date	Amendments	Issued to
29/10/2020 / S3-P01	Draft Report	Mayo County Council
25/11/2020 S3-P02	Draft with updated Impacts table to reflect amended Policies and Objectives (Appendix A)	Mayo County Council
21/12/2020 S3-P03	Draft with updated Impacts table to further amended Policies and Objectives (Appendix A)	Mayo County Council

Contract

This report describes work commissioned by Mayo County Council. Mayo representative for the contract was James Russell. Colm O'Leary, Hannah Mulcahy, Karen VanDorp, Ellis Hogan and Tanya Slattery of JBA Consulting carried out this work.

Prepared by Colm O'Leary BSc (Hons)

Assistant Ecologist

..... Ellis Hogan BSc (Hons) MSc

Assistant Ecologist

..... Karen Van Dorp BSc Msc

Ecologist

..... Hannah Mulcahy BSc MSc

Ecologist

..... Tanya Slattery BSc MSc MSc (Res)

Ecologist

Reviewed by Laura Thomas BA MRes PGCert CEcol MCIEEM

Principal Ecologist

Purpose

This document has been prepared as a NIR for Mayo County Council. JBA Consulting accepts no responsibility or liability for any use that is made of this document other than by the Client for the purposes for which it was originally commissioned and prepared.

JBA Consulting has no liability regarding the use of this report except to Mayo County Council.

Copyright

© JBA Consulting Engineers and Scientists Limited 2020.

Carbon footprint

A printed copy of the main text in this document will result in a carbon footprint of 58g if 100% post-consumer recycled paper is used and 73g if primary-source paper is used. These figures assume the report is printed in black and white on A4 paper and in duplex.

JBA is aiming to reduce its per capita carbon emissions.

DRAFT

Contents

1	Introduction	8
1.1	Background	8
1.2	Legislative Context	8
1.3	Appropriate Assessment Process	9
1.3.1	Stage 2 - AA	9
1.3.2	Stage 3 - Alternative Solutions	9
1.3.3	Stage 4 - IROPI	10
1.4	Methodology	10
1.4.1	Limitations and Constraints	11
2	Mayo County Development Plan	12
2.1	Background	12
2.2	Objectives of Mayo County Development Plan	12
2.3	Strategic Context	12
2.4	Format	12
2.5	Profile of County Mayo	14
2.5.1	Zone of Influence	15
3	Potential sources of impacts from the Plan	16
3.1	County	18
3.2	River Basin Level	19
3.3	Settlement Level	19
4	Alternatives to the Plan	20
5	Potential impacts of the Plan	21
5.1	Strategic Aims	21
5.2	Core Strategy	23
5.3	Location specific impacts	23
5.3.1	Balla	23
5.3.2	Ballinrobe	25
5.3.3	Ballyhaunis	26
5.3.4	Belmullet	27
5.3.5	Charlestown	27
5.3.6	Claremorris	28
5.3.7	Crossmolina	29
5.3.8	Foxford	29
5.3.9	Killala	30
5.3.10	Kiltimagh	30
5.3.11	Knock	31
5.3.12	Louisburgh	31
5.3.13	Newport	32
5.3.14	Swinford	32
5.4	Population, One-off rural Housing and Settlement plans	33
5.5	Retail Development	33
5.6	Renewable energy projects and telecommunications infrastructure	34
5.7	Water Quality	34
5.8	Climate Change	36
5.9	Air Quality	39
5.10	Agricultural development	39
5.11	Development of marine and coastal zone tourism and economic industry	40
5.12	Biodiversity, environment and green infrastructure	42
5.13	Tourism and Recreational pressures	43

5.14	Transport and related infrastructure	44
6	Cumulative and/or in-combination Impacts	46
6.1.1	River Basin Management Plan for Ireland 2018-2021	47
6.1.2	Water Services Strategic Plan 2015-2040 (Irish Water)	48
6.1.3	Plan for Forests & Freshwater Pearl Mussel in Ireland 2018 and The Pearl Mussel Project	48
6.1.4	Irish National Action Plan for the Sustainable Use of Pesticides	49
6.1.5	Ireland's Nitrates Action Programme	49
6.1.6	Climate Action Plan Ireland	49
6.1.7	Environmental River Enhancement Programme (EREP)	50
6.1.8	Green, Low-Carbon Agri-Environment Scheme (GLAS) and follow-on schemes	50
6.1.9	Food Wise 2025	50
6.1.10	National Biodiversity Action Plan 2017-2021	51
6.1.11	National Peatlands Strategy 2015	51
6.1.12	Project Ireland 2040	52
6.1.13	The National Spatial Strategy 2002-2020	52
6.1.14	Our Sustainable Future - Framework for Sustainable Development in Ireland	52
6.1.15	National Rural Development Programme 2014-2020	52
6.1.16	Sustainable Development Goals National Implementation Plan 2018 - 2020	53
6.1.17	Ireland Forestry Programme 2014-2020	53
6.2	Regional Plans and Projects with potential Cumulative and In-Combination Effects	53
6.2.1	Moy Water Management Unit Action Plan	54
6.2.2	Mayo County Council Renewable Energy Strategy 2011-2022	54
6.2.3	Destination Mayo - a strategy for the future development of tourism in County Mayo 2015-2020 (DRAFT)	54
6.2.4	County Development Plans Galway, Roscommon and Sligo	55
6.2.5	Shellfish Water Action Plans 2019	55
6.2.1	Mayo County Groundwater Protection Scheme	56
6.2.2	County Mayo Biodiversity Action Plan 2010-2015 / Strategy for the protection, conservation and promotion of Mayo's biodiversity 2020 - 2025	56
6.2.3	Mayo Local Economic and Community Plan 2015-2021	57
6.2.4	Northwest BAU (Business Area Unit) Strategic Plan 2016-2020	57
6.3	Local Plans and Projects with potential Cumulative and In-Combination Effects	57
6.3.1	Invasive Species Action Plans	57
6.3.2	Forthcoming Castlebar, Ballina and Westport Local Area Plans	58
6.3.3	All-Ireland Pollinator Plan	58
6.3.4	Planning Applications	58
6.3.5	Tourism and Leisure Activities and Associated Threats to Natura 2000 Sites in County Mayo	59
6.4	Climate Change	59
6.4.1	Climate change and biodiversity	60
6.5	Cumulative and in-combination impacts	60
6.6	Summary	61
7	Appropriate Assessment Screening	62
7.1	Introduction	62
7.2	Screening methodology	62
7.3	Study Area	62
7.4	Findings of the Screening Process	64
	Site Code	70
	Site Name	70
	Requirement for Appropriate Assessment	70

7.4.1	Qualifying Interests	73
7.4.2	Conservation Objectives	73
7.4.3	Cumulative Impact	74
7.5	Summary	74
8	Natura Impact Report	76
8.1	Potential significant effects	76
8.1.1	Loss or reduction of habitat area	76
8.1.2	Disturbance to key species	77
8.1.3	Fragmentation	86
8.1.4	Reduction in species density	87
8.1.5	Changes in key indicators of conservation value	87
8.2	Implementation routes for physical works	88
9	Mitigation Measures	90
9.1	Project Mitigation: Consenting Process	90
9.2	Project Mitigation: Pre-Construction / Detailed Design	90
9.2.1	Project Mitigation: Construction Stage	91
9.2.2	Project Monitoring	92
9.3	Summary and Conclusion	93
10	Influence of NIR on the Plan	94

DRAFT

List of Figures

Figure 1-1: The Appropriate Assessment Process (from: Appropriate Assessment of Plans and Projects in Ireland - Guidance for Planning Authorities, DEHLG, 2009).	9
Figure 5-1: Natura 2000 sites in the vicinity of Balla	25
Figure 5-2: Groundwater vulnerability	36
Figure 5-3: Location of coastal and marine Natura 2000 sites.	42

List of Tables

Table 2-1: Settlement tiers	13
Table 3-1 Summary of key impacts	16
Table 5-1: Natura 2000 sites within 15km from Balla	24
Table 5-2: Natura 2000 sites within 15km of Ballinrobe	25
Table 5-3: Natura 2000 sites within 15km of Ballyhaunis	26
Table 5-4: Natura 2000 sites within 15km of Belmullet	27
Table 5-5: Natura 2000 sites within 15km of Charlestown	28
Table 5-6: Natura 2000 sites within 15km of Claremorris	28
Table 5-7: Natura 2000 sites within 15km of Crossmolina	29
Table 5-8: Natura 2000 sites within 15km of Foxford	30
Table 5-9: Natura 2000 sites within 15km of Killala	30
Table 5-10: Natura 2000 sites within 15km of Kiltimagh	31
Table 5-11: Natura 2000 sites within 15km of Knock	31
Table 5-12: Natura 2000 sites within 15km of Louisburgh	32
Table 5-13: Natura 2000 sites within 15km of Newport	32
Table 5-14: Natura 2000 sites within 15km of Swinford	33
Table 5-15: Groundwater vulnerability in County Mayo	35
Table 5-16: Coastal and marine Natura 2000 sites	40
Table 5-17: Natura 2000 sites potentially impacted by tourism and recreation	43
Table 7-1: Natura 2000 sites screened and screening criteria	65
Table 7-2: Natura 2000 sites requiring further assessment	70
Table 8-1: Key species of the Screened-In Natura 2000 sites	78

Abbreviations

AA	Appropriate Assessment
CDP	County Development Plan
CIEEM	Chartered Institute of Ecology and Environmental Management
DoEHLG	Department of Environment, Heritage and Local Government
EC	European Communities
EclA	Ecological Impact Assessment
EIAR	Environmental Impact Assessment Report
EPA	Environmental Protection Agency
GIS	Geographical Information Systems
IROPI	Imperative Reason of Overriding Public Interest
NBDC	National Biodiversity Data Centre
NIR	Natura Impact Report
NPWS	National Parks and Wildlife Service

OPW..... Office of Public Works
QI Qualifying Interest
SAC Special Area of Conservation
SPA..... Special Protection Area
WWTP Wastewater Treatment Plant
ZOI Zone of Influence

DRAFT

1 Introduction

1.1 Background

JBA Consulting Ireland Ltd. has been commissioned by Mayo County Council to undertake a Natura Impact Report in relation to Mayo County Development Plan 2021 - 2027.

The Mayo County Development Plan 2021-2027 sets out the roadmap for the overall proper planning and sustainable development of County Mayo over the plan period. While the Plan is in place for a six-year period, it is framed having regard to the long-term development objectives of the County up until 2040, to align with national and regional spatial plans. The Plan governs the whole functional area of Mayo County Council, including the former Town Council functional areas of Castlebar, Ballina and Westport. The Plan, when adopted, will replace the Mayo County Development Plan 2014-2020.

This plan provides for, and manages, the physical, economic, and social development of the County, in the interests of the overall common good, and in compliance with environmental legislation. It includes a set of development objectives and standards, which set out where land is to be developed, and for what purposes (e.g. housing, retail, education, schools, employment, open space, amenity, conservation etc). It informs decisions on where public services such as roads and water infrastructure are to be provided and affects the type of buildings that can be constructed and how land is utilised. It influences many facets of daily economic and social life, in terms of where people can live, what services and facilities are available and where job opportunities are to be sited.

1.2 Legislative Context

Directive 92/43/EEC on the Conservation of Natural Habitats and Wild Fauna and Flora, known as the 'Habitats Directive' - provides legal protection for habitats and species of European importance. Article 2 of the Directive requires the maintenance or restoration of habitats and species of European Community interest, at a favourable conservation status. Articles 3 - 9 provide the legislative means to protect habitats and species of Community interest through the establishment and conservation of an EU-wide network of sites known as Natura 2000 sites. Natura 2000 sites are Special Areas of Conservation (SACs) designated under the Habitats Directive and Special Protection Areas (SPAs) designated under the Conservation of Wild Birds Directive (79 / 409 / EEC).

Articles 6(3) and 6(4) of the Habitats Directive set out the decision-making tests for plans or projects affecting Natura 2000 sites. Article 6(3) establishes the requirement for Appropriate Assessment:

"Any plan or project not directly connected with or necessary to the management of the site but likely to have a significant effect thereon, either individually or in combination with other plans or projects, shall be subject to appropriate assessment of its implications for the site in view of the site's conservation objectives. In the light of the conclusions of the assessment of the implications for the site and subject to the provisions of paragraph 4, the competent national authorities shall agree to the plan or project only after having ascertained that it will not adversely affect the integrity of the site concerned and, if appropriate, after having obtained the opinion of the general public."

Article 6(4) deals with the steps that should be taken when it is determined, as a result of Appropriate Assessment, that a plan/project will adversely affect a European site. Issues dealing with alternative solutions, imperative reasons of overriding public interest and compensatory measures need to be addressed in this case.

Article 6(4) states:

"If, in spite of a negative assessment of the implications for the site and in the absence of alternative solutions, a plan or project must nevertheless be carried out for imperative

reasons of overriding public interest, including those of a social or economic nature, the Member States shall take all compensatory measures necessary to ensure that the overall coherence of Natura 2000 is protected. It shall inform the Commission of the compensatory measures adopted.

Where the site concerned hosts a priority natural habitat type and / or a priority species, the only considerations which may be raised are those relating to human health or public safety, to beneficial consequences of primary importance for the environment or, further to an opinion from the Commission, to other imperative reasons of overriding public interest."

The requirements of Articles 6(3) and 6(4) of the Habitats Directive have been transposed into Irish legislation by means of the Habitats Regulations, 1997 (S.I. No. 94 of 1997) and the European Communities (Birds and Natural Habitats) Regulations 2011 (S.I. No. 477 / 2011).

1.3 Appropriate Assessment Process

Guidance on the Appropriate Assessment (AA) process was produced by the European Commission in 2002, which was subsequently developed into guidance specifically for Ireland by the Department of Environment, Heritage and Local Government (DEHLG) (2009). These guidance documents identify a staged approach to conducting an AA, as shown Figure 1-1.




Figure 1-1: The Appropriate Assessment Process (from: Appropriate Assessment of Plans and Projects in Ireland - Guidance for Planning Authorities, DEHLG, 2009).

The initial, screening stage of the Appropriate Assessment is to determine: whether the proposed plan or project is directly connected with or necessary for the management of the European designated site for nature conservation if it is likely to have a significant adverse effect on the European designated site, either individually or in combination with other plans or projects

For those sites where, potential adverse impacts are identified, either alone or in combination with other plans or projects, further assessment is necessary to determine if the proposals will have an adverse impact on the integrity of a European designated site, in view of the site's conservation objectives (i.e. the process proceeds to Stage 2).

1.3.1 Stage 2 - AA

This stage requires a more in-depth evaluation of the plan or project, and the potential direct and indirect impacts of them on the integrity and interest features of the European designated site(s), alone and in-combination with other plans and projects, taking into account the site's structure, function and conservation objectives. Where required, mitigation or avoidance measures will be suggested.

The competent authority can only agree to the plan or project after having ascertained that it will not adversely affect the integrity of the site(s) concerned. If this cannot be determined, and where mitigation cannot be achieved, then alternative solutions will need to be considered (i.e. the process proceeds to Stage 3).

1.3.2 Stage 3 - Alternative Solutions

Where adverse impacts on the integrity of Natura 2000 sites are identified, and mitigation cannot be satisfactorily implemented, alternative ways of achieving the objectives of the plan or project that avoid adverse impacts need to be considered. If none can be found, the process proceeds to Stage 4.

1.3.3 Stage 4 - IROPI

Where adverse impacts of a plan or project on the integrity of Natura 2000 sites are identified and no alternative solutions exist, the plan will only be allowed to progress if imperative reasons of overriding public interest can be demonstrated. In this case compensatory measures will be required.

The process only proceeds through each of the four stages for certain plans or projects. For example, for a plan or project, not connected with management of a site, but where no likely significant impacts are identified, the process stops at stage 1. Throughout the process, the precautionary principle must be applied, so that any uncertainties do not result in adverse impacts on a site.

This report is in support of a Natura Impact Report for Appropriate Assessment.

1.4 Methodology

The Screening for Appropriate Assessment and Natura Impact Report has been carried out with reference to the following documents:

- DoEHLG (2009 rev 2010) Appropriate Assessment of Plans and Projects in Ireland Guidance for Planning Authorities. Department of the Environment, Heritage and Local Government (DoEHLG, 2009)
- European Communities (EC) (2000) Managing Natura 2000 Sites: the provisions of Article 6 of the 'Habitats' Directive 92/43/EEC, Office for Official Publications of the European Communities, Luxembourg. European Commission (European Commission, 2000)
- EC (2002) Assessment of Plans and Projects Significantly Affecting Natura 2000 Sites: Methodological guidance on the provisions of Article 6(3) and (4) of the Habitats Directive 92/43/EEC, Office for Official Publications of the European Communities, Luxembourg. European Commission (European Commission et al., 2002)
- EC (2007) Guidance document on Article 6(4) of the 'Habitats Directive' 92/43/EEC – Clarification of the concepts of: alternative solutions, imperative reasons of overriding public interest, compensatory measures, overall coherence, opinion of the commission. European Commission (European Commission, 2007)
- CIEEM (2016). Guidelines for Ecological Impact Assessment in the UK and Ireland - Terrestrial, Freshwater and Coastal, Second Ed. (CIEEM, 2016)
- Fossitt, J., (2000). A Guide to Habitats in Ireland. The Heritage Council, Kilkenny (Fossitt, 2000)
- EC (2013) Interpretation Manual of European Union Habitats. Version EUR 27. European Commission
- National Parks and Wildlife Service (NPWS)(2019). The Status of EU Protected Habitats and Species in Ireland. National Parks and Wildlife Service, Department of the Environment, Heritage and Local Government, Dublin, Ireland

Data has been collected from a range of sources, including:

- NPWS website (www.npws.ie);

- River Basin Management Plans (RBMP) (www.wfdireland.ie);
- NBDC Biodiversity Maps (<http://maps.biodiversityireland.ie/#/Map>);
- Catchments (www.catchments.ie)
- Geological Survey Ireland (<https://www.gsi.ie>)
- The Pearl Mussel Project (www.pearlmusselproject.ie)
- Inland Fisheries Ireland (www.fisheriesireland.ie)

1.4.1 Limitations and Constraints

The screening assessment necessarily relies on some assumptions and it was inevitably subject to some limitations. These would not affect the conclusion, but the following points are recorded in order to ensure the basis of the assessment is clear:

- The County Development Plan is a higher-level plan. Therefore, this assessment cannot include any detailed assessment of any projects that may arise as a result of this Plan. Where broad level assessments are required to determine the potential for cumulative or in-combination impacts, the precautionary principle is used.
- Much of the implementation and monitoring of the plan is dependent on collaboration with stakeholders and a range of agencies. The implementation of the Plan will also depend on the economic climate, political support, council funding and availability of funding from other sources. It is not possible to predict with certainty where these can be relied upon or where unpredictable changes will occur. However, the assumption is made that in all instances legislation will be followed and further Appropriate Assessment will occur where needed by the Competent Authority.
- Any changes to the Plan will require re-assessment by a suitably qualified Ecologist to determine if further screening and assessment will be required. In all instances of this assessment, the precautionary principle is used.

2 Mayo County Development Plan

2.1 Background

The Plan consists of a number of aims and objectives, that are defined by key frameworks, strategies and policies as outlined below.

2.2 Objectives of Mayo County Development Plan

There are a significant number of objectives in the Mayo County Development Plan. These relate to each chapter of the Plan and include Housing, Economic Policy, Tourism and Recreation, Movement and Transport, Infrastructure, Sustainable Communities, Build Environment, Climate and Settlements. A full list of these and an assessment of the significance of any potential impacts that may arise as a result of the objectives is provided in Appendix A and Section 8.

2.3 Strategic Context

The Plan builds on the review of the 2014-2020 County Development Plan and is required to set out a strategy for the growth and development of County Mayo, consistent with national and regional spatial plans. The Plan must also have regard or be consistent with national development guidelines, local strategies and programmes, and must comply with both planning and environmental legislation. Since the adoption of the Mayo County Development Plan 2014-2020, significant changes have occurred in the policy and regulatory environment for land use planning in Ireland. A new national and regional hierarchy of plans are now in place.

Project Ireland 2040, which includes the National Planning Framework (NPF) and National Development Plan (NDP), sets out the Government’s policy on spatial planning over the period to 2040. The NPF influences all spatial plans from national to local levels. The Regional Spatial and Economic Strategy sets out the mechanism for delivering the NPF at a regional level. The County Development Plan will provide an overall strategy for development for the County, whilst local area plans provide more detailed planning guidance for urban areas over 5,000 populations.

The Plan is underpinned by a number of key frameworks, policies and strategies. These include The National Planning Framework, Regional Spatial and Economic Strategy (RSER), UN Sustainable Development Goals, Climate Action, Planning Guidelines and a number of Environmental Reports including this Nature Impact Report. This Natura Impact Report informs the Competent Authority for Appropriate Assessment of the overall Plan.

Further information on the background to the Plan can be found in the introduction of the Plan.

2.4 Format

The Plan consists 6 volumes. Volume 1 consists of a written statement and maps. The written statement comprises of 13 chapters. Chapters 2 to 12, inclusive, include an overarching strategic aim and a set of development objectives to achieve the strategic aim. The maps provide a graphic representation of the key elements of the Plan.

Volume 1 is set out as follows:

Chapter 1	Introduction
Chapter 2	Core and settlement strategy
Chapter 3	Housing

Chapter 4	Economic development
Chapter 5	Tourism development
Chapter 6	Movement and transport
Chapter 7	Infrastructure
Chapter 8	Sustainable communities
Chapter 9	Built environment
Chapter 10	Natural environment
Chapter 11	Climate action
Chapter 12	Renewable energy

Volume 2 contains the development management guidance document. The aim of the guidance document is to assist those wishing to develop in the County by outlining what will be required when applying for planning permission. The guidance document is based on the objectives in the Plan, government policy, planning guidance documents and established best practice.

Volume 3 comprises of Settlement Plans for towns and villages identified in Tiers 2 – 5 of the settlement hierarchy (see Table 2-1).

Table 2-1: Settlement tiers

Settlement Tiers	Settlements	Population 2016	Population increase to 2027	% Change	Area(ha.) of land required for new residential units to 2027
Tier I Strategic Growth Centres	Ballina	10,171	12,150	19.4%	33.68
	Castlebar	12,068	14,415	19.4%	39.96
	Westport	6,198	7,225	19.4%	20.48
Tier II Self-Sustaining Growth Towns	Ballinrobe	2,786	3,275	17.7%	8.42
	Ballyhaunis	2,366	2,785	17.7%	7.08
	Béal an Mhuirthead	1,019	1,200	17.7%	3.05
	Claremorris	3,687	4,340	17.7%	6.73
	Swinford	1,394	1,640	17.7%	4.15
Tier III Self-Sustaining Towns	Balla Charlestown Kiltimagh Killala Knock Louisburg Newport Foxford	7,824	8600	9.9%	13.25

Settlement Tiers	Settlements	Population 2016	Population increase to 2027	% Change	Area(ha.) of land required for new residential units to 2027
	Crossmolina				
Tier IV Rural Settlements	19 Settlements (see Table 2.x Settlement Hierarchy)	3931	4280	8.9%	6.85
Tier V Rural Villages	>25 Villages on the mainland and Islands (See Table 2.x Settlement Hierarchy)	N/A	330	N/A	6.4
Total		51,444	60,240	17.4%	
Open Countryside		79,063	85,460	7.9%	
County Total		130,507	145700	11.6%	

Volume 4 is the Plan for Ireland West Airport Knock (IWAK) Strategic Development Zone (SDZ).

Volume 5 comprises of the appendices for the plan and include Maps, the Housing Strategy, County Mayo Retail Strategy, List of Protected Views and Amenity Areas, Trees & Woodlands subject to Tree Preservation Orders, and Native Woodlands.

Volume 6 includes all three environmental reports undertaken in the preparation of the plan. The Strategic Environmental Assessment, Strategic Flood Risk Assessment and this report, the Natura Impact Report.

2.5 Profile of County Mayo

County Mayo is located on the west coast of Ireland in the province of Connacht. The County stretches from Lough Corrib and Killary Harbour in the south to the barony of Erris and Killala Bay in the north, and from the Atlantic Ocean on the west coast to the counties of Sligo and Roscommon on the east. Mayo is the third-largest County in area in the state and the second largest County in Connacht, in terms of size and population. Mayo has a population of 130,507, Castlebar being the largest town has a population of just over 12,000, followed by Ballina (10,171), Westport (6,198), Claremorris (3,687) and Ballinrobe (2,786). Over 71% of people in Mayo live in rural areas.

There is a distinct geological difference between the west and the east of the County. The west consists of large areas of extensive Atlantic blanket bog, whereas the east is largely a limestone landscape. Other key features of the County include:

- Mayo has the longest coastline in Ireland, at 1,168 km or approximately 21% of the total coastline of the State.
- The highest point in Mayo (and Connacht) is Mweelrea, at 814 m (2,671 ft).
- The River Moy in the northeast of the County is renowned for its salmon fishing.
- Ireland's largest island, Achill Island, lies off Mayo's west coast.
- Mayo has Ireland's highest cliffs at Croaghaun, Achill Island, while the Benwee Head cliffs in Kilcommon Erris drop almost perpendicularly 900 feet (270m) into the Atlantic Ocean.
- The northwest areas of County Mayo have some of the best renewable energy resources in Europe, if not the world, in terms of wind resources, ocean wave, tidal and hydroelectric resources.

The County possesses a strong community identity and a rich and diverse cultural heritage. Mayo's Gaeltacht region, comprising of three distinct areas (Erris; Achill Island; and Toormakeady), is the third largest in Ireland with 10,886 inhabitants, representing 11.5% of the total Gaeltacht population in the state. Significant progress continues to be made in the improvement of physical, social and community infrastructure in the towns and villages throughout the County, as it is recognised that social infrastructure and community development supports economic growth, provides employment opportunities and improves the well-being and quality of life for the people of Mayo.

The County contains an extensive national road network and rail network, which connects the towns of Ballina, Castlebar, Westport, Claremorris, Ballyhaunis and Foxford. The County also contains Ireland West Airport Knock (IWAK). The airport provides an international gateway to the west region, and its Strategic Development Zone (SDZ) has the potential to become a major enterprise and employment hub within the Atlantic Economic Corridor.

2.5.1 Zone of Influence

The Plan will primarily affect the County only, but a wider zone of influence (ZOI) is used for impacts relating to noise disturbance, air pollution, surface water and any supporting habitat for SAC/SPA species.

The ZOI was mainly determined to be within 15km of Mayo County. However, this was extended to 35km for surface water pathways where there are hydrological links to a Natura 2000 site with Freshwater Pearl Mussel as a Qualifying Feature. Further details of this are in Section 3.

3 Potential sources of impacts from the Plan

This plan provides for, and manages, the physical, economic, and social development of the County, in the interests of the overall common good, and in compliance with environmental legislation. It includes a set of development objectives and standards, which set out where land is to be developed, and for what purposes (e.g. housing, retail, education, schools, employment, open space, amenity, conservation etc). It informs decisions on where public services such as roads and water infrastructure are to be provided and affects the type of buildings that can be constructed and how land is utilised. It influences many facets of daily economic and social life, in terms of where people can live, what services and facilities are available and where job opportunities are to be sited.

However, this Plan does not in itself, include any specific projects or measures that could directly specify physical works to occur, without having to go through the normal consent process.

The key objectives, aims, strategies and policies from the Plan have been assessed and a summary of these are presented in Table 3-1. The potential impacts from the potential outcomes from each of these are identified in this table. A full assessment table of all objectives, aims, strategies and policies present in the overall Plan and their resulting potential impacts is provided in Appendix A. The results of the full assessment in the Appendix A does not differ from the below.

The summary objectives, aims, strategies and policies are assessed for significance of impact in Section 8.

Table 3-1 Summary of key impacts

Chapter	Summary of key impacts that may arise per Chapter as a result of their objectives, aims, strategies and/or policies
Chapter 2 – Core and strategy	Although some of the objectives can be considered to be a positive step, any works or projects arising from them could still have negative impacts upon the Natura 2000 sites and their Conservation Objectives (such as a reduction in water quality). Therefore, significantly adverse impacts upon the integrity of the Natura 2000 sites within the County and ZOI of any potential works, the works can still have negative impacts upon the Natura 2000 sites these objectives have not been screened out and developments that arise as a result of these Objectives will require further assessment.
Chapter 3 – Housing	Although some of the objectives can be considered to be a positive step in ensuring that development of Housing within the County does not significantly adversely impact upon the integrity of the Natura 2000 sites within the County and ZOI of any potential works, the works can still have negative impacts upon the Natura 2000 sites and their Conservation Objectives (such as causing species fragmentation). Therefore, these objectives have not been screened out and developments that arise as a result of these Objectives will require further assessment. Some positive impacts may through the retainment of existing buildings which may provide roosting sites for bats.
Chapter 4 – Economic Policy	Extraction of resources and further development of the County could result in direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes

Chapter	Summary of key impacts that may arise per Chapter as a result of their objectives, aims, strategies and/or policies
	in key indicators of conservation value, such as changes in water quality and quantity, and air quality.
Chapter 5 – Tourism and recreation	Any development could potentially have direct and/or indirect impact on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality. Potential negative impacts can occur by implementation of Plans that arise from this support. Therefore, these objectives have not been screened out and developments that arise as a result of these Objectives will require further assessment.
Chapter 6 – Movement and transport	Although some of the objectives can be considered to be a positive step in ensuring that development of movement and transport within the County does not significantly adversely impact upon the integrity of the Natura 2000 sites within the County and ZOI of any potential works, they works can still have negative impacts upon the Natura 2000 sites and their Conservation Objectives. Therefore, these objectives have not been screened out and developments that arise as a result of these Objectives will require further assessment.
Chapter 7 – Infrastructure	Development of infrastructure within the County could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality. Therefore, these objectives have not been screened out and developments that arise as a result of these Objectives will require further assessment.
Chapter 8 – Sustainable communities	Development within the County could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality. Therefore, these objectives have not been screened out and developments that arise as a result of these Objectives will require further assessment. Therefore, these objectives have not been screened out and developments that arise as a result of these Objectives will require further assessment.
Chapter 9 – Built environment	Development of the built environment within the County could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality. Therefore, these objectives have not been screened out and developments that arise as a result of these Objectives will require further assessment.

Chapter	Summary of key impacts that may arise per Chapter as a result of their objectives, aims, strategies and/or policies
Chapter 10 – Natural Environment	Although most objectives are determined to have a potential positive impact, some negative impacts are anticipated due to some objectives, namely NEO – 29 and NEO – 31. Therefore, these objectives have not been screened out and developments that arise as a result of these Objectives will require further assessment.
Chapter 11 - Climate	Development within the County could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality. Therefore, these objectives have not been screened out and developments that arise as a result of these Objectives will require further assessment.
Chapter 12 - Settlements	Development of settlements within the County could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality. Therefore, these objectives have not been screened out and developments that arise as a result of these Objectives will require further assessment.

Potential sources of impacts are determined based on the potential for implementation of the Plan through applicable measures, as a result of the Plan across various relevant Spatial Scales of Assessment (SSA):

- County
- The River Basin
- Each Sub-catchment of Coastal Area within the River Basin
- Settlements including towns, villages and community areas

3.1 County

Mayo County Development Plan is for the County of Mayo. Therefore, it can be assumed that it will have influences and impacts throughout the County. County Development Plan objectives, aims, strategies and policies applicable at this spatial scale include:

- Changes to land-use zoning
- Increase in one-off housing
- Potential for alternative renewable energy developments and associated infrastructure
- Quality of surface and groundwater leaving the County and discharging into Clew Bay
- Flood management and climate adaption issues
- Increase in tourism numbers and the development of amenities for tourists
- development of facilities for recreational activities including cycle lanes and walking routes

- new road infrastructure

3.2 River Basin Level

At this scale, objectives, aims, strategies and policies that could provide benefits to multiple settlements within the River Basin and other areas were considered, along with spatial and temporal coherence of measures being considered at smaller spatial scales. County Development Plan objectives, aims, strategies and policies applicable at this spatial scale include those listed above for impacts on a County level and also those that result in changes that will occur across County borders such as infrastructure developments, roads and transport routes, energy projects requiring connections and those that will cause changes to water quality and quantity and air quality.

3.3 Settlement Level

At this scale, objectives, aims, strategies and policies applicable to each particular Settlement in question were considered, even if the implementation of a given objectives, aims, strategies and policies includes the implication for works or activities outside of the Settlement, i.e., elsewhere in the County or catchment. Examples of where this might apply would be for General Settlement Objective GSO2 "Encourage and facilitate the development of the economic and tourism potential of towns in a manner that respects, builds on, protects and enhances the cultural, built heritage, natural heritage and local amenities of the town." which was assessed with each settlement in mind relevant to the Natura 2000 sites within the ZOI of each settlement.

Area specific Plan objectives, such as Objective KTP 1 for Kiltimagh "Support the reinstatement of the Western Rail Corridor, in particular actively seek and promote the re-opening of the Claremorris-Galway rail link, as well as a link to Sligo and to safeguard and protect these potential rail links from redevelopment for non-transport related purposes, in order not to preclude their future uses as an operational transportation network" were assessed on a more localised level.

4 Alternatives to the Plan

The development of the County Development Plan for Mayo included the development of a range of aims, strategies, objectives and policies and these have potential for effect at different spatial scales within the County. The SEA considered a range of alternative scenarios for accommodating future growth in Mayo provides alternatives to the aims, strategies, objectives and policies presented in the draft County Development Plan. The process of choosing the preferred aims, strategies, objectives and policies went through a number of steps starting off with a number of workshops during which alternatives were considered and questions were asked such as what will need to be provided for and what kind of environmental and ecological considerations will need to be made. During these workshops, the technical, social, economic and environmental impacts of a range of aims, objectives, policies and strategies were assessed. Notes were made on these and a multi-criteria assessment matrix was produced. Alternatives were considered at the various spatial scales. The 'Do Nothing' alternative, whereby the status quo remains and no County Development Plan, has been assessed. The aims, objectives, strategies and policies of the previous County Development Plan would remain in place as would the previous objectives dealing with spatial planning, adaptation to climate change etc. The Department of the Environment's requirements for Local Authorities to prepare climate change adaptation plans would, at a minimum, ensure that future risks to ecology and the environment through climate change would be considered in future planning. The impacts of the 'do nothing' alternative would be negative for the environmental objectives dealing with water and ecology and would have a long-term negative impact on humans and local economy particularly in the areas where Freshwater Pearl Mussel is present, or where one-off housing growth may be having an adverse impact upon the environment. This Plan directs development to town and village centres and for consideration of ecologically sensitive area. This approach allows for better protection of designated sites and achievement of WFD targets as serviced led development is directed to settlements. There are no other viable alternatives to the County Development Plan.

Section 6 of the SEA describes in technical detail, the alternatives considered at the various spatial scales.

5 Potential impacts of the Plan

As outlined in the EC guidance on the assessment of plans and projects affecting Natura 2000 sites (EC, 2002), impacts that could potentially occur through the implementation of the proposed Plan are as follows;

- Loss/ reduction of habitat area
- Disturbance to Key species
- Habitat or species fragmentation
- Reduction in species density
- Changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.

Potential impacts of the Plan are described relative to the Strategic Aims and key likely outcomes or actions of the Plan in the following sections. These impacts are then assessed in detail for significance, relevant to the Plan, in Section 8.

5.1 Strategic Aims

There are 10 Strategic Aims of the Plan, each of which relate to the advancement of the vision of the Plan. The Plan aims to build on previous successes and to strengthen Mayo's strategic advantage as a County, "to ensure that we utilise the strengths of our citizens, communities, built and natural environments, infrastructure and economic/employment development to their full potential, while combatting and adapting to climate change".

The Strategic Aims are as follows:

- **Housing** - To facilitate the sustainable growth of all rural areas, towns and villages throughout the county by seeking to accommodate, as far as possible, all persons in their choices to live in our rural areas, towns and villages; by supporting and strengthening the rural economy to sustain vibrant rural communities and by promoting consolidation and compact development of all urban and rural settlements in an attractive setting that provides a suitable mix of housing with supporting amenities; and by ensuring coordinated investment in infrastructure that will support economic competitiveness and create a high quality living and working environment.
- **Economic Development** - To promote and enhance Mayo's economic development potential through increased resilience in the county's enterprise, underpinned by talent and innovation, thereby ensuring that Mayo is best placed to excel in the long-term delivery of sustainable jobs and an enhanced standard of living for all.
- **Tourism Development** - To develop Mayo as a leading tourism destination through continued sustainable expansion of the tourism sector, with a focus on creating high-quality visitor services and the continued development and enhancement of visitor attractions and activities, capitalising on our natural and cultural heritage assets, whilst safeguarding these resources for future generations.
- **Movement and Transport** - To support increased use of sustainable modes of transport; the integration of spatial planning with transport planning; enhanced county and regional accessibility; the transition to a low carbon energy efficient transport system; and the development of a safer, more efficient, effective and connected transport system within Mayo.
- **Infrastructural Development** - To protect, improve and provide water, wastewater, surface water and flood alleviation services throughout the county, and to facilitate the provision of high quality information communication technology,

broadband, telecommunication information and electricity network required to support and enhance the key aims of best place to live and invest.

- **Sustainable Communities** - To develop and support vibrant sustainable communities in Mayo where people can live, work and enjoy access to a wide range of community, health, educational facilities and amenities, suitable to all ages and needs, in both urban and rural areas, thereby supporting a high quality of life for all to enjoy.
- **Built Environment** - To recognise and enhance the unique identity, character and built heritage of Mayo's towns, village and rural areas, to improve quality of life through the application of healthy placemaking, underpinned by good urban design with the creation of attractive public spaces that are vibrant, distinctive, safe and accessible and which promote and facilitate positive social interaction.
- **Natural Environment** - Continue to protect and enhance the county's natural heritage and biodiversity and ensure that networks of green and blue infrastructure are identified, created, protected and enhanced to provide a wide range of environmental, social and economic benefits to communities; To also improve the knowledge and understanding of the county's landscape and coast, and enhance the overall characteristics, qualities and diversity of landscape character, its sense of place and local distinctiveness in recognition of the amenity potential of the county.
- **Climate Action and Renewable Energy** - To transition to a low carbon and climate resilient county, with an emphasis on reduction in energy demand and greenhouse gas emissions, through a combination of effective mitigation and adaptation responses to climate change; in addition to maximising the opportunities to become a national leader in renewable energy generation, whilst increasing the resilience of our Natural and Cultural Capital to climate change by planning and implementing appropriate adaptation measure.
- **Settlements** - To develop Mayo's settlements as a network of attractive, livable towns and villages in the county with sustainable levels of population, employment activity and enhanced levels of amenity which encourage a high quality of life and well-being and support a sustainable synergy with the rural countryside.

While the Strategic Aims of the Plan do not directly specify any physical works that will occur, they guide the overall tone of the Plan. The Strategic Aims of the Plan are at a higher level and therefore, they do not directly result in any physical works that may cause adverse impacts. However, as they guide growth and development that requires physical works such as housing, retail and infrastructure, they will indirectly lead to projects that may potentially result in adverse impacts upon the overall integrity of any of the Natura 2000 sites that may be present within the Zone of Influence of the Plan. As this is a high-level Plan, the details to conduct assessments of such potential projects at this level and time are not possible and any such project would be required to go through the consenting process as outlined in Section 9.1.

Therefore, the Strategic Aims in themselves could result in indirect impacts upon the overall integrity of the Natura 2000 network, through loss / reduction of habitat area, disturbance of key species, habitat or species fragmentation, reduction in species density and/or changes in key indicators of conservation value such as changes in water quality and quantity, and air quality. Mitigation measures as outlined in Section 9.1 will be required.

5.2 Core Strategy

The Core Strategy is intended to set out high level policy to direct the future spatial development of County Mayo, consistent with the NPF, RSES and the Planning Acts. Under the Act, a Core Strategy focuses on:

- Defining a settlement hierarchy for the County that is consistent with the NPF and RSES.
- Transposing the prescribed NPF and RSES housing and population targets set at County level for the rural and urban centres identified within the settlement hierarchy.
- Providing an evidence-based rationale for the land proposed to be zoned for residential and mixed-use development, having regard to the capacity of existing zoned land and the phasing of development and taking account of the location of public transport and services.
- Demonstrating how the Planning Authority has had regard to the statutory Retail Planning Guidelines in setting out objectives for retail development.

As part of the Core Strategy, a settlement hierarchy has been developed, housing and population targets have been set within the hierarchy and objectives for retail development are also set. These are developed upon further in this report.

While the Core Strategy of the Plan does not directly specify any physical works that will occur, it gives structure to the overall Plan. The Core Strategies of the Plan are at a higher level and therefore, they do not directly result in any physical works that may cause adverse impacts. However, as they give structure to future plans for growth and development that will require physical works such as housing, retail and infrastructure, they will indirectly lead to projects that may potentially result in adverse impacts upon the overall integrity of any of the Natura 2000 sites that may be present within the Zone of Influence of the Plan. As this is a high-level Plan, the details to conduct assessments of any such project that may arise as a result of the Core Strategy at this level are not possible and any such project would be required to go through the consenting process as outlined in Section 9.1.

Therefore, the Core Strategies in themselves could result in indirect impacts upon the overall integrity of the Natura 2000 network, through loss / reduction of habitat area, disturbance of key species, habitat or species fragmentation, reduction in species density and/or changes in key indicators of conservation value such as changes in water quality and quantity, and air quality. The impacts of these, and of the Objectives of the Plan relative to each sector are discussed further in the below sections. These are divided into those impacts that are location specific (5.3), and those that are type specific that may incur impacts to the integrity of the Natura 2000 network throughout County Mayo and beyond the borders where relevant (Section 5.4 - 5.14).

5.3 Location specific impacts

Site specific impacts may occur to Natura 2000 sites within the ZOI of each settlement as listed in Section 5.4. These could be related to potential impacts that may arise as a result of any of the outcomes of the Plan including retail development, settlement development, increase in tourism numbers, climate change etc. These are summarised in the following sections.

5.3.1 Balla

There are 7 Natura 2000 sites within 15km from the town of Balla (See Table 5-1 and Figure 5-1). As a result of Mayo CPD there is potential for Balla Turlough SAC and the Moy River SAC to be impacted via ground and air pathways due to the close vicinity of the Natura

2000 site. Development that may occur within Balla may cause increased pressure on wastewater treatment plants and during high levels of rainfall, failing domestic wastewater treatment systems could cause a combined impact upon Balla Turlough SAC and the River Moy SAC by causing changes in in water quality, a key indicator of the conservation value of the site. Other Natura 2000 sites within the ZOI are unlikely to be impacted because they are not hydrologically connected to the site and/or are located too far from the settlement.

However, as any project that may arise as a result of the Plan will require individual appropriate assessment at a project level, this cannot be assessed at this level. Also, the Plan takes into consideration the requirements of the Habitats Directive and so, any project that is influenced by the Plan or follows the normal consent process should have sufficient assessment and mitigation in place to prevent such adverse impacts.

Table 5-1: Natura 2000 sites within 15km from Balla

Site Code	Natura site	Distance (km)
000463	Balla Turlough SAC	0.5
002081	Ballinafad SAC	4.2
002298	River Moy SAC	4.3
002179	Towerhill House SAC	9.3
000527	Moore Hall (Lough Carra) SAC	11.6
004051	Lough Carra SPA	11.4
001774	Lough Carra/Mask Complex SAC	11.4




Figure 5-1: Natura 2000 sites in the vicinity of Balla

5.3.2 Ballinrobe

There are 20 Natura 2000 sites within 15km from the town of Ballinrobe (See Table 5-2 and Appendix E.1). As a result of Mayo CPD there is potential for Lough Carra/Mask Complex SAC and Lough Mask SPA to be impacted via surface and groundwater pathways. QIs which could be impacted via these pathways include Wetland and waterbirds [A999] and Otter [1355]. Dependent on the nature and scale of any potential projects that may arise as a result of the Plan, other Natura 2000 sites are unlikely to be impacted via surface and groundwater pathways as they are located upstream of the settlement.

However, as any project that may arise as a result of the Plan will require individual Appropriate Assessment at a project level, this cannot be assessed at this level. Also, the Plan takes into consideration the requirements of the Habitats Directive and so, any project that is influenced by the Plan or follows the normal consent process should have sufficient assessment and mitigation in place to prevent such adverse impacts.

Table 5-2: Natura 2000 sites within 15km of Ballinrobe

Site Code	Natura site	Distance (km)
000461	Ardkill Turlough SAC	8.0
000474	Ballymaglancy Cave, Cong SAC	12.3

Site Code	Natura site	Distance (km)
000475	Carrowkeel Turlough SAC	10.2
000479	Cloughmoyne SAC	14.7
000480	Clyard Kettle-Holes SAC	4.8
000503	Greaghans Turlough SAC	9.2
002320	Kildun Souterrain SAC	7.7
000504	Kilglassan/Caheravoostia Turlough Complex SAC	6.9
001774	Lough Carra/Mask Complex SAC	2.9
000297	Lough Corrib SAC	10.7
001536	Mocorha Lough SAC	9.0
000527	Moore Hall (Lough Carra) SAC	9.9
002298	River Moy SAC	12.0
000525	Shrule Turlough SAC	11.1
000541	Skealaghan Turlough SAC	5.0
002179	Towerhill House SAC	10.3
004051	Lough Carra SPA	5.3
004042	Lough Corrib SPA	10.7
004062	Lough Mask SPA	4.9

5.3.3 Ballyhaunis

There are 10 Natura 2000 sites within 15km from the town of Ballyhaunis. (See Table 5-3 and Appendix E.2). Dependent on the nature and scale of any potential projects that may arise as a result of the Plan, Ballyhaunis is hydrologically connected to Lough Corrib SAC. Works that may arise as a result of the Plan has potential to impact Freshwater Pearl Mussel downstream of the settlement. Other Natura 2000 sites within the ZoI are unlikely to be impacted because they are not hydrologically connected to the site and/or are located too far from the settlement.

However, as any project that may arise as a result of the Plan will require individual appropriate assessment at a project level, this cannot be assessed at this time. Also, the Plan takes into consideration the requirements of the Habitats Directive and so, any project that is influenced by the Plan or follows the normal consent process should have sufficient assessment and mitigation in place to prevent such adverse impacts.

Table 5-3: Natura 2000 sites within 15km of Ballyhaunis

Site Code	Natura site	Distance (km)
000597	Carrowbehy/Caher Bog SAC	7.3
000600	Cloonchambers Bog SAC	11.2
000218	Coolcam Turlough SAC	10.7

Site Code	Natura site	Distance (km)
000255	Croaghill Turlough SAC	12.5
000604	Derrinea Bog SAC	9.6
000607	Errit Lough SAC	6.5
000297	Lough Corrib SAC	7.7
002298	River Moy SAC	2.5
001571	Urlaur Lakes SAC	9.1
002296	Williamstown Turloughs SAC	12.2

5.3.4 Belmullet

There are 11 Natura 2000 sites within 15km from the town of Belmullet (See Table 5-4 and Appendix E.3). As a result of Mayo CPD there is potential for Blacksod Bay/Broad Haven SPA, Mullet Peninsula SPA and Mullet/Blacksod Bay Complex SAC to be impacted via surface water, groundwater and land and air pathways due to the close vicinity of the Natura 2000 sites. Example QIs which could be impacted include Wetland and Waterbirds, Corncrake, and Otter. Other Natura 2000 sites within the ZoI are unlikely to be impacted because they are not hydrologically connected to the site and/or are located too far from the settlement.

However, as any project that may arise as a result of the Plan will require individual appropriate assessment at a project level, this cannot be assessed at this level. Also, the Plan takes into consideration the requirements of the Habitats Directive and so, any project that is influenced by the Plan or follows the normal consent process should have sufficient assessment and mitigation in place to prevent such adverse impacts.

Table 5-4: Natura 2000 sites within 15km of Belmullet

Site Code	Natura site	Distance (km)
000472	Broadhaven Bay SAC	0.0
000476	Carrowmore Lake Complex SAC	11.1
001501	Erris Head SAC	7.2
000507	Inishkea Islands SAC	14.4
000470	Mullet/Blacksod Bay Complex SAC	0.0
002998	West Connacht Coast SAC	4.6
004037	Blacksod Bay/Broad Haven SPA	0.0
004052	Carrowmore Lake SPA	11.1
004084	Inishglora and Inishkeeragh SPA	8.4
004227	Mullet Peninsula SPA	2.1
004093	Termoncarragh Lake and Annagh Machair SPA	4.4

5.3.5 Charlestown

There are 10 Natura 2000 sites within 15km from the town of Charlestown. (See Table 5-5 and Appendix E.5.) As a result of Mayo CPD there is potential for the River Moy SAC to be impacted via surface water, groundwater and land and air pathways due to the close vicinity

of the site. QIs which could potentially be impacted include Otter, Salmon, Brook Lamprey and Sea Lamprey. Due to the topography of the surrounding landscape other Natura 2000 sites are unlikely to be impacted. Other Natura 2000 sites within the ZoI are unlikely to be impacted because they are not hydrologically connected to the site and/or are located too far from the settlement.

However, as any project that may arise as a result of the Plan will require individual Appropriate Assessment at a project level, this cannot be assessed at this time. Also, the Plan takes into consideration the requirements of the Habitats Directive and so, any project that is influenced by the Plan or follows the normal consent process should have sufficient assessment and mitigation in place to prevent such adverse impacts.

Table 5-5: Natura 2000 sites within 15km of Charlestown

Site Code	Natura site	Distance (km)
001899	Cloonakillina Lough SAC	11.2
000604	Derrinea Bog SAC	14.3
000492	Dooastle Turlough SAC	11.5
000497	Flughany Bog SAC	12.2
000633	Lough Hoe Bog SAC	14.2
000634	Lough Nabrickkeagh Bog SAC	13.4
002006	Ox Mountains Bogs SAC	13.3
002298	River Moy SAC	0.0
000637	Turloughmore (Sligo) SAC	12.7
001571	Urlaur Lakes SAC	11.4

5.3.6 Claremorris

There are 11 Natura 2000 sites within 15km from the town of Claremorris (See Table 5-6 and Appendix E.4). Due to the location of Claremorris in relation to Natura 2000 sites and the surrounding topography there is unlikely to be impacts to Natura 2000 sites as a result of works associated with Mayo CPD. Other Natura 2000 sites within the ZoI are unlikely to be impacted because they are not hydrologically connected to the site and/or are located too far from the settlement.

However, as any project that may arise as a result of the Plan will require individual Appropriate Assessment at a project level, this cannot be assessed at this time. Also, the Plan takes into consideration the requirements of the Habitats Directive and so, any project that is influenced by the Plan or follows the normal consent process should have sufficient assessment and mitigation in place to prevent such adverse impacts.

Table 5-6: Natura 2000 sites within 15km of Claremorris

Site Code	Natura site	Distance (km)
000461	Ardkill Turlough SAC	14.2
000463	Balla Turlough SAC	11.2
002081	Ballinafad SAC	13.6
000475	Carrowkeel Turlough SAC	7.1
000503	Greaghans Turlough SAC	13.5

Site Code	Natura site	Distance (km)
000504	Kilglassan/Caheravoostia Turlough Complex SAC	11.2
001774	Lough Carra/Mask Complex SAC	14.1
000297	Lough Corrib SAC	11.4
000527	Moore Hall (Lough Carra) SAC	14.9
002298	River Moy SAC	8.2
002179	Towerhill House SAC	12.9

5.3.7 Crossmolina

There are 8 Natura 2000 sites within 15km from the town of Crossmolina (See Table 5-7 and Appendix E.7). As a result of Mayo CPD there is potential for impacts to occur to River Moy SAC via surface water, groundwater and land and air pathways due to the close vicinity of the Natura 2000 site. QIs which could potentially be impacted include Otter, Salmon, Brook Lamprey and Sea Lamprey. Due to the topography of the surrounding landscape other Natura 2000 sites are unlikely to be impacted. Other Natura 2000 sites within the ZOI are unlikely to be impacted because they are not hydrologically connected to the site and/or are located too far from the settlement.

However, as any project that may arise as a result of the Plan will require individual Appropriate Assessment at a project level, this cannot be assessed at this time. Also, the Plan takes into consideration the requirements of the Habitats Directive and so, any project that is influenced by the Plan or follows the normal consent process should have sufficient assessment and mitigation in place to prevent such adverse impacts.

Table 5-7: Natura 2000 sites within 15km of Crossmolina

Site Code	Natura site	Distance (km)
001922	Bellacorick Bog Complex SAC	11.9
000466	Bellacorick Iron Flush SAC	14.9
000458	Killala Bay/Moy Estuary SAC	13.1
002177	Lough Dahybaun SAC	13.5
002144	Newport River SAC	12.8
002298	River Moy SAC	0.0
004228	Lough Conn and Lough Cullin SPA	1.1
004036	Killala Bay/Moy Estuary SPA	11.9

5.3.8 Foxford

There are three Natura 2000 sites within 15km from the town of Foxford (See Table 5-8 and Appendix E.6). As a result of Mayo CPD there is potential for impacts to occur to River Moy SAC via surface water, groundwater and land and air pathways due to the close vicinity of the Natura 2000 site. QIs which could potentially be impacted include Otter, Salmon, Brook Lamprey and Sea Lamprey. Due to the topography of the surrounding landscape other Natura 2000 sites are unlikely to be impacted. Other Natura 2000 sites within the ZOI are unlikely to be impacted because they are not hydrologically connected to the site and/or are located too far from the settlement.

However, as any project that may arise as a result of the Plan will require individual Appropriate Assessment at a project level, this cannot be assessed at this time. Also, the Plan takes into consideration the requirements of the Habitats Directive and so, any project that is influenced by the Plan or follows the normal consent process should have sufficient assessment and mitigation in place to prevent such adverse impacts.

Table 5-8: Natura 2000 sites within 15km of Foxford

Site Code	Natura site	Distance (km)
000633	Lough Hoe Bog SAC	8.4
002298	River Moy SAC	0.0
004228	Lough Conn and Lough Cullin SPA	2.4

5.3.9 Killala

There are seven Natura 2000 sites within 15km from the town of Killala (See Table 5-9 and Appendix E.11). As a result of Mayo CPD there is potential for impacts to occur to Killala Bay/Moy Estuary SPA and Killala Bay/Moy Estuary SAC via surface water, groundwater and land and air pathways due to the close vicinity of the Natura 2000 site. QIs which could potentially be impacted include Otter and Harbour Seal. Due to the topography of the surrounding landscape other Natura 2000 sites are unlikely to be impacted. Other Natura 2000 sites within the ZoI are unlikely to be impacted because they are not hydrologically connected to the site and/or are located too far from the settlement.

However, as any project that may arise as a result of the Plan will require individual Appropriate Assessment at a project level, this cannot be assessed at this time. Also, the Plan takes into consideration the requirements of the Habitats Directive and so, any project that is influenced by the Plan or follows the normal consent process should have sufficient assessment and mitigation in place to prevent such adverse impacts.

Table 5-9: Natura 2000 sites within 15km of Killala

Site Code	Natura site	Distance (km)
004036	Killala Bay/Moy Estuary SPA	0.0
000458	Killala Bay/Moy Estuary SAC	0.0
004228	Lough Conn and Lough Cullin SPA	12.6
001922	Bellacorick Bog Complex SAC	11.2
000500	Glenamoy Bog Complex SAC	12.1
000516	Lackan Saltmarsh and Kilcummin Head SAC	5.2
002298	River Moy SAC	9.9

5.3.10 Kiltimagh

There are four Natura 2000 sites within 15km from the town of Kiltimagh (See Table 5-10 and Appendix E.9). As a result of Mayo CPD there is potential for impacts to occur to River Moy SAC via surface water, groundwater and land and air pathways due to the close vicinity of the Natura 2000 site. QIs which could potentially be impacted include Otter, Salmon, Brook Lamprey and Sea Lamprey. Due to the topography of the surrounding landscape other Natura 2000 sites are unlikely to be impacted. Other Natura 2000 sites within the ZoI are unlikely to be impacted because they are not hydrologically connected to the site and/or are located too far from the settlement.

However, as any project that may arise as a result of the Plan will require individual Appropriate Assessment at a project level, this cannot be assessed at this time. Also, the Plan takes into consideration the requirements of the Habitats Directive and so, any project that is influenced by the Plan or follows the normal consent process should have sufficient assessment and mitigation in place to prevent such adverse impacts.

Table 5-10: Natura 2000 sites within 15km of Kiltimagh

Site Code	Natura site	Distance (km)
000463	Balla Turlough SAC	8.8
002081	Ballinafad SAC	14.1
002298	River Moy SAC	0.5
001571	Urlaur Lakes SAC	13.9

5.3.11 Knock

There are five Natura 2000 sites within 15km from the town of Knock (See Table 5-11 and Appendix E.8). As a result of Mayo CPD there is potential for impacts to occur to River Moy SAC via surface water, groundwater and land and air pathways due to the close vicinity of the Natura 2000 site. QIs which could potentially be impacted include Otter, Salmon, Brook Lamprey and Sea Lamprey. Due to the topography of the surrounding landscape other Natura 2000 sites are unlikely to be impacted. Other Natura 2000 sites within the ZoI are unlikely to be impacted because they are not hydrologically connected to the site and/or are located too far from the settlement.

However, as any project that may arise as a result of the Plan will require individual Appropriate Assessment at a project level, this cannot be assessed at this time. Also, the Plan takes into consideration the requirements of the Habitats Directive and so, any project that is influenced by the Plan or follows the normal consent process should have sufficient assessment and mitigation in place to prevent such adverse impacts.

Table 5-11: Natura 2000 sites within 15km of Knock

Site Code	Natura site	Distance (km)
000463	Balla Turlough SAC	12.5
000607	Errit Lough SAC	13.4
000297	Lough Corrib SAC	13.6
002298	River Moy SAC	1.0
001571	Urlaur Lakes SAC	11.6

5.3.12 Louisburgh

There are nine Natura 2000 sites within 15km from the town of Louisburgh (See Table 5-12 and Appendix E.12). As a result of Mayo CPD there is potential for impacts to occur to West Connacht Coast SAC via surface water, groundwater and land and air pathways due to the close vicinity of the Natura 2000 site. Additionally, there is potential for impacts to Oldhead Wood SAC via land and air pathways. Other Natura 2000 sites within the ZoI are unlikely to be impacted because they are not hydrologically connected to the site and/or are located too far from the settlement.

However, as any project that may arise as a result of the Plan will require individual Appropriate Assessment at a project level, this cannot be assessed at this time. Also, the Plan takes into consideration the requirements of the Habitats Directive and so, any project

that is influenced by the Plan or follows the normal consent process should have sufficient assessment and mitigation in place to prevent such adverse impacts.

Table 5-12: Natura 2000 sites within 15km of Louisburgh

Site Code	Natura site	Distance (km)
004136	Clare Island SPA	11.8
002243	Clare Island Cliffs SAC	14.3
001482	Clew Bay Complex SAC	9.3
000485	Corraun Plateau SAC	12.2
000484	Cross Lough (Killadoon) SAC	8.5
001529	Lough Cahasy, Lough Baun And Roonah Lough SAC	5.6
001932	Mweelrea/Sheeffry/Erriff Complex SAC	4.5
000532	Oldhead Wood SAC	1.4
002998	West Connacht Coast SAC	1.3

5.3.13 Newport

There are five Natura 2000 sites within 15km from the town of Newport (See Table 5-13 and Appendix E.13). As a result of Mayo CPD there is potential for impacts to occur to Newport River SAC and Clew Bay Complex SAC via surface water, groundwater and land and air pathways due to the close vicinity of the Natura 2000 site. Example QIs that could be impacted include Salmon and Freshwater Pearl Mussel. Due to the topography of the surrounding landscape other Natura 2000 sites are unlikely to be impacted. Other Natura 2000 sites within the Zol are unlikely to be impacted because they are not hydrologically connected to the site and/or are located too far from the settlement.

However, as any project that may arise as a result of the Plan will require individual appropriate assessment at a project level, this cannot be assessed at this level. Also, the Plan takes into consideration the requirements of the Habitats Directive and so, any project that is influenced by the Plan or follows the normal consent process should have sufficient assessment and mitigation in place to prevent such adverse impacts.

Table 5-13: Natura 2000 sites within 15km of Newport

Site Code	Natura site	Distance (km)
004098	Owenduff/Nepin Complex SPA	5.3
000471	Brackloon Woods SAC	13.4
001482	Clew Bay Complex SAC	0.0
002144	Newport River SAC	0.0
000534	Owenduff/Nepin Complex SAC	5.3
002298	River Moy SAC	14.9

5.3.14 Swinford

There are 4 Natura 2000 sites within 15km from the town of Swinford (See Table 5-14 and Appendix E.10). As a result of Mayo CPD there is potential for impacts to occur to River Moy SAC via surface water, groundwater and land and air pathways due to the close vicinity of

the Natura 2000 site. QIs which could potentially be impacted include Otter, Salmon, Brook Lamprey and Sea Lamprey. Other Natura 2000 sites within the ZOI are unlikely to be impacted because they are not hydrologically connected to the site and/or are located too far from the settlement.

However, as any project that may arise as a result of the Plan will require individual Appropriate Assessment at a project level, this cannot be assessed at this time. Also, the Plan takes into consideration the requirements of the Habitats Directive and so, any project that is influenced by the Plan or follows the normal consent process should have sufficient assessment and mitigation in place to prevent such adverse impacts.

Table 5-14: Natura 2000 sites within 15km of Swinford

Site Code	Natura site	Distance (km)
004228	Lough Conn and Lough Cullen SPA	12.9
000633	Lough Hoe Bog SAC	10.2
002298	River Moy SAC	2.1
001571	Urlaur Lakes SAC	14.4

5.4 Population, One-off rural Housing and Settlement plans

As part of Mayo County Development Plan, Settlement Plans have been developed for a number of locations in Mayo. The below list includes several towns and villages that are scheduled to have settlement plans produced. The final list includes;

- Balla
- Ballinrobe
- Ballyhaunis
- Belmullet
- Charlestown
- Claremorris
- Crossmolina
- Foxford
- Kilalla
- Kiltimagh
- Knock
- Louisburgh
- Newport
- Swinford

Each settlement was assessed individually for potential impacts to Natura 2000 sites. Natura 2000 sites within 15km of each site was assessed, as well as Natura sites which are hydrologically connected to a site which have Freshwater Pearl Mussel as a Qualifying Interest up to 35km from a settlement. Impacts that may arise relative to each settlement are also assessed relative to the Objectives of the Plan in Section 8 and Appendix A.

5.5 Retail Development

The Mayo County Development Plan recognises a three-tier retail hierarchy in the County. This is consistent with the top three tiers of the Settlement Strategy and aims to concentrate higher order shopping functions in Castlebar, Ballina and Westport. It encourages more

generalised retail provision in the towns in Tier 2 and Tier 3 of the Settlement Strategy, while further providing for local shopping facilities in rural settlements and rural villages.

While the objectives, aims and policies for retail development in Plan do not directly specify any physical works that will occur, they do specify locations such as Castlebar, Ballina and Westport where specific recommendations are made. The encouragement for more generalised provision of retail facilities in towns in Tier 2 and Tier 3 of the Settlement Strategy as in Table 2-1, may effect Natura 2000 sites and their conservation objectives in those areas. These areas and Natura 2000 sites at risk are identified in Section 5.3.

The objectives and policies of the Plan related to retail development are at a higher level and therefore, they do not directly result in any physical works that may cause adverse impacts. However, as they give structure to future plans for growth and development that will require physical works to develop retail facilities, they will indirectly lead to projects that may potentially result in adverse impacts upon the overall integrity of any of the Natura 2000 sites that may be present within the Zone of Influence of the Plan. As this is a high-level Plan, the details to conduct assessments of any such project that may arise as a result of the objectives and policies at this level are not possible and any such project would be required to go through the consenting process as outlined in Section 9.1.

Therefore, the objectives and policies relating to retail development could result in direct and/or indirect impacts upon the overall integrity of the Natura 2000 network, through loss / reduction of habitat area, disturbance of key species, habitat or species fragmentation, reduction in species density and/or changes in key indicators of conservation value such as changes in water quality and quantity, and air quality.

5.6 Renewable energy projects and telecommunications infrastructure

Article 4 of Directive 2009/28/EC on renewable energy requires each Member State to adopt a national renewable energy action plan (NREAP). Ireland's NREAP sets out our national targets for the share of energy from renewable sources to be consumed in transport, electricity and heating and cooling in 2020. The plan demonstrates how the Member State will meet its overall national target established under the Directive. The National Energy Efficiency Action Plan (NEEAP) outlines how Ireland will achieve 20% energy efficiency savings, calculated on the basis of the average energy demand from 2001 to 2005. Central to this are the policies and measures identified by government to enable Ireland to achieve these targets.

Objective INP – 18 (chapter 7 of Mayo CPD), aims to “to support the provision of high-quality electricity infrastructure and development of enhanced electricity supply, to serve the existing and future needs of the County and to facilitate new transmission infrastructure projects that may be brought forward during the lifetime of the plan including the delivery and integration, including linkages of renewable energy proposals to the electricity transmission grid in a sustainable and timely manner, whilst seeking to protect and maintain bio-diversity, wildlife habitats, scenic amenities, including protected views and nature conservation.”

Therefore, the objectives and policies relating to renewable energy projects and telecommunications infrastructure could result in direct and/or indirect impacts upon the overall integrity of the Natura 2000 network, through loss / reduction of habitat area, disturbance of key species, habitat or species fragmentation, reduction in species density and/or changes in key indicators of conservation value such as changes in water quality and quantity, and air quality.

5.7 Water Quality

Under the Water Framework Directive (EU Directive 2000/60/EC), Mayo County Council is required as a public body, to co-ordinate their policies and operations to maintain water quality of unpolluted water bodies, and improve the status of polluted water bodies.

The main rivers in County Mayo and connected to the County are The Moy River, Clare River, Owenmore River Robe River and the River Deel.

These rivers are monitored by the EPA and Mayo County Council. Monitoring stations are located at several locations along The Moy River, Clare River, Owenmore River Robe River and the River Deel. The most recent water quality data show that the River Moy and Clare River is classified as having “Good” water quality and is “Not at Risk” from degradation. This Owenmore River and River Deel is classified as having “High” water quality. The River Robe is classified as having “Moderate” water quality and is “at risk” of further degradation.

Groundwater quality is protected under the European Communities Environmental Objectives Groundwater Regulations 2010, (S.I. No. 9 of 2010), which gives effect to the requirements of the Water Framework Directive (2000/60/EC) and the Groundwater Directive (2008/118/EC). Groundwater is important as a source of drinking water in Ireland and provides approximately 25% of all drinking water nationally. Groundwater connects wetlands, waterbodies and groundwater dependent terrestrial ecosystems throughout the County. It can be especially important during low-flow periods when groundwater forms a significant part of surface water flows.

Groundwater vulnerability in County Mayo is illustrated in Figure 5-2 and summarised in Table 5-15. The majority of groundwater in the County is classified as having “High” vulnerability (31.2%) or “Moderate” vulnerability (26.5%). 10.7% is classified as having “Low” vulnerability and 11.2% is classified as having “Extreme” vulnerability. The remaining area is covered by water (3.7%) and Rock at or near Surface or Karst (16.6%).

Table 5-15: Groundwater vulnerability in County Mayo

	Area (km ²)	Percentage (%)
Rock at or near Surface or Karst	924.5892	16.57027
Water	209.1013	3.747464
Extreme	627.5545	11.24688
High	1739.245	31.17033
Moderate	1483.048	26.57884
Low	596.2701	10.68621
Total	5579.808	100

The settlements of Kilalla and Newport are on the EPA’s priority list because they fall under one or more of the following criteria: discharging raw sewage because there is no treatment plant. Further information about EPA Priority Urban Areas can be found in the EPA’s Annual Urban Waste Water Report on the EPA website - www.epa.ie.




Figure 5-2: Groundwater vulnerability

Therefore, the objectives and policies that may give structure to, or direction for any development that may increase pressure on water quality or quantity in the area could result in direct and/or indirect impacts upon the overall integrity of the Natura 2000 network, through loss / reduction of habitat area, disturbance of key species, habitat or species fragmentation, reduction in species density and/or changes in key indicators of conservation value such as changes in water quality and quantity, and air quality.

5.8 Climate Change

The effects of climate change on the environment have economic and social implications, as well as implications for the environment themselves. Increasing temperatures, shifting seasons, changing rainfall patterns, increase in extremes of weather and sea level rise can all be caused by climate change. These changes can cause the spread of invasive species, changes in migration routes, loss or degradation of habitat and species and the overall loss of integrity of the Natura 2000 network. Overall, though, it can be difficult to predict with certainty the impacts of climate change as there are uncertainties in understanding how the earth's climatic system is changing.

Sites within the Natura 2000 network provide important localised space for nature to thrive, but they also form part of the wider network of green infrastructure across Europe. These

sites provide important refuges for biodiversity to allow species and habitats to adapt to climate change.

The effects of climate change on Natura 2000 sites can be considered as both direct impacts, and cumulative and in-combination impacts as a result of climate change. The cumulative and in-combination impacts of climate change are described in Section 6.

The following sections are guided by the decision framework (from (European Commission. Directorate General for the Environment. et al., 2013), to assist in deciding what actions may be required in County Mayo in order to mitigate against the impacts of climate change in the Mayo County Development Plan.

Temperature predictions in Ireland, show a clear west-to-east gradient, with the largest increases in the east of Ireland. Mid-century annual temperatures are projected to increase by 1-1.2°C to 1.3-1.6°C, dependant on future greenhouse gas emissions (Representative Concentration Pathway 4.5 (RCP4.5) or RCP8.5) (Nolan and Flanagan, 2020). Heatwaves are predicted to increase by the middle of the century and the numbers of frost and ice days are expected to decrease. Periods of frost and ice are important environmental drivers that trigger phenological responses in many plant and animal species. Species such as bats may not easily adapt to a reduction or loss of hibernation/torpor period and some plant seeds will not germinate without a cold spell to break dormancy. Extreme weather events such as heatwaves will occur more frequently and extreme temperatures are projected to increase by the middle of the century, causing increased stress to most species and the integrity of the Natura 2000 network.

Substantial decreases in rainfall are predicted for the summer months whilst during winter months increases in rainfall are predicted (Nolan and Flanagan, 2020), adding increased pressure on water dependant ecosystems. Bogs may be under threat as the water table drops, rivers and streams may dry up in total or in parts, fragmenting species from their spawning grounds. This could have particular impact to the Freshwater Pearl Mussel, as its lifecycle is dependent on Salmonids and a flow of highly oxygenated, clean water.

Increasing sea-levels are predicted worldwide as a result of climate change. Within County Mayo, there will be an increased likelihood of overtopping of flood defences during storm events in future years due to the impacts of climate change. Coastal towns such as, Ballina, have also been identified in the Regional Spatial and Economic Strategy as being at risk from storm surges and/or high tides emanating from the Atlantic Ocean. Low lying coastal settlements in County Mayo are particularly at risk from coastal flooding due to the impacts of climate change.

Natura 2000 sites and biodiverse habitats can mitigate and/or adapt the landscape against the predicted effects of climate change through the provision of valuable ecosystem services such as the storing carbon, reducing the amount in the atmosphere that contributes to increased temperatures. The Natura 2000 network in County Mayo, and in Ireland, encapsulates a broad variety of valuable habitats from marine and coastal wetlands, woodlands, grasslands and bogs.

Bogs and wetlands such as those at Bellacorick Bog Complex SAC, store water that could otherwise be increasing flood water levels. Trees such as those present in Brackloon Woods SAC absorb substantial amount of water and biodiverse habitats, or the diversity of habitats spread across the Natura 2000 network, provide a variety of trees and plant species that reduce available water, along with protecting the soil from erosion.

Any monoculture crop habitat, whether grown for food or other usage, is at risk from impacts that may be exacerbated by climate change such as disease, non-native invasive species and drought. Biodiverse habitats can provide reserves or refuges for species that may have adaptability to these and could be relied on in the future for genetic variability. Nature is an essential part of the solution to tackling climate change and its impacts.

Ecosystem services provided by the Natura 2000 network can reduce the impacts of sea level rise by providing natural coastal protection, they can assist in reducing overall temperature increases by increasing the capture of carbon and by climate regulation through provision of shade and storage of moisture in the land. They can also reduce the risk or impact, of extreme events that are caused or exacerbated by climate change such as flooding, fires and storms.

The vulnerability of habitats or species to climate change depends on their exposure, their sensitivity and their adaptive capacity. This varies for each habitat and species and is a complex assessment. All species and habitats in Mayo will be vulnerable to some extent. The habitats and species that occur within and around County Mayo will be susceptible to the effects of climate change in varied ways. The negative effects of climate change will impact upon habitats and species, those of which have an unfavourable conservation status, as well as deteriorate those with a favourable conservation status.

Marine and coastal habitats such as Atlantic salt meadows (*Glauco-Puccinellietalia maritima*) [1330], Reefs [1170] and Mudflats and sandflats not covered by seawater at low tide [1140] are amongst the most vulnerable habitats to climate change. The impact of climate change on Mayo's coastline is already evident, and further increases in relative sea levels, indicate that coastal areas will be increasingly susceptible to permanent inundation and erosion. These impacts are far reaching and are already changing the lives and livelihoods of coastal communities. As well as being vulnerable to increases in sea level, coastal and marine habitats and species are vulnerable to increasing temperatures changing species composition and altering spawning times or other natural patterns.

Marine and coastal species such as the Storm Petrel (*Hydrobates pelagicus*) [A014] or Puffin (*Fratercula arctica*) [A204] will be vulnerable to a loss of breeding, resting and feeding habitats due to climate change.

Ground and freshwater dependant terrestrial habitats and species, such as Turloughs [3180], Northern Atlantic wet heaths with *Erica tetralix* [4010], *Saxifraga hirculus* (Marsh Saxifrage) [1528], or *Vertigo geyeri* (Geyer's Whorl Snail) [1013] will be vulnerable from lowered precipitation and lower water table causing loss of suitable habitats and changes to key indicators of habitat quality such as water quantity.

Mammals including the *Rhinolophus hipposideros* (Lesser Horseshoe Bat) [1303], *Lutra lutra* (Otter) [1355] and *Phoca vitulina* (Harbour Seal) [1365] are vulnerable to changes in temperature and water quantity that may arise as a result of climate change.

Overall, of all the Natura 2000 habitats present, the most vulnerable habitat groups are the coastal habitats, freshwater habitats, bogs, mires and fens. However, in almost every habitat group, there are habitats and species present that are very vulnerable to climate change. The CDP also makes provisions for the management of green infrastructure for the regulation of micro-climates (green lung) and potentially, climate change adaptation. The Coastal Zone Policy NEP 16 specifically provides for the maintenance and enhancement of natural coastal defences to increase resilience to climate change.

Adaptive management and co-operation is key to managing the impacts that climate change may have on the vulnerabilities of the Natura 2000 network in Co Mayo and in Ireland. It is possible to change management regimes and policies to at least partially mitigate or adapt to impacts. The use of a structured, iterative process of optimal management decision-making based on system monitoring is needed, when faced with the uncertainty of climate change. Adaptive management builds in steps to assess progress towards achieving specified conservation goals and targets through monitoring actions, to see whether or not management actions are achieving species and/ or habitat targets. This is achieved in Ireland by National Parks and Wildlife Services, through the monitoring of Natura 2000 sites and the presentation of these results within the Article 17 reports. Under

Article 11 of the Habitats Directive, each member state is obliged to undertake surveillance of the conservation status of the natural habitats and species in the Annexes and under Article 17, to report to the European Commission every six years on their status and on the implementation of the measures taken under the Directive. The Mayo CDP recognises the importance of the Habitats Directive and seeks for all projects that may arise from the Plan to be compliant with the Directive. This is highlighted in the overarching strategic CDP objective SO 8 c) *To comply with the objectives and requirements of the Habitats Directive, specifically Article 6(3) and where necessary 6(4), Birds, Water Framework, and all other relevant EU Directives and all relevant transposing national legislation.*

As adaptive management is based on a learning process, it aims to improve management outcomes, by continuously adapting approaches, actions and measures on the basis of lessons learned. The development of the Mayo CDP is an iterative process which has considered, and will consider, the feedback from consultation and the outcome of the AA.

Mitigation measures within County Mayo and Ireland, that can mitigate against the impacts of climate change upon the Natura 2000 network include those that may occur on each site, those that must occur on a County level, and those that occur on a national and EU level. All mitigation measures required should be identified at the relevant levels through the AA process. Through assessing the potential impacts on all levels, including those that are cumulative and/or in-combination impacts and providing mitigation measures for these where relevant, significant adverse impacts should be mitigated against. Mitigation measures for those adverse impacts that relate to all potential adverse impacts including climate change are in Section 9.

5.9 Air Quality

The Air Framework Directive (96/62/EC) requires member states to divide countries into zones for the assessment and management of air quality

Ireland is divided into four zones which include:

- Zone A – Dublin Corporation
- Zone B – Cork Conurbation
- Zone C – Other Cities and large towns; and
- Zone D – Rural Ireland.

All of County Mayo is located within Zone D. Air quality is monitored by the EPA. Two air monitoring sites are located in Mayo, in Claremorris and Castlebar. The air quality index is calculated based on the information gathered from the monitoring stations using a Quality Index for Health, which is calculated every hour and indicates if air quality is good, fair, poor or very poor. The air quality in County Mayo is classified as “good”. with daily up to date information available for download from <http://www.epa.ie/air/quality/data/cm/>

5.10 Agricultural development

Agricultural development throughout County Mayo is an important economic driver and the Plan recognises this. The agricultural industry contributes to exports, provides raw materials for the food processing industry and the artisan food sector. The Council takes a positive approach to applications for sustainable agricultural developments generally, subject to the protection of ground waters, residential amenities, designated habitats and the landscape.

While considerations are made to designated habitats in general, they are more thoroughly considered in the overall objectives of the Plan. EOP 22 supports the implementation, at

county level, of the provisions set out in Food Harvest 2025, subject to environmental carrying capacity constraints. Food Harvest 2025 has undergone the AA process. It recommends monitoring across all sectors of the agri-food industry on the environmental impacts of the 2025 strategy at all levels. The results of this monitoring should result in directing any actions needed to prevent adverse impacts, however, this is unlikely to mitigate against any adverse impacts that may arise to the Natura network as a result of increased pressure caused by a growth in agriculture in the short term. The majority of the actions of Food Wise 2025 have been identified as having positive impacts and the overall conclusion of the AA report is that there would be no adverse impacts on the integrity of the Natura 2000 network as a result of Food Wise 2025.

While the Mayo CDP promotes the expansion of the Agri-Food Sector, it does so in consideration of the relevant guidelines and legislation including the Habitats Directive (See SO 8). Therefore, provided that the overarching Strategic Objectives of the Plan are considered in the expansion of the agricultural sector and that any development that occurs in the expansion of agriculture is subject to the normal processes and procedures for development including screening for AA is necessary, at this level, there is not likely to be any significant adverse impacts to the Natura 2000 network as a result of agricultural development in County Mayo.

5.11 Development of marine and coastal zone tourism and economic industry

County Mayo has extensive marine and coastal resources, and these resources can contribute to the economic stability of the county as a whole if developed appropriately and with sensitivity to the environment and Natura 2000 network. The marine industry can be extremely sensitive to change both on a social and environmental level. Development on land can lead to changes to water quality and have direct impacts upon coastal and marine resources. Climate change, tourism and recreational pressures also can put significant pressure on coastal and marine habitats and the species that are dependent on them.

The Natura 2000 sites that are dependent on marine and/or coastal habitats in County Mayo or within the ZOI of the CDP are listed below (see Table 5-16 and Figure 5-3). Qualifying interests of sites such as Clew Bay Complex SAC 001482, such as Otter and Common (Harbour) Seal are at risk of loss of breeding and resting habitats from coastal developments. The conservation objectives of habitats such as Atlantic Salt Meadows, or Shifting Dunes along the shoreline with *Ammophila arenaria* (white dunes) [21A0] are at risk from loss of habitat due to coastal development, erosion and climate change.

However, any development that may arise as a result of the Plan will be required to go through the consenting process and undergo AA screening if necessary. Therefore, it is unlikely that any coastal or marine developments that may arise as a result of the Plan will incur significant adverse impacts upon the Natura 2000 network. Although this is not likely to mitigate against any potential adverse impacts that may arise cumulatively as a result of the Plan with Climate Change and therefore, mitigation measure will be required.

Table 5-16: Coastal and marine Natura 2000 sites

SITECODE	Natura 2000 site
004004	Inishkea Islands SPA
004036	Killala Bay/Moy Estuary SPA
004036	Killala Bay/Moy Estuary SPA
004072	Stags of Broad Haven SPA
004074	Illanmaster SPA
004084	Inishglora and Inishkeeragh SPA
004093	Termoncarragh Lake and Annagh Machair SPA
004111	Duvillaun Islands SPA

SITECODE	Natura 2000 site
004136	Clare Island SPA
004144	High Island, Inishshark and Davillaun SPA
004170	Cruagh Island SPA
004177	Bills Rocks SPA
004212	Cross Lough (Killadoon) SPA
004221	Illaunnaon SPA
004227	Mullet Peninsula SPA
004231	Inishbofin, Omev Island and Turbot Island SPA
004235	Doogort Machair SPA
004037	Blacksod Bay/Broad Haven SPA
000278	Inishbofin And Inishshark SAC
000458	Killala Bay/Moy Estuary SAC
000458	Killala Bay/Moy Estuary SAC
000470	Mullet/Blacksod Bay Complex SAC
000472	Broadhaven Bay SAC
000484	Cross Lough (Killadoon) SAC
000500	Glenamoy Bog Complex SAC
000522	Lough Gall Bog SAC
001228	Aughrusbeg Machair And Lake SAC
001309	Omev Island Machair SAC
001311	Rusheenduff Lough SAC
001482	Clew Bay Complex SAC
001529	Lough Cahasy, Lough Baun And Roonah Lough SAC
001932	Mweelrea/Sheefry/Erriff Complex SAC
001955	Croaghaun/Slievemore SAC
002031	The Twelve Bens/Garraun Complex SAC
002118	Barnahallia Lough SAC
002130	Tully Lough SAC
001497	Doogort Machair/Lough Doo SAC
002265	Kingstown Bay SAC
000330	Tully Mountain SAC
000485	Corraun Plateau SAC
002243	Clare Island Cliffs SAC
002268	Achill Head SAC
000516	Lackan Saltmarsh and Kilcummin Head SAC
001501	Erris Head SAC
001513	Keel Machair/Menaun Cliffs SAC
002998	West Connacht Coast SAC
000495	Duvillaun Islands SAC
000507	Inishkea Islands SAC




Figure 5-3: Location of coastal and marine Natura 2000 sites.

5.12 Biodiversity, environment and green infrastructure

Provisions are made throughout the Plan, but in particular in Chapter 10 Natural Environment, *to protect and enhance Mayo's natural heritage and biodiversity, ensure that networks of green infrastructure are identified, created, protected and enhanced...*

While the aims, strategies, objectives and policies within the Plan relating to the environment and natural heritage are inherently of benefit to biodiversity and the Natura 2000 network, their execution must follow the same processes as those in all other sectors. Any projects that may arise as a result of the Plan must go through the normal consenting process and if necessary, may require screening for AA at a minimum. Considerations should always be made to consult with the relevant guidelines and best practise should always be followed.

Promotion, preservation and enhancement of biodiversity, treatment of Non-native Invasive Species or development of green infrastructure could impact upon the Natura 2000 network through various ways. Through reducing habitat quality, reduction in species density as a result of population fragmentation or direct mortalities and through changes to key indicators such as water quality or quantity.

The All Ireland Pollinator Plan has had a significant positive impact upon Ireland and in its promotion of biodiversity, many local groups and individuals have taken to seeding of wildflower meadows in previously unused or unappreciated land. While this can have a positive impact on biodiversity of brownfield sites or sites that have previously been fertilised and reseeded, wildflower seeds that are non-native are often used. This may not

be an issue in areas that are well managed or those that are not near to more natural Natura 2000 sites. But imported seeds that are spread within or adjacent to Natura 2000 sites, can reduce the quality of habitats present and replace native species impacting upon the overall integrity of the site.

Any works that may arise as a result of the Plan, that will occur within areas where non-native invasive species are present, will require biosecurity measures. This includes those works that will occur on land and/or in water. The spread of the Crayfish Plague is having devastating impacts upon populations of the White clawed Crayfish (*Austropotamobius pallipes*) with 100% mortalities. Therefore any projects or works that may occur as a result of the aims, objectives, policies and/or strategies of the Plan, whether for the protection of the natural environment or for any other means, should include provisions for biosecurity measures if necessary.

5.13 Tourism and Recreational pressures

A key element of the development plan is to promote tourism in County Mayo. Three key tourism pillars have been identified by the Plan, namely;

- The continued development and enhancement of tourism product and the various categories of tourism that bring visitors to Mayo;
- Ensuring that high quality services and accommodation are accessible and available for visitors;
- The identification of priority infrastructure designed to provide new and innovative activities and flagship products.

Tourism is of huge importance to the economy of County Mayo, particularly in coastal areas. The Wild Atlantic Way has been a huge boon to coastal areas in the west of Ireland. However, many Natura 2000 sites are under pressure from increased tourism and recreational activities; particularly sites with the following qualifying interests (Table 5-17):

Table 5-17: Natura 2000 sites potentially impacted by tourism and recreation

Qualifying features	Natura 2000 sites
Shifting dunes along the shoreline with <i>Ammophila arenaria</i> ("white dunes") [2130]	Killala Bay/Moy Estuary SAC
	Lough Cahasy, Lough Baun And Roonah Lough SAC
	Mullet/Blacksod Bay Complex SAC
	Mweelrea/Sheeffry/Erriff Complex SAC
Fixed coastal dunes with herbaceous vegetation ("grey dunes") [2150]	Killala Bay/Moy Estuary SAC
	Lackan Saltmarsh and Kilcummin Head SAC
	Lough Cahasy, Lough Baun And Roonah Lough SAC
Atlantic decalcified fixed dunes (Calluno-Ulicetea)	Mullet/Blacksod Bay Complex SAC
	Mweelrea/Sheeffry/Erriff Complex SAC
[21A0] Machairs (* in Ireland)	Clew Bay Complex SAC
	Doogort Machair/Lough Doo SAC

Qualifying features	Natura 2000 sites
	Glenamoy Bog Complex SAC
	Inishkea Islands SAC
	Keel Machair/Menaun Cliffs SAC
	Lough Cahasy, Lough Baun And Roonah Lough SAC
	Mullet/Blacksod Bay Complex SAC
	Mweelrea/Sheeffry/Erriff Complex SAC
	Omey Island Machair SAC
[1220] Perennial vegetation of stony banks	Clew Bay Complex SAC
	Cross Lough (Killadoon) SAC
	Keel Machair/Menaun Cliffs SAC
	Lough Cahasy, Lough Baun And Roonah Lough SAC
[1395] Petalophyllum ralfsii	Doogort Machair/Lough Doo SAC
	Glenamoy Bog Complex SAC
	Inishkea Islands SAC
	Keel Machair/Menaun Cliffs SAC
	Killala Bay/Moy Estuary SAC
	Mullet/Blacksod Bay Complex SAC
	Omey Island Machair SAC
[1310] Salicornia and other annuals colonizing mud and sand	Killala Bay/Moy Estuary SAC
	Lackan Saltmarsh and Kilcummin Head SAC
	Mullet/Blacksod Bay Complex SAC

Many sites with the above habitats in County Mayo have 'Human recreational activities, Golf links, walkways, increased human presence / walkways and recreational activities' listed as a threat on their Natura 2000 Standard Data forms.

Leisure fishing may also pose a threat to sites with aquatic habitats and species. For example, the River Moy is Ireland's premier Salmon river. Increased recreational fishing may result in increased pressure on this river and could result in detrimental impacts on Atlantic Salmon, one of the qualifying interests of this site.

5.14 Transport and related infrastructure

To enhance the local economy of County Mayo and to promote residential, commercial and recreational development, transport and its related infrastructure will likely increase. Currently the existing road network is deemed adequate for commercial transport of goods and to link population centres within the County and many roads have been upgraded to current best standard single carriageway. New or additional transport infrastructure to

support expanding development can have impacts, during construction and operation, on Natura 2000 sites within County Mayo. During construction of new infrastructure impacts can occur through habitat loss on habitats like:

- Machairs (21A0)
- Mediterranean salt meadows (*Juncetalia maritimī*) (1410)
- *Active raised bogs (7110)

The construction of new infrastructure may also impact on the following protected, and highly sensitive species, through pollution events or sediment runoff:

- Freshwater Pearl Mussel (*Margaritifera margaritifera*)
- White clawed Crayfish (*Austropotamobius pallipes*)
- Atlantic Salmon (*Salmo Salar*)

Furthermore, during operation of newly established infrastructure, the above QIs of Natura 2000 sites, including many other QIs (as identified in Table 8-1) have the potential to be impacted by habitat fragmentation, disturbance (visual and noise, polluted road runoff (hydrocarbons and microplastics) and vehicle emissions.

6 Cumulative and/or in-combination Impacts

As part of the Screening for an Appropriate Assessment, in addition to the proposed works, other relevant Projects and Plans in the region that may induce cumulative impacts must also be considered. As such, the list below contains Plans and Projects that are relevant to the Mayo County Development Plan and that may interact to cause cumulative and/or in-combination effects to Natura 2000 sites and Qualifying Interests. The Plans or Projects are listed according to a spatial hierarchy of National, Regional and Local. Potential cumulative and/or in-combination effects have been determined and are discussed in the section below, and are detailed in Appendix D. Mitigation measures that may be required are discussed in Section 9.

The following Projects and Plans were identified as potential sources of cumulative and in-combination impacts:

National

- River Basin Management Plan for Ireland 2018-2021
- Water Services Strategic Plan 2015-2040 (Irish Water)
- Local Authorities Water Programme
- Moy Water Management Unit Action Plan
- Plan for Forests & Freshwater Pearl Mussel in Ireland and The Pearl Mussel Project
- Irish National Action Plan for the Sustainable Use of Pesticides
- Ireland's Nitrates Action Programme
- Climate Action Plan Ireland
- Environmental River Enhancement Programme (EREP)
- Green, Low-Carbon Agri-Environment Scheme (GLAS)
- Food Wise 2025
- National Biodiversity Action Plan 2017-2021
- National Peatlands Strategy
- Project Ireland 2040 / National Development Plan 2018-2027 / National Planning Framework
- The National Spatial Strategy 2002-2020
- Our Sustainable Future - Framework for Sustainable Development in Ireland
- National Rural Development Programme 2014-2020
- Sustainable Development Goals National Implementation Plan 2018 - 2020
- Ireland Forestry Programme 2014-2020

Regional

- County Groundwater Protection Schemes
- Destination Mayo - a strategy for the future development of tourism in County Mayo 2015-2020 (DRAFT)
- Renewable Energy Strategy 2011-2022
- County Development Plan Galway, Roscommon and Sligo
- Shellfish Water Action Programmes / Pollution Reduction Programmes for shellfish
- County Mayo Biodiversity Action Plan 2010-2015 (new Plan for 2020-2025 in prep)

- Mayo local economic and community plan 2015-2021
- Northwest BAU (Business Area Unit) Strategic Plan 2016-2020

Local

- Invasive Species Action Plans
- Forthcoming Castlebar, Ballina and Westport Local Area Plans
- All-Ireland Pollinator Plan
- Planning Applications

6.1 National Plans and Projects with potential Cumulative and In-Combination Effects

National Plans and Projects that could have potential cumulative and in-combination effects were identified and are described below.

6.1.1 River Basin Management Plan for Ireland 2018-2021

The River Basin Management Plan (RBMP) (DoHLGH, 2018) for Ireland 2018-2021 sets out the actions that Ireland will take to improve water quality and achieve 'good' ecological status in water bodies (rivers, lakes, estuaries and coastal waters) by 2021 (DoHPLG, 2018a). Changes from previous River Basin Management Plans is that all River Basin Districts are merged as one national River Basin District. The Plan provides a more coordinated framework for improving the quality of our waters - to protect public health, the environment, water amenities and to sustain water-intensive industries, including agri-food and tourism, particularly in rural Ireland.

The first cycle of River Basin Management Plans included the Western River Basin District - River Basin Management Plan (ERBDMP) 2009 - 2015 (WFD, 2009). The plans summarised the water bodies that may not meet the environmental objectives of the WFD by 2015 and identified which pressures are contributing to the environmental objectives not being achieved. The plans described the classification results and identified measures that can be introduced in order to safeguard waters and meet the environmental objectives of the WFD:

- Prevent deterioration of water body status
- Restore good status to water bodies
- Achieve protected areas objectives
- Reduce chemical pollution of water bodies

With effective implementation of the RBMP, it can be expected to see the Plan's ambitious suite of measures translated into tangible improvements in water quality in over 700 water bodies around Ireland. The investment of €1.7 billion in waste-water infrastructure over the period of this Plan to 2021 will see over 250 projects in urban areas progressed and decisive steps towards sustainable and efficient water use will be taken. Assessment of risks to water quality in the planning processes will be enhanced and there will be more analyses of water quality carried out at a catchment level. Development related to the RBMP may have localised adverse impacts that will be assessed on a project level, however; overall impacts that may arise as a result of the RBMP are likely to be positive in nature. Due to the high level nature of this Plan, it is not possible to determine with confidence the likely impacts or mitigation measures required yet in detail.

Therefore, cumulative or in-combination impacts may arise as a result of projects that may occur as a result of this Plan. Any projects that may arise as a result of this plan will have a project level AA which will assess these in detail and provide

suitable mitigation measures where appropriate. Mitigation measures that may be required are discussed in Section 9.

6.1.2 Water Services Strategic Plan 2015-2040 (Irish Water)

The Water Services Strategic Plan 2015-2040 (WSSP) (Irish Water, 2015) by Irish Water sets out strategic objectives for the delivery of water services up to 2040. It details current and future challenges which affect the provision of water services and identifies the priorities to be tackled in the short and medium term. Development of infrastructure that maybe required for the implementation of the Plan may have localised adverse impacts.

Therefore, projects that may arise as a result of the WSSP will undergo AA at a project level, which will ensure that any cumulative or in-combination impacts are addressed. Due to the high-level nature of this Plan, it is not possible to determine with confidence the likely impacts or mitigation measures required yet in detail. Project level assessments will be required to determine any potential mitigation measures, as these will be dependent on the relevant project and any potential impacts that may arise as a result of this. Mitigation measures that may be required are discussed in Section 9.

6.1.3 Plan for Forests & Freshwater Pearl Mussel in Ireland 2018 and The Pearl Mussel Project

The Plan for Forests & Freshwater Pearl Mussel in Ireland (DoAFM, 2018) aims to eliminate, reduce or mitigate diffuse and point sources of sediment and nutrients and the disruption of the natural hydrological regime, arising from forests and regulated forestry activities, to ensure that these do not threaten the achievement of the conservation objectives for Freshwater Pearl Mussel set for each of the SACs involved. There is a potential for indirect impact on other species as a result of protection measures. It is likely that any impacts will be positive in nature, however, in order to determine cumulative impacts these will have to be screened against potential impacts that may occur as a result of any actions that will occur at a project level. Project level assessment will ensure that any potential cumulative impacts are addressed.

The Freshwater Pearl Mussel Project is a pilot agri-environment programme that seeks to improve the quality of watercourses to benefit the endangered FPM. Farmers will be rewarded for their environmental services whilst having the freedom and flexibility to farm. This will in turn improve the outcomes and long-term sustainability of agri-environment schemes for biodiversity, and for rural communities. The Programme is being run by the Pearl Mussel Project Team, based in counties Kerry and Mayo. It is likely that any impacts will be positive in nature, however, in order to determine cumulative impacts these will have to be screened against potential impacts that may occur as a result of any actions that will occur at a project level. Project level assessment will ensure that any potential cumulative impacts are addressed. Due to the high level nature of this Plan, it is not possible to determine with confidence the likely impacts or mitigation measures required yet in detail.

Therefore, cumulative or in-combination impacts may arise as a result of projects that may occur as a result of this Plan. Any projects that may arise as a result of this plan will have a project level AA which will assess these in detail and provide suitable mitigation measures where appropriate. Mitigation measures that may be required are discussed in Section 9.

6.1.4 Irish National Action Plan for the Sustainable Use of Pesticides

The EU Sustainable Use Directive requires each Member State to develop a National Action Plan for the Sustainable Use of Pesticides (NASUP) (DoAFM, 2019), which shall be reviewed at least every 5 years. The Plan sets out a national strategy to achieve a sustainable use of pesticides and aims to achieve a balance between ensuring human and environmental safety while maintaining continued viability of the farming and amenity sectors. Particular emphasis in this Plan is being placed on protecting drinking water from pesticides and no negative cumulative and/or in-combination effects are expected. However, actions that may arise as a result of the NAPSUP will be considered at a project level, which will ensure that any cumulative or in-combination impacts are addressed. Due to the high level nature of this Plan, it is not possible to determine with confidence the likely impacts or mitigation measures required yet in detail.

Therefore, cumulative or in-combination impacts may arise as a result of projects that may occur as a result of this Plan. Any projects that may arise as a result of this plan will have a project level AA which will assess these in detail and provide suitable mitigation measures where appropriate. Mitigation measures that may be required are discussed in Section 9.

6.1.5 Ireland's Nitrates Action Programme

Ireland's Nitrates Actions Programme (DoHLGH, 2017) is designed to prevent pollution of surface waters and groundwater from agricultural sources and to protect and improve water quality. Under the national NAP, each Member State's NAP must include 1) a limit on the amount of livestock manure applied to the land each year, 2) set periods when land spreading is prohibited due to risk, and 3) set capacity levels for the storage of livestock manure. Notwithstanding the economic success of the government's strategies to date, the agricultural sector faces a number of challenges, specifically relating to water quality standards and climate change commitments. As the industry embraces new levels of growth, it will also be required to show an absolute commitment to the principles of sustainability, recognising that gains in productivity must not be at the expense of the environment. As measures under the NAP are exclusively designed to prevent pollution of surface waters and groundwater, overall positive cumulative and/or in-combination effects are expected. However, due to the high level nature of this Plan, it is not possible to determine with confidence the likely impacts or mitigation measures required yet in detail.

Therefore, cumulative or in-combination impacts may arise as a result of projects that may occur as a result of this Plan. Any projects that may arise as a result of this plan will have a project level AA which will assess these in detail and provide suitable mitigation measures where appropriate. Mitigation measures that may be required are discussed in Section 9.

6.1.6 Climate Action Plan Ireland

Ireland's Climate Action Plan (Government of Ireland, 2019) sets out an ambitious course of action over the coming years to address this issue. The Plan identifies the nature and scale of the challenge, outlines the current state of play across key sectors including Electricity, Transport, Built Environment, Industry and Agriculture and charts a course towards ambitious decarbonisation targets. Reflecting the central priority climate change will have in Ireland's political and administrative systems into the future, the Plan sets out governance arrangements including carbon-proofing policies, establishment of carbon budgets, a strengthened Climate Change Advisory Council and greater accountability to the Oireachtas. It is not expected that measures proposed in this Plan will lead to significant cumulative and/or in-combination effects, although there might be a potential effect from alterations in land-use. Due to the high level nature of this Plan, it is not possible to determine with confidence the likely impacts or mitigation measures required yet in detail.

Therefore, cumulative or in-combination impacts may arise as a result of projects that may occur as a result of this Plan. Any projects that may arise as a result of this plan will have a project level AA which will assess these in detail and provide suitable mitigation measures where appropriate. Mitigation measures that may be required are discussed in Section 9.

6.1.7 Environmental River Enhancement Programme (EREP)

The Environmental River Enhancement Programme (EREP) (IFI, 2008) is an OPW funded project that is being coordinated and managed by Inland Fisheries Ireland. The Programme focuses on the enhancement of drained Salmonid rivers in Ireland. The EREP involves two different approaches to enhancement, these being capital enhancement and enhanced maintenance respectively. Infrastructural works that arise as a result of the EREP have the potential to impact other designated features, however, the effects will overall be positive. Actions that may arise as a result of the EREP will be considered at a project level, which will ensure that any cumulative or in-combination impacts are addressed. Due to the high level nature of this Programme, it is not possible to determine with confidence the likely impacts or mitigation measures required yet in detail.

Therefore, cumulative or in-combination impacts may arise as a result of projects that may occur as a result of this Programme. Any projects that may arise as a result of this plan will have a project level AA which will assess these in detail and provide suitable mitigation measures where appropriate. Mitigation measures that may be required are discussed in Section 9.

6.1.8 Green, Low-Carbon Agri-Environment Scheme (GLAS) and follow-on schemes

The GLAS Scheme is part of the Rural Development Programme 2014-2020 (DoAFM, 2014a). It provides funding to farmers in return for delivering environmental management on their land. Farmers must commit to the scheme for a minimum period of 5 years. Farmers participating in the GLAS must comply with several basic measures, including protecting and maintaining all watercourses and wells and cease using herbicides, pesticides and fertilisers in and around hedgerows, lakes, ponds, rivers and streams (except with consent). GLAS recognises the importance of the riparian zone in rural areas and any proposed works should make recommendations that are compatible with GLAS. Development of infrastructure may have localised adverse impacts, however; overall impacts that may arise as a result of this Scheme are likely to be positive in nature. Actions that may arise as a result of the Scheme will be considered at a project level, which will ensure that any cumulative or in-combination impacts are addressed. Due to the high level nature of this Plan, it is not possible to determine with confidence the likely impacts or mitigation measures required yet in detail.

Therefore, cumulative or in-combination impacts may arise as a result of projects that may occur as a result of this Scheme. Any projects that may arise as a result of this plan will have a project level AA which will assess these in detail and provide suitable mitigation measures where appropriate. Mitigation measures that may be required are discussed in Section 9.

6.1.9 Food Wise 2025

Food Wise 2025 (DoAFM, 2015) sets out a ten-year plan for Ireland's agri-food sector. It underlines the sector's unique and special position within the Irish economy, and it illustrates the potential which exists for this sector to grow even further. The Food Wise report was published in July 2015, with the Food Wise Implementation Plan and the final

Environmental Analysis on the strategy published in December 2015. Ireland's agri-food sector manages the vast majority of the natural resources in Ireland. In a global environment which is facing many challenges including an ever-increasing demand for food, increasing constraints on natural resources and the emerging challenges of Climate Change put pressure on the Irish agri-food sector. Ireland faces significant challenges in meeting some national and international environmental targets for air quality, biodiversity and water quality. Meeting Greenhouse Gas and ammonia emission reduction targets will be particularly challenging, but arresting biodiversity losses and continuing the improvement of water quality while increasing production will be equally demanding. Projects that may arise as a result of Food Wise 2025 plans have the potential to increase pressure on the environment through land management and therefore, could result in cumulative impacts to sites. However, due to the high level nature of this Plan, it is not possible to determine with confidence the likely impacts or mitigation measures required yet in detail.

Therefore, cumulative or in-combination impacts may arise as a result of projects that may occur as a result of Foodwise 2025. Any projects that may arise as a result of this plan will have a project level AA which will assess these in detail and provide suitable mitigation measures where appropriate. Mitigation measures that may be required are discussed in Section 9.

6.1.10 National Biodiversity Action Plan 2017-2021

The National Biodiversity Action Plan 2017-2021 (DoCHG, 2017) captures the objectives, targets and actions for biodiversity that will be undertaken by a wide range of government, civil society and private sectors to achieve Ireland's Vision for Biodiversity. It provides a framework to track and assess progress towards Ireland's Vision for Biodiversity over a five-year timeframe from 2017 to 2021. Implementation of proposed measures could help achieve the aims of the National Biodiversity Plan through optimisation of biodiversity benefits in planning and therefore overall impacts will be positive. However, due to the high level nature of this Plan, it is not possible to determine with confidence the likely impacts or mitigation measures required yet in detail.

Therefore, cumulative or in-combination impacts may arise as a result of projects that may occur as a result of the National Biodiversity Action Plan 2017-2021. Any projects that may arise as a result of this plan will have a project level AA which will assess these in detail and provide suitable mitigation measures where appropriate. Mitigation measures that may be required are discussed in Section 9.

6.1.11 National Peatlands Strategy 2015

The National Peatlands Strategy (NPWS, 2015) guides the Government's approach to peatlands management and conservation in the future, considering current and potential uses of this key resource. Specific outcomes of the Mayo CDP could impact the aims of the National Peatland Strategy, but implementation of specific measures could also contribute to the aim of the strategy in improving bogs. Projects that may arise as a result of the National Peatlands Strategy must be screened for in-combination and cumulative impacts that may occur with projects that arise as a result of the National Peatland Strategy at a project level. Due to the high level nature of this Plan, it is not possible to determine with confidence the likely impacts or mitigation measures required yet in detail.

Therefore, cumulative or in-combination impacts may arise as a result of projects that may occur as a result of the National Peatlands Strategy. Any projects that may arise as a result of this plan will have a project level AA which will assess these in detail and provide suitable mitigation measures where appropriate. Mitigation measures that may be required are discussed in Section 9.

6.1.12 Project Ireland 2040

Project Ireland 2040 (DoPER, 2018) is the government's long-term overarching strategy to make Ireland a better country for all of its people. Alongside the development of physical infrastructure, Project Ireland 2040 supports business and communities across all of Ireland in realising their potential. The National Development Plan and the National Planning Framework combine to form Project Ireland 2040. Development of infrastructure may have localised adverse impacts. Actions that may arise as a result of Project Ireland 2040 will be considered at a project level, which will ensure that any cumulative or in-combination impacts are addressed. Due to the high level nature of this Plan, it is not possible to determine with confidence the likely impacts or mitigation measures required yet in detail.

Therefore, cumulative or in-combination impacts may arise as a result of projects that may occur as a result of Project Ireland 2040. Any projects that may arise as a result of this plan will have a project level AA which will assess these in detail and provide suitable mitigation measures where appropriate. Mitigation measures that may be required are discussed in Section 9.

6.1.13 The National Spatial Strategy 2002-2020

The National Spatial Strategy 2002-2020 (DoHLGH, 2002) is a coherent national planning framework for Ireland, which aims to achieve a better balance of social, economic and physical development across Ireland, supported by more effective planning. In order to drive development in the regions, the NSS proposes that areas of sufficient scale and critical mass will be built up through a network of gateways and hubs. Development of infrastructure may have localised adverse impacts. However, due to the high level nature of this Plan, it is not possible to determine with confidence the likely impacts or mitigation measures required yet in detail.

Therefore, cumulative or in-combination impacts may arise as a result of projects that may occur as a result of the National Spatial Strategy. Any projects that may arise as a result of this plan will have a project level AA which will assess these in detail and provide suitable mitigation measures where appropriate. Mitigation measures that may be required are discussed in Section 9.

6.1.14 Our Sustainable Future - Framework for Sustainable Development in Ireland

This Framework (DoCCA, 2012) identifies some 70 measures to be implemented across government and tasks a high-level inter-departmental group with ensuring that the vision set out in the policy document is translated into clear and effective action. Development of infrastructure that may be required in order to achieve the measures of this Plan may have localised adverse impacts. Actions that may arise as a result of this Framework will be considered at a project level, which will ensure that any cumulative or in-combination impacts are assessed for significance. Due to the high level nature of this Plan, it is not possible to determine with confidence the likely impacts or mitigation measures required yet in detail.

Therefore, cumulative or in-combination impacts may arise as a result of projects that may occur as a result of this Framework. Any projects that may arise as a result of this plan will have a project level AA which will assess these in detail and provide suitable mitigation measures where appropriate. Mitigation measures that may be required are discussed in Section 9.

6.1.15 National Rural Development Programme 2014-2020

The National Rural Development Programme 2014-2020 (NRDP) (DoAFM, 2014b), as part of the Common Agricultural Policy (CAP), is a common set of objectives, principles and rules

through which the European Union coordinates support for European agriculture. The CAP is structured around two complementary pillars: Pillar 1 deals with direct payments to farmers and market management measures; Pillar 2 covers multi-annual rural development measures which include those that have beneficial impacts on the environment and climate change. The NRDP recognises that agriculture can have significant impacts on the environment, including biodiversity, flood and drought control, and as a carbon sink. Therefore, overall impacts that may arise as a result of the NRDP will likely be negative. However, any actions that may arise as a result of the NRDP will be considered at a project level, which will ensure that any cumulative or in-combination impacts are addressed. Due to the high level nature of this Plan, it is not possible to determine with confidence the likely impacts or mitigation measures required yet in detail.

Therefore, cumulative or in-combination impacts may arise as a result of projects that may occur as a result of this Programme. Any projects that may arise as a result of this plan will have a project level AA which will assess these in detail and provide suitable mitigation measures where appropriate. Mitigation measures that may be required are discussed in Section 9.

6.1.16 Sustainable Development Goals National Implementation Plan 2018 - 2020

The Sustainable Development Goals National Implementation Plan 2018 - 2020 (DoCCA, 2018) is in direct response to the 2030 Agenda for Sustainable Development and provides a whole-of-government approach to implement the 17 Sustainable Development Goals (SDGs). The Plan identifies four strategic priorities to guide implementation: 1) Awareness: raise public awareness of the SDGs; 2) Participation: provide stakeholders opportunities to engage and contribute to follow-up and review processes, and further develop national implementation of the Goals; 3) Support: encourage and support efforts of communities and organisations to contribute towards meeting the SDGs, and foster public participation; and 4) Policy alignment: develop alignment of national policy with the SDGs and identify opportunities for policy coherence. Actions that may arise as a result of this Plan will likely not result in cumulative and/or in-combination impacts.

Therefore, cumulative or in-combination impacts are unlikely to arise as a result of this Plan.

6.1.17 Ireland Forestry Programme 2014-2020

The Ireland Forestry Programme 2014-2020 (Forest Service/DoAFM, 2015) aims to develop a competitive and sustainable forest sector through the implementation of measures, in particular afforestation, woodland creation and the prevention and restoration of damage to forests. Development of the forest sector may have localised adverse impacts, however; some measures resulting from this Programme will be positive in nature. Actions that may arise as a result of the IFP will be considered at a project level, which will ensure that any cumulative or in-combination impacts are addressed. Due to the high level nature of this Plan, it is not possible to determine with confidence the likely impacts or mitigation measures required yet in detail.

Therefore, cumulative or in-combination impacts may arise as a result of projects that may occur as a result of this Plan. Any projects that may arise as a result of this plan will have a project level AA which will assess these in detail and provide suitable mitigation measures where appropriate. Mitigation measures that may be required are discussed in Section 9.

6.2 Regional Plans and Projects with potential Cumulative and In-Combination Effects

Regional Plans and Projects that could have potential cumulative and in-combination effects were identified and are described below.

6.2.1 Moy Water Management Unit Action Plan

The Moy Water Management Unit (WMU) Action Plan (Ordnance Survey Ireland/Galway County Council, 2010) maps the river and lake water bodies of the Moy Water Management Unit, states the status, impacts, pressures and risk for these water bodies and defines action programmes and objectives. In general, these tend to be positive in nature and designed for the overall management and preservation of water quality within the WMU, but at a high level. Due to the high level nature of this Plan, it is not possible to determine with confidence the likely impacts or mitigation measures required yet in detail.

Therefore, cumulative or in-combination impacts may arise as a result of projects that may occur as a result of this Plan. Any projects that may arise as a result of this plan will have a project level AA which will assess these in detail and provide suitable mitigation measures where appropriate. Mitigation measures that may be required are discussed in Section 9.

6.2.2 Mayo County Council Renewable Energy Strategy 2011-2022

The Mayo County Council Renewable Energy Strategy 2011-2022 (Mayo CoCo, 2011) sets out a path to allow County Mayo to contribute to meeting the national legally binding targets and sets out opportunities for individuals, communities and businesses to harness renewable energy in a sustainable manner and to assist in combating climate change. The Strategy also clarifies the approach Mayo County Council takes to renewable energy, and should assist direction and reduce uncertainty for the most regarding issues associated with renewable energy developments in Mayo. All major forms of renewable energy are considered in the Strategy, including micro renewables. This is a high-level Strategy and any projects that may arise as a result of this Strategy will be required to be assessed for significance of cumulative and/or in-combination effects at a project level. Development of infrastructure necessary to implement the Strategy may have localised adverse impacts. Due to the high-level nature of this Plan, it is not possible to determine with confidence the likely impacts or mitigation measures required yet in detail.

Therefore, cumulative or in-combination impacts may arise as a result of projects that may occur as a result of this Plan. Any projects that may arise as a result of this plan will have a project level AA which will assess these in detail and provide suitable mitigation measures where appropriate. Mitigation measures that may be required are discussed in Section 9.

6.2.3 Destination Mayo - a strategy for the future development of tourism in County Mayo 2015-2020 (DRAFT)

Destination Mayo, Mayo's Tourism Strategy (Mayo CoCo Enterprise & Investment Unit, 2015), seeks to build on the work of a previous study of tourism in Mayo called A Tourism Plan for Mayo. Some of the outcomes of the 1991 Plan included Mayo 5000, Mayo Naturally, designation of Ballycroy National Park and facilitated the development of a number of visitor attractions throughout the County, including Céide Fields, Mayo Naturally, National Museum of Ireland - Country Life and the Lough Lannagh Holiday Village. The vision for Mayo's tourism industry in 2020 is to be a high quality, inclusive and authentic destination. Key to the successful delivery of this vision are the availability of a skilled and enthusiastic workforce and good access to the County by road, rail and air. Destination Mayo seeks to promote tourism in Mayo and aims to attract more visitors to Mayo through destination marketing and other activities. It is expected that Projects that may result from this Plan

may have negative impacts, however; due to the high level nature of this Plan, it is not possible to determine with confidence the likely impacts or mitigation measures required yet in detail.

Therefore, cumulative or in-combination impacts may arise as a result of projects that may occur as a result of this Plan. Any projects that may arise as a result of this plan will have a project level AA which will assess these in detail and provide suitable mitigation measures where appropriate. Mitigation measures that may be required are discussed in Section 9.

6.2.4 County Development Plans Galway, Roscommon and Sligo

County Development Plans from counties sharing a border with County Mayo must also be considered, as proposed actions and measures from these Plans may induce cumulative and/or in-combination effects on SACs, SPAs and river systems in County Mayo.

A number of SACs and SPAs are overlapping Mayo’s borders and are shared with neighbouring counties. Any works in these areas may induce positive and/or negative cumulative in-combination impacts across the border into County Mayo. SACs and SPAs overlapping borders with neighbouring counties are:

Mayo/Galway border	Lough Carra/Mask Complex SAC
	Lough Corrib SAC
	Lough Corrib SPA
	Mweelrea/Sheeffry/Erriff Complex SAC
Mayo/Roscommon border	River Moy SAC
Mayo/Sligo border	River Moy SAC
	Lough Hoe Bog SAC
	Ox Mountains Bogs SAC
	Killala Bay/Moy Estuary SAC
	Killala Bay/Moy Estuary SPA

As County Mayo is bordered by County Galway, County Roscommon and County Sligo, the County Development Plans to be considered are:

- Galway County Development Plan 2015-2021 (Galway CoCo, 2015)
- Roscommon County Development Plan 2014-2020 (Roscommon CoCo, 2014)
- Sligo County Development Plan 2017-2023 (Sligo CoCo, 2014)

Actions that may arise as a result of the Galway, Roscommon and Sligo CDPs will be considered at a project level where appropriate, which will ensure that any cumulative or in-combination impacts are addressed. However, due to the high level nature of this Plan, it is not possible to determine with confidence the likely impacts or mitigation measures required yet in detail.

Therefore, cumulative or in-combination impacts may arise as a result of projects that may occur as a result of this Plan. Any projects that may arise as a result of this plan will have a project level AA which will assess these in detail and provide suitable mitigation measures where appropriate. Mitigation measures that may be required are discussed in Section 9.

6.2.5 Shellfish Water Action Plans 2019

Under the EU Shellfish Waters Directive, Ireland is obliged to establish measures to protect shellfish waters against pollution and to safeguard certain shellfish populations from various harmful consequences, resulting from the discharge of pollutant substances into the sea.

Shellfish Action Plans will next be reviewed on a priority basis starting in 2021 which is in line with the third River Basin Cycle under the Water Framework Directive. Any proposed works will have to ensure that the water quality of shellfish areas is not impacted. Any works that may arise as a result of these programmed will be designed to ensure that the water quality of shellfish areas is not adversely impacted and may even be designed to enhance water quality, causing positive impacts. Development of infrastructure required for these changes may have localised adverse impacts, however; overall impacts that may arise as a result of the Programmes are likely to be positive in nature. Actions that may arise as a result of the Programmes, must be assessed further at a project level. Due to the high level nature of these Programmes, it is not possible to determine with confidence the likely impacts or mitigation measures required yet in detail.

Therefore, cumulative or in-combination impacts may arise as a result of these projects that may occur as a result of these Programmes. Any projects that may arise as a result of this plan will have a project level AA which will assess these in detail and provide suitable mitigation measures where appropriate. Mitigation measures that may be required are discussed in Section 9.

6.2.1 Mayo County Groundwater Protection Scheme

The Mayo County Groundwater Protection Scheme (Geological Survey Ireland, 2018) provide guidelines for the planning and licensing authorities in carrying out their functions, and a framework to assist in decision-making on the location, nature and control of developments and activities in order to protect groundwater. County Groundwater Protection Schemes provide a framework to assist in decision-making on the location, nature and control of developments and activities in order to protect groundwater in a given location. Even though measures can improve groundwater, groundwater protection responses may involve hard structures or alterations to hydrology, and could therefore change water levels, quality or quantity, impacting sites and ground water dependent Qualifying Interests. Actions that may arise as a result of the County Groundwater Protection Schemes will be considered at a project level, which will ensure that any cumulative or in-combination impacts are addressed. Due to the high level nature of this Plan, it is not possible to determine with confidence the likely impacts or mitigation measures required yet in detail.

Therefore, cumulative or in-combination impacts may arise as a result of projects that may occur as a result of this Plan. Any projects that may arise as a result of this plan will have a project level AA which will assess these in detail and provide suitable mitigation measures where appropriate. Mitigation measures that may be required are discussed in Section 9.

6.2.2 County Mayo Biodiversity Action Plan 2010-2015 / Strategy for the protection, conservation and promotion of Mayo's biodiversity 2020 - 2025

The County Mayo Biodiversity Action Plan (BAP) (Mayo CoCo, 2010) aims to raise awareness of and promote the conservation of the natural heritage and biodiversity of the County. The Plan provides a framework for the conservation of biodiversity at a local level and helps ensure that national & international targets for biodiversity conservation can be achieved. A new strategy for the protection, conservation and promotion of Mayo's biodiversity is currently in preparation. This strategy will form part of the new County Mayo Heritage Plan 2020–2025 currently being developed and mirrors the National Biodiversity Action Plan. The County Mayo BAP provides a framework for the conservation of biodiversity at a local level and helps ensure that national and international targets for biodiversity conservation can be achieved, while at same time addressing local priorities. Therefore, overall impacts

will be positive. However, due to the high level nature of this Plan, it is not possible to determine with confidence the likely impacts or mitigation measures required yet in detail.

Therefore, cumulative or in-combination impacts may arise as a result of projects that may occur as a result of this Plan. Any projects that may arise as a result of this plan will have a project level AA which will assess these in detail and provide suitable mitigation measures where appropriate. Mitigation measures that may be required are discussed in Section 9.

6.2.3 Mayo Local Economic and Community Plan 2015-2021

The Mayo Local Economic and Community Plan 2015-2021 (Mayo Local Community Development Committee, 2015) identifies economic and local community issues in County Mayo and gives effect to the delivery of economic and community development in a manner that is consistent with higher level plans and strategies including the Regional Planning Guidelines for the West Region 2010-2022 and the Mayo County Development Plan 2014-2020. It is possible that any actions that may arise as a result of this Plan be assessed for significance of negative in-combination effects with the Mayo CDP. Development of infrastructure may have localised adverse impacts. However, due to the high level nature of this Plan, it is not possible to determine with confidence the likely impacts or mitigation measures required yet in detail.

Therefore, cumulative or in-combination impacts may arise as a result of projects that may occur as a result of this Plan. Any projects that may arise as a result of this plan will have a project level AA which will assess these in detail and provide suitable mitigation measures where appropriate. Mitigation measures that may be required are discussed in Section 9.

6.2.4 Northwest BAU (Business Area Unit) Strategic Plan 2016-2020

The Northwest Business Area Unit Strategic Plan (Coillte, 2016) sets out plans for the forest and non-forest business that will take place in the BAU during the plan period. In practicing sustainable forest management, Coillte's aim is to develop its forests in a way that is environmentally sustainable, socially sustainable and economically sustainable. Measures as a result of this Plan may have localised adverse impacts. Actions that may arise as a result of this Plan will be considered at a project level, which will ensure that any cumulative or in-combination impacts are addressed. However, due to the high level nature of this Plan, it is not possible to determine with confidence the likely impacts or mitigation measures required yet in detail.

Therefore, cumulative or in-combination impacts may arise as a result of projects that may occur as a result of this Plan. Any projects that may arise as a result of this plan will have a project level AA which will assess these in detail and provide suitable mitigation measures where appropriate. Mitigation measures that may be required are discussed in Section 9.

6.3 Local Plans and Projects with potential Cumulative and In-Combination Effects

Local Plans and Projects that could have potential cumulative and in-combination effects were identified and are described below.

6.3.1 Invasive Species Action Plans

An Invasive Species Action Plan (ISAP), like Mayo's Management of Invasive Alien Plant Species (IAPS) for Japanese Knotweed (Mayo CoCo, 2016) is a requiry under the National Biodiversity Action Plan to prepare strategies, in consultation with Northern Ireland, to control introduced species and to prevent, or minimise, future (accidental or deliberate)

introduction of alien species, which might threaten biodiversity. The purpose of an ISAP is to outline the work to be done and strategies to be followed for the control of invasive species. ISAPs are directly related to the control management of a species, however, there is potential to indirectly impact on other species as a result of measures that may be put in place. However; overall impacts will likely be positive. However, due to the high level nature of this Plan, it is not possible to determine with confidence the likely impacts or mitigation measures required yet in detail.

Therefore, cumulative or in-combination impacts may arise as a result of projects that may occur as a result of these Plans. Any projects that may arise as a result of this plan will have a project level AA which will assess these in detail and provide suitable mitigation measures where appropriate. Mitigation measures that may be required are discussed in Section 9.

6.3.2 Forthcoming Castlebar, Ballina and Westport Local Area Plans

Mayo County Council intends to prepare a Local Area Plans for the towns in the County which have a population exceeding 5000, these towns are Castlebar, Ballina and Westport. Castlebar and Ballina have been designated as Key Towns in the Regional Spatial and Economic Strategy (RSES), published by the Northern and Western Regional Assembly on 23 January 2020, whilst Westport was also identified as a regional driver. The formal preparation of the Local Area Plans for Castlebar, Ballina and Westport will commence after the Draft Mayo County Development Plan 2021-2027 has been published. It is possible that any actions that may arise as a result of Local Areas Plans are assessed for significance of negative in-combination effects with the Mayo CDP. However, due to the high level nature of this Plan, it is not possible to determine with confidence the likely impacts or mitigation measures required yet in detail.

Therefore, cumulative or in-combination impacts may arise as a result of projects that may occur as a result of these Plans. Any projects that may arise as a result of this plan will have a project level AA which will assess these in detail and provide suitable mitigation measures where appropriate. Mitigation measures that may be required are discussed in Section 9.

6.3.3 All-Ireland Pollinator Plan

The All-Ireland Pollinator Plan (NBDC, 2015) is a strategy that works to support the decline of pollinators including Ireland's 98 bee species, one-third of which are threatened with extinction. Mayo County Council signed up as a partner to the All-Ireland Pollinator Plan in May 2019 and works to support the All-Ireland Pollinator Plan through working with communities and local authority staff and hosting workshops and talks on the Plan. The AIPP is focused on taking measures in order to protect species and habitats and therefore overall positive impacts are expected. However, due to the high level nature of this Plan, it is not possible to determine with confidence the likely impacts or mitigation measures required yet in detail.

Therefore, cumulative or in-combination impacts may arise as a result of projects that may occur as a result of this Plan. Any projects that may arise as a result of this plan will have a project level AA which will assess these in detail and provide suitable mitigation measures where appropriate. Mitigation measures that may be required are discussed in Section 9.

6.3.4 Planning Applications

Planning and infrastructure may have localised adverse impacts. Due to the high level nature of this Plan, it is not possible to determine with confidence the likely impacts or

mitigation measures required yet in detail, and therefore actions that may arise as a result of these planning applications will be considered at a project level. This will ensure that any cumulative or in-combination impacts are addressed. Project level assessments will also be required to determine any potential mitigation measures, as these will be dependent on the relevant project and any potential impacts that may arise as a result of this.

6.3.5 Tourism and Leisure Activities and Associated Threats to Natura 2000 Sites in County Mayo

A key element of the development plan is to promote tourism in County Mayo. We recognise that tourism is of huge importance to the economy of County Mayo, particularly in coastal areas. The Wild Atlantic Way has been a huge boon to coastal areas in the west of Ireland. However, our review of the sites which would be considered within the CDP reveal that many sites are under pressure from increased tourism and recreational activities; particularly sites with the following qualifying interests:

- Shifting dunes along the shoreline with *Ammophila arenaria* ("white dunes") [2130]
- Fixed coastal dunes with herbaceous vegetation ("grey dunes") [2150]
- Atlantic decalcified fixed dunes (Calluno-Ulicetea)
- 21A0] Machairs (* in Ireland)
- [1220] Perennial vegetation of stony banks
- [1395] *Petalophyllum ralfsii*
- [1310] *Salicornia* and other annuals colonizing mud and sand
- [8330] Submerged or partly submerged sea caves

Many sites with the above habitats in County Mayo have 'Human recreational activities, Golf links, walkways, increased human presence / walkways and recreational activities' listed as a threat on their Natura 2000 Standard Data forms.

Leisure fishing may also pose a threat to sites with aquatic habitats and species. For example, the River Moy is Ireland's premier Salmon river. Increased recreational fishing may result in increased pressure on this river and could result in detrimental impacts on Atlantic Salmon, one of the qualifying interests of this site.

In order to ensure a balanced, pragmatic approach, we would work closely with the Forward Planning Team and would liaise with other bodies such as Fáilte Ireland and Inland Fisheries Ireland to ensure that increased tourism and recreational activity in County Mayo does not result in significant adverse effects on coastal, terrestrial and freshwater habitats. We also recognise the increasing level of importance being put on community-led approaches to planning and management of resources, through legislation such as the Water Framework Directive. For example, The Maharees Conservation Association in County Kerry has had great success in restoring sand dune ecosystems through a community-led initiative, supported by Kerry County Council, to tackle problems associated with increased tourism and leisure activities. We would therefore also propose to work with Mayo County Council to engage with communities early on in the development of the plan; such an approach may reduce the number of amendments required following the official consultation process.

6.4 Climate Change

There is a requirement in the AA process to consider the effects of climate change on Natura 2000 sites and how this may interact with the impacts and in-combination effects of the plan or project being considered. This can be somewhat challenging, given the uncertainties of climate change projections and the range of habitats that may be affected.

The Environmental Protection Agency's Climate Change Research Programme (CCRP) carries out relevant and up to date studies on climate change in Ireland. Analysis of the meteorological records shows that Ireland's climate is changing in line with global patterns.

The clearest trend is evident in the temperature records which show a mean temperature increase of 0.7°C between 1890 and 2008, i.e. an increase of 0.06°C per decade. The increase was 0.4°C during the period 1980-2008, i.e. equivalent to 0.14°C per decade. Other indicators of climate change in Ireland are 1) six of the ten warmest years in Ireland have occurred since 1990, 2) a reduction in the number of frost days and shortening of frost season length, and 3) an increase in annual rainfall in northern and western areas with decreases or small increases in the south and east.

These changes are reflected in Ireland's natural environment with an increase in the growing season and with greater number of animals suited to warmer temperatures being evident in Ireland and its surrounding waters.

Climate change impacts are projected to increase in the coming decades and during the rest of this century. Uncertainties remain in relation to the scale and extent of these impacts, particularly during the second half of the century. The greatest uncertainty lies in how effective global actions will be in reducing greenhouse gas emissions. Predicted adverse impacts include more intense storms and rainfall events, an increased likelihood and magnitude of river and coastal flooding, adverse impacts on water quality and changes in distribution of plant and animal species.

Mayo County Council has developed a Climate Adaptation Strategy named Climate Ready Mayo (Mayo CoCo, 2019). This Strategy sets out a vision for a County that will be climate ready, a County that understands how climate change will affect communities and businesses, and a County that works together to reduce the risk and avail of the opportunities that climate change will bring.

6.4.1 Climate change and biodiversity

Impacts that may occur to the Natura 2000 network of County Mayo as a result of climate change are discussed extensively in Section 8. However, climate change can also cause cumulative and in-combination impacts.

Impacts that will affect wetlands and water dependant ecosystems as a result of climate change, will be exacerbated by extractions of increasingly larger amounts of water to meet the needs of the population. Bogs, wetlands and groundwater dependant terrestrial ecosystems that are already under pressure from drainage and extraction will suffer as a result of the cumulative and in-combination impacts from the lowering of the water table due to reduced precipitation and extraction of water from the agricultural industry.

Rising sea levels, alongside increase in temperatures on land and in the water, will put cumulative pressure on marine and coastal habitats that are already under pressure from development. An increase in temperatures can lead to the direct impacts that have been discussed previously, but they can also cause the spread of non-native invasive species more suited to warmer temperatures. The impact of the spread of these species can cause cumulative impacts upon the integrity of the Natura 2000 network through loss and degradation of habitats present as well as loss of species such as those caused by the spread of the crayfish plague.

Mitigation measures are required to mitigate against the potential cumulative and in-combination impacts of climate change on the integrity of the Natura 2000 network in and around County Mayo. These are in Section 9.

6.5 Cumulative and in-combination impacts

Cumulative and in-combination impacts are assessed and listed in Appendix D. 017 in vicinity of proposed site.

6.6 Summary

Cumulative impacts have been assessed (see Appendix D). There may be an increase in pressures upon designated features as a result of actions that will take place in the implementation of these plans, any potential impacts must be determined through AA screening at a project level. Project level assessment will ensure that cumulative impacts are addressed and therefore, no negative effects were determined to result from cumulative or in-combination impacts, between this River Basins Plan and the plans and policies assessed.

Given the presence of pathways and the potential impacts posed by the proposed Plan on the Natura 2000 sites (Table 7-2), it is concluded that it is not possible to screen out likely significant effects on these Natura 2000 sites.

7 Appropriate Assessment Screening

7.1 Introduction

Assessment of the potential impacts of Mayo County Development Plans aims, objectives, strategies and policies within the Plan as described, are required under regulation 42 of the European Communities (Birds and Natural Habitats) Regulations 2011 (S.I. No. 477 of 2011).

This section aims to identify whether the proposed Plans aims, objectives, strategies and policies are likely to have a significant effect, either alone, or in-combination with other projects and plans, on the Natura 2000 sites within the zone of influence.

The 'screening' process addresses and records the reasoning and conclusions in relation to the first two tests of Article 6(3) of the Habitats Directive:

- Is the plan or programme directly connected to or necessary for the management of the site; and
- Will the plan or programme, alone or in-combination with other plans and projects, have a significant effect on a Natura 2000 site in view of its conservation objectives.

If the effects are deemed to be significant, potentially significant or uncertain, then the plan or programme that is under assessment is subject to a Stage 2 Appropriate Assessment, reported in the form of a Natura Impact Report.

The aims, objectives, strategies and policies of Mayo County Development Plan are not directly connected to the management of any Natura 2000 sites, however, they could have potential to cause significant effects on Natura 2000 sites.

7.2 Screening methodology

In accordance with DEHLG guidance, the key to determining if an Appropriate Assessment is required for a Plan, is in the assessment of whether the plan and its policies and objectives are likely to have a significant effect on a Natura 2000 site.

For this process, the screening of this plan has been broken down into 4 steps.

1. Description of the programme (Section 2);
2. Screening of Natura 2000 sites within the various zones of influence of the plan dependent on the presence of potential pathways and nature of the qualifying interests (Appendix B).
3. Assessing the measures to identify potential impacts. Determining the significance of these potential impacts and the requirement for follow up assessments. This is presented in Appendix A.
4. Screening Statement with conclusions. This is presented in Section 7.5.

7.3 Study Area

The following section describes the screening methods used at various scales to ensure inclusion of all Natura 2000 sites that may be potentially impacted by all objectives and measures of the Plan.

Identification of Relevant Natura 2000 sites

The relevant Natura 2000 sites were identified dependent on several factors. This included those within the River Basin, those within 15km of the River Basin and those connected hydrologically either through groundwater or surface water pathways as defined by the WFD and EPA. Natura 2000 sites containing Freshwater Pearl Mussel (FWPM) *Margaritifera margaritifera* or *Margaritifera durrovensis* within 35km were included, as were Natura 2000 sites that contained Groundwater Dependent Terrestrial Ecosystems (GWDTEs) that were

hydrologically connected. Consultation was carried out with EPA staff in relation to the method for screenings of GWDTEs.

Data for this process was obtained from the WFD and NPWS website and JBA derived data, then visualised and assessed using QGIS and Microsoft Excel. Further details of the relevant Natura 2000 sites within the likely zones of influence of the Plan are presented in tables in Section 7.4.

The zones of impact of consideration for potential impacts on Natura 2000 sites are discussed here:

County scale

Natura 2000 sites that are within the County, and therefore within the area of the Plan, are at risk of direct and indirect impacts as a result of the objectives and measures of the Plan.

15km Scale

A buffer area of 15km was used for selection of Natura 2000 sites, based on DEHLG Guidance (DEHLG, 2010), which is the distance considered appropriate for Plans. This 15km buffer also addresses the potential land and air pathways, as the distances defined in Ryan Hanley (2014) for land and air pathways, are well accommodated within 15km. This distance was evaluated on a case by case basis, dependent on the nature of the Qualifying Interests present. The method used in this current assessment is a slight variation on the method of Ryan Hanley (2014), as it uses more up to date information regarding potential pathways present in a catchment. The Ryan Hanley method, if it were used, would remove Natura 2000 sites with no surface water connectivity to a County, but which are situated within the catchment of a county and this may result in the exclusion of a Natura 2000 sites that may be potentially impacted by land and air pathways. Therefore, the use of the County and 15km buffer will, by default, include Natura 2000 sites potentially impacted by all relevant pathways.

Hydrologically connected

Hydrological connections between the River Basin and Natura 2000 sites were identified through the use of WFD defined active aquifers and EPA defined river network. These connections can increase or reduce the number of Natura 2000 sites at risk from potential impacts as a result of the objectives and measures of the Plan.

Groundwater pathways with the potential to transport impact to GWDTE of Natura 2000 sites, are defined by the use of active groundwater bodies shapefiles from the EPA Envision Maps, 2020. This method is based upon the WFD risk assessments and WFD GWDTE maps. According to the WFD assessment guidance, the impact of pollutants or nutrients within the zone of influence varies according to a number of factors including aquifer vulnerability.

Freshwater Pearl Mussel (FWMP) and Groundwater Dependent Terrestrial Ecosystems (GWDTE)

In order to take into account Freshwater Pearl Mussel (*Margaritifera margaritifera* and/or *Margaritifera durrovensis*) populations, all Natura 2000 sites within 35km of the River Basin (adapted from Ryan Hanley (2014)) were selected to include for Natura 2000 sites containing Freshwater Pearl Mussel populations. In relation to Freshwater Pearl Mussel, Natura 2000 sites that have FWPM as a qualifying interest were assessed and then on a case by case basis, the location of possible FWPM populations were examined.

Natura 2000 sites located outside of the River Basin's surface water and groundwater catchments were screened out on the basis that there would be no impact on Natura 2000 sites outside of the catchment. Those sites within River Basin then proceeded to a more detailed review by an ecologist in JBA and were either screened in or out on the basis of the following criteria:

Distance from the River Basin at various levels based upon an adapted methodology from Ryan Hanley (2014b) including the WFD surface, 15km buffer and groundwater catchments and the 35km downstream buffer for FWPM;

- Hydrological connectivity to River Basin;
- Qualifying interests and special conservation interests for which the site was selected and their sensitivities e.g. GWDTEs; and
- The conservation objectives for those sites.

7.4 Findings of the Screening Process

138 Natura 2000 sites were screened for potential impacts as they were determined to be within at least one of the previously described zones of influence (Section 7.3). 73 Natura 2000 sites are within County Mayo. These 73 Natura 2000 sites will require further assessment for potential impacts.

Of the 63 Natura 2000 sites outside County Mayo, 32 sites are within 15 km and are hydrologically connected. These 32 sites can be included with the original 73 sites that require further assessment for potential impacts totalling 105 Natura 2000 sites requiring further assessment. Another 2 Natura 2000 sites are within 15km but are not hydrologically connected, these 2 sites will not need to be considered further due to lack of pathway for transporting impacts.

Of the remaining 31 sites, 8 sites are connected only through Groundwater pathways but are greater than 15km away from the County. Ten sites are only connected through surface water pathways but are greater than 15km from the County and do not have Freshwater Pearl Mussel as a Qualifying Interest, therefore, do not require further assessment in keeping with the methodology as outlined in Section 1.4 and Section 2.5.1. Eight sites are connected via surface water and groundwater pathways but are greater than 15km from the County and so effects of dilution can be considered. Of the original 16 sites located greater than 15km from the County and are connected through groundwater pathways to the County, 15 have Groundwater Dependent Terrestrial Ecosystems. However, as they are beyond 15km they are unlikely to be adversely impacted by the project. The remaining four sites are not hydrologically connected to the County, therefore, do not require further assessment.

Of the 105 screened in Natura 2000 sites, Lough Corrib SAC, Mweelrea/Sheffry/Erriff Complex SAC and Newport River SAC have Freshwater Pearl Mussel as a Qualifying Interest and are located inside of the County so these have already been considered in the overall figure of 105 Natura 2000 sites. The Twelve Bens/Garraun Complex SAC has Freshwater Pearl Mussel as a Qualifying Interest but is located outside of the County. This SAC is not connected via surface water to the County but is connected by groundwater pathway.

In conclusion, a total of 105 Natura 2000 sites are within, or hydrologically connected to the County, may be impacted by measures implemented within the County and so may require further assessment depending on the nature of the potential impacts, as determined in Section 8 and Appendix A. As discussed in Section 5.3, as part of Chapter 12 of the Mayo CDP, settlement plans have been developed for a number of settlements. Works carried out as a result of Mayo CPD may result in impacts to Natura 2000 sites. For example, as a result of works associated with the Mayo CPD in Balla may result in impacts to Balla Turlough SAC via groundwater pathways due to the close vicinity of the Natura site to the settlement. Works associated with the settlement plan for Charlestown may result in impacts to the River Moy SAC which runs through the town via surface water pathways. Additionally, Natura 2000 sites may be impacted via land and air pathways as a result of the proposed settlement plants, for example the settlement plan for Newport may result in impacts to the Newport River SAC via land and air pathways due to the close vicinity of the Natura 2000 site. The 105 Natura 2000 sites, including the Natura 2000 sites potentially impacted

by proposed settlement plans, requiring further assessment are presented in Table 7-2 and their conservation objectives are listed in Appendix C.

Table 7-1: Natura 2000 sites screened and screening criteria

Site code	Natura 2000 sites listed	Within County	Distance (km)	FWPM	GWDTE	GW connected	SW connected
002268	Achill Head SAC	x			x	x	x
001626	Annaghmore Lough (Roscommon) SAC		35		x	x	
000461	Ardkill Turlough SAC	x			x	x	x
004133	Aughris Head SPA		19			x	x
001228	Aughrusbeg Machair And Lake SAC		19		x	x	x
000463	Balla Turlough SAC	x			x	x	x
002081	Ballinafad SAC	x				x	
000474	Ballymaglancy Cave, Cong SAC	x				x	
000622	Ballysadare Bay SAC		22		x	x	x
004129	Ballysadare Bay SPA		22		x	x	x
002118	Barnahallia Lough SAC		19		x	x	
001922	Bellacorick Bog Complex SAC	x			x	x	x
000466	Bellacorick Iron Flush SAC	x				x	x
002005	Bellacragher Saltmarsh SAC	x				x	x
000592	Bellanagare Bog SAC		13		x		x
004105	Bellanagare Bog SPA		12				x
004177	Bills Rocks SPA	x					x
004037	Blacksod Bay/Broad Haven SPA	x			x	x	x
000471	Brackloon Woods SAC	x				x	
001656	Bricklieve Mountains and Keishcorran SAC		11		x	x	x
000472	Broadhaven Bay SAC	x			x	x	x
000595	Callow Bog SAC		6		x	x	x
002347	Camderry Bog SAC		23		x		x

Site code	Natura 2000 sites listed	Within County	Distance (km)	FWPM	GWDTE	GW connected	SW connected
000597	Carrowbehy/Caher Bog SAC		1			x	x
000475	Carrowkeel Turlough SAC	x			x	x	x
000476	Carrowmore Lake Complex SAC	x			x	x	x
004052	Carrowmore Lake SPA	x				x	x
001242	Carrownagappul Bog SAC		26		x		x
002243	Clare Island Cliffs SAC	x				x	
004136	Clare Island SPA	x				x	x
001482	Clew Bay Complex SAC	x			x	x	x
001899	Cloonakillina Lough SAC	x			x	x	x
000600	Cloonchambers Bog SAC		5		x	x	x
000614	Cloonshanville Bog SAC		15		x		x
000479	Cloughmoyne SAC	x				x	
000480	Clyard Kettle-Holes SAC	x			x	x	x
002034	Connemara Bog Complex SAC		15		x	x	
004181	Connemara Bog Complex SPA		15			x	
000218	Coolcam Turlough SAC		6		x		x
002110	Corliskea/Trien/Cloonfelliv Bog SAC		10		x	x	x
000485	Corraun Plateau SAC	x				x	
001251	Cregduff Lough SAC		26		x		
001955	Croaghaun/Slievemore SAC	x				x	x
000255	Croaghill Turlough SAC		7		x		x
000484	Cross Lough (Killadoon) SAC	x				x	x
004212	Cross Lough (Killadoon) SPA	x				x	x
004170	Cruagh Island SPA		23				x
000627	Cummeen Strand/Drumcliff Bay (Sligo Bay) SAC		27				x

Site code	Natura 2000 sites listed	Within County	Distance (km)	FWPM	GWDTE	GW connected	SW connected
002350	Curraglehanagh Bog SAC		26		x		
000604	Derrinea Bog SAC		0.3		x	x	x
002197	Derrinlough (Cloonkeenleananode) Bog SAC		20		x	x	
001257	Dog's Bay SAC		27				x
000492	Doocastle Turlough SAC	x			x	x	x
004235	Doogort Machair SPA	x				x	x
001497	Doogort Machair/Lough Doo SAC	x				x	x
002338	Drumalough Bog SAC		4		x	x	x
000495	Duvillaun Islands SAC	x				x	x
004111	Duvillaun Islands SPA	x				x	x
001501	Erris Head SAC	x				x	
000607	Errit Lough SAC		0.5		x	x	x
000497	Flughany Bog SAC	x			x	x	x
000268	Galway Bay Complex SAC		23		x		x
000500	Glenamoy Bog Complex SAC	x			x		x
000503	Greaghans Turlough SAC	x			x	x	x
004144	High Island, Inishshark and Davillaun SPA		13				x
004074	Illanmaster SPA	x				x	x
004221	Illeannanooon SPA		11			x	
000278	Inishbofin And Inishshark SAC		7				x
004231	Inishbofin, Omev Island and Turbot Island SPA		7				
004084	Inishglora and Inishkeeragh SPA	x				x	x
000507	Inishkea Islands SAC	x				x	x
004004	Inishkea Islands SPA	x				x	x
004031	Inner Galway Bay SPA		24		x	x	x

Site code	Natura 2000 sites listed	Within County	Distance (km)	FWPM	GWDTE	GW connected	SW connected
001513	Keel Machair/Menaun Cliffs SAC	x				x	x
002320	Kildun Souterrain SAC	x				x	
000504	Kilglassan/Caheravoostia Turlough Complex SAC	x			x	x	x
002111	Kilkieran Bay And Islands SAC		22		x		
000458	Killala Bay/Moy Estuary SAC	x			x	x	x
004036	Killala Bay/Moy Estuary SPA	x			x	x	x
000285	Kilsallagh Bog SAC		14		x		x
002265	Kingstown Bay SAC		20		x	x	x
001669	Knockalongy and Knockachree Cliffs SAC		14			x	
000516	Lackan Saltmarsh and Kilcummin Head SAC	x				x	x
000295	Levally Lough SAC		17		x	x	
000296	Lisnageeragh Bog and Ballinastack Turlough SAC		15		x		x
001673	Lough Arrow SAC		16		x	x	x
004050	Lough Arrow SPA		16		x	x	x
001529	Lough Cahasy, Lough Baun And Roonah Lough SAC	x			x	x	x
004051	Lough Carra SPA	x				x	x
001774	Lough Carra/Mask Complex SAC	x			x	x	x
004228	Lough Conn and Lough Cullin SPA	x			x	x	x
000297	Lough Corrib SAC	x		x	x	x	x
004042	Lough Corrib SPA	x			x	x	x
002177	Lough Dahybaun SAC	x			x	x	x
000522	Lough Gall Bog SAC	x			x	x	x
004048	Lough Gara SPA		6			x	x
000633	Lough Hoe Bog SAC	x			x	x	x

Site code	Natura 2000 sites listed	Within County	Distance (km)	FWPM	GWDTE	GW connected	SW connected
000301	Lough Lurgheen Bog/Glenamaddy Turlough SAC		16		x		x
004062	Lough Mask SPA	x			x	x	x
000634	Lough Nabrickkeagh Bog SAC		2		x	x	x
002008	Maumturk Mountains SAC		0.4		x	x	
001536	Mocorha Lough SAC	x			x	x	x
002352	Monivea Bog SAC		31		x	x	
000527	Moore Hall (Lough Carra) SAC	x				x	x
004227	Mullet Peninsula SPA	x				x	x
000470	Mullet/Blacksod Bay Complex SAC	x			x	x	x
000612	Mullygollan Turlough SAC		31		x	x	
002129	Murvey Machair SAC		28				x
001932	Mweelrea/Sheefry/Erriff Complex SAC	x		x	x	x	x
002144	Newport River SAC	x		x		x	x
000532	Oldhead Wood SAC	x				x	
001309	Omey Island Machair SAC		28		x	x	
000534	Owenduff/Nepin Complex SAC	x			x	x	x
004098	Owenduff/Nepin Complex SPA	x				x	x
002006	Ox Mountains Bogs SAC	x			x	x	x
002298	River Moy SAC	x			x	x	x
001976	Rosroe Bog SAC		17		x		
001312	Ross Lake And Woods SAC		10		x		
001311	Rusheenduff Lough SAC		8		x	x	
000326	Shankill West Bog SAC		26		x		x
000525	Shrule Turlough SAC	x			x	x	x
000541	Skealaghan Turlough SAC	x			x	x	x

Site code	Natura 2000 sites listed	Within County	Distance (km)	FWPM	GWDTE	GW connected	SW connected
000542	Slieve Fyagh Bog SAC	x			x	x	x
000328	Slyne Head Islands SAC		27				x
002074	Slyne Head Peninsula SAC		23		x	x	
004072	Stags of Broad Haven SPA	x				x	x
000636	Templehouse And Cloonacleigha Loughs SAC		5		x	x	x
004093	Termoncarragh Lake and Annagh Machair SPA	x			x	x	x
002031	The Twelve Bens/Garraun Complex SAC		0.8	x	x	x	
002179	Towerhill House SAC	x				x	
002354	Tullaghanrock Bog SAC		6		x	x	x
002130	Tully Lough SAC		9		x	x	
000637	Turloughmore (Sligo) SAC		4		x	x	x
000638	Union Wood SAC		20			x	
001898	Unshin River SAC		14		x	x	x
001571	Urlaur Lakes SAC	x			x	x	x
002998	West Connacht Coast SAC	x					x
002296	Williamstown Turloughs SAC		7		x	x	x

Table 7-2: Natura 2000 sites requiring further assessment

Site Code	Site Name	Requirement for Appropriate Assessment
002268	Achill Head SAC	Required
000461	Ardkill Turlough SAC	Required
000463	Balla Turlough SAC	Required
002081	Ballinafad SAC	Required
000474	Ballymaglancy Cave, Cong SAC	Required
001922	Bellacorick Bog Complex SAC	Required
000466	Bellacorick Iron Flush SAC	Required
002005	Bellacragher Saltmarsh SAC	Required
004177	Bills Rocks SPA	Required

Site Code	Site Name	Requirement for Appropriate Assessment
004037	Blacksod Bay/Broad Haven SPA	Required
000471	Brackloon Woods SAC	Required
000472	Broadhaven Bay SAC	Required
000475	Carrowkeel Turlough SAC	Required
000476	Carrowmore Lake Complex SAC	Required
004052	Carrowmore Lake SPA	Required
002243	Clare Island Cliffs SAC	Required
004136	Clare Island SPA	Required
001482	Clew Bay Complex SAC	Required
001899	Cloonakillina Lough SAC	Required
000479	Cloughmoyne SAC	Required
000480	Clyard Kettle-Holes SAC	Required
000485	Corraun Plateau SAC	Required
001955	Croaghaun/Slievemore SAC	Required
000484	Cross Lough (Killadoon) SAC	Required
004212	Cross Lough (Killadoon) SPA	Required
000492	Doocastle Turlough SAC	Required
004235	Doogort Machair SPA	Required
001497	Doogort Machair/Lough Doo SAC	Required
000495	Duvillaun Islands SAC	Required
004111	Duvillaun Islands SPA	Required
001501	Erris Head SAC	Required
000497	Flughany Bog SAC	Required
000500	Glenamoy Bog Complex SAC	Required
000503	Greaghans Turlough SAC	Required
004074	Illanmaster SPA	Required
004084	Inishglora and Inishkeeragh SPA	Required
000507	Inishkea Islands SAC	Required
004004	Inishkea Islands SPA	Required
001513	Keel Machair/Menaun Cliffs SAC	Required
002320	Kildun Souterrain SAC	Required
000504	Kilglassan/Cahevavoostia Turlough Complex SAC	Required
000458	Killala Bay/Moy Estuary SAC	Required
004036	Killala Bay/Moy Estuary SPA	Required
000516	Lackan Saltmarsh and Kilcummin Head SAC	Required
001529	Lough Cahasy, Lough Baun And Roonah Lough SAC	Required
004051	Lough Carra SPA	Required
001774	Lough Carra/Mask Complex SAC	Required

Site Code	Site Name	Requirement for Appropriate Assessment
004228	Lough Conn and Lough Cullin SPA	Required
000297	Lough Corrib SAC	Required
004042	Lough Corrib SPA	Required
002177	Lough Dahybaun SAC	Required
000522	Lough Gall Bog SAC	Required
000633	Lough Hoe Bog SAC	Required
004062	Lough Mask SPA	Required
001536	Mocorha Lough SAC	Required
000527	Moore Hall (Lough Carra) SAC	Required
004227	Mullet Peninsula SPA	Required
000470	Mullet/Blacksod Bay Complex SAC	Required
001932	Mweelrea/Sheeffry/Erriff Complex SAC	Required
002144	Newport River SAC	Required
000532	Oldhead Wood SAC	Required
000534	Owenduff/Nephin Complex SAC	Required
004098	Owenduff/Nephin Complex SPA	Required
002006	Ox Mountains Bogs SAC	Required
002298	River Moy SAC	Required
000525	Shrule Turlough SAC	Required
000541	Skealaghan Turlough SAC	Required
000542	Slieve Fyagh Bog SAC	Required
004072	Stags of Broad Haven SPA	Required
004093	Termoncarragh Lake and Annagh Machair SPA	Required
002179	Towerhill House SAC	Required
001571	Urlaur Lakes SAC	Required
002998	West Connacht Coast SAC	Required
000614	Cloonshanville Bog SAC	Required
002034	Connemara Bog Complex SAC	Required
004181	Connemara Bog Complex SPA	Required
000296	Lisnageeragh Bog and Ballinastack Turlough SAC	Required
000285	Kilsallagh Bog SAC	Required
001669	Knockalongy and Knockachree Cliffs SAC	Required
001898	Unshin River SAC	Required
000592	Bellanagare Bog SAC	Required
004144	High Island, Inishshark and Davillaun SPA	Required
004105	Bellanagare Bog SPA	Required
001656	Bricklieve Mountains and Keishcorran SAC	Required
004221	Illeannan SPA	Required

Site Code	Site Name	Requirement for Appropriate Assessment
002110	Corliskea/Trien/Cloonfelliv Bog SAC	Required
002130	Tully Lough SAC	Required
001311	Rusheenduff Lough SAC	Required
000255	Croaghill Turlough SAC	Required
000278	Inishbofin And Inishshark SAC	Required
002296	Williamstown Turloughs SAC	Required
000595	Callow Bog SAC	Required
000218	Coolcam Turlough SAC	Required
004048	Lough Gara SPA	Required
002354	Tullaghanrock Bog SAC	Required
000600	Cloonchambers Bog SAC	Required
000636	Templehouse And Cloonacleigha Loughs SAC	Required
002338	Drumalough Bog SAC	Required
000637	Turloughmore (Sligo) SAC	Required
000634	Lough Nabrickkeagh Bog SAC	Required
000597	Carrowbehy/Caher Bog SAC	Required
002031	The Twelve Bens/Garraun Complex SAC	Required
000607	Errit Lough SAC	Required
002008	Maumturk Mountains SAC	Required
000604	Derrinea Bog SAC	Required

7.4.1 Qualifying Interests

The Qualifying interests for the Natura 2000 sites that were screened in are provided in Appendix B. Designated habitat types and species within the screened in SACs and SPAs have potential to be affected by the implementation of the proposed Plan.

7.4.2 Conservation Objectives

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Conservation objectives for SACs and SPAs (i.e. sites within the Natura 2000 network) are required for the habitats and species for which the sites are selected. Detailed site-specific conservation objectives have been provided for the majority of SACs and SPAs, which can be found within the Conservation Objectives document for each site on the NPWS website and Appendix C. Generic conservation objectives have been compiled for the remaining SAC and SPAs.

The overall aim of conservation objectives is for the maintenance or restoration of the favourable conservation conditions of the Annex I habitats and/or the Annex II species for which a SAC has been selected, under which the site-specific objectives contain more detailed attributes, measures and targets.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

The conservation objectives for SPAs are also to maintain or restore the favourable conservation condition of the bird species listed as Special Conservation Interests for SPAs, which are defined by the following list of attributes and targets;

- Population trend; Measure of percentage change and whether the long-term population trend stable or increasing.
- Distribution: Number, range, timing and intensity of use of areas. There is to be no significant decrease in the range, timing or intensity of use of areas by golden plover, other than that occurring from natural patterns of variation.

The conservation objective for non-breeding birds Special Conservation Interests for SPAs are as follows;

- To maintain the favourable conservation condition of the non-breeding water bird Special Conservation Interest species listed for a SPA.
- To maintain the favourable conservation condition of the wetland habitat for a SPA as a resource for the regularly occurring migratory water birds that utilise it.

The conservation objectives were considered when carrying out the AA screening process for the Mayo CDP that may potentially impact on Natura 2000 sites.

7.4.3 Cumulative Impact

Given the specificity of the relevant plans and project developments in Section 6 (see Appendix D), to the County, their connectivity in terms of surface water, groundwater and land & air pathways to the Natura 2000 sites is likely to be similar to the proposed site. With this in consideration, it can be stated that cumulative impacts are anticipated.

7.5 Summary

As outlined in Section 5 and Section 8, a number of objectives, aims, policies and strategies of the Plan are deemed to have a likely significant effect on Natura 2000 sites. Given that the implementation extent of these measures at Plan level is throughout the County and River Basins, those Natura 2000 sites listed in Table 7-2 could potentially be impacted as a result.

The proposed Plan is not directly connected with or necessary to the management of Natura 2000 sites. It is, accordingly, necessary for the competent authority to assess whether the proposed Plan, either individually or in combination with other plans or projects, would be likely to have significant effects on any Natura 2000 site.

Cumulative impacts have been assessed. There may be an increase in ecological and environmental pressures upon designated features as a result of actions that will take place in the implementation of these plans, however, any potential impacts must be determined through AA screening at a project level. Project level assessment will ensure that cumulative

impacts are addressed and therefore, no negative effects were determined to result from cumulative or in-combination impacts, because of the plans and policies assessed.

Given the presence of pathways and the potential impacts posed by the proposed Plan on the 105 Natura 2000 sites (Table 7-2), it is concluded by the authors of this report that it is not possible to rule out (screen out) likely significant effects on these Natura 2000 sites. Therefore, it is recommended by the authors of this report that the proposed Plan should be brought forward to the second stage of the Appropriate Assessment process. The resulting Natura Impact Statement follows.

8 Natura Impact Report

This Natura Impact Report (NIR) aims to determine whether the proposed Plan would result in significant adverse impacts on the integrity of any Natura 2000 site with respect to the site's structure, function, and/ or conservation objectives as defined in Appendix C. It also aims to provide supporting information for the competent authority to carry out an Appropriate Assessment of the proposed Plan.

Mayo County Development Plan is a high-level plan, which provides objectives, aims, strategies and policies for progression within Mayo. In general, this proposed Plan does not detail specifics of any potential projects that may be implemented that could cause significant effects on Natura 2000 sites although some specific locations are highlighted, and some specific projects are supported or guided as a result of the Plan. The sources of potential impacts are discussed in Section 3 and potential impacts are in Section 5. Thus, the potential significant effects from potential impacts as identified in Section 5 are discussed below in Section 8.1, in terms of the general measures being implemented throughout the County.

Overall, the Plan lays out a framework for development, promoting and guiding residential, commercial, recreational and infrastructural development within the County. There is likely to be direct and in-direct impacts upon the overall integrity of the Natura 2000 sites that have been screened in as in Table 7-2. The potential impacts identified in Section 3 are in general, related to the pressure that development places on nature and the environment.

Potential impacts that may arise are outlined below and discussed in further detail to determine significance based on their impact upon the overall integrity of the relevant Natura 2000 sites and their conservation objectives. Their conservation objectives are in Appendix C.

8.1 Potential significant effects

As outlined in Section 5, the potential impacts that could occur through the implementation of the Plan are;

- Loss/ reduction of habitat area
- Disturbance to key species
- Habitat or species population fragmentation
- Reduction in species density
- Changes in key indicators of conservation value, such as changes in water quality and quantity.

8.1.1 Loss or reduction of habitat area

Direct habitat loss is caused where there is complete removal of a habitat type. Loss of habitat can also occur through the deterioration of habitat quality and therefore a loss of the function of that habitat. This can be due to factors such as land take requirements, physical alterations and introduction of invasive species non-native as a result of development or works necessary to promote sustainable or any type of development. The outcomes of the objectives, aims, strategies and policies of the Plan, such as development leading to population increase, reduction in vacancy of existing buildings and the support of the growth of towns and villages could potentially cause these impacts.

There is potential that the Natura 2000 sites screened into this assessment may be adversely affected in this manner through the implementation of the proposed Plan.

Indirect habitat loss may also occur through disruption of hydrological links to a designated site that supports surface or ground water dependent habitats such as peatland and wetland habitats e.g. implementation of SuDs and diversion of water flows that may occur due to

new housing or retail development. The objectives of the proposed Plan, such as BAO1 which could result in the disruption of hydrological links in the settlement of Balla which could negatively impact the Qualifying Feature (Turloughs [3180]) of the nearby Balla Turlough SAC. The disruption of this hydrological link may result in a reduction of the water table directly impacting the site.

The implementation of environmental mitigation measures throughout County wide development and growth, although primarily beneficial in the provision of refuges for species and the protection and restoration of habitats including bog and wetland habitats, may result in changes of land use within the County. If measures are inappropriately implemented, habitat loss or degradation of habitat quality may occur. For example, the inappropriate implementation of measures may result in a reduction of water quality negatively impacting Qualifying Interests which rely on good water quality such as Freshwater Pearl Mussels in the Newport River SAC. A decline in water quality may also result in a decline in Salmonid abundance and therefore disrupting the life cycle of Freshwater Pearl Mussels.

As the objectives of the proposed Plan are set at a high level, loss or reduction of habitat area of the Natura 2000 sites screened into this assessment cannot be ruled out at this stage. Therefore, the significance of potential habitat loss will be assessed at project level and will be dependent on the final locations and detailed design of the proposed objectives.

Mitigation measures are detailed in Section 9 to ensure the potential for habitat loss will not adversely affect the integrity of a Natura 2000 site due to the implementation of the proposed Plan.

8.1.2 Disturbance to key species

Key species are those species listed within the annexes of the Habitats Directive for which Natura sites are designated. Disturbance to species supported by a Natura 2000 site may result due to physical disturbance of a habitat that may result in direct mortalities or displacement of the species. Sources of disturbance may also take the form of increased noise, visual presence of people, vibration and increased illumination of areas through increased or expanded development without mitigation measures in place. Increased disturbance levels within the sensitivity threshold of a species may cause the displacement or flight of a species from their respective habitat.

In relation to the proposed Plan, activities that may potentially disturb key species are those that will result in physical disturbance due to works being carried out, such as the implementation of the policies and/or objectives of the plan including increased tourism, population growth or development in general. These objectives may pose sources of disturbance during construction of works and their operation. The works may also cause disturbance to the resting and/ or breeding places and foraging areas of these key species.

The objectives of the proposed Plan have the potential to cause disturbance impacts to key species of the Natura sites screened into this assessment. As these objectives are set at a high level and the location has not been determined, the significance of disturbance to key species will be assessed at project level and will be dependent on the final locations and detailed design of the proposed objectives. As an example, objective FDO 2, exploring the creation of a new river walk, could cause disturbance to Otters, a Qualifying Interest of the River Moy SAC, may be disrupted due to increased noise, vibration and illumination if objectives are implemented in the settlement of Foxford.

Of the SACs and SPAs (Natura 2000 sites) screened into the assessment, the species that could be potentially be disturbed as a result of the Plan are listed below in Table 8-1.

Mitigation measures are detailed in Section 9 to ensure that disturbance to key species will not adversely affect the integrity of a Natura 2000 site due to the implementation of the proposed Plan.

Table 8-1: Key species of the Screened-In Natura 2000 sites

Species	Natura 2000 site
Arctic Tern	High Island, Inishshark and Davillaun SPA
	Inishglora and Inishkeeragh SPA
	Inishkea Islands SPA
	Lough Corrib SPA
Barnacle Goose	High Island, Inishshark and Davillaun SPA
	Termoncarragh Lake and Annagh Machair SPA
	Duvillaun Islands SPA
	Inishglora and Inishkeeragh SPA
	Inishkea Islands SPA
Bar-tailed Godwit	Blacksod Bay/Broad Haven SPA
	Killala Bay/Moy Estuary SPA
Brook Lamprey	River Moy SAC
	Lough Corrib SAC
Chough	Termoncarragh Lake and Annagh Machair SPA
	Clare Island SPA
Common Gull	Connemara Bog Complex SPA
	Clare Island SPA
	Inishkea Islands SPA
	Lough Carra SPA
	Lough Conn and Lough Cullin SPA
	Lough Corrib SPA

Species	Natura 2000 site
	Lough Mask SPA
Cormorant	Connemara Bog Complex SPA
	Inishglora and Inishkeeragh SPA
Dunlin	Termoncarragh Lake and Annagh Machair SPA
	Blacksod Bay/Broad Haven SPA
	Doogort Machair SPA
	Inishkea Islands SPA
	Killala Bay/Moy Estuary SPA
Fulmar	High Island, Inishshark and Davillaun SPA
	Clare Island SPA
	Duvillaun Islands SPA
Geyer's whorl snail	Ox Mountains Bogs SAC
	Bellacorick Bog Complex SAC
	Clew Bay Complex SAC
	Lough Hoe Bog SAC
	Mweelrea/Sheeffry/Erriff Complex SAC
Golden Plover	Connemara Bog Complex SPA
	Killala Bay/Moy Estuary SPA
	Owenduff/Nephin Complex SPA
	Lough Corrib SPA
Greenland white-	Lough Gara SPA
	Termoncarragh Lake and Annagh Machair SPA

Species	Natura 2000 site
fronted Goose	Lough Conn and Lough Cullin SPA
	Lough Corrib SPA
	Lough Mask SPA
Grey Plover	Killala Bay/Moy Estuary SPA
Harbour Seal	Clew Bay Complex SAC
	Inishkea Islands SAC
	Killala Bay/Moy Estuary SAC
Herring Gull	Inishglora and Inishkeeragh SPA
	Inishkea Islands SPA
Killarney Fern	Knockalongy and Knockachree Cliffs SAC
Leach's Storm-petrel	Stags of Broad Haven SPA
Light-belled Brent Goose	Blacksod Bay/Broad Haven SPA
Marsh Fritillary	Connemara Bog Complex SAC
	Bricklieve Mountains and Keishcorran SAC
Marsh Saxifrage	Owenduff/Nephin Complex SAC
	Ox Mountains Bogs SAC
	Bellacorick Bog Complex SAC
	Bellacorick Iron Flush SAC
	Carrowmore Lake Complex SAC

Species	Natura 2000 site
	Glenamoy Bog Complex SAC
Merlin	Connemara Bog Complex SPA
	Owenduff/Nephin Complex SPA
Narrow-mouthed Whorl Snail	Killala Bay/Moy Estuary SAC
	Mweelrea/Sheeffry/Erriff Complex SAC
Otter	Connemara Bog Complex SAC
	Owenduff/Nephin Complex SAC
	Mullet/Blacksod Bay Complex SAC
	River Moy SAC
	Clew Bay Complex SAC
	Lough Carra/Mask Complex SAC
	Lough Corrib SAC
	Mweelrea/Sheeffry/Erriff Complex SAC
	Unshin River SAC
	The Twelve Bens/Garraun Complex SAC
Petalwort	Mullet/Blacksod Bay Complex SAC
	Doogort Machair/Lough Doo SAC
	Glenamoy Bog Complex SAC
	Inishkea Islands SAC
	Keel Machair/Menaun Cliffs SAC
	Mweelrea/Sheeffry/Erriff Complex SAC
Redshank	Killala Bay/Moy Estuary SPA

Species	Natura 2000 site
Salmon	Connemara Bog Complex SAC
	Owenduff/Nephin Complex SAC
	Newport River SAC
	River Moy SAC
	Glenamoy Bog Complex SAC
	Lough Corrib SAC
	Mweelrea/Sheeffry/Erriff Complex SAC
	Unshin River SAC
	The Twelve Bens/Garraun Complex SAC
	Maumturk Mountains SAC
Sanderling	Blacksod Bay/Broad Haven SPA
	Inishkea Islands SPA
	Killala Bay/Moy Estuary SPA
Sandwich Tern	Illaunnaon SPA
	Blacksod Bay/Broad Haven SPA
	Carrowmore Lake SPA
	Cross Lough (Killadoon) SPA
Sea Lamprey	River Moy SAC
	Killala Bay/Moy Estuary SAC
	Lough Corrib SAC
Shag	Clare Island SPA
	Inishglora and Inishkeeragh SPA

Species	Natura 2000 site
	Inishkea Islands SPA
Slender Green Feather-moss	Owenduff/Nephin Complex SAC
	Carrowmore Lake Complex SAC
	Glenamoy Bog Complex SAC
	Lough Carra/Mask Complex SAC
	Lough Corrib SAC
Slender naiad	Connemara Bog Complex SAC
	Lough Corrib SAC
	Lough Dahybaun SAC
	Mweelrea/Sheeffry/Erriff Complex SAC
	Tully Lough SAC
	Rusheenduff Lough SAC
	The Twelve Bens/Garraun Complex SAC
	Maumturk Mountains SAC
Storm Petrel	Stags of Broad Haven SPA
	Bills Rocks SPA
	Duvillaun Islands SPA
	Illanmaster SPA
	Inishglora and Inishkeeragh SPA
Tufted Duck	Lough Conn and Lough Cullin SPA
	Lough Corrib SPA
	Lough Mask SPA

Species	Natura 2000 site
White-clawed Crayfish	River Moy SAC
	Lough Corrib SAC
	Lough Hoe Bog SAC
	Bricklieve Mountains and Keishcorran SAC
Whooper Swan	Lough Gara SPA
	Bellanagare Bog SPA
	Termoncarragh Lake and Annagh Machair SPA
Corncrake	Termoncarragh Lake and Annagh Machair SPA
	Mullet Peninsula SPA
Lapwing	Termoncarragh Lake and Annagh Machair SPA
Freshwater Pearl Mussel	Newport River SAC
	Lough Corrib SAC
	Mweelrea/Sheeffry/Erriff Complex SAC
	The Twelve Bens/Garraun Complex SAC
Lesser Horseshoe Bat	Ballinafad SAC
	Ballymaglancy Cave, Cong SAC
	Kildun Souterrain SAC
	Lough Carra/Mask Complex SAC
	Lough Corrib SAC
	Moore Hall (Lough Carra) SAC
	Towerhill House SAC
Puffin	Bills Rocks SPA

Species	Natura 2000 site
Red-throated Diver	Blacksod Bay/Broad Haven SPA
Great Northern Diver	Blacksod Bay/Broad Haven SPA
Slavonian Grebe	Blacksod Bay/Broad Haven SPA
Common Scoter	Blacksod Bay/Broad Haven SPA
	Lough Conn and Lough Cullin SPA
	Lough Corrib SPA
Red-breasted Merganser	Blacksod Bay/Broad Haven SPA
Ringed Plover	Blacksod Bay/Broad Haven SPA
	Inishkea Islands SPA
	Killala Bay/Moy Estuary SPA
Curlew	Blacksod Bay/Broad Haven SPA
	Killala Bay/Moy Estuary SPA
Kittiwake	Clare Island SPA
Guillemot	Clare Island SPA
Razorbill	Clare Island SPA
Grey Seal	Duvillaun Islands SAC
	Inishbofin and Inishshark SAC
Common Bottlenose Dolphin	Duvillaun Islands SAC
	West Connacht Coast SAC

Species	Natura 2000 site
Lesser Black-backed Gull	Inishglora and Inishkeeragh SPA
Purple Sandpiper	Inishkea Islands SPA
Turnstone	Inishkea Islands SPA
Little Tern	Inishkea Islands SPA
Gadwall	Lough Corrib SPA
Shoveler	Lough Corrib SPA
Pochard	Lough Corrib SPA
Hen Harrier	Lough Corrib SPA
Coot	Lough Corrib SPA
Black-headed Gull	Lough Corrib SPA
	Lough Mask SPA
Common Tern	Lough Corrib SPA

8.1.3 Fragmentation

Habitat and species population fragmentation occurs when the connectivity between habitats and species is restricted or lost and impedes the natural movement of a species. This can result in singular habitat units being formed and isolated species communities. Fragmentation can arise as a result of barriers to migratory movements where important corridors exist, e.g. river, riparian and coastal corridors. The inappropriate installation of measures that may act as a barrier to free movement of a species and/ or physically separate a habitat, will result in fragmentation.

The implementation of the policies and/or objectives of the plan including increased tourism, population growth or development in general may pose sources of disturbance during construction of works and their operation that can lead to fragmentation. They may result in the installation of structures that may impeded movement of a species. For example, the unmitigated installation of a bridge needed for access to a housing estate may impede the movement of Salmon within a river catchment and act as a migratory barrier. This may in turn prevent Salmon reaching Freshwater Pearl Mussel located upstream, thus

breaking the lifecycle of Freshwater Pearl Mussel and creating an isolated population. Long term this could result in the end of this population if not mitigated against.

As the objectives of the proposed Plan are not specific regarding location of implementation, fragmentation of the Natura 2000 sites screened into this assessment cannot be ruled out. Also, the significance of potential fragmentation will be required to be assessed at project level and will be dependent on the final locations and detailed design of the resulting projects that arise from this Plan. An example of where species fragmentation could occur is if objective GSO 11 is implemented in the settlement of Newport, where increased development and tourism could impact the movement of Salmon into the Newport River SAC either due to the installation of structures which may impede Salmon migration or degradation of the habitat due to increased pollution making the SAC unsuitable for the QI. In this case, the fragmentation of Salmon could impact the ability of Freshwater Pearl Mussel to complete its lifecycle.

Mitigation measures are detailed in Section 9 to ensure that fragmentation will not adversely affect the integrity of a Natura 2000 site due to the implementation of the proposed Plan.

8.1.4 Reduction in species density

Reduction in the density of species may arise due to the factors discussed above; habitat loss or degradation, disturbance and fragmentation. As outlined above, the objectives of the proposed Plan have the potential to cause these impacts. The implementation of the objectives for rural development could also result in development in rural areas where species are sensitive to human activity.

As this is a high level plan and the objectives, policies, strategies and aims of the proposed Plan are not specific regarding location of implementation, reduction in species density of the Natura 2000 sites screened into this assessment cannot be ruled out. Therefore, the significance of potential reduction of species density will be assessed at project level and will be dependent on the final locations and detailed design of the proposed objectives. A decline in water quality and/or species fragmentation of the River Moy SAC due to the implementation of objectives (such as FDO 3) for the settlement of Foxford may impede migratory species (Salmon, Brook Lamprey, Sea Lamprey) from travelling, potentially impacting the reproductive success of these species, resulting in an decrease in species density in the Natura 2000 site in the long term.

Mitigation measures are detailed in Section 9 to ensure that reduction in species density will not adversely affect the integrity of a Natura 2000 site due to the implementation of the proposed Plan.

8.1.5 Changes in key indicators of conservation value

The key indicators of conservation value for the Natura sites likely to be affected by the implementation of the proposed Plan include surface water and groundwater quality and quantity. Waterbodies are one of the key pathways for impact and are easily affected by impacts themselves. As outlined in the screening process and Table 7-2, the Natura sites either fall within the area of the River Basin or are hydrologically connected to the area of the proposed Plan. Any deterioration in water quality could potentially result in adverse impacts, either directly or indirectly, on a range of habitats and species that are surface water or groundwater dependent, such as lakes, rivers, coastal lagoons and bays, seals, Salmon, Otter and Freshwater Pearl Mussel. Maintaining the hydrological regime, e.g. recharge, flow rates and direction, of these habitats is also very important for the function and structure of the habitats and the species that they support.

The outcomes of the objectives, aims, strategies and policies of the Plan, such as development leading to population increase, reduction in vacancy of existing buildings and

the support of the growth of towns and villages could potentially cause impacts to water quality and quantity. These pressures would mainly arise from the activity of both land-based and in-stream works, where there is the potential for discharge of silt laden runoff and pollutants such as hydrocarbons to coastal, freshwater and groundwater bodies. Hydrological alterations may also occur through the potential diversion of flows and implementation of SuDs and natural flood risk management measures that are used for sustainable development and growth. Therefore, all projects that arise as a result of the objectives, aims and strategies of this Plan must be appropriately assessed for their suitability at a given location.

As the objectives of the proposed Plan are not specific regarding location of implementation, changes in key indicators of conservation value of the Natura 2000 sites screened into this assessment cannot be ruled out. Therefore, the significance of potential changes in key indicators will be assessed at project level and will be dependent on the final locations and detailed design of the proposed objectives. Implementation of objectives around the settlement of Foxford has potential to cause an increase in dust emissions and release of pollutants via surface water runoff resulting in a decline in water quality of the River Moy SAC which divides the settlement. This would affect the conservation value of the Natura 2000 site, impacting QIs such as Salmon, Otter, White-clawed Crayfish, Sea Lamprey and Brook Lamprey which rely on good water quality.

Mitigation measures are detailed in Section 9 to ensure that changes in key indicators of conservation value will not adversely affect the integrity of a Natura 2000 site due to the implementation of the proposed Plan.

8.2 Implementation routes for physical works

Measures or projects arising from the objectives, aims, strategies or policies of the Plan requiring physical works may either require planning consent or confirmation, or will be an exempted development.

Works that will require planning consent or confirmation, will be carried out by either a private developer or the Local Authority. Works may progress to construction stage as one of the following:

- Project led by private developer in line with the Aims, policies, objectives or strategies of the Plan.
- Project led by the Local Authority under the Planning and Development Regulations.
- Project led by the Local Authority under the Strategic Infrastructure Act.

Project level assessments that may be required for all types of project include:

- Environmental Impact Assessment: For a project above the thresholds specified under Article 24 of the European Communities (Environmental Impact Assessment) Regulations, 1989 as amended or a project likely to have significant effects on the environment, having regard to the criteria specified for under Article 27 of the same EIA Regulations 1989 as amended.
- Appropriate Assessment: All projects will be screened for Appropriate Assessment and, where there is a potential for a significant effect on a European (Natura 2000) site, an Appropriate Assessment will be undertaken in accordance the European Communities (Birds and Natural Habitats) Regulations 2011.

Exempted developments include those of limited scale and scope, that may fall under the category of flood mitigation works or housing protection schemes. Exempted developments may be carried out by Local Authorities under funding by the OPW, will be exempted in

accordance with the Planning and Development Act 2000 (as amended) and will comply with all relevant environmental legislation. This could require the undertaking of an EIA or AA screening for physical works. Local Authorities must supply written confirmation of legislative compliance under condition of funding.

9 Mitigation Measures

Mayo County Development Plan will not in itself, cause any adverse impacts to the overall integrity of the Natura 2000 sites within the ZOI of the Plan. However, Projects that arise as a result of the Plan may have adverse effects.

Projects stemming from the Mayo County Development Plan will apply a range of standard processes and measures that will mitigate potential environmental impacts. While the applicability of processes and particular measures will be dependent on the nature and scale of each project, examples of typical processes and measures that will be implemented where applicable at the different stages of project implementation are set out in the below sections.

It must also be noted that some Aims, Objectives, Strategies and policies of the Plan will increase the levels of environmental protection afforded to Natura 2000 sites and their conservation objectives e.g. Strategic Objective SO 8:

“Appropriate Assessment, Strategic Environmental Assessment and Strategic Flood Risk Assessment

a) To ensure the assessment of all planning applications in the Plan area have regard to the information, data and requirements of the Appropriate Assessment Natura Impact Report, SEA Environmental Report and Strategic Flood Risk Assessment Report contained in Appendices XX of the Mayo CDP 2021-2027

b) To require project planning to be fully informed by ecological and environmental constraints at the earliest stage of project development and any necessary assessment to be undertaken, including assessments of disturbance to species protected under the Wildlife Act and/or the Flora Protection Act.

c) To comply with the objectives and requirements of the Habitats Directive, specifically Article 6(3) and where necessary 6(4), Birds, Water Framework, and all other relevant EU Directives and all relevant transposing national legislation.

d) Ensure that proposals for developments located within identified or potential flood risk areas, or which may exacerbate the risk of flooding elsewhere, are assessed in accordance with the provisions of the Flood Risk Management Guidelines (DoEHLG/OPW 2009) and Circular PL2/2014 (or any updated/superseding document), the relevant policies, objectives and guidelines within this plan and shall also take account of the National CFRAM Programme Flood Hazard Mapping and Flood Risk Management Plans when they become available.”

9.1 Project Mitigation: Consenting Process

As set out in Section 8.2 above, the consenting process for the progression of measures, actions or projects supported, guided or recommended within the Plan involving physical works, will require the applicable environmental assessments. Also, the consenting authorities may set out specific environmental conditions as part of the project approval.

9.2 Project Mitigation: Pre-Construction / Detailed Design

For the detailed design of projects that may arise as a result of the Plan, where options are available, the design should use a hierarchy to mitigation measures along the following principles:

- Avoidance: avoid creating the potential impact where feasible.
- Mitigation: minimise the potential impact through mitigating measures

- Enhancement: Enhance the environment to better than pre-project conditions, where reasonably possible

The progression of any projects that may arise as a result of the Plan, through the detailed design phase can entail a series of surveys to inform the design, where the scale of surveys would be proportionate to the complexity and potential impacts of the project. These can include:

- engineering structure surveys,
- topographical surveys,
- habitat and species surveys¹
- ornithological surveys,
- bat surveys,
- fish surveys,
- water quality surveys,
- archaeological surveys,
- landscape and visual assessments,
- land valuation surveys and
- other surveys as deemed necessary to prepare a project.

Where necessary, Wildlife Derogation Licences will be sought from Department of Arts, Heritage, Regional, Rural and Gaeltacht Affairs.

The scope of any necessary EIS will contain a WFD assessment, which will include a hydro-morphological assessment, to more clearly consider and support the Water Framework Directive (WFD) objectives. This WFD assessment will inform the project level AA regarding likely significant effects and adverse impacts on the site integrity of Natura 2000 sites in respect of their conservation objectives and if necessary, appropriate mitigation measures will be implemented at project level to ensure adverse effects will not occur.

9.2.1 Project Mitigation: Construction Stage

For large and complex projects and sites, where environmental management may entail multiple aspects, a project specific Construction Environmental Management Plan (CEMP) may be developed. This will form a framework for all environmental management processes, mitigation measures and monitoring and will include other environmental requirements such as invasive species management measures, if applicable.²

¹ In the context of ecological mitigation, the habitat and species surveys are conducted as required to assess the various aspects for the project, such as ecological surveys for:

- protected or notable habitats and species, including Annex 1 habitats, Annex II and Annex IV species,
- species protected under the Wildlife Acts,
- species protected under the Flora Protection Order,
- the resting and breeding places of relevant species and,
- invasive species, both plant and animal.

² There are a range standard type mitigation measures consisting of good construction practices and good planning of works, that are used within construction projects such as for example: Refuelling of plant and vehicles away from watercourses, Installation of wheel-wash and plant washing facilities, working in-channel or on specific works only within environmental windows e.g. in-stream works in Salmonid channels from May to September.

A designated environmental officer and project ecologist will be appointed, as appropriate for the project.

Biosecurity measures may be required and should be considered depending on the location and conditions on-site.

9.2.2 Project Monitoring

The Plan, with its associated SEA and plan-level AA, sets out a series of monitoring requirements, in connection with the SEA objectives and the predicted effects of the Plan. For measures involving physical works, the project-level EIA and AA, where conducted, will set out the specific monitoring required for each measure.

9.3 Summary and Conclusion

A number of aims, objectives, strategies and policies have been developed as a result of the Plan that are designed to promote the sustainable development and growth of County Mayo. This NIR has determined the potential impacts that may occur through the implementation of the Plan and has assessed each one of these potential impacts in as much detail as is possible with the level of detail that is available at this high level. General mitigation measures have been provided in this NIR, however, where actions may occur in the implementation of these measures, project level assessment of works will be required.

Mitigation measures that are provided in the NIR will be implemented throughout the entire extent of the execution of projects that may stem from the Plan. This includes mitigation measures to be implemented at the Consenting Process Stage, Pre-Construction, Construction and Monitoring for any projects that may arise as a result of the Plan.

Providing the recommended mitigation measures are implemented, it can be concluded that this Plan will not adversely impact on the Natura 2000 sites, either alone or in-combination with other plans, projects or policies.

10 Influence of NIR on the Plan

This NIR informs both the Draft Mayo County Development Plan and the Draft SEA Statement and Monitoring Programme. The Plan template has changed in response to direct observations made in the NIR and may change further following Public Consultation of the NIR.

The Plan may be revised and the outcome of the revision of the Plan, depending on the result of statutory consultation and the findings of this NIR. These revisions may include clarifications and/or refinement of processes or any measures to be implemented.

The approval / adoption of the Plan has not and does not confer approval or permission for the installation or construction of any physical works, without AA Screening and Appropriate Assessment as necessary. Implementation of the Plan to the level as described at this stage, will require the inclusion of mitigation measures that have been detailed in this NIR. Following the approval of the Plan, the next stage is to progress the proposed Plan by undertaking more detailed assessment and design at project level for any structural or physical works, before submitting the proposal for Public Exhibition or planning permission as required depending on the nature of the works. For measures involving physical works, the project-level EIA and AA, where conducted, will set out the specific monitoring and mitigation measures required for each measure. The need for on-site specific mitigation measures at a project level will be a requirement of the planning consent for physical project works. This NIR supports the recommendations of the Plan in this regard.

Appendices

A Impact Assessment

A.1 Chapter 2 – Core Strategy

Core Strategy			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective				
Strategic County Development Plan Objectives	SO 1 Population growth	To plan for a population increase in County Mayo during the plan period of 15,200 persons	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Strategic County Development Plan Objectives	SO 2 Quality of Life for Citizens and Communities	To promote the well-being and quality of life of all citizens and communities in Mayo and to enhance the attractiveness of the county as a place in which to live, work, visit, invest in and enjoy.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Strategic County Development Plan Objectives	SO 3 Employment and Investment	To support employment, encourage enterprise, maximise investment and create an environment that will establish Mayo as a premier investment location, capitalising on the county's existing and emerging key economic drivers, such as the Coastal Corridor and Marine Environment, the Atlantic Economic Corridor, Ireland	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Core Strategy			Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective	Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		West Airport Knock (IWAK), the IWAK Strategic Development Zone and the Economic Growth Clusters of Ballina/ North Mayo, Castlebar-Westport, Ballinrobe, Ballyhaunis and Claremorris and Ballina-Killala.			
Strategic County Development Plan Objectives	SO 4 Low Carbon and Climate Resilient County	To transition to a low carbon and climate resilient county, by promoting sustainable settlement patterns, the integration of land-use and sustainable modes of transport, encouraging walking, cycling and public transport, increasing reliance on green energy sources, encouraging urban and rural communities to facilitate effective change and by building climate change resilience and climate action into all services and functions of Mayo County Council.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy	
Strategic County Development Plan Objectives	SO 5 Development of Sustainable Communities	To support and develop sustainable communities and a competitive county that enhances the health and wellbeing of our people and places from rural to urban, with access to employment opportunities, accessibility to high quality housing and physical, social and community infrastructure, including quality transport infrastructure and new digital technologies, where people of	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy	

Core Strategy				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective		Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red		Mitigation
		all ages can enjoy a good quality of life and a sense of pride in their place.		
Strategic County Development Plan Objectives	SO 6 Inclusive Society	To promote and assist an inclusive county, to ensure that all people of Mayo have access to the range of resources to allow them to fully participate in their local communities, to tackle poverty and social exclusion.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Strategic County Development Plan Objectives	SO 7 Protection of Natural and Built Environment	To create a sustainable Mayo that protects and enhances the natural and built environment, which underpins Mayo's unique identity, environmental character and quality of life and also creates a vibrant county that is an attractive destination for business and enterprise, tourists, residents of the county and creative industries alike.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Strategic County Development Plan Objectives	SO 8 Supporting Rural Areas	To promote the role of Mayo's rural countryside, by developing a sustainable synergy between the rural area and network of settlements, enhancing the rural economy with improved connectivity, broadband, rural economic development opportunities and smarter working opportunities, all within the context of the sustainable management of land and resources	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Core Strategy			Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective	Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
Strategic County Development Plan Objectives	SO 9 Appropriate Assessment, Strategic Environmental Assessment and Strategic Flood Risk Assessment	<p>a) To ensure the assessment of all planning applications in the Plan area have regard to the information, data and requirements of the Appropriate Assessment Natura Impact Report, SEA Environmental Report and Strategic Flood Risk Assessment Report contained in Volume 5 of the Mayo CDP 2021-2027.</p> <p>b) To require project planning to be fully informed by ecological and environmental constraints at the earliest stage of project development and any necessary assessment to be undertaken, including assessments of disturbance to species protected under the Wildlife Act and/or the Flora Protection Act.</p> <p>c) To comply with the objectives and requirements of the Habitats Directive, specifically Article 6(3) and where necessary 6(4), Birds, Water Framework, and all other relevant EU Directives and all relevant transposing national legislation. d) Ensure that proposals for developments located within identified or potential flood risk areas, or which may exacerbate the risk of flooding elsewhere, are assessed in accordance with the provisions of the Flood Risk</p>	Policies or proposals which steer change in such as way as to protect international nature conservation sites from adverse effects	N/A	

Core Strategy			Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective	Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		Management Guidelines (DoEHLG/OPW 2009) and Circular PL2/2014 (or any updated/superseding document), the relevant policies, objectives and guidelines within this plan and shall also take account of the National CFRAM Programme Flood Hazard Mapping and Flood Risk Management Plans when they become available.			
Strategic County Development Plan Objectives	SO 10 Implementation of National and Regional Objectives	To contribute and progress, as practicable, towards achievement of the National Strategic Objectives of Project 2040, the Regional Growth Ambitions of the Northern and Western Regional Assembly's RSES, and the Sustainable Development Goals of the 2030 Agenda for Sustainable Development	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy	
Strategic County Development Plan Objectives	SO 11 Urban Renewal and Regeneration	Continue to enhance the towns and villages of County Mayo, through renewal and regeneration, improvements to public realm infrastructure, healthy place-making and by improving the visual amenity, urban design, viability, vibrancy of these areas so that people can live, work and invest in the these areas.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy	
Strategic County Development	SO 12 Integrated Land Use and	Integrate land use planning and sustainable transportation planning, promote the consolidation of	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura	Refer to Chapter 9 of this NIR for mitigation measures for any plan or	

Core Strategy				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective		Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red		Mitigation
Plan Objectives	Transportation Planning	development, encourage sustainable travel patterns by reducing the need to travel particularly by private transport, while prioritising walking, cycling and public transport.	2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	project arising from this objective/ policy
Core Strategy Policies	CSP1	To promote and facilitate the development of sustainable communities in the county, by managing the level of growth in each settlement to ensure future growth is in accordance with the Core Strategy and County Settlement Hierarchy, in order to deliver sustainable and vibrant rural and urban communities.	This objective could lead to increased development of sustainable communities. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Core Strategy Policies	CSP2	To support the implementation of the Core Strategy for Mayo in a manner that is consistent with policies at a national and regional level, in particular population targets and distribution.	This objective could lead to population increase. This may put pressures on the natural environment within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Core Strategy Policies	CSP3	To promote measures to reduce vacancy and the underuse of existing building stock and support initiatives that promote the reuse, refurbishment and retrofitting of existing buildings within all settlements in the Settlement Hierarchy and the rural countryside	This objective could lead to development of derelict buildings. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Core Strategy			Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective	Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
Core Strategy Policies	CSP4	To support the compact growth of towns and villages to ensure that development proceeds sustainably and at an appropriate scale, density and sequence and in line with the Core Strategy Table.e and in line with the Core Strategy Table.	This objective could lead to population increase and development. This may put pressures on the natural environment within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy	
Core Strategy Objectives	CSO1	To secure the implementation of the population and housing growth household allocation set out in the Core Strategy and Settlement Strategy, in so far as practicable, by facilitating rural housing, while allowing for the accommodation of further residential growth in our designated settlements, subject to the availability of infrastructure and services.	This objective could lead to population increase. This may put pressures on the natural environment within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy	
Core Strategy Objectives	CSO2	To ensure that sufficient zoned lands are available to satisfy the housing requirements of the designated settlements in the county over the lifetime of the Plan.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy	
Core Strategy Objectives	CSO3	To adopt Local Area Plans for Ballina, Castlebar and Westport that align with the NPF, RSES and this Core Strategy. During the transition period between	This objective could indirectly lead to increased development through objectives of the Local Area Plans. Any development within the county could potentially have direct and/or indirect impacts on	Refer to Chapter 9 of this NIR for mitigation measures for any plan or	

Core Strategy				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective		Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red		Mitigation
		adoption of this County Development Plan and the adoption of the Local Area Plans for Ballina, Castlebar and Westport, the objectives (including zoning objectives), policies and standards in this County Development Plan shall apply to these towns.	the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	project arising from this objective/ policy
Core Strategy Objectives	CSO4	To move towards more compact towns by promoting the development of infill and brownfield/consolidation/regeneration sites and the redevelopment of under-utilised land within and close to the existing built up footprint of existing settlements in preference to edge of centre locations.	This objective could lead to increased development on brownfield sites. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Core Strategy Objectives	CSO5	To deliver at least 30% of all new homes in urban areas within the existing built up footprint of settlements	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Core Strategy Objectives	CSO6	To deliver at least 20% of all new homes in rural towns and villages within the existing built up footprint of settlements	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Core Strategy				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective		Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red		Mitigation
			as changes in water quality and quantity, and air quality.	
Core Strategy Objectives	CSO7	To implement all land use planning policies and objectives in a manner which takes account of and is consistent with the Core Strategy, in order to accelerate a transition to a greener, low carbon and climate resilient county with a focus on reduced travel demand through the promotion of sustainable settlement patterns	Policies or proposals which steer change in such as way as to protect international nature conservation sites from adverse effects	N/A
Core Strategy Objectives	CSO8	Monitor development for compliance with the objectives of the Core Strategy and adjust, where necessary, the approach taken to the consideration of development proposals, in order to ensure effective alignment with National and Regional policy and objectives	Policies or proposals which steer change in such as way as to protect international nature conservation sites from adverse effects	N/A
Core Strategy Objectives	CSO9	Monitor and maintain a record of residential development permitted in settlements designated under the Settlement Hierarchy in accordance with any Departmental guidance that may arise.	General statement of policy/general aspiration which will not lead to development	N/A
SETTLEMENT STRATEGY POLICIES	SSP 1	Support the appropriate growth of the Rural Countryside by offering a sustainable choice for people to live in	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of	Refer to Chapter 9 of this NIR for mitigation measures for any plan or

Core Strategy				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective		Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red		Mitigation
		order to maintain vibrant Rural Communities.	habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	project arising from this objective/ policy
SETTLEMENT STRATEGY POLICIES	SSP 2	Support the continued growth and sustainable development of Ballina, Castlebar and Westport, as designated Strategic Growth Towns in the Settlement Strategy, capitalising on Ballina’s designation as a Key Town in the context of the Sligo Regional Growth Centre and Castlebar/Westport as a linked growth driver in the region.	This objective could lead to increased development in Ballina, Castlebar and Westport. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
SETTLEMENT STRATEGY POLICIES	SSP 3	Promote consolidation in Self-Sustaining Growth Towns coupled with targeted investment where required, to improve local employment, services and sustainable transport options and to become more self-sustaining settlements.	This objective could lead to population increase. This may put pressures on the natural environment within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
SETTLEMENT STRATEGY POLICIES	SSP 4	Promote commensurate population and employment growth in the designated Self-Sustaining Towns, providing for natural increases and to become more economically self-sustaining, in line with the quality and capacity of public transport, services and infrastructure available.	This objective could lead to increased development of infrastructure . Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Core Strategy				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
SETTLEMENT STRATEGY POLICIES	SSP 5	To promote and encourage the sustainable, balanced development of the Rural Settlements and Rural Villages in an incremental manner, with the emphasis on small scale development over a medium to long-term period, in keeping with the character of the settlement.	This objective could lead to increased development in rural areas. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
SETTLEMENT STRATEGY POLICIES	SSP 6	To support the creation of healthy and sustainable communities that encourages and facilitates walking and cycling and general physical activity through the implementation of best practices in urban design that promotes permeability and interconnecting spaces.	This objective could lead to increased development of infrastructure . Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
SETTLEMENT STRATEGY POLICIES	SSP 7	To promote the integration of land use and transportation policies and to prioritise provision for cycling and walking travel modes and the strengthening of public transport.	This objective could lead to increased development of infrastructure . Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
SETTLEMENT STRATEGY POLICIES	SSP 8	To develop the county in accordance with the Settlement Hierarchy and to require future residential development to locate at and be of a scale appropriate to the settlement tiers and levels identified in the Core Strategy	General statement of policy/general aspiration which will not lead to development	N/A

Core Strategy				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective		Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red		Mitigation
		Table, subject to the availability of infrastructure and services.		
SETTLEMENT STRATEGY OBJECTIVES	SSO 1	To facilitate the appropriate growth of the Rural Countryside by offering a sustainable choice for people to live in order to maintain vibrant Rural Communities.		
SETTLEMENT STRATEGY OBJECTIVES	SSO 2	To facilitate the development of Ballina, Castlebar and Westport to underpin their role as designated Strategic Growth Towns in the Settlement Hierarchy and to ensure that the growth of these towns takes place in an orderly and sustainable fashion that will not detract from the vitality and viability of their town centres.	This objective could lead to increased development in Ballina, Castlebar and Westport. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
SETTLEMENT STRATEGY OBJECTIVES	SSO 3	To require sustainable, compact, sequential growth and urban regeneration in Ballina, Castlebar and Westport by consolidating the built-up footprints of these towns through a focus on regeneration and development of town centre infill and brownfield sites, and encouraging regeneration of underutilised, vacant and derelict lands for residential development and mixed use to facilitate population growth.	This objective could lead to increased development on brownfield sites in Ballina, Castlebar and Westport. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Core Strategy				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
SETTLEMENT STRATEGY OBJECTIVES	SSO 4	To apply higher densities to the higher order settlements of Ballina, Castlebar and Westport (see DM Standards), to align with their roles within the settlement hierarchy, subject to good design and development management standards being met.	This objective could lead to population increase in Ballina, Castlebar and Westport. This may put pressures on the natural environment within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
SETTLEMENT STRATEGY OBJECTIVES	SSO 5	To apply graded densities in towns and villages having regard to their role in the Settlement Hierarchy and where they are commensurate to the existing built environment.	This objective could lead to population increase in Ballina, Castlebar and Westport. This may put pressures on the natural environment within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
SETTLEMENT STRATEGY OBJECTIVES	SSO 6	To strengthen the core of settlements and encourage the compact growth of settlements by way of the development of infill sites, brownfield lands, under-utilised land / buildings, vacant sites, and derelict sites within the existing built-up footprint of the settlements, and develop outwards from the centre in a sequential manner	This objective could lead to increased development on brownfield sites. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
SETTLEMENT STRATEGY OBJECTIVES	SSO 7	To promote measures to reduce vacancy and the underuse of existing building stock and support initiatives	This objective could lead to development of derelict buildings. Any development within the county could potentially have direct and/or indirect impacts on	Refer to Chapter 9 of this NIR for mitigation measures for any plan or

Core Strategy				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective		Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red		Mitigation
		that promote the reuse, refurbishment and retrofitting of existing buildings and the regeneration of under-utilised lands within all settlements in the Settlement Hierarchy and the rural countryside.	the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	project arising from this objective/ policy
SETTLEMENT STRATEGY OBJECTIVES	SSO 8	To ensure that settlements grow in a manner that is self-sustaining with sufficient social and economic infrastructure, and to a scale which aligns with the Settlement Hierarchy prescribed in the Core Strategy	General statement of policy/general aspiration which will not lead to development	N/A
SETTLEMENT STRATEGY OBJECTIVES	SSO 9	Ensure that all applications for retail developments are appropriate to the role, function and capacity of settlements within the settlement hierarchy and have regard to the Retail Planning Guidelines for Planning Authorities (DoEHLG 2012).	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
SETTLEMENT STRATEGY OBJECTIVES	SSO 10	To support and facilitate the delivery of consolidation and targeted 'catch up' investment in services, infrastructure, amenities and local employment to settlements, where appropriate.	This objective could lead to increased development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
SETTLEMENT STRATEGY OBJECTIVES	SSO 11	To seek to progress the list of Municipal District Projects over the plan period, as set out in Appendix IV.	This objective could lead to increased development of infrastructure. Any development within the county could potentially have direct and/or indirect	Refer to Chapter 9 of this NIR for mitigation measures for any plan or

Core Strategy				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective		Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red		Mitigation
			impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	project arising from this objective/ policy
SETTLEMENT STRATEGY OBJECTIVES	SSO 12	Ensure the promotion of sustainable settlement and transportation strategies in urban and rural areas including the promotion of measures to— (i) reduce energy demand in response to the likelihood of increases in energy and other costs due to long-term decline in non-renewable resources, (ii) reduce anthropogenic greenhouse gas emissions, and (iii) address the necessity of adaptation to climate change; in particular, having regard to location, layout and design of new development	General statement of policy/general aspiration which will not lead to development	N/A

A.2 Chapter 3 – Housing

Strategic Aim: To facilitate the sustainable growth of all rural areas, towns and villages throughout the county by seeking to accommodate, as far as possible, all persons in their choices to live in our rural areas, towns and villages; by supporting and strengthening the rural economy to sustain vibrant rural communities and by promoting consolidation and compact development of all urban and rural settlements in an attractive setting that provides a suitable mix of housing with supporting amenities; and by ensuring coordinated investment in infrastructure that will support economic competitiveness and create a high quality living and working environment.

Housing				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
Housing Strategy Objective	HSO 1	To reserve sufficient lands to facilitate and implement the Housing Strategy and its policies, as informed by the HNDA undertaken as part of this development plan	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Housing Strategy Objective	HSO 2	To secure the implementation of the Part V Housing Strategy, in particular, through the reservation of 10% of all land zoned solely for residential use, or for a mixture of residential or other uses, to be made available for the provision of social and affordable housing referred to in section 94(4)(c) of the Planning & Development Act 2000 (as amended) and shall be provided in accordance with an Agreement referred to in section 96 of the Planning & Development Act 2000 (as amended) and in accordance with Part V Ministerial guidance or any future revised guidance.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Housing				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
Housing Strategy Objective	HSO 3	To increase the stock of social housing within the county in order to meet the social housing needs identified in this Housing Strategy as well as the long-term housing needs of existing households on the local authority housing waiting list.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Housing Strategy Objective	HSO 4	To seek to procure vacant homes where needed and appropriate, to ensure their continued use for residential purposes and develop housing on infill / brownfield sites within town and village centres for social and affordable housing provision.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Housing Strategy Objective	HSO 5	To seek to procure vacant sites or derelict/substandard sites where needed and appropriate, to facilitate the regeneration and repopulation within town and village centres for social and affordable housing provision.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Housing				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
Housing Strategy Objective	HSO 6	Secure the implementation of the Council’s Traveller Accommodation Programme 2019-2024 and to review this programme if required and/or deemed to be necessary, during the plan period.	General statement of policy/general aspiration which will not lead to development	N/A
Housing Strategy Objective	HSO 7	To co-operate with Voluntary Housing Associations and other providers of social housing within County Mayo to secure the delivery of new housing of appropriate design and at appropriate locations over the lifetime of the Plan.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Housing Strategy Objective	HSO 9	To support the ongoing monitoring and review of the HNDA, in accordance with the forthcoming guidance on HNDA methodology to be issued by the Department of Housing, Planning and Local Government.	General statement of policy/general aspiration which will not lead to development	N/A
Housing Strategy Objective	HSO 10	To review, as may be necessary, the County Mayo Housing Strategy should a variation to the Mayo County Council Development Plan 2021-2027 be made or if planning legislation (notably ‘Part V’) is amended during the lifetime of plan	General statement of policy/general aspiration which will not lead to development	N/A
Housing Strategy Policy	HSP1	To promote social integration and the provision of a range of dwelling types in residential developments that would encourage a mix of tenure, particularly in any state funded house building programmes	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Housing				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective		Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red		Mitigation
			as changes in water quality and quantity, and air quality.	
Housing Strategy Policy	HSP2	To promote a mixture of house types, tenures and sizes to reasonably match the requirements of different categories of households and ensure that the special requirements of older persons, persons with disabilities and persons with learning difficulties, are developed in convenient, easily accessible and permeable locations	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Housing Strategy Policy	HSP3	To support the provision of accommodation for older people and for people with disabilities that would allow for independent and semi-independent living, in locations that are proximate to town and village centres and services and amenities such as shops, local healthcare facilities, parks and community centres.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Housing Strategy Policy	HSP4	To promote new social and affordable housing developments designed and constructed on the principles of universal design and life-long adaptability, energy efficient, good design and healthy placemaking	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Housing				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
			as changes in water quality and quantity, and air quality.	
Housing Strategy Policy	HSP 5	To support the work of voluntary and cooperative housing associations in County Mayo	General statement of policy/general aspiration which will not lead to development	N/A
Housing Strategy Policy	HSP 6	To support a balanced supply of private, social and affordable housing in residential development, in order to promote social integration in towns and villages in County Mayo.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Rural Housing Objectives	RHO 1	To facilitate single houses in the open countryside to applicants with a demonstrable economic or social need in <i>Rural Areas under Strong Urban Influence</i> .	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Rural Housing Objectives	RHO 1	An economic need would include applicants who are functionally dependent on the local rural area for employment, where they seek to build their first home i.e. employment is rural based.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Housing				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
			fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	
Rural Housing Objectives	RHO 1	A social need would include applicants who have long standing local intrinsic links to the rural area, where they seek to build their first home i.e. growing up in the area, educated in the area and continue to have strong social links to the rural area.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Rural Housing Objectives	RHO 1	Note: An occupancy clause will be attached to any grant of planning permission.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Rural Housing Objectives	RHO 2	In rural areas not classified as in <i>Rural Areas under Strong Urban Influence</i> , there is a presumption in favour of facilitating the provision of single housing in the countryside, based on siting and design criteria for rural housing in statutory guidelines and plans, except in the case of single houses seeking to locate along	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Housing				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective		Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red		Mitigation
		Mayo's <i>Scenic Routes/Scenic Routes with Scenic Views or Coastal Areas/Lakeshores</i> (See RHO 3).	key indicators of conservation value, such as changes in water quality and quantity, and air quality.	
Rural Housing Objectives	RHO 3	To consider facilitating single houses along <i>Scenic Routes/Scenic Routes with Scenic Views or Coastal Areas/Lakeshores</i> to applicants who can demonstrate a social need to locate in the area concerned, whilst ensuring that it:	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Rural Housing Objectives	RHO 3	· Does not impinge in any significant way on the character, integrity and distinctiveness of the area;	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Rural Housing Objectives	RHO 3	· Meets high standards in siting and design;	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Housing				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective		Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red		Mitigation
			as changes in water quality and quantity, and air quality.	
Rural Housing Objectives	RHO 3	· Satisfies all other criteria with regard to, inter alia, servicing, public safety, and environmental considerations;	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Rural Housing Objectives	RHO 3	· Demonstrates enhancement to local landscape character and ecological connectivity	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Rural Housing Objectives	RHO 3	Note: An occupancy clause will be attached to any grant of planning permission.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Housing				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
			as changes in water quality and quantity, and air quality.	
Rural Housing Objectives	RHO 4	<p>Housing applications, within Mayo’s Coastal Areas and Lakeshores and within areas along scenic routes with designated scenic views, will considered where the applicants can demonstrate a long-standing social link to the area concerned, whilst ensuring that it:</p> <ul style="list-style-type: none"> • Does not impinge in any significant way on the character, integrity and distinctiveness of the area • Cannot be considered at an alternative location • Meets high standards in siting and design • Satisfies all other criterial with regard to, inter alia, servicing, public safety, and environmental considerations • Demonstrates enhancement to local landscape character and ecological connectivity <p>Note: An occupancy clause will be attached to any grant of planning permission.</p>	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Rural Housing Objectives	RHO 5	To advise all rural housing applicants to utilise the Design Guidelines for Rural Housing (Mayo County Council) and core principles of same.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Housing				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
Rural Housing Objectives	RHO 6	To review the existing Design Guidelines for Rural Housing during the lifetime of this plan.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Rural Housing Objectives	RHO 7	That there will be a general presumption against allowing ribbon development i.e. greater than 5 houses in a row over 250m of road frontage, in any area outside of the development boundaries of all settlements listed in the Settlement Hierarchy of this plan.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Rural Housing Objectives	RHO 8	Applicants seeking to replace or reuse an existing house or other structure such as a disused barn, church, schoolhouse or other substantial building in any rural area will not be required to demonstrate a housing need and will be assessed under normal planning considerations only.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Housing				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
Rural Housing Objectives	RHO 9	To discourage the demolition and replacement of traditional or vernacular rural houses in order to protect the varied types of housing stock in rural areas of the County and to preserve the rural built heritage. Demolition and replacement will only be considered, on a case by case basis, where it is clearly demonstrated by way of a suitably qualified structural engineer's report that the dwelling/structure is not reasonably capable of being made structurally sound or otherwise improved.	General statement of policy/general aspiration which will not lead to development	N/A
Rural Housing Objectives	RHO 10	To require that any proposal to extend/refurbish an existing rural dwelling house, occupied or otherwise, takes account of the siting and size of the existing building and endeavours to ensure that the design, scale and materials used in the refurbishment and/or extension are in keeping and sympathetic with the existing structure and that mature landscape features are retained and enhanced, as appropriate.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Rural Housing Objectives	RHO 11	Buildings which are ancillary to existing rural dwelling(s), such as self-isolation units/granny flats/independent living unit or remote working office unit will be considered on their individual merits, subject to compliance with the criteria outlined in Section 2.9 of Volume 2 (Development Management Standards) of the Plan.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Housing				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
Rural Housing Policy	RHP 1	To support and promote strong vibrant sustainable rural communities in County Mayo.	General statement of policy/general aspiration which will not lead to development	N/A
Rural Housing Policy	RHP 2	To support a balanced approach to the development of rural areas to retain vibrancy, to accommodate within the rural area people who are functionally or socially part of the rural community, and to direct urban generated housing demand into established rural settlements.	General statement of policy/general aspiration which will not lead to development	N/A
Rural Housing Policy	RHP 3	To endeavour to accommodate the housing needs of the population, as projected in the Core Strategy, while at all times seeking to facilitate, as far as possible, all persons in their choices to live in our rural areas, towns and villages, subject to relevant development management controls and standards and carrying capacity of natural resources	General statement of policy/general aspiration which will not lead to development	N/A
Rural Housing Policy	RHP 4	To ensure that future housing in rural areas have regard to the Sustainable Rural Housing Guidelines for Planning Authorities 2005 (DOEHLG) or any amended or superseding guidelines.	General statement of policy/general aspiration which will not lead to development	N/A
Rural Housing Policy	RHP 5	To ensure that rural housing applications employ site specific design solutions to provide for proposals that integrate into the landscape, reflect and enhance local landscape character and that respect their location in terms of siting, design, materials, finishes and landscaping.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Housing				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
Rural Housing Policy	RHP 6	To encourage the reuse of an existing rural building/structure other than a house for residential development subject to proper planning and sustainable development.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Rural Housing Policy	RHP 7	To consider replacement dwellings or development of other structures to habitable homes in all rural areas, subject to normal planning considerations such as availability of services, adequacy of ground conditions for disposal of effluent from the development, traffic safety, residential amenity, visual amenity etc. Where it is proposed to replace a dwelling, the replacement dwelling may require to be located on the footprint of the existing structure and the scale and character of the existing building may require replication or be of similar scale and design, depending on the location of the development (e.g. sensitive or vulnerable locations such as coastal, the shorelines of large lakes or upland areas).	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Rural Housing Policy	RHP 8	To require that new houses in the rural areas ensure the protection of water quality in the arrangements for on-site waste water disposal, ensure provision of a safe means of access in relation to road and public safety, avoid flood risk and ensure the conservation of sensitive areas such as natural habitats, the environs of protected structures and other aspects of heritage.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Housing				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
			key indicators of conservation value, such as changes in water quality and quantity, and air quality.	
TOWN AND VILLAGE HOUSING POLICIES	TVHP 1	To support the development of quality residential schemes having regard to and being consistent with the standards and principles set out in the Sustainable Residential Development in Urban Areas Guidelines for Planning Authorities (2009) and any relevant specific planning policy requirements (SPPRs) in the ‘Urban Development and Building Heights Guidelines for Planning Authorities’ (2018) and the ‘Sustainable Urban Housing: Design Standards for New Apartments, Guidelines for Planning Authorities’ (2018) (and as updated).	General statement of policy/general aspiration which will not lead to development	N/A
TOWN AND VILLAGE HOUSING OBJECTIVES	TVHP 1	To ensure that a suitable variety and mix of dwelling types and sizes is provided in developments to meet different needs, having regard to demographic and social changes.	General statement of policy/general aspiration which will not lead to development	N/A
TOWN AND VILLAGE HOUSING POLICIES	TVHP 2	To support the creation of attractive residential developments with a range of housing options and appropriate provision of functional public and private open space that is consistent with the standards and principles set out in the Guidelines for Planning Authorities on Sustainable Residential Development in Urban Areas and the associated Urban Design Manual – A Best Practice Guide, DEHLG (2009) and any subsequent guidelines.	General statement of policy/general aspiration which will not lead to development	N/A
TOWN AND VILLAGE HOUSING OBJECTIVES	TVHP 2	To require residential development to demonstrate that a housing density appropriate to its context is achieved, providing for a sustainable pattern of	General statement of policy/general aspiration which will not lead to development	N/A

Housing			
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective		Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		development, whilst ensuring a high-quality living environment.	
TOWN AND VILLAGE HOUSING POLICIES	TVHP 3	To encourage and foster the creation of attractive, mixed use, sustainable communities that include a suitable mix of housing types and tenure with supporting facilities, amenities and services that meet the needs of the community and are in accordance with the principles of universal design, life-long adaptability and energy efficiency, and urban greening in as far as practicable.	N/A
TOWN AND VILLAGE HOUSING OBJECTIVES	TVHP 3	Ensure all apartments are designed having regard to the standards principles and any specific planning policy requirements (SPPRs) outlined in the Design Standards for New Apartments - Guidelines for Planning Authorities (March 2018) (or as updated).	N/A
TOWN AND VILLAGE HOUSING POLICIES	TVHP 4	To encourage the reuse of upper floors above commercial premises for residential accommodation.	N/A
TOWN AND VILLAGE HOUSING OBJECTIVES	TVHP 4	To ensure the provision of childcare facilities as an integral part of proposals for new residential developments, having regard to the DEHLG's Childcare Facilities Guidelines for Planning Authorities, 2001 (as may be updated) in relation to the provision of childcare facilities, where appropriate.	N/A
TOWN AND VILLAGE HOUSING POLICIES	TVHP 5	To promote higher residential densities in appropriate locations and in particular close to town centres and along public transport corridors, in accordance with the Guidelines for Planning Authorities on Sustainable Residential Development in Urban Areas, DEHLG (2009).	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Housing				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
			conservation value, such as changes in water quality and quantity, and air quality.	
TOWN AND VILLAGE HOUSING OBJECTIVES	TVHP 5	To facilitate higher and increased building heights at suitable locations within the higher order settlements in accordance with the Sustainable Residential Development in Urban Areas Guidelines for Planning Authorities (2009); Urban Development and Building Heights Guidelines for Planning Authorities’ (2018) and the ‘Sustainable Urban Housing: Design Standards for New Apartments, Guidelines for Planning Authorities’ (2018) (or as updated), including any relevant specific planning policy requirements (SPPRs), where appropriate.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
TOWN AND VILLAGE HOUSING POLICIES	TVHP 6	To promote innovation in architectural design that delivers buildings of a high quality that positively contribute to the built environment and local streetscape.	General statement of policy/general aspiration which will not lead to development	N/A
TOWN AND VILLAGE HOUSING OBJECTIVES	TVHP 6	To require that applications for residential development take an integrated and balanced approach to movement, place making, and streetscape design in accordance with the requirements of the Design Manual for Urban Roads and Streets, DTTS and DECLG (2013 as amended).	Policy or proposal that could not have any conceivable effect on a site	N/A
TOWN AND VILLAGE HOUSING POLICIES	TVHP 7	To promote the development of vacant residential and regeneration sites in all development centres in the county, as appropriate, in accordance with the requirements of the Urban Regeneration and Housing Act 2015 (as amended).	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Housing				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
			as changes in water quality and quantity, and air quality.	
TOWN AND VILLAGE HOUSING OBJECTIVES	TVHP 7	To ensure the provision of adequate areas of high quality, safe and overlooked open space within residential developments and support the provision of play and recreational areas in all new large residential developments.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
TOWN AND VILLAGE HOUSING POLICIES	TVHP 8	To direct multiple residential developments to those settlements identified in the Core Strategy and to require that the scale of such development is in accordance with growth projected within that specific settlement, except where there is otherwise a demonstrable need.	General statement of policy/general aspiration which will not lead to development	N/A
TOWN AND VILLAGE HOUSING OBJECTIVES	TVHP 8	To require that development proposals for new residential developments in settlements demonstrate a high-quality design process including layout, specification and external finishes.	Policy or proposal that could not have any conceivable effect on a site	N/A
TOWN AND VILLAGE HOUSING OBJECTIVES	TVHP 9	To guide multiple residential development in a sequential manner outward from the core area of settlements, to maximise the utility of existing and future infrastructure provision, to promote sustainability, to make more efficient use of underutilised lands, and to avoid the extension of services and utilities to more remote areas.	Policy on avoiding development in remote areas will reduce the impact on Natura 2000 sites	N/A

Housing				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
TOWN AND VILLAGE HOUSING OBJECTIVES	TVHP 10	To prepare design guidelines for new residential developments in towns and villages during the lifetime of the plan. In the interim, the design guidance and principles set out in the Village Design Statements for Louisburgh, Mulranny and Belcarra and the Sustainable Residential Development in Urban Areas and accompanying Urban Design Manual (DoEHLG) 2009 (or as updated) shall apply, where appropriate.	General statement of policy/general aspiration which will not lead to development	N/A

DRAFT

A.3 Chapter 4 – Economy

Strategic Aim: The aim of this economic chapter is to provide for the future well-being of the residents of the county and the region by facilitating economic development; to promote the growth of employment opportunities in all sectors, in accordance with the principles of sustainable development; to achieve a reduction in the unsustainable levels of commuting from the county; to provide a greater focus on community building and improve the quality of life for all.

Economy				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
Economic Development Policy	EDP1	Support and promote economic opportunities identified in Mayo County Council’s economic strategy – Mayo: <i>Sustaining Jobs, Supporting Growth & Winning Investment</i> or any amended or replacement strategy.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Economic Development Objectives	EDO 1	To facilitate and support the continued growth of the economy in the county in a sustainable manner and in accordance with the National Planning Framework (NPF) and the Regional Spatial and Economic Strategy.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Economic Development Objectives	EDO 2	To support and facilitate the economic development of the county in a manner which is consistent with the economic pillars identified in the Enterprise and Investment Units Economic Strategy.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats,	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Economy				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
			disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	
Economic Development Objectives	EDO 3	Continue to promote the county to attract enterprise and investment into Mayo through the Enterprise & Investment Unit and/or Local Enterprise Office, with a focus on a number of established and emerging sectors including tourism, manufacturing, marine, renewable energy, ICT, food and agri-food.	This objective could indirectly lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Economic Development Objectives	EDO 4	Continue to support new and existing enterprises in the county through the provision of training, mentoring, financial supports, advice and networking events.	General policy statement which will, in itself, not lead to development	N/A
Economic Development Objectives	EDO 5	Encourage enterprise and employment development to locate in brownfield sites or unoccupied buildings in town centres or where appropriate in existing industrial/retail parks or other brownfield industrial sites in preference to undeveloped zoned or unzoned lands.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Economic Development Objectives	EDO 6	Facilitate the economic development of Mayo to create a viable and favourable economic environment for business and enterprise, whilst delivering sustainable	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats,	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Economy				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective		Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red		Mitigation
		jobs, employment opportunities and an enriched standard of living for all.	disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	
Economic Development Objectives	EDO 7	To identify and promote a range of locations within the County for different types of enterprise activity including international business and technology parks, small and medium enterprises (SME) and micro enterprise centre.	This objective could indirectly lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Economic Development Objectives	EDO 8	To co-operate with local and national development agencies and engage with existing and future employers in order to maximise job opportunities in the County.	General policy statement which will, in itself, not lead to development	N/A
Economic Development Objectives	EDO 9	Facilitate agri-industry and other rural enterprise activities that are dependent on their locality in rural locations, where it can be demonstrated that the development will not have significant adverse effects on the environment, including the integrity of the Natura 2000 network, residential amenity or visual amenity. Preference will be given to occupying vacant structures which the planning authority consider appropriate for the use proposed.	Policies or proposals which steer change in such as way as to protect international nature conservation sites from adverse effects	N/A

Economy				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
Economic Development Objectives	EDO 10	Encourage and facilitate home-based employment of appropriate type, size and scale, where it can be demonstrated that the development will not have significant adverse effects on the environment, including the integrity of the Natura 2000 network, residential amenity or visual amenity.	Policies or proposals which steer change in such as way as to protect international nature conservation sites from adverse effects	N/A
Economic Development Objectives	EDO 11	To work in partnership with relevant stakeholders, to ensure that a sustainable approach is taken to enterprise development and employment creation across all sectors of the Mayo economy, in accordance with the Green Economy national frameworks relevant to each sector.	General policy statement which will, in itself, not lead to development	N/A
Economic Development Objectives	EDO 12	Engage with all relevant government stakeholders, enterprise agencies and sectoral representatives in pursuing 'green' approaches to economic development and actively collaborate with key industry and educational bodies to promote Mayo based initiatives across the economic sectors.	General policy statement which will, in itself, not lead to development	N/A
Economic Development Objectives	EDO 13	Promote quality employment and residential developments in proximity to each other in order to reduce the need to travel.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value,	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Economy				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
			such as changes in water quality and quantity, and air quality.	
Economic Development Objectives	EDO 14	To encourage the provision of 'live work' communities, in which employment, residency and sustainable transport facilities are located in close proximity to each other, to reduce long distance commuter trends and congestion, as well as reducing outward migration from the county.	General policy statement which will, in itself, not lead to development	N/A
Economic Development Objectives	EDO 15	To ensure that people intensive developments are located close to the strategic public transport network.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Economic Development Objectives	EDO 16	To address the rate of out bound commuting, with the provision of 'live work' communities in strategic settlements served by sustainable transport, thereby improving quality of life, encouraging volunteerism and community engagement.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Economic Development Objectives	EDO 17	Encourage mixed use settlement forms and sustainable centres, in which employment, residency, education and	This objective could lead to increased development. Any development within the county could potentially have direct and/or	Refer to Chapter 9 of this NIR for mitigation measures for any plan

Economy				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		local services / amenities are located in close proximity to each other.	indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	or project arising from this objective/ policy
Economic Development Objectives	EDO 18	To work with Irish Water and other infrastructure providers, to support the provision of services and facilities to accommodate the future economic growth of the County and to seek to reserve infrastructure capacity for employment generating uses.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Economic Development Objectives	EDO 19	Ensure that there is sufficient quantum of zoned lands to facilitate a range of enterprise across the county in line with the settlement hierarchy.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Economic Development Objectives	EDO 20	To support start-up businesses and small-scale industrial enterprise at appropriate locations throughout the County, subject to the principles of proper planning and sustainable development.	This objective could indirectly lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value,	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Economy				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
			such as changes in water quality and quantity, and air quality.	
Economic Development Objectives	EDO 21	To further linkages and partnerships with GMIT including the branding of the area as a centre of excellence in the knowledge-based economy.	General policy statement which will, in itself, not lead to development	N/A
Economic Development Objectives	EDO 22	To encourage and facilitate small indigenous industries, at appropriate locations with good communication infrastructure, in recognition of their increasing importance in providing local employment and helping to stimulate economic activity within small communities.	General policy statement which will, in itself, not lead to development	N/A
Economic Development Objectives	EDO 23	To support the use of town centre core locations for new service focused enterprises.	This objective could indirectly lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Economic Development Objectives	EDO 24	Support the continued development of Galway Mayo Institute of Technology (GMIT), Castlebar Campus including the development of a shared approach to enterprise development with the Local Enterprise Office.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value,	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Economy				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
			such as changes in water quality and quantity, and air quality.	
Economic Development Objectives	EDO 25	To acquire suitable land (subject to the availability of funding), including where appropriate, disused sites in State ownership, for creative and innovative entrepreneurial initiatives and the provision of clustered incubator units.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Economic Development Objectives	EDO 26	To promote the development of the key strategic employment sites identified in the Economic Development Strategy for County Mayo.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Economic Development Objectives	EDO 27	To support and promote the equine industry in the County as an economic and employment provider.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Economy				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
Economic Development Objectives	EDO 28	To work with Eirgrid, as far as practicable, to ensure power infrastructure is available for the development of zoned employment lands within the N5 corridor.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Economic Development Objectives	EDO 29	Review the Development Contribution Scheme.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Ireland West Airport Knock Policies	EDP 2	Promote and support the strategic role of IWA Knock SDZ as a significant regional economic driver and promote the SDZ location as a regional economic business and enterprise hub.	Increased air traffic within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Ireland West Airport Knock Policies	EDP 3	Support the development of the IWA Knock SDZ in tandem with the required infrastructure for both the expansion of the airport and in its delivery as a regional business and enterprise hub.	This objective could lead to increased development of the airport. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Economy				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
			conservation value, such as changes in water quality and quantity, and air quality.	
Ireland West Airport Knock Objectives	EDO 30	Promote and facilitate the growth of IWAK as a creative and innovative area that is globally competitive, internationally linked, attractive and open.	General policy statement which will, in itself, not lead to development	N/A
Ireland West Airport Knock Objectives	EDO 31	Support, promote and facilitate the implementation of the approved Planning Scheme for the designated SDZ at IWA Knock (including the mitigation measures of the statutory environmental assessment carried out for the planning scheme), which provides the framework for the expansion of the Airport in terms of its transport and business operations and as a new business and enterprise destination.	This objective could indirectly lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Ireland West Airport Knock Objectives	EDO 32	Support and recognise the important national and Regional role of IWAK in the economic life of the county and region and to facilitate its activities and development, having regard to the IWAK SDZ.	This objective could indirectly lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Atlantic Economic Corridor Policies	EDP 4	Promote, support and facilitate the Atlantic Economic Corridor initiative in a sustainable manner to secure the long-	This objective could indirectly lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network	Refer to Chapter 9 of this NIR for mitigation measures for any plan

Economy				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective		Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red		Mitigation
		term economic and employment growth for Mayo.	through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	or project arising from this objective/ policy
Atlantic Economic Corridor Policies	EDP 5	Promote, support and facilitate the appropriate development of Mayo's towns, villages and rural areas that function as part of the AEC.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Atlantic Economic Corridor Objectives	EDO 33	To advance and promote the AEC in order to maximise the advantages of Western Region links and the development of an All-Ireland economy.	This objective could indirectly lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Atlantic Economic Corridor Objectives	EDO 34	To support the sustainable development of a linked Castlebar-Westport Hub as a driver of economic development for the county and region, building on the dual capacity, complementarity and combined strengths of both towns in terms of population, established commercial, social and cultural links and proximity,	This objective could indirectly lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value,	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Economy				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		thereby strengthening the urban structure of the region and enabling the realisation of the Atlantic Economic Corridor.	such as changes in water quality and quantity, and air quality.	
Atlantic Economic Corridor Objectives	EDO 35	Facilitate the development of the Western Rail Corridor, in order to support and underpin the development of the AEC, to encourage the economic growth of Mayo and the wider region and to support vibrant local communities.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Atlantic Economic Corridor Objectives	EDO 36	Promote and facilitate in cooperation with relevant stakeholders, the provision of AEC enterprise hubs within the towns and villages in Mayo.	This objective could indirectly lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Atlantic Economic Corridor Objectives	EDO 37	Work with adjoining counties to secure the AEC, as a model of balanced regional development.	General policy statement which will, in itself, not lead to development	N/A
Growth Clusters Policies	EDP 6	To support and promote the sustainable development of the Castlebar-Westport Economic Growth Cluster as a driver of economic development for the County and Region, building on the dual	This objective could indirectly lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats,	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Economy				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		capacity, complementarity and combined strengths of both towns in terms of population, established commercial, social and cultural links and proximity, thereby strengthening the urban structure of the region and enable the realisation the Atlantic Economic Corridor.	disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	
Growth Clusters Policies	EDP 7	To support and grow the role of Castlebar as the main administrative, healthcare and education centre in the county, the main centre for commerce and enterprise, capitalising on its designation as a Key Town within the context of the Galway Metropolitan City and Sligo Regional Growth Centre and its location in relation to the Atlantic Economic Corridor, in order to facilitate long-term economic growth within the context of a high quality environment supporting a wide range of services and amenities.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Growth Clusters Policies	EDP 8	To support and grow the role of Ballina as the key economic driver in the north-west of the county, capitalising on its designation as a Key Town within the context of the Sligo Regional Growth Centre and its location in relation to the Atlantic Economic Corridor, in order to facilitate long-term economic growth within the context of a high quality	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Economy				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		environment, supporting a wide range of services and amenities.		
Growth Clusters Policies	EDP 9	To support and grow the role of Westport as a key economic driver in the county and as a national tourism hub, capitalising on its significant tourism related assets, including its designation as a Heritage Town and its status as a Key Town within the context of Galway Metropolitan City, Sligo Regional Growth Centre and its location in relation the Atlantic Economic Corridor, in order to facilitate long-term economic growth within the context of a high quality environment, supporting a wide range of services and amenities.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Growth Cluster Objectives	EDO 38	Seek to support the implementation of the recommendations of the Castlebar-Westport Economic Growth (CWEG) Cluster 2040.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Growth Cluster Objectives	EDO 39	Prepare an economic growth strategy for Ballina and its supporting catchment to advance the economic development of North Mayo.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Economy				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
			changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	
Growth Cluster Objectives	EDO 40	Investigate the potential for the Tier 2 urban settlements of Ballinrobe, Ballyhaunis and Claremorris to function as an economic growth cluster in South Mayo, capitalising on their proximity to the IWAK and AEC	General policy statement which will, in itself, not lead to development	N/A
Retail Policies	EDP 10	Support the vitality and viability of existing town and village centres and facilitate a competitive and healthy environment for the retailing sector into the future, by ensuring that future growth in retail floorspace responds to the identified settlement hierarchy, the sequential approach, the appropriate protection of the built environment and the needs of the projected population of the settlement area.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Retail Policies	EDP 11	Encourage local partnership, town teams or community organisations to develop and implement local economic initiatives which will enhance town and village centres.	General policy statement which will, in itself, not lead to development	N/A
Retail Policies	EDP 12	Promote the reuse of vacant floorspace. Alternative uses shall be assessed on their own merits against the requirements of the proper planning and sustainable development of the area.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Economy				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
			changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	
Retail Policies	EDP 13	Support and promote the retail sector in the county and ensure compliance with the Retail Planning Guidelines for Planning Authorities DoECLG (April 2012) and Retail Design Manual DoECLG (April 2012) or any amended or superseding version of the guidelines and design manual.	General policy statement which will, in itself, not lead to development	N/A
Retail Policies	EDP 14	Promote the reuse or reactivation of vacant underutilised properties/shop units in order to assist with the regeneration of streets and settlements in the county.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Retail Objectives	EDO 41	To implement/review the Mayo County Retail Strategy.	General policy statement which will, in itself, not lead to development	N/A
Retail Objectives	EDO 42	Promote and reinforce all town centres in the county as primary shopping areas.	General policy statement which will, in itself, not lead to development	N/A
Retail Objectives	EDO 43	To adhere to the principle of 'sequential approach' in the consideration of retail applications located outside of core retail areas.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value,	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Economy				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
			such as changes in water quality and quantity, and air quality.	
Retail Objectives	EDO 44	Ensure proposals for retail development in towns and villages make a positive contribution to the general townscape through the promotion of excellence in urban design, signage, consideration of the built heritage and designed to a scale appropriate to the settlement.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Retail Objectives	EDO 45	To continue to implement and facilitate environmental, amenity and recreational improvements to the public realm, in existing town and village core retail areas.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Retail Objectives	EDO 46	Strictly control the location and range of goods sold from retail warehouse units in order to ensure that the retail primacy of the town centre is not unduly prejudiced.	General policy statement which will, in itself, not lead to development	N/A
Retail Objectives	EDO 47	Promote and facilitate on-street activities including street markets and farmers / country markets in all existing retail centres or any event that adds to the vitality and viability of existing town and village centres, whilst not unduly impacting on the existing retail function.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value,	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Economy				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
			such as changes in water quality and quantity, and air quality.	
Smart Economy Policies	EDP 15	Support and facilitate the delivery of the National Broadband Plan and Mayo County Councils Digital Strategy as a means of developing further opportunities for enterprise, employment, education, innovation and skills development, for those who live and work in rural areas throughout Mayo, where appropriate	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Smart Economy Policies	EDP 17	Develop the ICT sector in Mayo, along with key stakeholders and relevant agencies to ensure that the economic, potential of the sector is secured for the benefit of the local, regional and national economy.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Smart Economy Policies	EDP 16	Support and facilitate the development of a Smart County to benefit the economic growth, quality of life and wellbeing of the people of Mayo and advance the NWRA 'Smart Region' Growth Ambition in the RSES.	General policy statement which will, in itself, not lead to development	N/A
Smart Economy Policies	EDP 18	Support the development of a Smart County to develop and diversify the rural economy, to build on local enterprise and infrastructure assets, to drive innovations	General policy statement which will, in itself, not lead to development	N/A

Economy				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		around energy, transport, agri-food, tourism, e-services and remote working.		
Smart Economy Objectives	EDO 48	Support the role out of actions listed in the Mayo County Council Digital Strategy.	General policy statement which will, in itself, not lead to development	N/A
Smart Economy Objectives	EDO 49	Support the development of sites where data centres, ICT related development and high potential start-ups can thrive.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Smart Economy Objectives	EDO 50	To identify suitable locations and support the provision of co-working facilities, digital hubs/eHubs and eWorking centres throughout the county that function as outreach hubs for out of county employers and that promote flexible working arrangements and strengthen the AEC hub network.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Rural Economy Policies	EDP 19	Support, in conjunction with Mayo LEO and other agencies, the development of indigenous industry and business start-ups in rural employment centres (villages and settlements) in the county, subject to compliance with siting, design and environmental considerations.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Economy				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
Rural Economy Policies	EDP 20	To support the implementation of the new LEADER Rural Development Programme 2014-2020 and any subsequent amended/updated programme for the county..	This objective could lead to increased development of the airport. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Rural Economy Objectives	EDO 51	Facilitate agri-industry and other rural enterprise activities that are not suitable in a settlement and are dependent on their locality in rural locations, where it can be demonstrated that the development will not have significant adverse effects on the environment, including the integrity of the Natura 2000 network, residential amenity or visual amenity. Preference will be given to occupying vacant structures which the planning authority consider appropriate for the use proposed. Where proposals demonstrate measures to promote environmental enhancement through improved ecological connectivity such as measures in the Pollinator Plan, additional native species planting or blue and green infrastructure measures, these will be favourably considered.	Policies or proposals which steer change in such as way as to protect international nature conservation sites from adverse effects	N/A
Rural Economy Objectives	EDO 52	To support rural entrepreneurship and the development of micro businesses	This objective could lead to increased development. Any development within the	Refer to Chapter 9 of this NIR for mitigation

Economy				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		(generally less than 10 no. employees) in rural areas where environmental and landscape impact is minimal and such developments do not generate significant or undue traffic. This objective shall not apply to sites accessed from the National Road Network	county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	measures for any plan or project arising from this objective/ policy
Rural Economy Objectives	EDO 54	Facilitate remote working in the rural area, at an appropriate scale, for enterprise/businesses that do not require visiting members of the public, subject to normal planning considerations.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Agriculture Policies	EDP 21	Support the implementation of the Mayo County Council Agricultural Strategy to promote the continued development and expansion of the Agri-Food Sector subject to the measures and environmental objectives of the forthcoming Common Agricultural Policy Strategy for Ireland.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Agriculture Policies	EDP 22	Support the implementation, at county level, of the provisions set out in Food Harvest 2025, subject to environmental carrying capacity constraints.	This objective could lead to an increase in agricultural pressure. Increase in agricultural pressure could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Economy				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
			changes in key indicators of conservation value, such as changes in water quality and quantity and air quality.	
Agriculture Policies	EDP 23	Maintain a vibrant and healthy agricultural sector based on the principles of sustainable development, whilst at the same time supporting alternative employment in or close to rural areas to sustain rural communities.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Agriculture Policies	EDP 24	Work with the Department of Agriculture, Food and the Marine, Teagasc and all other stakeholders to support the agricultural and agri-business sector, as appropriate and in line with environmental commitments under Climate Action, Biodiversity and Water Quality	This objective could lead to an increase in agricultural pressure. Increase in agricultural pressure could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Agriculture Policies	EDP 25	Work with the Northern and Western Regional Assembly and other relevant stakeholders in identifying areas of high value agricultural land in the county to address the need for sustainable food supplies. The consideration of future climate scenarios, and water availability, climate change adaption and environmental enhancement for	This objective could lead to an increase in agricultural pressure. Increase in agricultural pressure could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Economy				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		agricultural purposes shall form part of this assessment.		
Agriculture Policies	EDP 26	Support and promote the equine industry in the county as an economic and employment provider.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Agriculture Policies	EDP 27	Support the development of appropriately located allotments, in areas which have good access to and are proximate to built-up and residential areas.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Agriculture Objectives	EDO 55	To support, promote and consider, on their individual merits, the reuse of redundant agricultural buildings and the development of new buildings to accommodate farm diversification / enterprise within an overall farmyard complex.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Agriculture Objectives	EDO 56	Support, promote and facilitate the implementation of the actions of Mayo	This objective could lead to increased development. Any development within the	Refer to Chapter 9 of this NIR for mitigation

Economy				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		County Council's agricultural strategy - <i>A Sustainable Agricultural Strategy for Mayo</i> or any amended/replacement strategy to secure economic and employment growth in the county in a sustainable manner.	county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	measures for any plan or project arising from this objective/ policy
Agriculture Objectives	EDO 57	Encourage proposals for farm shops where it can be clearly demonstrated that:	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Agriculture Objectives	EDO 57	(a) the products to be sold are primarily produce grown on the farm holding;	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
		(b) the scale and scope of the retailing proposed will not harm the viability or retail facilities in any nearby town or village;	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Economy				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
			changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	
		(c) and the proposed shop is operated by the owner of the farm and is ancillary to the main use of the property for agricultural activities.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Agriculture Objectives	EDO 58	Ensure that equine based developments are located on suitable and viable landholdings and are subject to normal planning, siting and design considerations.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Agriculture Objectives	EDO 59	Facilitate the development of allotments of an appropriate scale, subject to normal planning considerations and which meet the following criteria:	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Economy				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		(i) The lands are situated within or immediately adjacent to the edge of towns/villages or are easily accessible to the residents of a particular town or village; and	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
		(ii) Adequate water supply and adequate parking facilities can be provided.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Extractive Industry Policy	EDP 28	Support and facilitate adequate supplies of aggregate resources to meet the future growth needs of the county and the wider region where there is a proven need for a certain mineral/aggregate and to exercise appropriate control, while addressing key environmental, traffic and social impacts and details of rehabilitation.	Extracting resources within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Extractive Industry Policy	EDP 29	Ensure that the development of aggregate resources (stone and sand/gravel deposits) is carried out in a manner which minimises effects on the	Extracting resources within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or	Refer to Chapter 9 of this NIR for mitigation measures for any plan

Economy				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		environment, including the Natura 2000 network, amenities, infrastructure and the community, and has full regard to the principles of sustainability.	fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	or project arising from this objective/ policy
Extractive Industry Objectives	EDO 60	Ensure that the development of aggregate resources (stone and sand/gravel deposits) is carried out in a manner which minimises effects on the environment, including the Natura 2000 network, amenities, infrastructure and the community, and has full regard to the principles of sustainability.	Policies or proposals which steer change in such as way as to protect international nature conservation sites from adverse effects	N/A
Extractive Industry Objectives	EDO 61	Have regard to the Quarry and Ancillary Activities Planning Guidelines for Planning Authorities DoEHLG (April 2004) or any new or subsequent quarry guidance.	General policy statement which will, in itself, not lead to development	N/A
Green Economy Policies	EDP 34	Support the implementation of the Green Economy national frameworks through encouraging enterprise development and employment creation across all sectors of the Mayo economy.	General policy statement which will, in itself, not lead to development	N/A
Green Economy Policies	EDP 35	Support rural diversification through sustainable rural development practices, investment in rural towns and villages, providing for access to technology and skills- development networks.	General policy statement which will, in itself, not lead to development	N/A
Green Economy Objectives	EDO 64	Support and facilitate the Green Economy in County Mayo, as appropriate.	General policy statement which will, in itself, not lead to development	N/A

Economy				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
Green Economy Objectives	EDO 65	Facilitate the development of industries that create and employ green technologies and take measures to accelerate the transition towards a low carbon economy and circular economy.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Green Economy Objectives	EDO 66	Facilitate homebased economic activity that allows employers, enterprises and entrepreneurs the option of working from home or local hubs to reduce commuting and congestion, subject to normal planning considerations.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Green Economy Objectives	EDO 67	Support Renewable energy initiatives that facilitate a low carbon transition.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Major Accidents & Seveso Objectives	EDO 68	To ensure that appropriate distances are maintained between any proposed development and any existing Seveso II establishment, in the interest of the	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network	Refer to Chapter 9 of this NIR for mitigation measures for any plan

Economy				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		health and safety of the occupiers of the proposed development.	through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	or project arising from this objective/ policy
Major Accidents & Seveso Objectives	EDO 69	To require that, the siting of new establishments, or modification of existing establishments classified under the Seveso II Directive as listed in Appendix and new development in the vicinity of existing establishments shall take into account the need to prevent major accidents involving hazardous substances and safeguard both the public and the environment.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

A.4 Chapter 5 – Tourism

The strategic aim of this chapter is to promote and facilitate a sustainable and well-managed year-round, high-quality tourism industry that generates economic benefits to all areas of the county, thereby contributing to the wider tourism industry of the region.

Tourism				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
General Tourism Policies	TRP 1	To work in partnership with local, national and international agencies/bodies to promote County Mayo as a tourist destination and to support and encourage cohesion and linkages between the relevant agencies/ bodies to implement the key tourism objectives of this Plan	General statement of policy/general aspiration which will not lead to development	N/A
General Tourism Policies	TRP 2	Support and promote sustainable tourism development, accessible to all throughout the county and work in partnership with tourism organisations and adjoining Local Authorities where necessary, in securing the development of tourism enterprises and infrastructure. Subject to suitable locations where it can be demonstrated that the development will not have significant adverse effects on the environment, including the integrity of the Natura 2000 network, residential amenity or visual amenity.	This Policy could promote development. Although sustainable and in consideration of the Natura 2000 network, impacts may still arise as a result of this Policy. Any developments can potentially have direct and/or indirect impact on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality. Therefore, mitigation measures will be required.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
General Tourism Policies	TRP 3	Support the work of Fáilte Ireland and the strategic growth of tourism development in the county through:	This Policy could promote development for tourism, particularly in the Clew Bay and North Mayo/Erris areas. Any developments can potentially have direct and/or indirect	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

		<p>(a) The implementation of Fáilte Irelands designations - ‘Always On’ Hubs, ‘Seasonal’ Service Centres and ‘Attractions’ within the county.</p> <p>(b) The implementation of Fáilte Irelands Visitor Experience Development Plans (VEDP’s) for Clew Bay and North Mayo/Erris.</p>	<p>impact on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality. Therefore, mitigation measures will be required.</p>	
General Tourism Policies	TRP 4	<p>Co-operate with Fáilte Ireland, Tourism Ireland, and any other relevant bodies in the implementation of Destination Mayo 2016-2021 by:</p>	<p>This Policy could promote development for tourism, particularly in the areas around the Great Western Greenway at Castlebar to Westport and to The Wild Atlantic Way. Any developments can potentially have direct and/or indirect impact on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality. Therefore, mitigation measures will be required.</p>	<p>Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy</p>
General Tourism Policies	TRP 4	<p>(a) Encouraging investment in the tourism industry in the county with specific reference to leisure activities (including walking, cycling, equestrian and family focused activities), including connectivity to the Great Western Greenway at Castlebar to Westport and to the Wild Atlantic Way.</p>	<p>This Policy could promote development for tourism, particularly in the areas around the Great Western Greenway at Castlebar to Westport and to The Wild Atlantic Way. Any developments can potentially have direct and/or indirect impact on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality. Therefore, mitigation measures will be required.</p>	<p>Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy</p>
General Tourism Policies	TRP 4	<p>(b) Encouragement and support of the upgrading of public transport facilities in</p>	<p>This Policy could promote development for tourism. Any developments can potentially</p>	<p>Refer to Chapter 9 of this NIR for mitigation measures for any</p>

		Destination Towns, including the provision of Transport Hubs/Links.	have direct and/or indirect impact on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality. Therefore, mitigation measures will be required.	plan or project arising from this objective/ policy
General Tourism Policies	TRP 4	(c) Supporting the development of new and emerging tourism products and facilities or upgrading/extension of existing tourist facilities at tourist sites within the county, within proper planning and sustainable development principles.	This Policy could promote development for tourism. Any developments can potentially have direct and/or indirect impact on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality. Therefore, mitigation measures will be required.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
General Tourism Policies	TRP 5	Support the delivery of high quality 'destination town' experiences for visitors by supporting expansion in accommodation and facilities within Destination Towns and supporting infrastructural investment, including improvements to the public realm, transport links, accommodation, the night-time economy and the sustainable development of our natural and built heritage.	This Policy could promote development for tourism. Any developments can potentially have direct and/or indirect impact on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality. Therefore, mitigation measures will be required.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
General Tourism Policies	TRP 6	Promote and support the continued strategic development of Westport, Ballina and Castlebar as tourist destinations through:	Proposed developments like the extension of the GWG and NCP have the potential to directly and/or indirectly negatively impact the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

			value, such as changes in water quality and quantity, and air quality.	
General Tourism Policies	TRP 6	(a) Supporting the enhancement of Westport as a major tourism centre in the West of Ireland, building on its reputation as one of Ireland's premier visitor destinations along the Wild Atlantic Way, as a heritage town and gateway to areas of outstanding natural and built heritage.	General statement of policy/general aspiration which will not lead to development	N/A
General Tourism Policies	TRP 6	(b) Supporting the development of Westport's key assets e.g. Westport House Demesne, Quay area, extension of the Great Western Greenway and additional greenway linkages e.g. Westport-Cong Trail and National Coastal Path.	Proposed developments like the extension of the GWG and NCP have the potential to directly and/or indirectly negatively impact the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
General Tourism Policies	TRP 6	(c) Exploration of the development potential of Westport and Ballina harbour areas in terms of marine related tourism and extensive marine resources.	Developments like marina, pontoon and mooring facilities and other infrastructural developments, ranging from car park enhancements, addition of toilet, changing and shower facilities to more comprehensive Activity Service Centres have the potential to directly and/or indirectly negatively impact the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
General Tourism Policies	TRP 6	(d) Promote the significant tourism potential of Ballina and its location as a prominent stop on the Wild Atlantic Way, a gateway to	Promoting tourism potential of these sites has the potential to lead to increased pressure from increased footfall, which has	Refer to Chapter 9 of this NIR for mitigation measures for any

		Northwest Mayo, the Céide Fields, and internationally renowned salmon fishing, through investment in tourism related infrastructure, including Monasteries of the Moy Greenway from Beleek to Killala, incorporating EuroVelo1 Atlantic Coastal Route, Mountain Biking Trail via Ballina connecting the Wild Nephin Ballycroy National Park to the National Mountain Bike Trail at Coolaney and ongoing development of the Wild Atlantic Way, including Discovery Points at Crockets Town.	the potential to directly and/or indirectly negatively impact the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	plan or project arising from this objective/ policy
General Tourism Policies	TRP 6	(e) Promote the development of Castlebar as a nationally important Sports Tourism & Adventure Hub, including leveraging existing sports assets and existing natural and built facilities at Lough Lannagh Holiday Village.	This Policy could promote development for tourism and sports. Any developments can potentially have direct and/or indirect impact on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality. Therefore, mitigation measures will be required.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
General Tourism Objectives	TRO 1	To work with all relevant stakeholders and Fáilte Ireland to facilitate the erection of standardised signage for tourism facilities and tourist attractions as part of national and regional initiatives.	This Policy is promoting the erection of signage which, although small in size, still has potential to have direct and/or indirect impact on the Natura 2000 network through direct loss of habitats, disturbance of species, or changes in key indicators of conservation value, such as changes in water quality and quantity dependant on the location of the signage an its installation. Therefore, mitigation measures will be required.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
General Tourism Policies	TRO 2	Enable, facilitate and encourage the growth and sustainability of the tourism sector,	Promoting growth through developments in rural areas can have the potential to	Refer to Chapter 9 of this NIR for mitigation measures for any

		through supporting the provision of tourism enterprise developments in rural areas including open farms, subject to the provision of adequate infrastructure and compliance with normal planning considerations.	directly and/or indirectly negatively impact the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality. Potential negative impacts.	plan or project arising from this objective/ policy
General Tourism Objectives	TRP 7	To encourage the clustering of tourism products and services within identified hubs, to facilitate the sharing of infrastructure and services where possible, to increase linkages within and reduce leakage from the local economy	This Policy could promote infrastructural development for linking services. Any developments can potentially have direct and/or indirect impact on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality. Therefore, mitigation measures will be required.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Activity, Adventure & Sports Tourism Policies	TRP 8	To support the implementation of the opportunities and actions identified in the Mayo Tourism Strategy and Action Plan – Destination Mayo 2015-2021, regarding the development of activity, adventure and sports tourism at strategic locations throughout the County in conjunction with the Tourism section of Mayo County Council and other relevant stakeholders.	Any development could potentially have direct and/or indirect impact on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality. Potential negative impacts can occur by implementation of Plans that arise from this support.	Further development of walking routes and trails could potentially have direct and/or indirect impact on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality. Potential negative impacts can occur by implementation of Plans that arise from this support.

Activity, Adventure & Sports Tourism Policies	TRP 9	<p>To support the implementation of the strategy for Greenway development in Ireland - “Strategy for the Future Development of National and Regional Greenways” in relation to the development and extension of greenways throughout the county, in conjunction with the Tourism section of Mayo County Council and other relevant stakeholders.</p>	<p>Proposed developments like the extension of the GWG have the potential to directly and/or indirectly negatively impact the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.</p>	<p>Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy</p>
Activity, Adventure & Sports Tourism Policies	TRP 10	<p>To promote Mayo as a premier walking/cycling destination in the Country and support the further development of walking routes and trails within the county and the integration and linkage of these with other existing / proposed routes and trails both within and outside of County Mayo, in accordance with national walking strategy guidance and conjunction with the Tourism Section of Mayo County Council, Fáilte Ireland and other relevant stakeholders.</p>	<p>Further development of walking routes and trails could potentially have direct and/or indirect impact on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality. Potential negative impacts can occur by implementation of Plans that arise from this support.</p>	<p>Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy</p>
Activity, Adventure & Sports Tourism Policies	TRP 11	<p>To support the development of a regional water-based leisure sector in a sustainable manner, making the best use of existing and planned infrastructure and resources, in a manner that is sensitive to the natural and cultural heritage resources, in conjunction with relevant regional partners.</p>	<p>Even though a statement is made about making the best use of existing infrastructure and a sensitive manner of development, this support will likely lead to increased development and increased pressure on sites. Any development could potentially have direct and/or indirect impact on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality. Potential negative impacts can occur by implementation of Plans that arise from this support.</p>	<p>Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy</p>

<p>Activity, Adventure & Sports Tourism Policies</p>	<p>TRP 12</p>	<p>To support local, national and international tourism agencies and bodies in their work to promote Mayo as a unique tourism destination and build on the 'Wild Mayo' adventure brand and to continue to position itself as 'The Heartbeat of the Wild Atlantic Way' through an interagency approach, utilising available funding streams to implement the tourism objectives of this plan.</p>	<p>Promoting tourism potential of County Mayo has the potential to lead to increased pressure from increased footfall, which has the potential to directly and/or indirectly negatively impact the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.</p>	<p>Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy</p>
<p>Activity, Adventure & Sports Tourism Policies</p>	<p>TRP 13</p>	<p>To work with the National Transport Authority, Fáilte Ireland, Waterways Ireland and all stakeholders to develop a coordinated approach to the selection, delivery and servicing of future greenways, blueways, trails and routes throughout the county.</p>	<p>No negative impacts anticipated from implementation of objective/policy at this stage. Any future greenways, blueways, trails or routes that may be selected as a result of this Policy will require project level assessment.</p>	<p>N/A</p>
<p>Activity, Adventure and Sports Tourism Objectives</p>	<p>TRO 3</p>	<p>To explore a feasibility study, including the development of route options, for the creation of a coastal walking/cycling route along the western seaboard, in collaboration and partnership with the NWRA, other relevant local authorities and stakeholders, including the public, to enhance access to our tourist assets along the route of then Wild Atlantic Way.</p>	<p>Further development of walking routes and trails could potentially have direct and/or indirect impact on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality. Potential negative impacts can occur by implementation of Plans that arise from this support.</p>	<p>Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy</p>
<p>Activity, Adventure and Sports Tourism Objectives</p>	<p>TRO 4</p>	<p>To investigate the development of bike trails and sites at Keenagh / Ballycastle/Bunnyconnellan Bike Trail and the Mountain Bike Centre at the Wild Nephin site, in conjunction with the Tourism Section of Mayo County Council</p>	<p>Further development of walking routes and trails could potentially have direct and/or indirect impact on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.</p>	<p>Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy</p>

			<p>Potential negative impacts can occur by implementation of Plans that arise from this support.</p>	
<p>Spiritual and Pilgrimage Tourism Policy</p>	<p>TRP 14</p>	<p>(a) To exploring the development of a long-distance Spiritual Trail linking Croagh Patrick, Ballintubber Abbey and Knock Shrine. Investigate the potential of linking this trail to Burriscarra, Cong, Turlough and the Monasteries of the Moy through the Tourism Section of Mayo County Council and other relevant stakeholders.</p> <p>(b) To explore a range of spiritual walking and pilgrimage events to showcase Mayo as a spiritual tourism destination, with particular emphasis on generating overnight stays for visitors to Knock Shrine.</p> <p>(c) To support and facilitate the development of Knock as a world-renowned religious tourist destination</p> <p>(d) To investigate the delivery of pilgrim trail improvements and upgrades on the Croagh Patrick trail and Tochar Padraig trail. Opportunities to enhance ecological connectivity should be integrated as part of any linking of routes to strengthen and support green infrastructure. SEA and AA measures relating to the above were identified in the Destination Mayo Plan and should be adhered to and integrated to any project level assessments. tourism in Mayo, in conjunction with the Fáilte Ireland, the Tourism section of Mayo County Council and other relevant stakeholders through:</p>	<p>The development of such a trail could potentially have direct and/or indirect impact on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality. Negative impacts can potentially occur by development of this trail.</p>	<p>Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy</p>

Cultural and Amenity Tourism Policies	TRP 15	To support the implementation of the opportunities and actions identified in the Mayo Tourism Strategy and Action Plan – Destination Mayo 2015-2021 regarding the development of cultural and amenity tourism and the Wild Atlantic Way in Mayo, in conjunction with the Tourism section of Mayo County Council and other relevant stakeholders.	General statement of policy/general aspiration which will not lead to development	N/A
Cultural and Amenity Tourism Policies	TRP 16	Support the protection and enhancement of Mayo’s historic buildings, gardens and museums as cultural tourist attractions.	General statement of policy/general aspiration which will not lead to development	N/A
Cultural and Amenity Tourism Policies	TRP 17	Support developments which will enable and encourage countryside recreation and an increased appreciation of, and access to, the natural environment.	Supporting these developments can potentially have direct and/or indirect impact on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality. Potential negative impacts can occur through supporting these developments.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Cultural and Amenity Tourism Policies	TRP 18	To work in partnership with Coillte to identify opportunities for tourism and recreation facility development within commercially managed forests, where appropriate, and promote and continue to be involved in the Neighbour Wood Scheme (2017) and to identify areas at local level that are suitable for such schemes.	Supporting these developments can potentially have direct and/or indirect impact on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality. Potential negative impacts can occur through supporting these developments.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Cultural and Amenity	TRO 5	Encourage access to forestry and woodlands, including private forestry, in cooperation with stakeholders for walking routes, bridle paths,	Encouraging access to forestry and woodlands can potentially have direct and/or indirect impact on the Natura 2000	Refer to Chapter 9 of this NIR for mitigation measures for any

Tourism Objectives		mountain biking, nature walks, orienteering, hiking, recreational areas and other similar facilities and to retain existing public rights of way through forest lands.	network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality. Potential negative impacts can occur supporting these developments.	plan or project arising from this objective/ policy
Cultural and Amenity Tourism Objectives	TRO 6	To investigate the development of Moore Hall Estate and walled garden in conjunction with the Tourism Section of Mayo County Council and relevant working groups.	Unlikely to have any significant adverse impacts upon the integrity of any Natura 2000 site	N/A
Cultural and Amenity Tourism Objectives	TRO 7	To investigate the further potential of the Wild Nephin Ballycroy National Park and Dark Sky initiative in a sustainable manner. This approach should demonstrate good practice in terms of eco-tourism and wilderness and seek to gain recognised accreditation on same.	Even though it is stated that further development will be in a sustainable manner, this can potentially have direct and/or indirect impact on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality. Potential negative impacts can occur from further development and investigating the tourism potential of the Park.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Cultural and Amenity Tourism Objectives	TRO 8	Explore the development of community walks, off road trails/rural trail developments, parks, other outdoor amenities and recreational infrastructure, and work with relevant landholders and recreational/tourism agencies to increase access to the countryside and our coastal areas, subject to proper planning and sustainable development principles.	Even though the increased access will be subject to proper planning and sustainable development principles, this policy statement will indirectly lead to increased development, which can potentially have direct and/or indirect impact on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

<p>Cultural and Amenity Tourism Objectives</p>	<p>TRO 9</p>	<p>Encourage sensitively designed and located development which provides for the appreciation, interpretation, upgrade and provision of access to natural habitats, scenic vistas and heritage features for the benefit of rural tourism, subject to normal planning and nature conservation consideration.</p>	<p>Regardless of the fact that it will be subject to nature conservation consideration, any development can potentially have direct and/or indirect impact on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.</p>	<p>Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy</p>
<p>Cultural and Amenity Tourism Objectives</p>	<p>TRO 10</p>	<p>To utilise Mayo’s natural and heritage resources to foster the development of tourism as a viable sustainable sector of the economy, in a sustainable manner, which complements the scale, quality and unique features of the county</p>	<p>Increased tourism can have direct and/or indirect impact on the Natura 2000 network through pressure on sites created by more footfall. Potential negative impacts can therefore occur.</p>	<p>Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy</p>
<p>Food and Culinary Tourism Policy</p>	<p>TRP 19</p>	<p>To support the development of Mayo as a ‘Foodie Destination’ through the implementation of the Fáilte Ireland ‘Food Tourism Development Strategy 2018-2023’ and the Mayo County Council Food and Drink Strategy 2020-2025 and the opportunities and actions identified in the Mayo Tourism Strategy and Action Plan – Destination Mayo 2015-2021 regarding the development of food tourism in Mayo, in conjunction with the Tourism section of Mayo County Council and other relevant stakeholders.</p>	<p>General statement of policy/general aspiration which will not lead to development</p>	<p>N/A</p>
<p>Food and Culinary Tourism Objectives</p>	<p>TRO 11</p>	<p>Explore the development of a Mayo food network and market signature food experiences around the county, including artisan food trails such as the Gourmet Greenway and Edible Greenway and other food trails, at appropriate locations throughout the county.</p>	<p>Development of trails can potentially have direct and/or indirect impact on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.</p>	<p>Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy</p>

<p>Food and Culinary Tourism Objectives</p>	<p>TRO 12</p>	<p>To facilitate Feile na Tuaithe and the Westport and Ballina food festivals and other food and drinks tourism events at appropriate locations throughout the county.</p>	<p>One off events that may be held in rural locations could occur within the footprint of a Natura 2000 site. Any events facilitated within the ZOI of a Natura 2000 site could cause direct and/or indirect adverse impacts to the Natura 2000 network causing fragmentation or loss of habitats, disturbance or fragmentation of key species, or changes to indicators of conservation value such as changes in water quality.</p>	<p>Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy</p>
<p>Festivals and Events Policies</p>	<p>TRP 20</p>	<p>To support the implementation of the opportunities and actions identified in the Mayo Tourism Strategy and Action Plan – Destination Mayo 2015-2021 regarding the development festivals and events in Mayo, in conjunction with the Tourism section of Mayo County Council and other relevant stakeholders by:</p>	<p>General statement of policy/general aspiration which will not lead to development</p>	<p>N/A</p>
<p>Festivals and Events Policies</p>	<p>TRP 20</p>	<p>(a) supporting community groups and festival committees to identify and access new sources of funding for festivals and events in the county;</p>	<p>General statement of policy/general aspiration which will not lead to development</p>	<p>N/A</p>
<p>Festivals and Events Policies</p>	<p>TRP 20</p>	<p>(b) promoting the development of a variety of new festivals, including Mayo Dark Skies International and sporting events, to appeal to a wide range of visitors and to increase the profile of the county as a key tourism destination.</p>	<p>Festivals or events that may be held throughout Mayo could occur within the footprint of a Natura 2000 site. Any events facilitated within the ZOI of a Natura 2000 site could cause direct and/or indirect adverse impacts to the Natura 2000 network causing fragmentation or loss of habitats, disturbance or fragmentation of key species, or changes to indicators of conservation value such as changes in water quality.</p>	<p>Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy</p>

<p>Festivals and Events Policies</p>	<p>TRP 21</p>	<p>Support and promote existing festivals and sporting events, in conjunction with the relevant tourism and sporting agencies to increase the cultural, heritage and lifestyle profile of the county, and to facilitate the establishment of new events where viable, subject to the satisfactory location, access, parking provision and protection of the surrounding environment.</p>	<p>New or existing festivals or events that may be held throughout Mayo could occur within the footprint of a Natura 2000 site. Any events facilitated within the ZOI of a Natura 2000 site could cause direct and/or indirect adverse impacts to the Natura 2000 network causing fragmentation or loss of habitats, disturbance or fragmentation of key species, or changes to indicators of conservation value such as changes in water quality.</p>	<p>Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy</p>
<p>Tourism and Infrastructure Policies</p>	<p>TRP 22</p>	<p>To support the implementation of priority infrastructural developments and tourism facilities identified by the Tourism Section of Mayo County Council, including the development of new tourist facilities or upgrading/extension of existing tourist facilities at tourist sites within the county, within proper planning and sustainable development principles.</p>	<p>Development of trails can potentially have direct and/or indirect impact on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.</p>	<p>Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy</p>
<p>Tourism and Infrastructure Policies</p>	<p>TRP 23</p>	<p>Support and promote sustainable tourism development, accessible to all throughout the County and to work in partnership with tourism organisations and adjoining Local Authorities where necessary, in securing the development of tourism enterprises and infrastructure. Subject to suitable locations where it can be demonstrated that the development will not have significant adverse effects on the environment, including the integrity of the Natura 2000 network, residential amenity or visual amenity</p>	<p>Any development can potentially have direct and/or indirect impact on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.</p>	<p>Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy</p>
<p>Tourism and Infrastructure Policies</p>	<p>TRP 24</p>	<p>To encourage the clustering of tourism products and services within identified hubs, to facilitate the sharing of infrastructure and</p>	<p>This Policy could promote infrastructural development for linking services. Any developments can potentially have direct</p>	<p>Refer to Chapter 9 of this NIR for mitigation measures for any</p>

		services where possible, to increase linkages within and reduce leakage from the local economy.	and/or indirect impact on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality. Therefore, mitigation measures will be required.	plan or project arising from this objective/ policy
Tourism and Infrastructure Policies	TRO 13	Identify opportunities for funding for infrastructural projects and facilities, including LEADER, national and European funding schemes and seek to maximise the benefit of such funding opportunities to the county.	Any development can potentially have direct and/or indirect impact on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Tourism Accommodation Policy	TRP 25	To promote the sustainable development of the tourism sector in appropriate locations throughout the county.	Increased tourism can have direct and/or indirect impact on the Natura 2000 network through pressure on sites created by more footfall. Potential negative impacts can therefore occur.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Tourism Accommodation Objectives	TRP 26	Encourage proposals to reinstate, conserve and/or replace existing ruinous or disused dwellings for holiday home purposes, subject to normal planning considerations including design, safe access and provision of any necessary wastewater disposal facilities.	Any development can potentially have direct and/or indirect impact on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Tourism Accommodation Objectives	TRO 14	To ensure that tourism related accommodation such as holiday homes, hotels, caravan/camping parks, glamping etc. are primarily located within existing settlements where there is existing infrastructure provision to service the development and where they can contribute to maintenance of essential rural services.	This Policy could promote development for tourism. Although sustainable and in consideration of the Natura 2000 network, impacts may still arise as a result of this Policy. Any developments can potentially have direct and/or indirect impact on the	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

		<p>Certain forms of low-impact tourist accommodation such as Camping and Glamping/Pod sites may also be considered outside of existing settlements where it is:</p> <ul style="list-style-type: none"> • proposed to incorporate the reuse an existing structure as an integral part of the development • adjacent to, and capable of availing of, an existing appropriate commercial enterprise or community facility or located on an existing farm <p>In all cases the facility shall be of an appropriate scale for the location and shall have a high standard of design, layout, landscape, including Sections 6.2 and 6.3 of the Development Management Standards (Volume 2), and environmental protection so as not to impact negatively on the visual and residential amenity of the area or have significant adverse effects on the environment, including the integrity of the Natura 2000 network.</p>	<p>Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality. Therefore, mitigation measures will be required.</p>	
Tourism Accommodation Objectives	TRO 15	<p>To facilitate, where appropriate, the conversion of former demesnes or estate dwellings and their outbuildings into tourism facilities, subject to good planning principles and architectural practice.</p>	<p>Any development can potentially have direct and/or indirect impact on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.</p>	<p>Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy</p>
Tourism Accommodation Objectives	TRO 16	<p>Facilitate the development of a variety of quality tourist accommodation types, at suitable locations, throughout the county.</p>	<p>Any development can potentially have direct and/or indirect impact on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.</p>	<p>Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy</p>
Tourism Accommodation Objectives	TRO 17	<p>To positively consider the development of new hotels in existing settlements, with particular preference for locations in larger settlements.</p>	<p>Any development can potentially have direct and/or indirect impact on the Natura 2000 network through fragmentation or</p>	<p>Refer to Chapter 9 of this NIR for mitigation measures for any</p>

		In rural locations (i.e villages / rural nodes), it must be demonstrated that: (i) the area proposed to be served by the new development has high visitor numbers associated with an existing attraction / facility; (ii) a need for new / additional hotel type accommodation for these visitors has been identified having regard to the profile of the visitor and the availability and proximity of existing hotels in the area; and (iii) the distance of the location from a significant settlement is such that visitors to the area/attraction are unlikely to avail of existing hotel facilities.	loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	plan or project arising from this objective/ policy
Tourism Accommodation Objectives	TRO 18	To positively consider the (part) conversion of existing dwellings to Bed & Breakfasts (B&Bs) and Guesthouses, to be operated by the owner-occupier of the dwelling. Applications for new build B&Bs /guesthouses will in the first instance be evaluated as private dwellings and the objectives and standards applicable in that area type (e.g. large town, rural town, rural area etc) will be applied.	Any development can potentially have direct and/or indirect impact on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Tourism Accommodation Objectives	TRO 19	To ensure that holiday home / self-catering developments on a farm holding shall be provided by farmhouse extension or by the utilisation of other existing dwellings / structures on the property. Only where it has been demonstrated that these are not viable options, will permission be considered for new build development. Any new build development shall be in close proximity to the existing farmhouse.	Any development can potentially have direct and/or indirect impact on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Tourism Accommodation Objectives	TRO 20	To facilitate the development of hostels along established walking / hiking routes and	Any development can potentially have direct and/or indirect impact on the Natura 2000 network through fragmentation or	Refer to Chapter 9 of this NIR for mitigation measures for any

		adjacent to existing tourism / recreation facilities, subject to normal planning criteria.	loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	plan or project arising from this objective/ policy
Flagship Infrastructure Projects Policy	TRO 21	To strategically facilitate the development of infrastructural flagship projects through the implementation of the National Tourism Strategy 'People, Place and Policy: Growing Tourism to 2025' Department of Transport, Tourism and Sport and the Mayo County Council Tourism Strategy 'Destination Mayo'	Any development can potentially have direct and/or indirect impact on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	
Flagship Infrastructure Projects Objectives	TRO 22	Identify all opportunities for funding for Flagship Projects, including LEADER, national and European funding schemes and seek to maximise the benefit of such funding opportunities to the county.	General statement of policy/general aspiration which will not lead to development	N/A
Wild Atlantic Way Policy	TRP 27	To support the promotion of the Wild Atlantic Way in its role to grow the economic contribution of tourism along its route, through the upgrade and improvement of the touring network, facilities and visitor attractions through:	Any development could potentially have direct and/or indirect impact on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality. Potential negative impacts can occur by implementation of Plans that arise from this support.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Wild Atlantic Way Policy	TRP 27	(a) Exploration of the development of signature discovery points to assist and secure Mayo's position as 'The Heartbeat of the Wild Atlantic Way'.	Any development can potentially have direct and/or indirect impact on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

			changes in water quality and quantity, and air quality.	
Wild Atlantic Way Policy	TRP 27	(b) Working with all relevant stakeholders and Fáilte Ireland to facilitate the erection of standardised signage for tourism facilities and tourist attractions along the Wild Atlantic Way.	This Policy is promoting the erection of signage which, although small in size, still has potential to have direct and/or indirect impact on the Natura 2000 network through direct loss of habitats, disturbance of species, or changes in key indicators of conservation value, such as changes in water quality and quantity dependant on the location of the signage an its installation. Therefore, mitigation measures will be required.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Wild Atlantic Way Policy	TRP 27	(c) Working with relevant landholders and recreational/tourism agencies to increase access to the countryside and our coastal areas, and to ensure maintenance and access to the existing network of trails, paths, and tourist sites along the Wild Atlantic Way.	Increased access can potentially have direct and/or indirect impact on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Wild Atlantic Way Policy	TRP 27	(d) Supporting the exploration of the development of new tourist services, parking and facilities or upgrading/extension of existing tourist services, parking and facilities at tourist sites along the Wild Atlantic Way	Any development can potentially have direct and/or indirect impact on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Walking & Greenway Policies	TRP 28	To support the implementation of the NWRA Regional Outdoor Recreation Strategy, through the extension of greenways, walking routes, tracks and trails within the county and the integration and linkage of them with other existing / proposed greenways, walking routes,	Developments like the extension of greenways can potentially have direct and/or indirect impact on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

		tracks and trails, both within and outside County Mayo.	conservation value, such as changes in water quality and quantity, and air quality.	
Walking & Greenway Policies	TRP 29	To support the progression of long-distance linear greenways and greenway town networks, and the further development and improvements of the following walking trails and coastal paths in the county: Westport/Cong Walking trail; Croagh Patrick trail upgrade; Tourmakeady Trail; Bangor trail; Ceide /Downpatrick Coastal Path; and Slievemor Trail, in conjunction with the Tourism Section and other relevant stakeholders.	Developments like the extension of greenways can potentially have direct and/or indirect impact on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Marine Facilities Policy	TRP 30	Support the protection and enhancement of our islands, coastline and waterways as tourism products and attractions.	General statement of policy/general aspiration which will not lead to development	N/A
Marine Facilities Objectives	TRO 23	Identify strategic marine locations, in line with the Mayo Tourism Strategy and Action Plan, for the development of pontoon and mooring facilities.	Developments like pontoon and mooring facilities and other infrastructural developments, ranging from car park enhancements, addition of toilet, changing and shower facilities to more comprehensive Activity Service Centres have the potential to directly and/or indirectly negatively impact the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Marine Facilities Objectives	TRO 24	Explore the development of activity service centres in accordance with the Mayo Tourism Strategy and Action Plan.	Any development could potentially have direct and/or indirect impact on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

			indicators of conservation value, such as changes in water quality and quantity, and air quality. Potential negative impacts could occur through the development of activity service centres	
Marine Facilities Objectives	TRO 25	Investigate facility upgrades at Ronagh Pier.	Any development could potentially have direct and/or indirect impact on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality. Potential negative impacts could occur through the development of activity service centres	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Marine Facilities: Adventure Policy	TRP 31	Support the development of an activity centre at Keel and activity hub at Lough Lannagh, Castlebar	Any development could potentially have direct and/or indirect impact on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality. Potential negative impacts could occur through the development of activity service centres	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Adventure Policy	TRO 26	Support the protection and enhancement of our counties lakes as visitor attractions, through the identification of infrastructure requirements and development of masterplans including:	Any development could potentially have direct and/or indirect impact on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality. Potential negative impacts could occur through the development of activity service centres	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Adventure Policy	TRO 26	(a) The exploration of the development of the Pontoon Lakes masterplan and construction of a boardwalk in conjunction with the Tourism Section of Mayo County Council;	Developments like an activity centre or an activity hub can potentially have direct and/or indirect impact on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Adventure Policy	TRO 26	(b) Working with relevant landholders and recreational/tourism agencies to increase access to Mayo's lakes.	Increasing access to Mayo's lakes can potentially have direct and/or indirect impact on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Adventure Policy	TRP 32	Preserve and protect verified public rights of way which give access to seashore, mountain, lakeshore, riverbank or other places of natural beauty or recreational utility	Preservation and/or protection of public rights of way could involve changes to substrate, fencing or other physical works. Any works that may occur within the ZOI of a Natura 2000 site could potentially cause impacts to the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Public Rights of Way Objectives	TRO 27	The Council, within the lifetime of this plan, to:		
Public Rights of Way Objectives	TRO 27	a. Identify, preserve and enhance existing accesses and public rights of way to recreational areas including the coast, upland areas, lakeshores, river-bank areas and heritage sites	Preservation and/or protection of public rights of way could involve changes to substrate, fencing or other physical works. Any works that may occur within the ZOI of a Natura 2000 site could potentially cause impacts to the Natura 2000 network	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

			through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	
Public Rights of Way Objectives	TRO 27	b. Where necessary, establish new accesses and public rights of way to recreational areas including the coast, upland areas, lakeshores, river-bank areas and heritage sites in co-operation with landowners and the local community. When public rights of way are identified, the owners of the public rights of way shall be notified in accordance with the Planning & Development Acts 2000-2010	Establishing new accesses can potentially have direct and/or indirect impact on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Public Rights of Way Objectives	TRO 27	c. Map public rights of way in accordance with the Planning & Development Acts 2000-2010	General statement of policy/general aspiration which will not lead to development	N/A
Public Rights of Way Objectives	TRO 27	a. and b. shall be carried out in accordance with best sustainable management practices, any guidance from the Department of the Environment, Community and Local Government and where it can be demonstrated that the development will not have significant adverse effects on the environment including the integrity of the Natura 2000 network.	General statement of policy/general aspiration which will not lead to development	N/A
Public Rights of Way Objectives	TRO 28	To seek to identify and protect over the lifetime of the Plan further existing <u>verified</u> rights of way which give access to seashore, mountain, lakeshore, riverbank or other place of natural beauty or recreational utility, as they become available to the Planning Authority over the lifetime of the plan. (accompanied by mapping showing public rights of way).	Protection of public rights of way could involve changes to substrate, fencing or other physical works. Any works that may occur within the ZOI of a Natura 2000 site could potentially cause impacts to the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

			value, such as changes in water quality and quantity, and air quality.	
--	--	--	--	--

DRAFT



A.5 Chapter 6 – Movement and transport

Strategic Aim: To achieve a sustainable, integrated and low carbon transport system with excellent connectivity within and to Mayo, by enhancing existing strategic transportation infrastructure in the county. To work with other agencies in the provision of infrastructure to attract new business investment and people into the county.

Movement and transport				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
Integrated Land Use and Transportation Policies	MTP – 1	Support sustainable travel in the County by ensuring future population and employment growth predominantly takes place in urban areas that will warrant provision of public transport services	This objective could lead to increased development of public transport. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Integrated Land Use and Transportation Policies	MTP – 2	To support and facilitate the integration of land use with transportation infrastructure, through the development of sustainable compact settlements which are well served by public transport.	This objective could lead to increased development of public transport. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Integrated Land Use and Transportation Policies	MTP – 3	Support and facilitate any ‘Smarter Travel’ initiatives that will improve sustainable transportation within the County, including public transport, electric and hybrid vehicles, car clubs, public bike schemes, improved pedestrian and cycling facilities, as appropriate	This objective could lead to increased development of public transport. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Movement and transport				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
Integrated Land Use and Transportation Objective	MTO - 1	To prepare and commence implementation of, Local Transport Plans (LTP), in conjunction with the National Transport Authority and relevant stakeholders, for Ballina, Castlebar and Westport, other settlements where appropriate.	General statement of policy/general aspiration which will not lead to development	N/A
Land Use Integration & Sustainable Transport Objectives	MTO - 2	Ensure that planning applications for large scale developments, that are significant trip intensive generators are accompanied by a Mobility Management Plan	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Sustainable Mobility Policy	MTP - 4	Support sustainable mobility, enhanced regional accessibility and connectivity within County Mayo in accordance with the National Policy Outcomes of the National Planning Framework 2040, National Development Plan and the Regional Planning Objectives of the Regional & Spatial, Economic Strategy.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Sustainable Mobility Policy	MTP - 5	Promote the transition to a low carbon integrated transport system by firstly reducing the need for travel through the use of design solutions and innovative approaches with regards to the Design Manual for Urban Roads and Streets, and subsequently to shift to environmentally sustainable modes of transport	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Movement and transport				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
Sustainable Mobility Objectives	MTO - 3	To liaise and collaborate with relevant agencies to support and encourage the growth of electric vehicles and EBikes with support facilities/infrastructure, through a roll-out of additional electric charging points in collaboration with relevant agencies at appropriate locations with a particular emphasis in public parking areas and employment locations (including along the Wild Atlantic Way) including retrofit of charging points in existing urban centres	This objective could lead to development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Sustainable Mobility Objectives	MTO - 4	To increase cycling usage in Tier I and Tier II settlements in line with the national average (2016).	This objective could lead to development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Pedestrians & Cyclists Policy	MTP – 6	Support safer cycling/walking routes to encourage people to be more physically active for transport and leisure purposes.	This objective could lead to development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Pedestrians & Cyclists Policy	MTP – 7	To promote the design and construction of new developments to create low carbon, walkable neighbourhoods and workplaces containing high quality green and blue infrastructure.	This objective could lead to development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation	Refer to Chapter 9 of this NIR for mitigation measures for any plan

Movement and transport				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
			or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	or project arising from this objective/ policy
Pedestrians & Cyclists Policy	MTO 5	To encourage and facilitate the maintenance and further development of the public footpath network, walking and cycling routes and associated infrastructure and where possible the retrofitting of cycle and pedestrian routes into the existing urban road network.	This objective could lead to development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Pedestrians & Cyclists Policy	MTO 6	To support the establishment of a network of interlinked cycle ways and walk ways in the county and the adjoining counties, having regard to best practice standards including the Design Manual for Urban Roads and Streets and the NTA Cycle Manual or any amending/superseding national guidance or manuals	The development of cycling infrastructure within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Pedestrians & Cyclists Policy	MTO 7	To endeavour to identify, preserve and enhance existing accesses and public rights of way to recreational areas including the coast, upland areas, lakeshores, river-bank areas and heritage sites over the lifetime of the plan, subject to any forthcoming departmental guidance.	General statement of policy/general aspiration which will not lead to development	N/A
Pedestrians & Cyclists Policy	MTO 8	To encourage, where appropriate, the incorporation of safe and efficient cycleways, accessible footpaths and pedestrian routes into the design schemes for town centres/neighbourhood centres, residential, educational, employment, recreational	This objective could lead to development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Movement and transport				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		developments and other uses, with the design informed by published design manuals, including the Design Manual for Urban Roads and Streets and the NTA Cycle Manual or any amending/superseding national guidance or manuals.	value, such as changes in water quality and quantity, and air quality.	
Pedestrians & Cyclists Policy	MTO 9	To continue the development of a network of Greenways in the County in accordance with best practice and where it can be demonstrated that the development will not have significant adverse effects on the environment including the integrity of the Natura 2000 network.	Policies or proposals which steer change in such as way as to protect international nature conservation sites from adverse effects therefore positive impacts are anticipated from the implementation of this objective/policy	N/A
Pedestrians & Cyclists Policy	MTO 10	Investigate the potential of providing loped Greenways / Walkways for the Tier 1, 2 and 3 Settlements of the County, where it can be demonstrated that such schemes will not have a significant adverse effect on the environment including the integrity of the Natura 2000 network	Policies or proposals which steer change in such as way as to protect international nature conservation sites from adverse effects therefore positive impacts are anticipated from the implementation of this objective/policy	N/A
Pedestrians & Cyclists Policy	MTO 11	To investigate the protentional of providing a number of 'Park and Stride' facilities in appropriate locations in the county, subject to an environmental assessment of identified locations.	General statement of policy/general aspiration which will not lead to development	N/A
Pedestrians & Cyclists Policy	MTO 12	Seek to advance the walking/ cycling projects listed in the table below	This objective could lead to development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Movement and transport				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
Pedestrians & Cyclists Objectives	MTO 13	Protect open spaces, with multifunctional green and blue infrastructure in developments, with connections to the wider network of open spaces and habitats	Policies or proposals which steer change in such as way as to protect international nature conservation sites from adverse effects therefore positive impacts are anticipated from the implementation of this objective/policy	N/A
Bus Policy	MTP – 8	Support and encourage public transport providers and rural community transport initiatives and programmes, such as the Local Link Rural Transport Programme, to enhance to provision of public transportation services linking rural villages to the main towns of Mayo	This objective could lead to development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Bus Policy	MTO – 14	Support the operation of existing bus services by facilitating the provisions of improved facilities and services for bus users in towns and villages including the provision of set down areas for coaches and bus shelters at all bus stops where feasible.	This objective could lead to development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Rail Policy	MTP - 9	To Support the enhancement of rail services to Dublin and commuter services between Ballina, Castlebar, Westport and Claremorris with connectivity to Galway and Limerick Metropolitan Cities and major international ports such as Shannon / Foynes with the realisation of re-opening the Western Rail Corridor	This objective could lead to development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Rail Policy	MTP - 10	To support the re-opening of the Western Rail Corridor in order to deliver the Tuam –	This objective could lead to development of infrastructure. Any development within the county	Refer to Chapter 9 of this NIR for mitigation

Movement and transport				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		Claremorris – Sligo Rail to an appropriate level of service and to a standard capable of facilitating passenger and freight transport, and would present an opportunity to provide an integrated rail linkage to the IWA Knock SDZ.	could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	measures for any plan or project arising from this objective/ policy
Rail Policy	MTP - 11	Support and encourage the provision of a high quality rail network and service (including commuter services) and ancillary works for passenger and freight carriage to, from and within the County, where it can be demonstrated that the development will not have significant adverse effects on the environment including the integrity of the Natura 2000 network	Policies or proposals which steer change in such as way as to protect international nature conservation sites from adverse effects therefore positive impacts are anticipated from the implementation of this objective/policy	N/A
Rail Policy	MTO 15	To liaise with and encourage Iarnród Éireann to:	This objective could lead to development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Rail Policy	MTO 15	A. Continue investment in rail freight facilities at Ballina & Claremorris.	This objective could lead to development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Movement and transport				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
Rail Policy	MTO 15	Increase frequency of commuter services on mainline rail network between Westport, Castlebar & Ballina	This objective could lead to development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Rail Objectives	MTO 16	Support, and encourage the upgrading of existing railway stations, and protect, as required, lands necessary for the upgrading of existing railway lines or stations.	This objective could lead to development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Rail Objectives	MTO 17	To work with the National Transport Authority & Iarnród Éireann, to safeguard and protect all existing or historic rail lines and associated facilities from redevelopment for non-transport related purposes in order to not preclude their future use as an operational transportation network	This objective could lead to development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Rail Objectives	MTO 18	Support and facilitate the velo rail project on the Western Rail Corridor as an interim use for the rail line pending its reopening for passenger and rail freight.	This objective could lead to development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Movement and transport				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
			value, such as changes in water quality and quantity, and air quality.	
National Roads Policy	MTP 12	to enhance Regional Accessibility between key urban centres of population and their regions through the protection of the capacity, efficiency and safety of the National Road Network (listed in appendix) in Mayo	This objective could lead to development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
National Roads Policy	MTP 13	To support the upgrading to a 'High Quality Road' of both the N5 & N17 National Primary Routes, within the county, as part of the Tran-European Transport Network (TEN-T).	This objective could lead to development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
National Roads Policy	MTP 14	To support Transport Infrastructure Ireland in the provision of service and rest area facilities that may be proposed by the TII, and have regard to the provisions of Section 2.8 of the DoECLG Spatial Planning and National Roads Guidelines and the NRA Service Policy (August 2014), with regard to any other proposals for roadside service facilities or off-line service facilities along national roads and junctions that may be promoted by private developers.	This objective could lead to development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
National Roads Policy	MTP 15	To protect the capacity, efficiency and safety of the national road network in Mayo by complying	This objective could lead to development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts	Refer to Chapter 9 of this NIR for mitigation measures for any plan

Movement and transport				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		with the 'Spatial Planning and National Roads - Guidelines for planning authorities' (2012).	on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	or project arising from this objective/ policy
National Roads Policy	MTP-16	To avoid the creation of any additional access points from new development or the generation of increased traffic from existing accesses to national roads to which speed limits greater than 60 km/h apply.	Policy that cannot lead to development or other change therefore no negative impacts anticipated from implementation of objective/policy.	N/A
National Roads Policy	MTP-17	In relation to sections of national roads on the approaches to or exit from urban centres that are subject to a speed limit of 60 kph before a lower 50 kph limit is encountered – otherwise known as transitional zones - to provide for a limited level of direct access to facilitate orderly urban development. Any such proposal must, however, be subject to a road safety audit carried out in accordance with the TII requirements and a proliferation of such entrances, which would lead to a diminution in the role of such zones, must be avoided	This objective could lead to development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
National Roads Objectives	MTO 19	To apply an less restrictive approach to non-residential development of strategic or national importance or extensions to such developments accessing onto the National Road Network in accordance with the provisions of Section 2.6 of the 'Spatial Planning and National Roads - Guidelines for planning authorities' (2012).	This objective could lead to development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Movement and transport				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
National Roads Objectives	MTO 20	To ensure that developments which have the potential to generate significant traffic movements, as per the TII traffic and transportation guidelines which will be required to carry out the assessments set out 7.5 of the Development Management (Volume 2).	This objective could lead to development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
National Roads Objectives	MTO 21	To seek to progress the National Road projects, listed in Table 6.5 subject, to required environmental assessments.	This objective could lead to development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
National Roads Objectives	MTO 22	To seek to review, in conjunction with TII, a reduction of the maximum speed limit along National Routes, where such routes pass through identified settlements in the Settlement Strategy of this Plan.	Policy that cannot lead to development or other change therefore no negative impacts anticipated from implementation of objective/policy.	N/A
Non-National Roads Policy	MTP – 18	to enhance Regional Accessibility between key settlements of Mayo and their regions and to safeguard existing and future capital investment through the protection of the capacity, efficiency and safety of Strategically Important Regional Roads (listed in appendix	This objective could lead to development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Non-National Roads Policy	MTP – 19	The Council, in co-operation with the NTA, TII, Department of Transport, Tourism and Sport and	Policies or proposals which steer change in such as way as to protect international nature conservation	N/A

Movement and transport				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		other stakeholders, to continue with the strengthening and improvement of the local road network including links, by-passes and relief roads, with priority given to those serving the Key Towns and interconnection between such settlements, where it can be demonstrated that the development will not have significant adverse effects on the environment or Natura 2000 network	sites from adverse effects therefore positive impacts are anticipated from the implementation of this objective/policy	
Non-National Roads Policy	MTP 20	It is an objective of Mayo County Council, in relation to lands adjoining Strategically Important Regional Roads to which to which speed limits greater than 60 km/h apply, to avoid the creation of any additional access points from new development or the generation of increased traffic from existing accesses to Strategically Important Regional Roads to which speed limits greater than 60 km/h apply, unless it can be demonstrated the development is required for economic or social reasons and cannot be accessed from a non-Strategically Important Regional Road.	This objective could lead to development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Non-National Roads Policy	MTP 21	Implement the recommendations of the Design Manual for Urban Roads and Streets (DMURS) in relation to urban streets and roads within the 50/60 kph zone	This objective could lead to development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Movement and transport				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
Non-National Roads Objectives	MTO 23	To improve and maintain regional and county roads in line with the annual roads programme and allocated budgets.	This objective could lead to development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Non-National Roads Objectives	MTO 24	Liaise with the TII and DTTAS regarding the revision of speed limits in the County.	General statement of policy/general aspiration which will not lead to development	N/A
Non-National Roads Objectives	MTO 25	Facilitate the continued improvement and upgrading of all roads, should their status be re-graded, under the national roads programme and / or the Council's road programme. The Council will seek and support the upgrading of the status of regional roads in the county which perform functions akin to National Secondary Routes.	This objective could lead to development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Non-National Roads Objectives	MTO 26	Promote road safety measures throughout the County, including traffic calming, road signage and parking.	This objective could lead to development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Non-National Roads Objectives	MTO 27	To seek to progress the non-National Road projects, listed in Table No. 6.5 subject, to required environmental assessments.	This objective could lead to the development of roads. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of	Refer to Chapter 9 of this NIR for mitigation measures for any plan

Movement and transport				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
			habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	or project arising from this objective/ policy
Air Transport Policy	MTP - 22	Support the development of Ireland West Airport Knock as a strategic driver of economic development in the area and as a vital asset in maintaining & enhancing the connectivity of the region	This objective could lead to development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Air Transport Policy	MTO 28	Safeguard public safety in the area around Ireland West Airport Knock by ensuring all development within 15km of Ireland West Airport Knock be subject to safe-guarding restrictions outlined in the Development Guidance document of this Plan.	Safe-guarding of land to prevent development therefore no negative impacts anticipated from implementation of objective/policy.	N/A
Air Transport Policy	MTO 29	Ensure any development associated with light aircraft/helicopter activity is approximately located in areas that avoid significant adverse effects on the environment, the integrity of the Natura 200 network and residential amenity	Policies or proposals which steer change in such as way as to protect international nature conservation sites from adverse effects therefore positive impacts are anticipated from the implementation of this objective/policy	N/A
Ports, Harbours & Piers Policies	MTP 23	Support the development and improvements of ports, harbours, piers, slipways and associated shore facilities and access, where appropriate, where it can be demonstrated that the development will not have significant adverse effects on the environment including the integrity of the Natura 2000 network	Policies or proposals which steer change in such as way as to protect international nature conservation sites from adverse effects therefore positive impacts are anticipated from the implementation of this objective/policy	N/A

Movement and transport				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
Ports, Harbours & Piers Objectives	MTO 30	To investigate the feasibility of create a deep sea harbour/port at Kilcummin	This objective could lead to development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Ports, Harbours & Piers Objectives	MTO 31	Facilitate the implementation of the Marine infrastructural projects listed in Table 6.7	This objective could lead to development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

A.6 Chapter 7 – Infrastructure

The strategic aim of this chapter is to protect, improve and provide water, wastewater, surface water and flood alleviation services throughout the county, and to facilitate the provision of high quality information communication technology, broadband, telecommunication information and electricity network required to support and enhance the key aims of best place to live/invest.

Infrastructure				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
Drinking Water	INP - 1	To liaise and work in conjunction with Irish Water in the delivery of an adequate level of water services infrastructure through the Capital Investment Plan 2017-2021 and Rural Water Programme 2019-2021 and any subsequent Plans or Programmes, to ensure a sufficient capacity of water supply is available for the settlements of the County Settlement Hierarchy	This objective could lead to development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Drinking Water	INP - 2	To liaise and work in conjunction with Irish Water to promote the sustainable development of water supply and drainage infrastructure in the county and the region, in accordance with the objectives and recommendations set out in the Irish Water's Water Services Strategic Plan.	This objective could lead to development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Drinking Water	INP - 3	To liaise with Irish Water to develop and implement Water Safety Plans to protect sources of public water supply and their contributing catchment.	This objective could lead to development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Infrastructure				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
			value, such as changes in water quality and quantity, and air quality.	
Drinking Water Objectives	INO - 1	To implement the Rural Water Programme 2019-2021 and any subsequent plans.	This objective could indirectly lead to development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Drinking Water Objectives	INO - 2	Provide guidance and advice regarding the protection of water supply to private wells with the overall responsibility for protection remaining with the householder.	General statement of policy/general aspiration which will not lead to development	N/A
Drinking Water Objectives	INO - 3	To ensure that any new development connects to a public water supply or Group Water Scheme, where available. Connections to wells for individual housing units in unserved rural areas will only be considered where there is no public water main or Group Water Scheme serving the site and where it can be demonstrated that connection to the proposed well will not have significant adverse effects on water quality or water quantity in the area and can provide a potable water supply in accordance with EU Drinking Water standards.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Drinking Water Objectives	INO 4	To advance key Capital Projects as outlined in the 5-year Capital Programme.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network	Refer to Chapter 9 of this NIR for mitigation measures for any plan or

Infrastructure				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
			through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	project arising from this objective/ policy
Drinking Water Objectives	INO 5	To support and facilitate key upgrades to the Achill water supply and the provision of drinking water in the Murrisk area.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Drinking Water Objectives	INO 6	To support and facilitate upgrades to the water schemes listed in Table 7.1 below.	This objective could indirectly lead to development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Wastewater	INP - 4	To liaise and work in conjunction with Irish Water in the delivery of an adequate level of wastewater services infrastructure to ensure a sufficient capacity of wastewater supply is available for the settlements of the County Settlement Hierarchy; to upgrade capacity in settlements that have an identified capacity shortfall such as Hollymount, Louisburgh, Ballindine and Doogort;	This objective could lead to development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Infrastructure				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
			value, such as changes in water quality and quantity, and air quality.	
Wastewater	INP - 5	Collaborate with Irish Water in contributing towards compliance with the relevant provisions of the Urban Wastewater Treatment Regulations 2001 and 2004 and the Waste Water Discharge (Authorisation) Regulations 2007 as amended.	Compliance reducing pollutants into river systems would steer change in such as way as to protect international nature conservation sites from adverse effects	N/A
Wastewater	INP - 6	Encourage and support a changeover from septic tanks/private wastewater treatment plants to public collection networks wherever feasible, subject to connection agreements with Irish Water and to ensure that any future development connects to the public wastewater infrastructure where it is available.	This objective could lead to development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Wastewater Objectives	INO 7	to require development in serviced areas to connect to the public foul sewer network, where available.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Wastewater Objectives	INO 8	to require development in an unsewered area which includes a septic tank/proprietary effluent treatment unit and percolation area to be rigorously assessed in accordance with the accepted EPA Code of Practice for single houses or small communities, business, leisure centres	Policies or proposals which steer change in such as way as to protect international nature conservation sites from adverse effects	N/A

Infrastructure				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		and hotels, taking into account the cumulative effects of existing and proposed developments in the area.		
Wastewater Objectives	INO 9	To actively endeavour to upgrade capacity in settlements that have an identified capacity shortfall, such as Hollymount, Louisburgh, Ballindine and Doogort, through Irish Water's Small Towns and Villages Growth Programme or any superseding programmes.		
Waste Management	INP - 7	Support the Implementation of the Connacht Ulster Regional Waste Management Plan 2015-2021(as amended) or replacement plan with particular emphasis on reuse, recycling and disposal of residual waste in the most appropriate manner where it can be demonstrated that the development will not will not have significant adverse effects on the environment, the integrity of the Natura 2000 network, traffic safety, residential or visual amenity.	Policies or proposals which steer change in such as way as to protect international nature conservation sites from adverse effects therefore positive impacts are anticipated from the implementation of this objective/policy	N/A
Waste Management	INP - 8	To promote the sustainable management of waste generation and investment in different types of waste treatment and support a healthy environment, circular economy and society.	This objective could indirectly lead to development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Waste Management Objectives	INO 10	Promote prioritising prevention, reuse, recycling and recovery, and to sustainably manage residual waste. New developments will be expected to	General statement of policy/general aspiration which will not lead to development	N/A

Infrastructure				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		take account of the provisions of the Connacht Ulster Regional Waste Management Plan 2015-2021(as amended) and observe those elements of it that relate to waste prevention and minimisation, waste recycling facilities, and the capacity for source segregation.		
Waste Management Objectives	INO 11	To provide and support the provision of bring banks or other appropriate recycling facilities throughout the County	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Waste Management Objectives	INO 12	To continue to expand environmental awareness initiatives designed to create increased public awareness of waste prevention, minimisation, reuse and resource efficiency.	General statement of policy/general aspiration which will not lead to development	N/A
Waste Management Objectives	INO 13	To encourage community/voluntary groups to establish additional waste services or facilities (e.g. small-scale facilities for recycling, reuse, repair) in their area and assist them to develop a strategy to provide such facilities for and with members of their community.	This objective could indirectly lead to development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Waste Management Objectives	INO 14	To continue to support with local and Tidy Towns initiatives in the maintenance and conservation of	This objective could indirectly lead to development. Any development within the county could potentially have direct and/or	Refer to Chapter 9 of this NIR for mitigation measures

Infrastructure				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		our local urban and rural communities throughout the county.	indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	for any plan or project arising from this objective/ policy
Waste Management Objectives	INO 15	To seek the effective engagement of local communities in the county to promote their role in recycling waste and tackling the problem of illegal dumping within the county through liaison with the environmental Awareness Officer.	General statement of policy/general aspiration which will not lead to development	N/A
Surface Water	INP 9	To liaise and work in conjunction with Irish Water in the implementation of the Memorandum of Understanding (MOU) for surface water drainage and flood management, including the separation of foul and surface water drainage networks where feasible and undertake drainage network upgrades to help remove surface water misconnection and infiltration.	This objective could lead to development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Surface Water	INP 10	Support, in conjunction with Irish Water, the improvement of storm water infrastructure to improve sustainable drainage and reduce the risk of flooding in urban environments	This objective could lead to development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Surface Water Objectives	INO 16	Support & promote the use of green infrastructure, for example green roofs, green	This objective could lead to development of infrastructure. Any development within the	Refer to Chapter 9 of this NIR for

Infrastructure				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		walls, planting and green spaces for surface water run-off retention purposes, in the interests of flood mitigation and climate change adaptation.	county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	mitigation measures for any plan or project arising from this objective/ policy
Surface Water Objectives	INO 17	To require the use of SuDS to minimise and limit the extent of hard surfacing and paving and require the use of sustainable drainage techniques where appropriate, for the new development or for extensions to existing developments, in order to reduce the potential impact of existing and predicted flooding risks.	This objective could lead to development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Surface Water Objectives	INO 18	Ensure new development is adequately serviced with surface water drainage infrastructure, which meets the requirements of the Water Framework Directive, associated River Basin Management Plans and Catchment Flood Risk Assessment Management (CFRAM) Plans.	This objective could lead to development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Flood Risk Management	INP 11	Have regard to the Guidelines for Planning Authorities on the Planning System and Flood Risk Management (DoEHLG/OPW 2009) and Circular PL2/2014, in the preparation of plans and strategies related to development and in the assessment of projects.	General statement of policy/general aspiration which will not lead to development	N/A

Infrastructure				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
Flood Risk Management	INP 12	Support the implementation of recommendations in the Flood Risk Management Plans (FRMP's), including planned investment measures for managing and reducing flood risk.	Development of flood defences could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Flood Risk Management	INP 13	Support the implementation of recommendations in the CFRAM Programme to ensure that flood risk management policies and infrastructure are progressively implemented.	General statement of policy/general aspiration which will not lead to development	N/A
Flood Risk Management Objectives	INO 19	Ensure that a flood risk assessment is carried out for any development proposal where a flood risk is identified in accordance with the Planning System and Flood Risk Management (DoEHLG/OPW 2009) and Circular PL2/2014. This assessment shall be appropriate to the scale and nature of risk to the potential development.	General statement of policy/general aspiration which will not lead to development	N/A
Flood Risk Management Objectives	INO 20	Consult with the OPW in relation to proposed developments in the vicinity of drainage channels and rivers for which the OPW are responsible, and to retain a strip on either side of such channels where required, to facilitate maintenance access thereto.	Any works occurring in river systems could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Flood Risk Management Objectives	INO 21	Assist the OPW in developing catchment-based Flood Risk Management Plans for rivers in County Mayo and have regard to their provisions/recommendations.	Any works occurring in river systems could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance	Refer to Chapter 9 of this NIR for mitigation measures for any plan or

Infrastructure				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
			or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	project arising from this objective/ policy
Flood Risk Management Objectives	INO 22	Protect the integrity of any formal (OPW or Mayo County Council) flood risk management infrastructure, thereby ensuring that any new development does not negatively impact any existing defence infrastructure or compromise any proposed new infrastructure.	Any works occurring in river systems could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Flood Risk Management Objectives	INO 23	Ensure that where flood risk management works take place that the natural and cultural heritage, rivers, streams and watercourses are appropriately protected.	Any works occurring in river systems could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Flood Risk Management Objectives	INO 24	Consult, where necessary, with Inland Fisheries Ireland, the National Parks and Wildlife Service and other relevant agencies in the provision of flood alleviation measures in the County	General statement of policy/general aspiration which will not lead to development	N/A
Flood Risk Management Objectives	INO 25	Ensure each flood risk management activity is examined to determine actions required to embed and provide for effective climate change adaptation as set out in the OPW Climate Change Sectoral Adaptation Plan Flood Risk Management applicable at the time.	General statement of policy/general aspiration which will not lead to development	N/A

Infrastructure				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
Flood Risk Management Objectives	INO 26	To cooperate with the Office of Public works in the delivery of the Crossmolina Flood Relief scheme and other schemes that may be brought forward in the lifetime of this Plan	Any works occurring in river systems could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Flood Risk Management Objectives	INO 27	To identify and preserve vulnerable floodplains, wetlands and coastal areas to the maximum possible extent in both urban and rural areas.	Policies or proposals which protects habitats and thus protects international nature conservation sites from adverse effects	N/A
Broadband	INP 14	Support and facilitate the implementation of the National Broadband Plan and the Mayo Digital Strategy as a means of developing further opportunities for enterprise, employment, education, innovation and skills development for those who live and work in rural areas.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Broadband	INP 15	Support the delivery of high capacity Information Communications Technology Infrastructure, broadband connectivity and digital broadcasting, throughout the County, in order to ensure economic competitiveness for the enterprise and commercial sectors and in enabling more flexible work practices e.g. teleworking/homeworking.	This objective could lead to development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Infrastructure				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
Broadband Objectives	INO 28	To require all new development to provide specific ducting to enable broadband infrastructure, where appropriate.	This objective could lead to development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Broadband Objectives	INO 29	Support and facilitate the European Commission’s “Wifi4EU project” through the provision of free Wifi Hotspots at appropriate publicly accessible locations throughout the County.	This objective could lead to development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Broadband Objectives	INO 30	To identify suitable locations and support the provision of co-working facilities, digital hubs/eHubs and eWorking centres throughout the County that function as outreach hubs for employers and promote flexible working arrangements	This objective could lead to development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Telecommunications Policy	INP 16	To support the delivery of Telecommunication Infrastructure in the County having regard to the Government Guidelines ‘Telecommunications Antennae and Support Structures-Guidelines for	Policies or proposals which steer change in such as way as to protect international nature conservation sites from adverse effects	N/A

Infrastructure				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		Planning Authorities' 1996 (DoEHLG), the 'Guidance on the potential location of overground telecommunications infrastructure on public roads', (Dept of Communications, Energy & Natural Resources, 2015) and Circular Letter PL 07/12 and where it can be demonstrated that the development will not have significant adverse impacts on communities, public rights of way and on the built or natural environment including the integrity of the Natura 2000 network.		
Telecommunications Policy	INP 17	To promote Mayo as a sustainable international destination for ICT infrastructures such as data centres and associated economic activities at appropriate locations.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Telecommunications Objectives	INO 31	To maximise and widely promote connectivity of Mayo based on/building on existing ICT infrastructure and sub-sea fibre optic cables where possible.	This objective could lead to development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Telecommunications Objectives	INO 32	Encourage the location of any telecommunications structures have regard to the Landscape Appraisal of County Mayo, and where	This objective could lead to increased development. Any development within the county could potentially have direct and/or	Refer to Chapter 9 of this NIR for mitigation measures

Infrastructure				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		possible, advise on a less intrusive location in areas where they are unlikely to intrude on the setting of, or views of/from, national monuments or protected structures.	indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	for any plan or project arising from this objective/ policy
Telecommunications Objectives	INO 33	To maintain and update the council’s register of approved ducting and telecommunication structures in the County, to assist in the assessment of future telecommunication developments. The Council will encourage co-location of antennae on existing support structures and to require documentary evidence as to the non-availability of this option in proposals for new structures. The shared use of existing structures will be required where the numbers of masts located in any single area is considered to have an excessive concentration.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Telecommunications Objectives	INO 34	To work with statutory undertakers to make the most efficient use of infrastructure in the delivery of broadband in the county, particularly encouraging the use of existing telecommunications ducting where it is available.	General statement of policy/general aspiration which will not lead to development	N/A
Telecommunications Objectives	INO 35	To actively engage with telecommunication service providers to help identify, improve and/or eliminate mobile phone signal blackspots within the county, including an examination of the feasibility and suitability of council owned lands/assets to support economic development and social inclusion, in accordance with section	This objective could indirectly lead to development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Infrastructure				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		211 of the Planning and Development Act 2000 (as amended).	value, such as changes in water quality and quantity, and air quality.	
Electricity Policies	INP 18	To support the provision of high-quality electricity infrastructure and development of enhanced electricity supply, to serve the existing and future needs of the County and to facilitate new transmission infrastructure projects that may be brought forward during the lifetime of the plan including the delivery and integration, including linkages of renewable energy proposals to the electricity transmission grid in a sustainable and timely manner, whilst seeking to protect and maintain bio-diversity, wildlife habitats, scenic amenities, including protected views and nature conservation.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Electricity Policies	INP 19	To co-operate and liaise with statutory and other energy providers in relation to power generation in order to ensure adequate power capacity for the existing and future business and enterprise needs of the County.	This objective could indirectly lead to development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Electricity Policies	INP 20	Support the statutory providers of national grid infrastructure by safeguarding such strategic corridors from encroachment by other developments that might compromise the provision of energy networks where strategic route corridors have been identified;	This objective could indirectly lead to development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Infrastructure				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
			value, such as changes in water quality and quantity, and air quality.	
Electricity Objectives	INO 36	Facilitate the progression of and implement improvements to the existing electricity networks and facilitate the development of new transmission infrastructure projects in accordance with EirGrid’s Implementation Plan Strategy 2020-2025 (or any superseding strategies) that might be brought forward during the lifetime of this plan, subject to relevant Irish planning and European environmental legislation including Article 6 of the Habitats Directive and/or other environmental assessment.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Electricity Objectives	INO 37	Ensure the provision, where feasible, of electricity cables been located underground, especially in the urban environment, and generally within a reason of public open space.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Electricity Objectives	INO 38	To seek the delivery of the necessary integration of transmission network requirements to facilitate linkages of renewable energy proposals to the electricity transmission grid in a sustainable and timely manner.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Infrastructure				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
Natural Gas	INP 21	To support and facilitate the improvement and extension of the gas grid network in County Mayo to serve existing and envisaged future residential, commercial and industrial development.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Natural Gas	INP 22	To support the development of enhanced gas supplies, which do not negatively impact on environmental quality, landscape, wildlife, habitats or residential amenity and which are critical to the economic development of the County.	This objective could lead to development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Natural Gas	INP 23	To support the continued use of the Bellinaboy gas terminal as the primary hub to bring ashore any future gas reserves utilising the existing gas grid connection.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Natural Gas Objectives	INO 39	To seek the extension of the Gas Network to other towns in the County and to Ireland West Airport Knock.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network	Refer to Chapter 9 of this NIR for mitigation measures for any plan or

Infrastructure			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective				
			through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	project arising from this objective/ policy

DRAFT

A.7 Chapter 8 – Sustainable Communities

Strategic aim: To develop and support vibrant sustainable communities in Mayo where people can live, work and enjoy access to a wide range of community, health and educational facilities and amenities, suitable for all ages and needs, in both urban and rural areas, thereby supporting a high quality of life for all to enjoy.

Sustainable Communities				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
Community development policies	SCP 1	To support empowerment and capacity building in communities and support participation in community development initiatives.	General statement of policy/general aspiration which will not lead to development	N/A
Community development policies	SCP 2	Recognise and support the Mayo diaspora worldwide through the implementation of the Mayo Diaspora strategy.	General statement of policy/general aspiration which will not lead to development	N/A
Community Development Objectives	SCO 1	To identify and facilitate the development of suitable sites for community facilities within the county, particularly in newly developing areas. These sites should be easily accessible (walking and cycling) and promote the use of public transport.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/policy
Community Development Objectives	SCO 2	To support and assist communities to plan for the future of their villages and towns through a bottom up, participative and inclusive approach of the Mayo Community Futures process of community action planning.	General statement of policy/general aspiration which will not lead to development	N/A
Community Development Objectives	SCO 3	Facilitate voluntary and community groups through the ongoing development of the Mayo Public Participation Network.	General statement of policy/general aspiration which will not lead to development	N/A

Sustainable Communities				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
Community Development Objectives	SCO 4	To promote and advance the attractiveness of the towns and villages throughout the county by assisting in the enhancement of the natural and built environment, through initiatives such as Pride of Place and the coordinated efforts of Mayo County Council, tidy towns organisations, community groups and local development companies.	General statement of policy/general aspiration which will not lead to development	N/A
Community Development Objectives	SCO 5	Retain existing community facilities and public open spaces, unless a sustainable alternative can be provided in the immediate locality.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Community Facilities and Social Infrastructure Policies	SCP 3	Ensure that County Mayo is equipped with physical and other infrastructure necessary to allow communities the capacity to develop and flourish.	This objective could lead to development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Community Facilities and Social Infrastructure Policies	SCP 4	Support the provision of a modern and effective fire and rescue services for the county.	This objective could indirectly lead to development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from

Sustainable Communities				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
			habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	this objective/ policy
Community Facilities and Social Infrastructure Policies	SCP 5	Support and facilitate the development of places of worship and multi-faith facilities at appropriate locations, such as town and village centres.	This objective could lead to development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Community Facilities and Social Infrastructure Policies	SCP 6	Ensure that new social infrastructure developments are accessible and inclusive for a range of users, by adopting a universal design approach and providing for an age friendly society in which people of all ages can live full active, valued and healthy lives.	This objective could lead to development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Community Facilities and Social Infrastructure Policies	SCP 7	Protect the cultural heritage of historical burial grounds within the county and to encourage their management and maintenance in accordance with conservation principles.	General statement of policy/general aspiration which will not lead to development	N/A
Community Facilities and Social	SCP 8	Facilitate the development of new or extended burial grounds and crematoria by the zoning of land at suitable locations, where appropriate, and providing local authority burial grounds, subject to	This objective could lead to development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network	Refer to Chapter 9 of this NIR for mitigation measures for any plan or

Sustainable Communities				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
Infrastructure Policies		appropriate safeguards with regard to environmental, noise and traffic impacts.	through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	project arising from this objective/ policy
Community Facilities and Social Infrastructure Policies	SCP 9	Ensure that all buildings, public open spaces, recreational and amenity areas are accessible for people with disabilities.	General statement of policy/general aspiration which will not lead to development	N/A
Community Facilities and Social Infrastructure Policies	SCP 10	Direct residential care homes for older people, retirement homes, nursing homes, independent living units, assisted living units, retirement villages and sheltered accommodation are located within urban and rural towns and villages, ensuring that they are appropriate in scale to the size of the settlement.	General statement of policy/general aspiration which will not lead to development	N/A
Community Facilities and Social Infrastructure Objectives	SCO 6	To work with public service delivery bodies to improve the level of accessibility and facilitate the delivery of more effective and integrated services in communities.	General statement of policy/general aspiration which will not lead to development	N/A
Community Facilities and Social Infrastructure Objectives	SCO 7	To support and assist the provision of a broad range of community facilities within settlements or in close proximity to existing facilities or multi-purpose centres or public transport routes to meet the changing needs of all sectors of the community.	This objective could lead to development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Sustainable Communities				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
Community Facilities and Social Infrastructure Objectives	SCO 8	Support and promote the development of an inter-county coastal path linking Mayo, Galway and Sligo, subject to no significant adverse impacts on the environment including the integrity of any Natura 2000 site.	Policies or proposals which steer change in such as way as to protect international nature conservation sites from adverse effects therefore positive impacts are anticipated from the implementation of this objective/policy	N/A
Social Inclusion Policies	SCP 11	Support initiatives that foster social inclusion amongst groups that are vulnerable to poverty and exclusion.	General statement of policy/general aspiration which will not lead to development	N/A
Social Inclusion Policies	SCP 12	Support the implementation of the Mayo Local Economic and Community Plan in collaboration with the Local and Economic Development Committee to reduce the number of people in or at risk of social exclusion.	General statement of policy/general aspiration which will not lead to development	N/A
Social Inclusion Policies	SCP 13	Support and promote a more diverse and socially inclusive society that targets equality of opportunity and a better quality of life for all citizens, through improved integration and greater accessibility in the delivery of sustainable communities and the provision of associated services.	General statement of policy/general aspiration which will not lead to development	N/A
Social Inclusion Policies	SCP 14	To support the implementation of the Mayo Age Friendly County Programme 2016-2020 as implemented by the Mayo Age Friendly Alliance (and any updated editions) and any key actions in relation to the physical environment.	General statement of policy/general aspiration which will not lead to development	N/A
Social Inclusion Policies	SCP 15	Promote Universal Design and well-designed lifetime, adaptable and age friendly housing in accordance with best practice and with the policies and principles contained in Building for Everyone: A Universal Design Approach (National Disability Authority, 2012) and Sustainable Residential	This objective could lead to development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from

Sustainable Communities				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		Development in Urban Area: Guidelines for Planning Authorities and its companion document Urban Design Manual (DEHLG, 2009) and particularly on infill and brownfield sites, walkable to existing services and facilities.	changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	this objective/ policy
Social Infrastructure Objectives	SCO 9	Facilitate the provision of community and resource centres and youth clubs/ cafes, recreational amenities and other facilities for younger people by the identification of suitably located sites, including Council landbanks and by assisting in the provision of finance, where possible.	This objective could lead to development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Social Infrastructure Objectives	SCO 10	Secure the implementation of the Council's Traveller Accommodation Programme (2019-2024) and to review this programme if required and/or deemed to be necessary, during the Plan period.	This objective could lead to development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Health and Wellbeing Policies	SCP 16	To promote the improvement of health and wellbeing services throughout the county, in particular to encourage the integration of health and wellbeing services and facilities with new and existing community facilities, where feasible.	General statement of policy/general aspiration which will not lead to development	N/A
Health and Wellbeing Policies	SCP 17	To look favourably upon the development of primary care centre, clinics and facilities for the specific needs of an ageing population, in	This objective could indirectly lead to development of infrastructure. Any development within the county could potentially have direct	Refer to Chapter 9 of this NIR for mitigation measures

Sustainable Communities				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		accessible locations, particularly town centres, which comprise / provide access to a range of healthcare services.	and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	for any plan or project arising from this objective/ policy
Health and Wellbeing Policies	SCP 18	To support the Health Service Executive and other statutory and voluntary agencies in the provision of appropriate healthcare facilities, including the development of both the system of hospital care and the provision of community-based primary care facilities.	This objective could indirectly lead to development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Health and Wellbeing Policies	SCP 19	To support the key priorities of the Mayo Healthy and Wellbeing Plan 2018-2022 (as updated in the future) to promote health and well-being.	General statement of policy/general aspiration which will not lead to development	N/A
Health and Wellbeing Policies	SCP 20	To support the key priorities and actions of the Mayo Children and Young People's Plan 2018-2020 (as updated in the future) to secure better outcomes for children and young people through more effective integration of existing services and interventions at local level.	General statement of policy/general aspiration which will not lead to development	N/A
Health and Wellbeing Policies	SCP 21	To support health and wellbeing initiatives and healthcare provision so that Mayo is a healthy and caring county for all.	General statement of policy/general aspiration which will not lead to development	N/A
Health and Wellbeing Objective	SCO 11	Support the provision of improved health services on suitably zoned lands in the Tier 1 and 2 towns in the settlement hierarchy, on lands in the town centres or immediately adjacent to town centres	This objective could indirectly lead to development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000	Refer to Chapter 9 of this NIR for mitigation measures for any plan or

Sustainable Communities				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		(based on the sequential approach) which are appropriately serviced (water service, footpaths, lighting etc.)	network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	project arising from this objective/ policy
Recreation Facilities and Provision Policy	SCP 22	To promote the improvement of health and wellbeing services throughout the county, in particular to encourage the integration of health and wellbeing services and facilities with new and existing community facilities, where feasible.	General statement of policy/general aspiration which will not lead to development	N/A
Recreation Facilities and Provision Objectives	SCO 12	Support and facilitate the implementation of the recommendations of Mayo County Council Play and Recreation Strategy or any subsequent strategy, in conjunction with all relevant agencies.	This objective could indirectly lead to development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Recreation Facilities and Provision Objectives	SCO 13	To support and facilitate the provision for the development of an indoor all-weather multi-sport dome for Ballina to serve clubs and organisations in North Mayo/ West Sligo.	This objective could lead to development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Education, Training and Skills Policies	SCP 23	To support initiatives which provide opportunities for people in Mayo to access appropriate	General statement of policy/general aspiration which will not lead to development	N/A

Sustainable Communities				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		education and training provision necessary to allow them to realise their full potential.		
Education, Training and Skills Policies	SCP 24	Ensure the provision of new educational facilities are located within existing settlements.	General statement of policy/general aspiration which will not lead to development	N/A
Education, Training and Skills Objectives	SCO 14	Support and facilitate the establishment of co-working/ remote working hubs and creative hubs as either standalone facilities themselves or ancillary to public buildings, libraries and community centres or in towns and village centres, as appropriate.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Education, Training and Skills Objectives	SCO 15	Support and promote the growth, development and success of the Connacht Ulster Alliance consortium's (including GMIT Mayo) ambition towards becoming a Technological University serving the west and northwest.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Education, Training and Skills Objectives	SCO 16	Support the provision of childcare facilities and new and refurbished schools on well located sites, within or close to existing built-up areas, that meet the diverse needs of local populations.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value,	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Sustainable Communities				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
			such as changes in water quality and quantity, and air quality.	
Education, Training and Skills Objectives	SCO 17	Assist the County Childcare Committee in identifying priority areas within the county for the provision of childcare facilities.	General statement of policy/general aspiration which will not lead to development	N/A
Arts and Libraries Policies	SCP 25	To implement the Creative Ireland Programme 2017-22, Council's Arts Strategy 2018-2022 and the Mayo Culture and Creative Strategy 2018-2022 or any subsequent strategy over the lifetime of the plan.	General statement of policy/general aspiration which will not lead to development	N/A
Arts and Libraries Policies	SCP 26	To recognise and project Mayo's unique identity in an appropriate manner and to promote and market the county to derive optimum social, cultural and economic benefits.	This objective could indirectly lead to development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/policy
Arts and Libraries Policies	SCP 27	Continue to promote the use of the library service and further develop each library as a community gathering place and learning hub.	General statement of policy/general aspiration which will not lead to development	N/A
Arts and Libraries Objective	SCO 18	To implement the Mayo County Library Development Plan 2015-2020 (as extended) and any subsequent Library Development Plans.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value,	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/policy

Sustainable Communities				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
			such as changes in water quality and quantity, and air quality.	
Arts and Libraries Objective	SCO 19	To support the implementation of the Mayo County Council Strategic Arts Plan 2018-2022 (and any superseding version) and to ensure comprehensive provision of arts infrastructure incorporating spaces suitable for the arts and a variety of arts practices throughout the County.	General statement of policy/general aspiration which will not lead to development	N/A
Islands and Gaeltacht Policies	SCP 28	To support initiatives at promoting Gaeilge, the Irish language, and the need for access to social supports in terms of language, education and employment for refugees, asylum seekers and migrants.	General statement of policy/general aspiration which will not lead to development	N/A
Islands and Gaeltacht Policies	SCP 29	To promote and protect Mayo's heritage and culture and the advancement of the Irish Language.	General statement of policy/general aspiration which will not lead to development	N/A
Islands and Gaeltacht Policies	SCP 30	Support the inhabited islands in County Mayo and recognise the special planning and development needs of islands and islands communities, particularly access, infrastructure and services.	This objective could indirectly lead to development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/policy
Islands and Gaeltacht Objectives	SCO 20	To ensure that the Irish language and our unique linguistic heritage becomes more visible and audible and integrated into all activities in Mayo.	General statement of policy/general aspiration which will not lead to development	N/A
Islands and Gaeltacht Objectives	SCO 21	To strengthen the unique linguistic and cultural heritage of Gaeltacht Mhaigh Éo, by supporting and facilitating improved physical, social and	This objective could indirectly lead to development of infrastructure. Any development within the county could potentially have direct	Refer to Chapter 9 of this NIR for mitigation measures

Sustainable Communities				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective		Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red		Mitigation
		economic infrastructure at appropriate locations throughout the Gaeltacht areas and require a Language Impact Statement in respect of significant developments within or close to Gaeltacht areas, or where deemed appropriate, to determine their impact on the usage of Irish as the community language.	and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	for any plan or project arising from this objective/ policy
Islands and Gaeltacht Objectives	SCO 22	To maintain and support the communities of Mayo's inhabited islands by facilitating sustainable social and economic development through the provision of, or facilitating the provision of, infrastructure (social and physical), housing and services, having regard to the traditional building patterns of the Islands and the need to protect the cultural and natural heritage of the Islands.	This objective could indirectly lead to development of infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

A.8 Chapter 9 – Built Environment

Strategic aim: To recognise the importance of identifying, valuing and safeguarding our archaeological, architectural and cultural heritage for future generations and aim to do so by means of proper management, sensitive enhancement and/or appropriate development of this resource. To also protect and enhance the unique identity and character of Mayo's towns and villages and improve quality of life and well-being through the application of Healthy Placemaking, underpinned by good urban design, by the creation of attractive public spaces that are vibrant, distinctive, safe and accessible and which promote and facilitate positive social interaction.

Built environment				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
Archaeological Heritage Policies	BEP 1	To support and promote the protection, appropriate management and sympathetic enhancement of the county's archaeological heritage within the Plan area, in particular by implementing the Planning and Development Act 2000 (as amended) and the National Monuments Act 1930 (as amended).	General statement of policy/general aspiration which will not lead to development	N/A
Archaeological Heritage Policies	BEP 2	To protect archaeological sites, monuments, underwater archaeology and archaeological objects in their setting, which are listed on the Record of Monuments and Places for Mayo.	General statement of policy/general aspiration which will not lead to development	N/A
Archaeological Heritage Policies	BEP 3	To protect all sites and features of archaeological interest discovered subsequent to the publication of the Record of Monument and Places, in situ (or at a minimum preservation by record) , having regard to the advice and recommendations of the National Monuments Service of the Department of Culture, Heritage and the Gaeltacht.	General statement of policy/general aspiration which will not lead to development	N/A
Archaeological Heritage Objectives	BEO 1	Protect the archaeological heritage and sites identified in the Record of Monuments and Places, National Monuments in the ownership or guardianship of the State in addition to	General statement of policy/general aspiration which will not lead to development	N/A

Built environment				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		National Monuments that are the subject of Preservation Orders, and to safeguard the integrity of the archaeological sites in their setting.		
Archaeological Heritage Objectives	BEO 2	Protect the tentative World Heritage Site in Mayo on the UNESCO Tentative List - Ireland 2010, The Céide Fields, from inappropriate development and support its nomination to World Heritage Status.	General statement of policy/general aspiration which will not lead to development	N/A
Archaeological Heritage Objectives	BEO 3	To implement, in partnership with the County Mayo Heritage Forum, relevant stakeholders and the community, the County Mayo Heritage Plan and any revisions thereof.	General statement of policy/general aspiration which will not lead to development	N/A
Archaeological Heritage Objectives	BEO 4	To ensure that development in the vicinity of a Recorded Monument or Zone of Archaeological Potential is sited and designed in a sensitive manner, with a view to minimal detracting from the monument or its setting.	General statement of policy/general aspiration which will not lead to development	N/A
Archaeological Heritage Objectives	BEO 5	To encourage the management and maintenance of the county's archaeological heritage, including historic burial grounds, in accordance with best conservation practice that considers the impact of climate change.	General statement of policy/general aspiration which will not lead to development	N/A
Archaeological Heritage Objectives	BEO 6	To promote awareness of and encourage the provision of access to, the archaeological resources of the county.	General statement of policy/general aspiration which will not lead to development	N/A
Architectural Heritage Policies	BEP 4	To protect the architectural heritage of County Mayo which is a unique and special resource.	General statement of policy/general aspiration which will not lead to development	N/A

Built environment				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
Architectural Heritage Policies	BEP 5	To promote best conservation practice and encourage the use of appropriately qualified professional advisors, tradesmen and craftsmen with recognised conservation expertise, for works to protected structures or historic buildings in an Architectural Conservation Area.	General statement of policy/general aspiration which will not lead to development	N/A
Architectural Heritage Policies	BEP 6	To encourage the conservation of Protected Structures, and where appropriate, the adaptive re-use of existing buildings and sites in a manner compatible with their character and significance.	This objective could lead to development/retention of abandoned or derelict building which could contain Annex II species. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Architectural Heritage Policies	BEP 7	Protect buildings and structures included in the Record of Protected Structures (RPS) which forms part of this Plan.	General statement of policy/general aspiration which will not lead to development	N/A
Architectural Heritage Objectives	BEO 7	To review and update the Record of Protected Structures on an on-going basis and to make additions and deletions, as appropriate.	General statement of policy/general aspiration which will not lead to development	N/A
Architectural Heritage Objectives	BEO 8	To ensure the protection and sympathetic enhancement of buildings and structures included and proposed for inclusion in the Record of Protected Structures (RPS) that are of special architectural, historical, archaeological, artistic, cultural, scientific, social or technical interest, together with the integrity of their character and setting.	General statement of policy/general aspiration which will not lead to development	N/A

Built environment				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
Architectural Heritage Objectives	BEO 9	To protect the setting of protected structures and seek to prevent the demolition or inappropriate alteration of Protected Structures, which would adversely impact on the character and special interest of the structure, where appropriate.	General statement of policy/general aspiration which will not lead to development	N/A
Architectural Heritage Objectives	BEO 10	Ensure that any new development or alteration to a building within or adjoining an Architectural Conservation Area positively enhances the character of the area and is appropriate in terms of the proposed materials, scale, density, layout, proportions, plot ratio and building lines.	General statement of policy/general aspiration which will not lead to development	N/A
Architectural Heritage Objectives	BEO 11	Identify places of special architectural, historical, archaeological, artistic, cultural, scientific, social or technical interest, and to define them as Architectural Conservation Areas and to undertake an assessment to inform the potential ACA designation for the following areas:- Ballinrobe, Killala, Pontoon and Dugort.	Policies or proposals which steer change in such as way as to protect international nature conservation sites from adverse effects therefore positive impacts are anticipated from the implementation of this objective/policy	N/A
Historic Building Stock and Vernacular Architecture Policies	BEP 8	To encourage the retention, sympathetic maintenance and sustainable re-use of historic buildings, including vernacular dwellings or farm buildings and the retention of historic streetscape character, fabric, detail and features, where appropriate.	This objective could lead to development/retention of abandoned or derelict building which could contain Annex II species. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Built environment				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
			value, such as changes in water quality and quantity, and air quality.	
Historic Building Stock and Vernacular Architecture Policies	BEP 9	To actively promote the retention and restoration of thatched dwellings as a key component of the built heritage of the county.	This objective could lead to development/retention of abandoned or derelict building which could contain Annex II species. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Historic Building Stock and Vernacular Architecture Policies	BEP 10	To encourage the protection, retention, appreciation and appropriate revitalisation of the vernacular heritage of Mayo.	General statement of policy/general aspiration which will not lead to development	N/A
Historic Building Stock and Vernacular Architecture Policies	BEP 11	To promote the sympathetic maintenance refurbishment and re-use of vernacular built heritage and to support the retention of original fabric such as windows, doors, renders/pub/shop-fronts, roof coverings and interiors.	General statement of policy/general aspiration which will not lead to development	N/A
Historic Building Stock and Vernacular Architecture Policies	BEP 12	To support proposals to appropriately refurbish and extend vernacular structures in a semi-derelict or derelict condition.	This objective could lead to development/retention of abandoned or derelict building which could contain Annex II species. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project

Built environment				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
			through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	arising from this objective/ policy
Historic Building Stock and Vernacular Architecture Policies	BEP 13	To encourage the protection, conservation, promotion and enhancement of Country Houses, Gardens and Demesnes in the county and support public awareness, enjoyment of and access to these sites, where appropriate.	This objective could lead to development/retention of abandoned or derelict building which could contain Annex II species. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Historic Building Stock and Vernacular Architecture Policies	BEP 14	To discourage development that would lead to a loss of, or cause damage to, the character, the principle components of, or the setting of Country Houses, Gardens and Demesnes.	General statement of policy/general aspiration which will not lead to development	N/A
Historic Building Stock and Vernacular Architecture Objectives	BEO 12	To identify and retain good examples of vernacular architecture and historic street furniture in situ, for example, cast-iron post boxes, water pumps, signage, street lighting, kerbing and traditional road and street surface coverings.	General statement of policy/general aspiration which will not lead to development	N/A
Historic Building Stock and	BEO 13	To ensure that conversions or extensions of traditional buildings or the provision of new adjoining buildings, are sensitively designed and	This objective could lead to development/retention of abandoned or derelict building which could contain Annex	Refer to Chapter 9 of this NIR for mitigation

Built environment				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
Vernacular Architecture Objectives		do not detract from the character of the historic building.	II species. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	measures for any plan or project arising from this objective/ policy
Historic Building Stock and Vernacular Architecture Objectives	BEO 14	To update the survey of surviving thatched structures in the county and to promote available grant schemes in order to assist owners with their retention and repair.	This objective could lead to development/retention of abandoned or derelict building which could contain Annex II species. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Historic Building Stock and Vernacular Architecture Policies	BEO 15	To preserve the character and setting (for example, gates, gate piers and courtyards) of historic building and vernacular buildings, where deemed appropriate by the planning authority.	General statement of policy/general aspiration which will not lead to development	N/A
Sustainable Buildings and Structures Policies	BEP 15	Support and promote retaining built urban fabric/structures in towns and villages, in the interest of sustainable development in the national and global context of locking in carbon.	General statement of policy/general aspiration which will not lead to development	N/A
Sustainable Buildings and	BEP 16	Promote and support the re-use and re-purposing of extant building stock, in the first	This objective could lead to development/retention of abandoned or	Refer to Chapter 9 of this NIR for

Built environment				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
Structures Policies		instance, over demolition and rebuilding building, where practical, with reference to the loss of our historic building stock, sense of place and the environmental cost.	derelict building which could contain Annex II species. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	mitigation measures for any plan or project arising from this objective/ policy
Sustainable Buildings and Structures Policies	BEP 17	Support and promote the development and use of passive solar design principles in all new developments taking account of national guidelines.	This objective could lead to development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Sustainable Buildings and Structures Policies	BEP 18	Encourage the use of green roofs and green walls particularly on apartment, industrial, commercial, leisure and educational buildings.	This objective could lead to development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Sustainable Buildings and Structures Policies	BEP 19	Encourage improved energy efficiency of existing building stock and promote energy efficiency and conservation in the design and	This objective could lead to development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network	Refer to Chapter 9 of this NIR for mitigation measures for any

Built environment				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective		Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation	
		development of all new buildings, including local authority dwellings.	through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	plan or project arising from this objective/ policy
Sustainable Buildings and Structures Objectives	BEO 16	To seek to reduce reliance on fossil fuels in the county by reducing the energy demand of existing buildings, in particular residential dwellings.	General statement of policy/general aspiration which will not lead to development	N/A
Sustainable Buildings and Structures Objectives	BEO 17	To require, where feasible and practicable, the provision of green roof technology for all new public buildings (Council buildings, school buildings, hospitals, community centres, sports facilities, libraries, Garda stations etc.) to assist in flood alleviation, insulation and improved biodiversity and to actively promote these measures where appropriate in new commercial and industrial buildings.	This objective could lead to development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Sustainable Buildings and Structures Objectives	BEO 18	To seek to improve the energy efficiency of the county's existing building stock in line with good architectural conservation practice and to promote energy efficiency and conservation in the design and development of all new buildings in the county, in accordance with the Building Regulations Part L (Conservation of Fuel and Energy).	General statement of policy/general aspiration which will not lead to development	N/A
Sustainable Buildings and Structures Objectives	BEO 19	To consider the adaptability of buildings over time and seek to improve the efficiency of existing building stock, promote energy efficiency, high levels of energy conservation and the use of renewable energy sources in the	General statement of policy/general aspiration which will not lead to development	N/A

Built environment				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		design and development of all new buildings in the county.		
Sustainable Buildings and Structures Objectives	BEO 20	Inform and encourage new developments to mitigate against, and adapt to, where possible the impacts of climate change through the location, layout and design of the development.	General statement of policy/general aspiration which will not lead to development	N/A
Sustainable Buildings and Structures Objectives	BEO 21	To ensure that new development proposals maximise energy efficiency through siting, layout, design and incorporate best practice in energy technologies, conservation and smart technology.	General statement of policy/general aspiration which will not lead to development	N/A
Placemaking Policies	BEP 20	Provide for a high-quality public realm and public spaces by promoting quality urban design that accommodates creative patterns of use, having regard to the physical, cultural, and social identities of individual settlements.	This objective could lead to development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Placemaking Policies	BEP 21	To encourage the continued vitality and viability of town and village centres by promoting ongoing environmental improvements to the public realm, including blue and green infrastructure measures.	This objective could lead to development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Placemaking Policies	BEP 22	To encourage high quality and well-designed buildings, structures, public spaces and streets	This objective could lead to development. Any development within the county could	Refer to Chapter 9 of this NIR for

Built environment				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		and support and promote healthy place-making and quality of life.	potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	mitigation measures for any plan or project arising from this objective/ policy
Placemaking Policies	BEP 23	To encourage and facilitate improvements to the physical fabric and environment of town and village centres, including streetscape, street furniture, landscaping (hard and soft), signage and wirescape, while recognising that both private and public developments can contribute to effective public realm.	This objective could lead to development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Placemaking Policies	BEP 24	To be flexible in terms of enabling brownfield / infill development within settlements, focusing on design-led and performance-based outcomes, rather than specifying absolute requirements in all cases.	This objective could lead to development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Placemaking Policies	BEP 25	To support the consolidation and aggregation of land where required in order to enable regeneration and proper planning and sustainable development.	This objective could lead to development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project

Built environment				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
			changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	arising from this objective/ policy
Placemaking Policies	BEO 22	To apply the following key attributes when considering public realm and public space enhancements: Accessible - connected and linked permeable spaces to ensure ease of movement. Functional - safe, adaptable and social environments to attract and foster activity. Attractive - visually pleasing spaces with high quality design, materials and installations (lighting, furniture and signage) based on a singular common design theme. Distinctive - reference to local context and building on the character and identity of place. Where appropriate, recreational considerations and access to blue and greens space should be underpinned by the Green Space Principles[1] including: <ul style="list-style-type: none"> Enhance urban greening through planting strategies that mitigate noise and air pollution and maximise local biodiversity gain and facilitate sustainable drainage (e.g. deciduous wooded and wildflower meadow areas). A networked approach: emphasising green infrastructure networks (rather than isolated parks) can provide new opportunities for connecting existing and new green spaces and creating linkages between urban and rural areas. Examples include greenways and linear parks, local greenways or cycleways that link to 	This objective could lead to development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Built environment				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective		Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation	
		<p>regional and national greenways and de-culverting watercourses to provide new blue corridors.</p> <ul style="list-style-type: none"> Well managed and maintained, creating a high-quality environment: poorly managed spaces or vandalism lead to negative perceptions among potential users. Multifunctional uses: examples include spaces that encourage active mobility, physical activity and sports, relaxation and tranquillity, and opportunities for social exchange (e.g. that incorporate community gardens or encourage park runs). Create multisensory restorative environments that help mitigate the psychological stresses of modern living through the provision of “restive places for rejuvenation”. 		
Placemaking Objectives	BEO 23	<p>To facilitate, promote and encourage the development of Town Centre Consolidation Sites and Opportunity Sites, identified in Chapter 12 (Settlement, or any such regeneration sites in the Local Area Plans Ballina, Castlebar and Westport for appropriate development, that contributes positively to the character of the settlement. Any proposal brought forward on Opportunity Sites shall be in accordance with the Development Principles for Town Centre Consolidation Sites/Opportunity Sites should include of an urban design statement, site brief/masterplan and shall demonstrate the rationale for the</p>	<p>This objective could lead to development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.</p>	<p>Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy</p>

Built environment				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		proposal and how it will interact within its context and the wider urban area.		
Placemaking Objectives	BEO 24	Require that all new developments, including public open spaces, cater for disability needs by way of appropriate design of the built environment.	This objective could lead to development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Placemaking Objectives	BEO 25	To make a Place-Making Strategy for towns and implement Town Centre Renewal Plans.	General statement of policy/general aspiration which will not lead to development	N/A
Placemaking Objectives	BEO 26	To support the preparation of Design Guidelines to provide for improvements in the appearance of streetscapes.	This objective could lead to development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Placemaking Objectives	BEO 27	To support the revitalisation of vacant spaces for example with cost effective, temporary uses that build on the longer-term vision for space.	This objective could lead to development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Built environment				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
			value, such as changes in water quality and quantity, and air quality.	
Placemaking Objectives	BEO 28	Require proposals for public realm enhancements to include inclusive universal design principles.	General statement of policy/general aspiration which will not lead to development	N/A
Placemaking Objectives	BEO 29	Seek funding to support the preparation of site-specific Public Realm Strategies to enhance the unique characteristics and assets of Mayo's towns and villages.	This objective could lead to development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Placemaking Objectives	BEO 30	Prepare and deliver Place-making Strategies for Castlebar and Ballina.	General statement of policy/general aspiration which will not lead to development	N/A
Placemaking Objectives	BEO 31	Prepare a Placemaking Strategy for the Tier 2 towns of Ballinrobe, Ballyhaunis, Belmullet, Claremorris and Swinford.	General statement of policy/general aspiration which will not lead to development	N/A
Regeneration Policies	BEP 26	Promote the regeneration of settlements by making better use of underutilised land and buildings, particularly within the existing built-up areas to achieve compact growth.	This objective could lead to development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Built environment				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
Regeneration Policies	BEP 27	To support initiatives that promote the reuse, refurbishment and retrofitting of existing buildings within town and village centres.	This objective could lead to development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Regeneration Policies	1111BEP 28	To apply for funding under various funding streams to facilitate the enhancement, revitalisation, renewal and regeneration of communities and town/village centres, and the delivery of innovative and transformational regeneration proposals, for example, under the Urban and Rural Regeneration and Development Funds and Town and Village Renewal Schemes.	This objective could lead to development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Regeneration Policies	BEP 29	Promote the consolidation of town and village centres with a focus on the regeneration of underused buildings and strategic sites and on the establishment of a mix of uses to encourage greater vibrancy outside of business hours.	This objective could lead to development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Regeneration Policies	BEP 30	Promote regeneration and revitalisation of small towns and villages and support local enterprise and employment opportunities to	This objective could lead to development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network	Refer to Chapter 9 of this NIR for mitigation measures for any

Built environment				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		ensure their viability as service centres for their surrounding rural areas.	through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	plan or project arising from this objective/ policy
Regeneration Objectives	BEO 32	Require all development proposals for strategic opportunity, brownfield and infill sites to be accompanied by a site brief and/or masterplan that sets out a phased programme for the regeneration of the site and demonstrates how the proposal will comply with national guidelines that seek to integrate principles of good urban design and placemaking	General statement of policy/general aspiration which will not lead to development	N/A
Regeneration Objectives	BEO 33	Support the regeneration of rural towns and villages through identification of regeneration projects for rural towns, villages and rural areas and promoting the utilisation of investment opportunities such as the Rural Regeneration and Development Fund.	This objective could lead to development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Regeneration Objectives	BEO 34	Support the viability of small towns and villages, through sustainable targeted measures that address vacant premises and deliver sustainable reuse and regeneration outcomes targeted in core areas.	This objective could lead to development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Built environment				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
Regeneration Objectives	BEO 35	Use specific powers, such as the Vacant Sites register to address issues of vacancy and underutilisation of strategic lands in town centres, including the implementation of the Vacant Sites Levy in accordance with the Urban Regeneration and Housing Act 2015.	This objective could lead to development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Regeneration Objectives	BEO 36	To establish a database of strategic brownfield and infill sites so that brownfield land re-use can be managed and co-ordinated across multiple stakeholders, as part of an active land management process.	This objective could lead to development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Regeneration Objectives	BEO 37	To identify derelict sites and vacant sites which are suitable for redevelopment and to maintain the respective registers	This objective could lead to development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

A.9 Chapter 10 – Natural Environment

Strategic aim: to continue to protect and enhance Mayo’s natural heritage and biodiversity; ensure that networks of green infrastructure are identified, created, protected and enhanced to provide a wide range of environmental, social and economic benefits to communities; improve the knowledge and understanding of the county’s landscape and coast; enhance the overall characteristics, qualities and diversity of landscape character, its sense of place and local distinctiveness in recognition of the amenity potential of the county.

Natural Environment				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
Biodiversity, Designated and Non-designated sites	NEP - 1	To support the protection, conservation and enhancement of natural heritage of County Mayo, including the protection of the integrity of European sites, that form part of the Natura 2000 Network, the protection of Natural Heritage Areas, proposed Natural Heritage Areas Ramsar Sites, Nature Reserves and Wild Fowl Sanctuaries (and other designated sites including any future designations).	Policies or proposals which steer change in such as way as to protect international nature conservation sites from adverse effects	N/A
Biodiversity, Designated and Non-designated sites	NEP - 2	To support the implementation of the National Biodiversity Action Plan 2017-2021, the National Pollination Plan 2015-2020 and County Mayo Biodiversity Plan 2015-2020 and any future editions, in partnership with relevant stakeholders, subject to available resources.	Policies or proposals which steer change in such as way as to protect international nature conservation sites from adverse effects	N/A
Biodiversity, Designated and Non-designated sites	NEP - 3	To support the implementation, in partnership with the County Mayo Heritage Forum, relevant stakeholders and the community, of the objectives and associated actions the County Mayo Heritage Plan and future editions thereof, which relate to the remit and functions of Mayo County Council.	General statement of policy/general aspiration which will not lead to development	N/A

Natural Environment				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
Biodiversity, Designated and Non-designated sites	NEP - 4	To conserve and enhance the county's biodiversity and ecological connectivity identified areas of local biodiversity importance (Local Biodiversity Areas) in the town and villages in Mayo.	Protection of native habitats will positively impact on Natura 2000 Sites	N/A
Biodiversity, Designated and Non-designated sites	NEP - 5	To promote and support increased public participation in biodiversity conservation by supporting and encouraging community-led initiatives.	General statement of policy/general aspiration which will not lead to development	N/A
Biodiversity, Designated and Non-designated sites	NEP - 6	To support the maintenance of geological and geomorphological heritage values of County Geological Sites and through consultation with the Geological Survey of Ireland and seek to promote access to such sites where possible.	This Policy could promote development for tourism. Any developments can potentially have direct and/or indirect impact on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality. Therefore, mitigation measures will be required.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Biodiversity, Designated and Non-designated sites	NEP - 7	To encourage the effective management of native and semi-natural woodlands, groups of trees and individual trees in the discharge of development management functions.	Protection of native habitats will positively impact on Natura 2000 Sites	N/A
Biodiversity, Designated and Non-designated sites Objectives	NEO - 1	To review the County Mayo Heritage Plan and County Mayo Biodiversity Plan as appropriate.	Policies or proposals which steer change in such as way as to protect international nature conservation sites from adverse effects	N/A

Natural Environment				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
Biodiversity, Designated and Non-designated sites Objectives	NEO - 2	To create a Wetland Database for County Mayo of known and potential wetland sites in the county and to develop a corresponding GIS dataset as an important tool for future biodiversity and natural heritage conservation planning in the county.	Protection of native habitats will positively impact on Natura 2000 Sites	N/A
Biodiversity, Designated and Non-designated sites Objectives	NEO - 3	To ensure the unique ecological, scenic, recreational and environmental character of Wild Nephin Ballycroy National Park is protected and enhanced, and developed appropriately.	This Policy could promote development for tourism. Any developments can potentially have direct and/or indirect impact on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality. Therefore, mitigation measures will be required.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Biodiversity, Designated and Non-designated sites Objectives	NEO - 4	To protect and enhance biodiversity and ecological connectivity in County Mayo, including woodlands, trees, hedgerows, semi-natural grasslands, rivers, streams, natural springs, wetlands, stone - walls, geological and geo-morphological systems, other landscape features and associated wildlife where these form part of the ecological network.	Protection of native habitats will positively impact on Natura 2000 Sites	N/A
Biodiversity, Designated and Non-designated sites Objectives	NEO - 5	To actively increase awareness of the importance of the natural heritage of the County and to promote education, knowledge and pride in our natural heritage.	Protection of native habitats will positively impact on Natura 2000 Sites	N/A

Natural Environment				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
Biodiversity, Designated and Non-designated sites Objectives	NEO - 6	To protect surface waters, aquatic and wetland habitats and freshwater and water- dependent species through the implementation of all appropriate and relevant Directives and transposed legislation and seek to protect and conserve the quality, character and features of inland waterways by controlling developments close to navigable and non-navigable waterways.	Protection of native habitats will positively impact on Natura 2000 Sites	N/A
Biodiversity, Designated and Non-designated sites Objectives	NEO - 7	To seek the protection, where possible the riparian zones of watercourses throughout the County, recognising the benefits they provide in relation to flood risk management and their protection of the ecological integrity of watercourse systems.	Protection of native habitats will positively impact on Natura 2000 Sites	N/A
Biodiversity, Designated and Non-designated sites Objectives	NEO - 8	To maintain, protect and where possible enhance bogs, fens and turloughs, where appropriate, in County Mayo.	Protection of native habitats will positively impact on Natura 2000 Sites	N/A
Biodiversity, Designated and Non-designated sites Objectives	NEO - 9	Recognise the importance of woodlands, tree lines hedgerows, stone - walls, watercourses and associated riparian vegetation to support bat populations and where possible developments will be encouraged to retain such features.	Protection of native habitats will positively impact on Natura 2000 Sites	N/A
Biodiversity, Designated and Non-designated	NEO - 10	to install nest boxes in all new and existing Municipal buildings, as appropriate.	Protection of native habitats will positively impact on Natura 2000 Sites	N/A

Natural Environment				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
sites Objectives				
Biodiversity, Designated and Non-designated sites Objectives	NEO - 11	To have regard to the policies and guidance of National Parks and Wildlife Service of the DoEHLG in respect of proposed development where it is possible that such development may impact on a designated European or national site or a site proposed for designation.	Policies or proposals which steer change in such as way as to protect international nature conservation sites from adverse effects	N/A
Biodiversity, Designated and Non-designated sites Objectives	NEO - 12	To support the Joyce Country and Western Lakes Geopark Project aim of establishing a NEP - w UNEP - SCO Global Geopark in the South Mayo and North Connemara area of Galway.	This Policy could promote development for tourism. Any developments can potentially have direct and/or indirect impact on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality. Therefore, mitigation measures will be required.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Biodiversity, Designated and Non-designated sites Objectives	NEO - 13	To ensure the protection of trees or groups of trees protected under Tree Preservation Orders, as well as recognise the value and encourage the retention and management of other trees and woodlands which make a valuable contribution to the character of the landscape, a settlement or its setting.	Protection of native habitats will positively impact on Natura 2000 Sites	N/A
Invasive Species	NEP - 8	To support measures for the prevention and/or eradication of invasive species as appropriate within the county	Although this Policy/objective will likely lead to a sustainable and positive impact, this policy/objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Natural Environment				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
			network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	
Invasive Species Objectives	NEO - 14	To ensure that where the presence of invasive species is identified at the site of any proposed development or where the proposed activity has an elevated risk of resulting in the presence of these species, details of how these species will be managed and controlled will be required.	Although this Policy/objective will likely lead to a sustainable and positive impact, this policy/objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/policy
Peatlands policy	NEP - 9	To support the protection and restoration of peatlands in County Mayo, where appropriate, in order to transition towards a low-carbon and circular economy.	Protection of native habitats will positively impact on Natura 2000 Sites	N/A
Peatlands policy	NEP - 10	To recognise the role of peatlands as carbon sinks to combat climate change, and ensure that peatland areas, including those designated or proposed for designation (pNHA, NHA or SAC), are conserved for their ecological, climate regulation, archaeological, cultural and educational significance.	Protection of native habitats will positively impact on Natura 2000 Sites	N/A
Peatlands Objectives	NEO - 15	As part of the implementation of Climate Ready Mayo, Climate Adaption Strategy, to develop and implement a Peatland Management Strategy for County Mayo that will: (a) Identify	Protection of native habitats will positively impact on Natura 2000 Sites	N/A

Natural Environment				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		damaged Peatlands in the county and those at risk from climate change and becoming carbon emitters. b) Initiate conservation and management of Mayo's peatlands, particularly those sites nominated for designation as Special Areas of Conservation and Natural Heritage Areas, to preserve the habitat and their unique ecosystems, managing flood risk and other environmental benefits.		
Peatlands Objectives	NEO - 16	To actively increase public awareness of the importance of peatlands as carbon sinks to combat climate change.	Protection of native habitats will positively impact on Natura 2000 Sites	N/A
Green and blue Infrastructure policy	NEP - 11	To recognise the economic, social, environmental and physical value of green infrastructure	General statement of policy/general aspiration which will not lead to development	N/A
Green and blue Infrastructure policy	NEP - 12	To seek to protect and expand the green infrastructure within the county, and ensure their connectivity to provide additional green infrastructure, where possible, and to encourage green infrastructure to be spatially connected to facilitate the extension or establishment of ecological corridors.	Although this Policy/objective will likely lead to a sustainable and positive impact, this policy/objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Green and blue Infrastructure policy	NEP - 13	To promote and enhance green and blue infrastructure and seek to integrate the provision of green infrastructure with infrastructure provision and replacement,	Although this Policy/objective will likely lead to a sustainable and positive impact, this policy/objective could lead to increased development. Any development within the	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising

Natural Environment				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		including walking and cycling routes, as appropriate, while protecting and enhancing natural heritage and improving ecological corridors.	county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	from this objective/policy
Green and Blue Infrastructure Objectives	NEO - 17	To prepare a Green and blue Infrastructure Strategy for the County over the plan period, which will be incorporated into local area plans and non-statutory plans over the lifetime of the plan.	General statement of policy/general aspiration which will not lead to development	N/A
Green and Blue Infrastructure Objectives	NEO - 18	To identify green infrastructure and sustainable design that supports biodiversity and natural systems to adapt to climate change and contribute to climate adaptation in the built and natural environment.	Although this Policy/objective will likely lead to a sustainable and positive impact, this policy/objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/policy
Green and Blue Infrastructure Objectives	NEO - 19	To ensure the design and construction of new developments create low carbon, walkable neighbourhoods and workplaces, provides for high quality green infrastructure, where appropriate.	Although this Policy/objective will likely lead to a sustainable and positive impact, this policy/objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/policy

Natural Environment				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
			species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	
Green and Blue Infrastructure Objectives	NEO -20	To protect open spaces, with multifunctional green and blue infrastructure in developments, with connections to the wider Network of open spaces and habitats.	Although this Policy/objective will likely lead to a sustainable and positive impact, this policy/objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Green and Blue Infrastructure Objectives	NEO - 21	Protect and enhance the County's floodplains and wetlands as 'green infrastructure' which provides space for storage and conveyance of floodwater, enabling flood risk to be more effectively managed and reducing the NEP - ed to provide flood defenses in the future, subject to normal planning and environmental criteria.	Although this Policy/objective will likely lead to a sustainable and positive impact, this policy/objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Landscape policy	NEP - 14	To protect the physical, visual and scenic character of County Mayo and to preserve the unique landscape character of the area	Although this Policy/objective will likely lead to a sustainable and positive impact, this policy/objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Natural Environment				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
			network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	
Landscape Objectives	NEO - 22	Consider development, along Mayo's Scenic routes, that can demonstrate a clear need to locate in the area concerned, whilst ensuring that it: <ul style="list-style-type: none"> Does not impinge in any significant way on the character, integrity and distinctiveness of the area, Meets high standards in siting and design Satisfies all other criteria with regard to, inter alia, servicing, public safety, and environmental considerations Rural housing applications along Coastal Areas and Lakeshores must comply with the requirements set out in Objective RHO 4 (Chapter 3).	Although this Policy/objective will likely lead to a sustainable and positive impact, this policy/objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/policy
Landscape Objectives	NEO -23	Consider Development, within Mayo's Coastal Areas and Lakeshores and within areas along scenic routes with designated scenic views, that can demonstrate a long-standing social link to the area concerned, whilst ensuring that it <ul style="list-style-type: none"> Does not impinge in any significant way on the character, integrity and distinctiveness of the area Cannot be considered at an alternative location Meets high standards in siting and design Satisfies all other criteria with regard to, inter alia, servicing, public safety, and environmental considerations 	Although this Policy/objective will likely lead to a sustainable and positive impact, this policy/objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/policy
Landscape Objectives	NEO -24	To ensure all development proposals are consistent with the Landscape Appraisal of	Although this Policy/objective will likely lead to a sustainable and positive impact,	Refer to Chapter 9 of this NIR for mitigation

Natural Environment				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		County Mayo and the associated Landscape Sensitivity Matrix and future editions thereof.	this policy/objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	measures for any plan or project arising from this objective/policy
Landscape Objectives	NEO -25	To review the Landscape appraisal for Mayo and update this plan as appropriate, following publication of the statutory guidelines for Planning Authorities on local Landscape Character Assessments as detailed in the National Landscape Strategy 2015-2025.	General statement of policy/general aspiration which will not lead to development	N/A
Landscape Objectives	NEO -26	Require a Landscape/Visual Impact Assessment to accompany significant proposals, located within or adjacent to sensitive landscapes, where appropriate.	General statement of policy/general aspiration which will not lead to development	N/A
Coastal Zone Policy	NEP - 15	To protect the character, visual, recreational, ecological and amenity value of the coast and provisions for public access, while recognising the NEP - eds of coastal communities to live, work and interest with the coast.	Construction of coastal protection structures within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/policy
Coastal Zone Policy	NEP - 16	To maintain and enhance our natural coastal defences to increase resilience to climate change	Construction of coastal protection structures within the county could potentially have direct and/or indirect	Refer to Chapter 9 of this NIR for mitigation measures for any plan

Natural Environment				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
			impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	or project arising from this objective/ policy
Coastal Zone Objectives	NEO - 27	To ensure that the County's natural coastal defences, such as beaches, sand dunes, coastal wetlands and estuaries are not compromised by inappropriate works or development.	Construction of coastal protection structures within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Coastal Zone Objectives	NEO - 28	To ensure that any conservation works on the coastal dune - systems shall be carried out in accordance with best practice, subject to ecological impact assessment and Appropriate Assessment, as appropriate	Provision of Appropriate Assessment and protection of native habitats will positively impact on Natura 2000 Sites	N/A
Coastal Zone Objectives	NEO - 29	To investigate how the County's natural coastal defences, can be enhanced to increase climate resilience of our coastal communities.	Construction of coastal protection structures within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Natural Environment				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
Coastal Zone Objectives	NEO - 30	To ensure any new development within areas liable to coastal flooding and assessed and developed in accordance with the Flood Risk Management Guidelines for Planning Authorities (Department of the Environment, Heritage and Local Government and Office of Public Works, 2009).	General statement of policy/general aspiration which will not lead to development	N/A
Coastal Zone Objectives	NEO - 31	To protect the coastal zone - through the protection, enhancement and maintenance of the current status of the designated Blue Flag beaches and Green Coasts and seek to increase the number of beaches and coasts holding this status in the future	Construction of coastal protection structures within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Coastal Zone Objectives	NEO - 32	As part of the implementation of Climate Ready Mayo, Climate Adaption Strategy, to develop, in consultation with key stakeholders, an Integrated Coastal Zone - Management Plan for County Mayo to preserve, enhance and develop protective habitats of coastal ecosystems, dunes and wetlands, as well as protect critical infrastructure and assets from damaging storm surges, and to ensure new developments take account of future risk from coastal erosion/storm surges and sea level rise, including the identification and restriction of development in coastal erosion zones where appropriate, and ecosystem based adaptation	Construction of coastal protection structures within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Natural Environment				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		actions to manage climate risk and build resilience to climate change.		
Coastal Zone Objectives	NEO - 33	Ensure new developments take account of future risk from coastal erosion/storm surges and sea level rise, including the identification and restriction of development in coastal erosion zones where appropriate, and ecosystem-based adaptation actions to manage climate risk and build resilience to climate change.	General statement of policy/general aspiration which will not lead to development	N/A
Water Quality Policy	NEP - 17	To promote public awareness of water quality issues and the measures required to protect surface water and groundwater bodies for inappropriate and damaging development.	Protection or improvement to water quality will positively impact on Natura 2000 Sites	N/A
Water Quality Policy	NEP - 18	To co-operate with the EPA and other authorities in the continued implementation of the EU Water Framework Directive.	Protection or improvement to water quality will positively impact on Natura 2000 Sites	N/A
Water Quality Policy	NEP - 19	To protect existing groundwater sources and aquifers in the County and to manage development in a manner consistent with the protection of these resources. to meet the requirements of the Water Framework Directive	Protection or improvement to water quality will positively impact on Natura 2000 Sites	N/A
Water Quality Policy	NEP - 20	To meet our targets to achieve 'good status' in all water bodies in compliance with the Water Framework Directive and to cooperate with the implementation of the National River Basin Management Plan 2018-2021.	Protection or improvement to water quality will positively impact on Natura 2000 Sites	N/A
Water Quality Policy	NEP - 21	To manage, protect and enhance surface water and ground water quality to meet the	Protection or improvement to water quality will positively impact on Natura 2000 Sites	N/A

Natural Environment				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		requirements of the Water Framework Directive.		
Water Quality Policy	NEP - 22	To encourage the use of catchment-sensitive farming practices, in order to meet Water Framework Directive targets and comply with the River Basin Management Plan.	Protection or improvement to water quality will positively impact on Natura 2000 Sites	N/A
Water Quality Objectives	NEO - 34	To ensure that the Water Framework Directive, the River Basin Management Plan and any subsequent Water Management Plans are fully considered throughout the planning process.	Protection or improvement to water quality will positively impact on Natura 2000 Sites	N/A
Water Quality Objectives	NEO - 35	To ensure, through the implementation of the River Basin Management Plan(s) and the associated Programmes of Measures and any other associated legislation or revised plans with all relevant stakeholders, the protection and improvement of all drinking water, surface water and ground waters throughout the County.	Protection or improvement to water quality will positively impact on Natura 2000 Sites	N/A
Water Quality Objectives	NEO - 36	To manage in a sustainable manner the existing groundwater sources and aquifers in the County and manage development in a manner consistent with the sustainable management of these resources in conformity with the EU Environmental objectives (Groundwater) Regulations 2010 and the second cycle National River Basin Management Plan 2018-2021, and any subsequent plan and the Groundwater Protection Scheme.	Protection or improvement to water quality will positively impact on Natura 2000 Sites	N/A
Water Quality Objectives	NEO - 37	To protect groundwater sources through the implementation of the Groundwater Protection Scheme and Source Protection Zones.	Protection or improvement to water quality will positively impact on Natura 2000 Sites	N/A

Natural Environment				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		Development proposals within these zones which have the potential to pose a risk to groundwater will be required to demonstrate that no reasonable alternative site is available and that groundwater quality will be protected to the satisfaction of the Council		
Water Quality Objectives	NEO - 38	To protect both ground and surface water resources and to work with Irish Water to develop and implement Drinking Water Safety Plans, to protect sources of public water supply and their contributing catchment, and to work with the National Federation of Group Water Schemes, in respect of Source Protection Plans for Group Water Schemes to protect these sources.	Protection or improvement to water quality will positively impact on Natura 2000 Sites	N/A
Water Quality Objectives	NEO - 39	To comply with the Blue Dot Catchments Programme and protect high status water bodies in County Mayo, and ensure all proposed development which may have an impact on a high status water quality site will require site specific assessment to demonstrate localised pressures and demonstrate suitable mitigation measures in order to protect these sites.	Protection or improvement to water quality will positively impact on Natura 2000 Sites	N/A
Water Quality Objectives	NEO - 40	To protect through its regulatory controls and in conjunction with the Local Authority Waters Programme, water bodies with 'high ecological status', to restore water bodies that have fallen below 'high ecological status', to maintain water bodies at 'Good Status' and to mitigate	Protection or improvement to water quality will positively impact on Natura 2000 Sites	N/A

Natural Environment				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		threats to water bodies identified as 'At Risk' i.e. 'Moderate and Poor Status'		
Water Quality Objectives	NEP - 23	To support and facilitate the implementation of the Air Quality Regulations.	Improvement to air quality will positively impact on Natura 2000 Sites	N/A
Water Quality Objectives	NEP - 24	To promote the implementation of the Noise Directive 2002/49/EC and associated Environmental Noise Regulations 2006, as amended.	Positive impacts through reduction in Noise pollution under Noise Directive are generally anticipated from implementation of objective/policy	N/A
Water Quality Objectives	NEO - 41	Promote the achievement of best ambient air quality, compatible with sustainable development, in accordance with the EU Ambient Air Quality and Cleaner Air for Europe (CAFE) Directive (2008/50/EC) and by ensuring that all air emissions associated with new developments are within Environmental Quality Standards as set out in the Air Quality Standards Regulations 2011 (SI No. 180 of 2011) (or any updated/superseding documents).	Improvement to air quality will positively impact on Natura 2000 Sites	N/A
Water Quality Objectives	NEO - 42	To raise awareness of artificial light pollution and identify where lighting improvements or adjustments could be made to reduce its impact, where appropriate.	Positive impacts through reduction in light pollution are generally anticipated from implementation of objective/policy	N/A
Water Quality Objectives	NEO - 43	To protect Mayo Dark Sky Park at Wild Nephin Wilderness Park from adverse levels of artificial light pollution and encourage the use of Dark Sky Friendly lighting for all new lighting and lighting upgrades.	Positive impacts through reduction in light pollution are generally anticipated from implementation of objective/policy	N/A

A.10 Chapter 11 – Climate Action and Renewable Energy

Strategic Aim: to transition to a low carbon and climate resilient county, with an emphasis on reduction in energy demand and greenhouse gas emissions, through a combination of effective mitigation and adaptation responses to climate change; in addition to maximising the opportunities to become a national leader in renewable energy generation, whilst safeguarding the environment and other amenities.

Climate				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
Climate Action Policies	CAP 1	To support and enable the implementation and achievement of European and national objectives for climate adaptation and mitigation as detailed in the following documents, taking into account other provisions of the Plan (including those relating to land use planning, energy, sustainable mobility, flood risk management and drainage);	General statement of policy/general aspiration which will not lead to development	N/A
Climate Action Policies	CAP 1	<ul style="list-style-type: none"> Climate Action Plan (2019 and any subsequent versions); National Climate Change Adaptation Framework (2018 and any subsequent versions); Relevant provisions of any Sectoral Adaptation Plans prepared to comply with the requirements of the Climate Action and Low Carbon Development Act 2015, including those seeking to contribute towards the National Transition Objective, to pursue, and achieve, the transition to a low carbon, climate resilient and environmentally sustainable economy by the end of the year 2050; and Mayo Council Climate Change Adaptation Strategy (2019-2024 and any subsequent versions). 	General statement of policy/general aspiration which will not lead to development	N/A

Climate				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
Climate Action Policies	CAP 2	To support the National Climate Change Strategy and methods of reducing anthropogenic greenhouse gases on an ongoing basis through implementation of supporting objectives in this Plan, particularly those supporting use of alternative and renewable energy sources, sustainable transport, air quality, coastal zone management, flooding and soil erosion and promotion of the retention of, and planting of trees, hedgerows and afforestation, subject to no significant adverse effects on the environment including the integrity of the Natura 2000 network.	Policies or proposals which steer change in such way as to protect international nature conservation sites from adverse effects	N/A
Climate Action Policies	CAP 3	To support, promote and facilitate the advancement of climate action at the local and community level in County Mayo and to raise general awareness of issues associated with climate action and climate change mitigation and adaptation.	General statement of policy/general aspiration which will not lead to development	N/A
Climate Action Policies	CAP 4	To support local, regional, national and international initiatives for climate adaptation and mitigation and to limit emissions of greenhouse gases through energy efficiency and the development of renewable energy sources, which make use of all natural resources, including publicly owned lands, in an environmentally acceptable manner.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Climate Action Policies	CAP 5	To support the National Dialogue on Climate Action, in an effort to increase awareness of climate change, behavioural change and adaptation actions and in doing so provide an	General statement of policy/general aspiration which will not lead to development	N/A

Climate				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		ongoing platform for planning climate resilience with a focus on personal responsibility at all levels.		
Climate Action Policies	CAP 6	To support the transition to a competitive, low carbon, climate-resilient and environmentally sustainable economy by 2050, by way of reducing greenhouse gases, increasing renewable energy, and improving energy efficiency.	General statement of policy/general aspiration which will not lead to development	N/A
Climate Action Policies	CAP 7	To support and promote the enhancement of carbon sinks such as peatlands, appropriate afforestation and permanent grasslands, with consideration of afforestation on cut away peatlands.	Protection of native habitats will positively impact on Natura 2000 Sites	N/A
Climate Action Policies	CAP 8	To cooperate with the Climate Action Regional Office (CARO) in respect of the implementation of existing and future climate change adaption and mitigation strategies.	General statement of policy/general aspiration which will not lead to development	N/A
Climate Action Policies	CAP 9	To support Ireland's renewable energy commitments outlined in national policy by facilitating the development and exploitation of all appropriate renewable energy sources at suitable locations within the county, where such development does not have a negative impact on the surrounding environment (including water quality), landscape, biodiversity or local amenities, so as to provide for further residential and enterprise development within the county.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Climate Action Objectives	CAO 1	To support and advance the provision of renewable energy resources and programmes in line with the Government's National Renewable Energy Action Plan (NREAP), the Governments' Energy White Paper "Ireland's Transition to a Low	General statement of policy/general aspiration which will not lead to development	N/A

Climate				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		Carbon Energy Future (2015-2030) and any other relevant policy adopted during the lifetime of this plan.		
Climate Action Objectives	CAO 2	To support, facilitate and advance the implementation of Climate Ready Mayo, the Climate Change Adaptation Strategy for County Mayo.	General statement of policy/general aspiration which will not lead to development	N/A
Climate Action Objectives	CAO 3	To carry out a carbon emissions baseline for County Mayo over the lifetime of the plan.	General statement of policy/general aspiration which will not lead to development	N/A
Climate Action Objectives	CAO 4	To develop and implement a climate change screening checklist and guidance document over the lifetime of the plan, to ensure new development takes account of climate change over the lifetime of a development, in particular with regard to its location, site layout, building, ventilation and cooling, drainage, water, outdoor spaces and connectivity.	General statement of policy/general aspiration which will not lead to development	N/A
Climate Action Objectives	CAO 5	To identify and develop Decarbonising Zone(s) in Mayo, as per Action 165 of the Climate Action Plan 2019 and to promote the use of the Decarbonisation Zone(s) as an exemplary example for best practice within the county.	General statement of policy/general aspiration which will not lead to development	N/A
Climate Action Objectives	CAO 6	To increase the resilience of Natural and Cultural Capital, as per Goal 3 of Climate Ready Mayo, by: (a) Building awareness of Nature Based Adaptation Solutions and Green Infrastructure. (b) Support biodiversity for its intrinsic value within the natural environment and its importance in climate change adaptation.	General statement of policy/general aspiration which will not lead to development	N/A

Climate				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		(c) Develop a database of impacts of climate change on Mayo's Natural Environment. (d) Identify Cultural and Heritage Sites vulnerable to climate change and develop adaptation and management policies. (e) Encourage adaptation in Agriculture and Local Food Supply		
Climate Action Objectives	CAO 7	To recognise, support and facilitate Ballina to become Ireland's Greenest Town by 2025	General statement of policy/general aspiration which will not lead to development	N/A
Renewable Energy Policies	REP 1	To support Ireland's renewable energy commitments outlined in national policy by facilitating the development and exploitation of a range of renewable energy sources at suitable locations within the county, where such development does not have a negative impact on the surrounding environment (including water quality), landscape, biodiversity or local amenities to ensure the long term sustainable growth of the county	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Renewable Energy Policies	REP 2	To support, within the context of the Offshore Renewable Energy Development Plan (OREDPP) and its successors, the progressive development of Ireland's offshore renewable energy potential, including domestic and international grid connectivity enhancements.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Renewable Energy Policies	REP 3	To actively encourage and support the sustainable development, renewal and maintenance of energy generation infrastructure in order to maintain a	Policies or proposals which steer change in such as way as to protect international nature conservation sites from adverse effects	N/A

Climate				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		secure energy supply, while protecting the landscape, archaeological and built heritage and having regard to the provisions of the Habitats Directive.		
Renewable Energy Policies	REP 4	To ensure that developers of proposed large-scale renewable energy projects carry out community consultation in accordance with best practice and commence the consultation at the initiation of project planning.	General statement of policy/general aspiration which will not lead to development	N/A
Renewable Energy Policies	REP 5	To promote the use of efficient energy storage systems and infrastructure that supports energy efficiency and reusable energy system optimization, subject to the proper planning and sustainable development of the area and consideration of environmental and ecological sensitivities.	Policies or proposals which steer change in such a way as to protect international nature conservation sites from adverse effects	N/A
Renewable Energy Policies	REP 6	To work with relevant stakeholders and industry to establish Mayo as a centre of excellence for renewable energy research and development activities.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Renewable Energy Policies	REP 7	To promote the harnessing of wind energy to contribute toward decarbonising County Mayo, including new emerging by-product markets.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Climate				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
			indicators of conservation value, such as changes in water quality and quantity, and air quality.	
Renewable Energy Objectives	REO 1	To co-operate with the Northern and Western Regional Assembly in identifying Strategic Energy Zones as areas suitable for larger, energy generating projects, community and micro energy production, whilst ensuring environmental constraints and a regional landscape strategy are considered.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Renewable Energy Objectives	REO 2	To examine options to ensure that community benefits are derived from renewable energy development in the County	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Renewable Energy Objectives	REO 3	To encourage and facilitate, where possible, the production of energy from established and emerging renewable technologies.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Renewable Energy Objectives	REO 4	To support and implement the recording and monitoring of renewable energy potential in the county in partnership with other stakeholders including the Sustainable Energy Authority of Ireland (SEAI).	General statement of policy/general aspiration which will not lead to development	N/A

Climate				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
Renewable Energy Objectives	REO 5	To support and work in partnership with local communities in the development of energy efficient and renewable energy projects.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Renewable Energy Objectives	REO 6	To ensure all renewable energy proposal comply with the provisions of the Mayo County Council Renewable Energy Strategy 2011-2022 (or as updated).	General statement of policy/general aspiration which will not lead to development	N/A
Renewable Energy Objectives	REO 7	To review the Mayo County Renewable Energy Strategy 2011-2022 in accordance with future legislative guidelines.	General statement of policy/general aspiration which will not lead to development	N/A
Renewable Energy Objectives	REO 8	To encourage the development of wind energy, in accordance with Government policy and having regard to the <i>Landscape Appraisal of County Mayo</i> and the Wind Energy Development Guidelines (2006) and Mayo Renewable Energy Strategy, or any revisions thereof or future guidelines.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Renewable Energy Objectives	REO 9	To support Ireland's renewable energy commitments outlined in national policy by promoting the development of solar energy.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Climate				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
Renewable Energy Objectives	REO 10	To encourage solar energy in commercial and residential developments, subject to the proper planning and sustainable development of the area and consideration of environmental and ecological sensitivities.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Renewable Energy Objectives	REO 11	To ensure that solar farm development proposals in the vicinity of major road networks & transport nodes, such as Ireland West Airport Knock (15km Radius), do not create a traffic hazard or endanger aircraft safety by reason of glint and glare.	General statement of policy/general aspiration which will not lead to development	N/A
Renewable Energy Objectives	REO 12	To support offshore and tidal renewable energy developments subject to environmental considerations and the protection of the amenities of the surrounding areas in accordance with the OREDP, subject to proper planning and environmental considerations.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Renewable Energy Objectives	REO 13	To recognise the important role of the Atlantic Marine Energy Test Site (AMETS) tidal wave test site off Belmullet.	General statement of policy/general aspiration which will not lead to development	N/A
Renewable Energy Objectives	REO 14	To support the development of appropriate land-based infrastructure at suitable locations, in order to facilitate the transition between the land and sea necessary for off- shore renewable energy projects.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Climate				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
			indicators of conservation value, such as changes in water quality and quantity, and air quality.	
Renewable Energy Objectives	REO 15	To facilitate large and smaller scale geothermal energy generating developments both standalone and in conjunction with other renewable energy projects.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Renewable Energy Objectives	REO 16	To promote the use of geothermal heat pumps for space heating and cooling as well as water heating in domestic, commercial and recreational buildings.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Renewable Energy Objectives	REO 17	To promote on-site wind/solar energy development or other emerging energy technologies, where energy generated is primarily required to meet the needs of households, communities, agriculture and other businesses to reduce their carbon emissions.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Renewable Energy Objectives	REO 18	Support, promote and facilitate community energy-based initiatives such as the Sustainable Energy Authority of Ireland's (SEAI) 'Sustainable Community Energy' scheme or similar community	General statement of policy/general aspiration which will not lead to development	N/A

Climate				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		energy initiatives to help achieve low carbon communities.		
Renewable Energy Objectives	REO 19	To support Ireland's renewable energy commitments outlined in national policy by promoting the use of district heating systems in new residential and commercial developments.	General statement of policy/general aspiration which will not lead to development	N/A
Renewable Energy Objectives	REO 20	To consider using heat mapping to support developments which deliver energy efficiency and the recovery of energy that would otherwise be wasted.	General statement of policy/general aspiration which will not lead to development	N/A
Renewable Energy Objectives	REO 21	To carry out a feasibility assessment for district heating in County Mayo and identify local waste heat sources or renewable energy sources.	General statement of policy/general aspiration which will not lead to development	N/A
Renewable Energy Objectives	REO 22	To promote the use of efficient energy storage systems and infrastructure that supports energy efficiency and reusable energy system optimisation, in accordance with proper planning and sustainable development	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

A.11 Chapter 12 – Settlement Plans

Strategic Aim: to create a network of attractive, liveable towns and villages in the County, with increased levels of population, employment activity and enhanced levels of amenity, which support a high quality of life and wellbeing, along with providing an alternative residential choice for those who may not wish to live in the rural countryside.

Policies and objectives have been devised for Tier II towns: Ballinrobe, Ballyhaunis, Belmullet, Claremorris & Swinford; and Tier III towns: Balla, Charlestown, Crossmolina, Foxford, Killala, Kiltimagh, Knock, Louisburgh, and Newport.

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
General Settlement Policies	GSP 1	Support and promote commensurate population, service and employment growth, to enable the Tiers 2 and 3 settlements to fulfil their role as a self-sustaining growth towns and self-sustaining towns, respectively.	General statement of policy/general aspiration which will not lead to development	N/A
General Settlement Policies	GSP 2	Support and encourage the development/redevelopment of identified Town Centre Consolidation Sites in Tier II settlements and Opportunity Sites in Tier III settlements.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
General Settlement Policies	GSP 3	Support the provision of mixed-use developments in the town centre which create opportunities to live, work, shop, etc., within the town and reduce the propensity to travel by private car.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
			conservation value, such as changes in water quality and quantity, and air quality.	
General Settlement Policies	GSP 4	Encourage re-development of all derelict buildings within all settlement plan areas listed on the Derelict Sites Register.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
General Settlement Policies	GSP 5	Support and promote the re-use and regeneration of derelict land and other buildings in town centre areas for retail and other appropriate uses.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
General Settlement Policies	GSP 6	To encourage the appropriate use of unoccupied/derelict buildings in the town for start-up businesses and community facilities.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
General Settlement Policies	GSP 7	Co-operate with relevant agencies to secure improvements to the public	This objective could lead to increased development. Any development within the	Refer to Chapter 9 of this NIR for mitigation

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		transport system serving Tier 2 and 3 settlement towns and to support initiatives designed to improve bus interchange facilities.	county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	measures for any plan or project arising from this objective/ policy
General Settlement Policies	GSP 8	Support the establishment of green routes and an accessible walking / cycling network throughout the settlement plan areas and surrounding areas that provides safe and attractive circulation routes for pedestrians and cyclists for the enjoyment and recreational use of the entire community by linking residential areas, community facilities, amenities and the town centre.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
General Settlement Policies	GSP 9	Support and encourage Irish Water in increasing the provision of adequate wastewater and water infrastructure, to ensure that services are delivered in line with the further development and growth of settlements.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
General Settlement Policies	GSP 10	Support and encourage key stakeholders/providers in increasing the provision of	This objective could lead to increased development. Any development within the county could potentially have direct and/or	Refer to Chapter 9 of this NIR for mitigation measures for any plan

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		adequate key physical infrastructure (i.e. transportation, parking, communications, energy etc.) and to support the provision of key social infrastructure (health care services, education facilities, burial grounds, fire and emergency services, recreational, cultural facilities etc.) in Tier II and Tier III settlements.	indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	or project arising from this objective/ policy
General Settlement Objectives	GSO 1	To ensure appropriate development occurs in a sequential manner outward from the core area, to maximise the utility of existing and future infrastructure provision, to promote sustainability and active travel, to make more efficient use of underutilised lands, and to avoid the inappropriate extension of services and utilities.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
General Settlement Objectives	GSO 2	Encourage and facilitate the development of the economic and tourism potential of towns in a manner that respects, builds on, protects and enhances the cultural, built heritage, natural heritage and local amenities of the town.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
General Settlement Objectives	GSO 3	Ensure that the town centre is accessible to all members of the community, including people with	This objective could lead to increased development. Any development within the county could potentially have direct and/or	Refer to Chapter 9 of this NIR for mitigation measures for any plan

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		mobility issues, the elderly and people with young children.	indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	or project arising from this objective/ policy
General Settlement Objectives	GSO 4	Work with Irish Water and landowners on the “New Homes in Small Towns and Villages” initiative to augment the delivery of actions by Local Authorities, Irish Water, communities and other stakeholders in the provision of services and serviced sites to create “build your own home” opportunities within the existing footprint of rural settlements, to meet housing demand.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
General Settlement Objectives	GSO 5	Require proposals for new development to integrate with existing Green Infrastructure networks and contribute to the development and protection of overall Green Infrastructure assets.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
General Settlement Objectives	GSO 6	To use active land management measures, such as the Vacant Site Levy and Derelict Site Levy, to ensure the delivery of the projected	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000	Refer to Chapter 9 of this NIR for mitigation measures for any plan

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		housing units in the identified Town Centre Consolidation Sites and residential zoned lands in Tier II Self-Sustaining Growth Towns, as set out in the Core Strategy or any subsequent Town Centre Consolidation Sites identified over the lifetime of the plan.	network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	or project arising from this objective/ policy
General Settlement Objectives	GSO 7	To promote and facilitate the delivery of multiple residential development in the identified Opportunity Sites for Tier III Self-Sustaining Towns or any subsequent Opportunity Sites identified over the lifetime of the plan.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
General Settlement Objectives	GSO 8	Ensure applications for development within the settlement boundaries on lands identified as flood risk areas including benefitting lands, shall be subject to a Specific Flood Risk Assessment and Justification Test, in accordance with the Planning System and Flood Risk Management – Guidelines for Planning Authorities, 2009 or any superseding guidelines and circulars.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
General Settlement Objectives	GSO 9	To facilitate, identify, support and secure a strategic footpath and cycleway network throughout the	This objective could lead to increased development. Any development within the county could potentially have direct and/or	Refer to Chapter 9 of this NIR for mitigation measures for any plan

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		plan areas, as appropriate and as resources allow.	indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	or project arising from this objective/ policy
General Settlement Objectives	GSO 10	Support and facilitate pedestrian mobility and safety in the town by introducing traffic calming measures and pedestrian crossings.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
General Settlement Objectives	GSO 11	Support and where possible, implement measures to create interpretative walking routes in and around the town, linking the town's special features of built and natural heritage interest.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
General Settlement Objectives	GSO 12	Develop the local economy by encouraging additional commercial businesses and industries and to promote the clustering of such industries on suitably zoned land in Tier II settlements and at	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		appropriate locations within Tier II settlements.	conservation value, such as changes in water quality and quantity, and air quality.	
General Settlement Objectives	GSO 13	Protect the role of the town centre as the dominant retailing and commercial area.	General statement of policy/general aspiration which will not lead to development	N/A
General Settlement Objectives	GSO 14	Implement the land use zoning objectives in each Tier 2 settlement plans.	General statement of policy/general aspiration which will not lead to development	N/A
General Settlement Objectives	GSO 15	To actively provide or facilitate the provision/upgrade of identified infrastructural deficiencies in settlement plan areas.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Ballinrobe Settlement Plan Policies	BEP 1	Promote the development of Ballinrobe as a driver of economic growth for the south region of county Mayo and fulfil its role as a designated Self-Sustaining Growth Town and support the potential for the creation of an Economic Growth Cluster in tandem with the towns of Ballyhaunis and Claremorris.	General statement of policy/general aspiration which will not lead to development	N/A
Ballinrobe Settlement Plan Policies	BEP 2	Promote and support the re-development and refurbishment of Bridge Street/High Street whilst safeguarding the protected	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		structures and their curtilages along the street.	habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	
Ballinrobe Settlement Plan Policies	BEP 3	Promote and support the appropriate refurbishment of Cranmore House and the Military Barracks for amenity and cultural purposes.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Ballinrobe Settlement Plan Policies	BEP 4	Promote and support the re-establishment of the Ballinrobe to Claremorris dismantled railway line as a walking and cycling route or as a spur line of the Western Rail corridor.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Ballinrobe Settlement Plan Policies	BEP 5	Promote and support the enhancement and extension of the Bowers Walk River walkway and safeguard the value of the river as an ecological “green corridor”. Riverside walkway provisions should be incorporated, where appropriate, into development proposals bounding the river.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
Ballinrobe Settlement Plan Policies	BEP 6	Promote and support the implementation of the projects listed within the Ballinrobe Public Realm Plan during the plan period, to improve attractiveness and permeability of the public realm.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Ballinrobe Settlement Plan Objectives	BEO 1	Ensure infill development respects the unique height-to-width ratio with respect to building design in Ballinrobe town centre and maintain the existing sense of enclosure on town centre streets.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Ballinrobe Settlement Plan Objectives	BEO 2	To consider the designation in the town centre of an Architectural Conservation Area, including all or parts of Bowgate Street, Main Street, High Street/Bridge Street, Glebe Street and Abbey Street.	Policies or proposals which steer change in such as way as to protect international nature conservation sites from adverse effects	N/A
Ballinrobe Settlement Plan Objectives	BEO 3	Support and facilitate the development of the town centre consolidation site located to the rear of New Street/Convent Road for appropriate uses, as outlined in the land uses generally permitted on town centre consolidation sites in	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		the Land Use Zoning Matrix, with pedestrian links to New Street/Convent Road.	conservation value, such as changes in water quality and quantity, and air quality.	
Ballinrobe Settlement Plan Objectives	BEO 4	Seek and encourage ways to make more use of the Cornmarket area, including making the area pedestrian and market-place friendly and encourage and support the provision of a weekly market in the Cornmarket area.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Ballinrobe Settlement Plan Objectives	BEO 5	Protect the water quality and riparian zone of the rivers Robe and Bulkan. Any proposed developments adjacent to or close to watercourses shall be carefully assessed to ensure that there is no adverse impact to the water course, its riparian zone or to any waterbody into which it flows.	Policies or proposals which steer change in such as way as to protect international nature conservation sites from adverse effects	N/A
Ballinrobe Settlement Plan Objectives	BEO 6	Seek to make Tree Preservation Orders for the tree groups along the River Robe and important tree groups in the town.	Policies or proposals which steer change in such as way as to protect international nature conservation sites from adverse effects	N/A
Ballinrobe Settlement Plan Objectives	BEO 7	To encourage development in the town of Ballinrobe in accordance with the Land Use Zoning Map (Map BE1).	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
			species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	
Ballinrobe Settlement Plan Objectives	BEO 8	To encourage and facilitate the re-use and regeneration of derelict land and buildings for retail and other town centre uses with due cognisance to the Sequential Approach prescribed in the Retail Planning Guidelines 2012. (Refer to Map BE2)	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Ballinrobe Settlement Plan Objectives	BEO 9	To use active land management measures, such as the vacant site levy and derelict site levy to ensure the delivery of the projected housing units for Ballinrobe, as set out in the Core Strategy, on town centre consolidation sites and residential zoned lands (Map BE3).	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Ballyhaunis Settlement Plan Policies	BSP 1	Promote the development of Ballyhaunis as a driver of economic growth for the southeast region of County Mayo and fulfil its role as a designated Self-Sustaining Growth Town and to further investigate the potential for the creation of an Economic Growth Cluster, in tandem with the towns of Ballinrobe and Claremorris.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
Ballyhaunis Settlement Plan Policies	BSP 2	Protect St. Mary's Abbey and its associated lands as one of the town's key tourist attractions.	General statement of policy/general aspiration which will not lead to development	N/A
Ballyhaunis Settlement Plan Policies	BSP 3	Support community-led developments, where appropriate, including the implementation of Community Action Plan for Ballyhaunis (2016-2021) or any amended or superseding Community Futures Plan for the town.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Ballyhaunis Settlement Plan Policies	BSO 4	Promote and support the enhancement and extension of the Dalgan River walkway and to safeguard the value of the river as an ecological "green corridor". Riverside walkway provisions should be incorporated, where appropriate, into development proposals bounding the river.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Ballyhaunis Settlement Plan Policies	BSO 5	Promote the town's strategic location in relation to Ireland West Airport Knock.	General statement of policy/general aspiration which will not lead to development	N/A
Ballyhaunis Settlement Plan Objectives	BSO 1	Improve the attractiveness of the built fabric of the town centre, including the encouragement of appropriate redevelopment and renewal of vacant sites and buildings, in particular Knox St, and	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		to ensure high architectural quality in all new developments.	conservation value, such as changes in water quality and quantity, and air quality.	
Ballyhaunis Settlement Plan Objectives	BSO 2	Support and facilitate the development of the four identified town centre consolidation sites for appropriate uses, as outlined in the land uses generally permitted on town centre consolidation sites in the Land Use Zoning Matrix.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Ballyhaunis Settlement Plan Objectives	BSO 3	Continue to facilitate the further development of the Abbey Walk an amenity / recreation area in Ballyhaunis.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Ballyhaunis Settlement Plan Objectives	BSO 4	Provide a civic amenity facility in Ballyhaunis.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Ballyhaunis Settlement Plan Objectives	BSO 5	Continue to encourage rail use as a sustainable mode of transport and	This objective could lead to increased development. Any development within the	Refer to Chapter 9 of this NIR for mitigation

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		to promote the upgrading of the rail services to the town and connecting bus services.	county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	measures for any plan or project arising from this objective/ policy
Ballyhaunis Settlement Plan Objectives	BSO 6	Ensure the future protection of lands adjacent to the railway station from inappropriate development that could jeopardize the long-term viability and improvement of the transport facility.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Ballyhaunis Settlement Plan Objectives	BSO 7	Protect the water quality and riparian zone of the Dalgan River and to encourage appropriate developments that enhance its landscape setting and public benefit.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Ballyhaunis Settlement Plan Objectives	BSO 8	Facilitate the construction of the town's outer bypass.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
			species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	
Ballyhaunis Settlement Plan Objectives	BSO 9	Ensure that the strategic importance, capacity and safe operation of the Knock Link Road is maintained by restricting the number of access points entering and exiting onto the road.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Ballyhaunis Settlement Plan Objectives	BSO 10	To encourage development in the town of Ballyhaunis in accordance with the Land Use Zoning Map (Map BS1).	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Ballyhaunis Settlement Plan Objectives	BSO 11	To encourage and facilitate the re-use and regeneration of derelict land and buildings for retail and other town centre uses with due cognisance to the Sequential Approach prescribed in the Retail Planning Guidelines 2012. (Refer to Map BS2)	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
Ballyhaunis Settlement Plan Objectives	BSO 12	To use active land management measures, such as the vacant site levy and derelict site levy to ensure the delivery of the projected housing units for Ballyhaunis, as set out in the Core Strategy, on town centre consolidation sites and residential zoned lands (Map BS3).	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Béal an Mhuirthead Settlement Plan Policies	BTP 1	Promote the development of Béal an Mhuiread as a driver of economic growth for the Erris region and fulfil its role as a designated Self-Sustaining Growth Town.	General statement of policy/general aspiration which will not lead to development	N/A
Béal an Mhuirthead Settlement Plan Policies	BTP 2	Strengthen and protect the linguistic heritage of the town and to promote use of the Irish language in the public realm through signage and other measures throughout the town centre area.	General statement of policy/general aspiration which will not lead to development	N/A
Béal an Mhuirthead Settlement Plan Policies	BTP 3	Support the provision of a swimming pool in Béal an Mhuirthead through public private partnership, assistance to a private promoter or any other means.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
Béal an Mhuirthead Settlement Plan Policies	BTP 4	Support community-led development, where appropriate, including the implementation of Community Action Plan for Belmullet (2019-2024) or any amended or superseding Community Futures Plan for the town.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Béal an Mhuirthead Settlement Plan Objectives	BTO 1	To work with local agencies, in particular Údarás na Gaeltachta, to encourage the development and growth of Béal an Mhuirthead as an employment centre for the wider rural area.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Béal an Mhuirthead Settlement Plan Objectives	BTO 2	Prioritise the construction of a civic amenity site in Béal an Mhuirthead.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Béal an Mhuirthead Settlement Plan Objectives	BTO 3	Create a more pedestrian friendly environment in the town centre of Béal an Mhuirthead.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000	Refer to Chapter 9 of this NIR for mitigation measures for any plan

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective		Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red		Mitigation
			network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	or project arising from this objective/ policy
Béal an Mhuirthead Settlement Plan Objectives	BTO 4	To support and facilitate the development of the three identified town centre consolidation sites in Béal an Mhuirthead for appropriate uses, as outlined in Section 12.6.11, and as generally permitted on town centre consolidation sites in the Land Use Zoning Matrix.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Béal an Mhuirthead Settlement Plan Objectives	BTO 5	To encourage development in the town of Béal an Mhuirthead in accordance with the Land Use Zoning Map (Map BT1).	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Béal an Mhuirthead Settlement Plan Objectives	BTO 6	To encourage and facilitate the re-use and regeneration of derelict land and buildings for retail and other town centre uses with due cognisance to the Sequential Approach prescribed in the Retail Planning Guidelines 2012. (Refer to Map BT2)	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
			conservation value, such as changes in water quality and quantity, and air quality.	
Béal an Mhuirthead Settlement Plan Objectives	BTO 7	To use active land management measures, such as the vacant site levy and derelict site levy to ensure the delivery of the projected housing units for Béal an Mhuirthead, as set out in the Core Strategy, on town centre consolidation sites and residential zoned lands (Map BT3).	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Claremorris Settlement Plan Policies	CSP 1	To promote the development of Claremorris as a driver of economic growth for the southeast region and fulfil its role as a designated Self-Sustaining Growth Town and to further investigate the potential for the creation of an Economic Growth Cluster in tandem with the towns of Ballinrobe and Ballyhaunis.	General statement of policy/general aspiration which will not lead to development	N/A
Claremorris Settlement Plan Policies	CSP 2	To support and promote the development potential of Claremorris as a transportation hub given its trans-nodal infrastructural location and siting along the Atlantic Economic Corridor.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Claremorris Settlement Plan Policies	CSP 3	To support the reinstatement of the Western Rail Corridor, in particular	This objective could lead to increased development. Any development within the	Refer to Chapter 9 of this NIR for mitigation

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective		Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red		Mitigation
		to actively seek and promote the re-opening of the Claremorris-Galway rail link as well as a link to Sligo and to safeguard and protect these potential rail links from redevelopment for non-transport related purposes, in order not to preclude their future uses as an operational transportation network.	county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	measures for any plan or project arising from this objective/ policy
Claremorris Settlement Plan Policies	CSP 4	To support the development of Claremorris station for passenger and freight services and as a distribution depot.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Claremorris Settlement Plan Policies	CSP 5	To support and develop Claremorris as a low carbon town, in conjunction with relevant stakeholders, including Claremorris and Western District Energy Co-Operative.	General statement of policy/general aspiration which will not lead to development	N/A
Claremorris Settlement Plan Objectives	CSO 1	To develop an attractive, serviced industrial land bank to the north of the inner relief road, on lands zoned for industrial development (refer to Map xxx), suitable for a range of high class and environmentally acceptable business and technology / industrial park type developments.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
			conservation value, such as changes in water quality and quantity, and air quality.	
Claremorris Settlement Plan Objectives	CSO 2	To continue to facilitate the further development of the Mayfield Lough and McMahon Park as an amenity / recreation area in Claremorris	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Claremorris Settlement Plan Objectives	CSO 3	To support and facilitate the development of the six identified town centre consolidation sites in Claremorris for appropriate uses, as outlined in Section 12.7.11, and as generally permitted on town centre consolidation sites in the Land Use Zoning Matrix.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Claremorris Settlement Plan Objectives	CSO 4	To provide a civic amenity facility in Claremorris.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Claremorris Settlement Plan Objectives	CSO 5	To examine the feasibility of, with the aim of constructing, the	This objective could lead to increased development. Any development within the	Refer to Chapter 9 of this NIR for mitigation

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		following road projects within the plan area:	county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	measures for any plan or project arising from this objective/ policy
Claremorris Settlement Plan Objectives	CSO 5	(i) construction of an inner relief road from the N60 to the old N17	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Claremorris Settlement Plan Objectives	CSO 5	(ii) construction of a link road from the L-5572-0 (Cloonconner Road) to the old N17 Claremorris/Ballindine Road, with the intention of extinguishing the vehicular right of way on the Cloonconner Road to/from the N17 By-Pass	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Claremorris Settlement Plan Objectives	CSO 5	The above road projects will be subject to the normal planning and environmental considerations, including an assessment in accordance with Article 6 of the EU Habitats Directive, as appropriate.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
			species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	
Claremorris Settlement Plan Objectives	CSO 6	To safeguard and protect the Claremorris By-Pass from development for non-infrastructure related purposes that would prejudice its future use as part of the Western Infrastructural Corridor.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Claremorris Settlement Plan Objectives	CSO 7	To protect, maintain and enhance, in conjunction with the relevant agencies, the conservation value of Mayfield and Clare Loughs, their associated wetland habitats and rich diversity of plant and animal species and the extent, quality and connectivity of associated surface waters and wetlands as well as to promote the educational role of the lakes.	Policies or proposals which steer change in such as way as to protect international nature conservation sites from adverse effects	N/A
Claremorris Settlement Plan Objectives	CSO 8	To protect and enhance the conservation value of the disused railway lines in the town and their setting / margins as wildlife habitats and corridors notwithstanding that such disused railway lines may be developed at some future date as part of the County's infrastructure /	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		greenway network, in a manner that has regard to their inherent conservation value.	conservation value, such as changes in water quality and quantity, and air quality.	
Claremorris Settlement Plan Objectives	CSO 9	To comply with the requirements of the objectives in the Economic Development Chapter of this Plan regarding Seveso II site 'Calor Gas Teo'.	General statement of policy/general aspiration which will not lead to development	N/A
Claremorris Settlement Plan Objectives	CSO 10	To encourage development in the town of Claremorris in accordance with the Land Use Zoning Map (Map CS1).	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Claremorris Settlement Plan Objectives	CSO 11	To encourage and facilitate the re-use and regeneration of derelict land and buildings for retail and other town centre uses with due cognisance to the Sequential Approach prescribed in the Retail Planning Guidelines 2012. (Refer to Map CS2)	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Claremorris Settlement Plan Objectives	CSO 12	To use active land management measures, such as the vacant site levy and derelict site levy to ensure the delivery of the projected housing units for Claremorris, as set out in	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		the Core Strategy, on town centre consolidation sites and residential zoned lands (Map CS3).	habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	
Swinford Settlement Plan Policies	SDP 1	To promote the development of Swinford as a driver of economic growth for the east region of Mayo and fulfil its role as a designated Self-Sustaining Growth Town.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Swinford Settlement Plan Policies	SDP 2	To support and promote the reinstatement of the Western Rail Corridor.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Swinford Settlement Plan Policies	SDP 3	Support community-led developments, where appropriate, including the implementation of Community Action Plan for Swinford (2013-2018) or any amended or superseding Community Futures Plan for the town.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
Swinford Settlement Plan Objectives	SDO 1	To protect and enhance existing amenity facilities in Swinford, particularly the golf course, tennis courts, Amenity Park and Brabazon Woods.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Swinford Settlement Plan Objectives	SDO 2	To protect and enhance areas considered unique and important in the town such as the Courthouse, the Famine Graveyard and Church of Ireland Graveyard, the Church, the Railway Bridge, the Signal Box and the Water Tower.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Swinford Settlement Plan Objectives	SDO 3	To support and facilitate the development of the six identified town centre consolidation sites in Swinford for appropriate uses, as outlined in Section 12.8.11, and as generally permitted on town centre consolidation sites in the Land Use Zoning Matrix.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Swinford Settlement Plan Objectives	SDO 4	To develop the local economy by encouraging additional healthcare and pharmaceutical industries in the town and to promote the clustering	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000	Refer to Chapter 9 of this NIR for mitigation measures for any plan

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective		Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red		Mitigation
		of such industries on suitably zoned land.	network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	or project arising from this objective/ policy
Swinford Settlement Plan Objectives	SDO 5	To ensure infill development respects the unique height-to-ratio building design in Swinford town centre along Market Street and Main Street.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Swinford Settlement Plan Objectives	SDO 6	To protect the Spaddagh and Derryronan Rivers and all water courses and surface waters within the area, their water quality, ecology and function as ecological corridors, in this regard developers are required to show that any proposed development in the town will not adversely affect the integrity of the River Moy candidate Special Area of Conservation.	Policies or proposals which steer change in such as way as to protect international nature conservation sites from adverse effects	N/A
Swinford Settlement Plan Objectives	SDO 7	To develop and maintain a pedestrian access to Pound Street car park from the town centre	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
			species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	
Swinford Settlement Plan Objectives	SDO 8	To examine the feasibility of enhancing the link between the town centre and the Tesco development site.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Swinford Settlement Plan Objectives	SDO 9	To support and facilitate the local community in measures to commemorate Swinford 2050 including oak tree planting on site to the rear of the former vocational school.	General statement of policy/general aspiration which will not lead to development	N/A
Swinford Settlement Plan Objectives	SDO 10	To encourage and facilitate the re-use and regeneration of derelict land and buildings for retail and other town centre uses with due cognisance to the Sequential Approach prescribed in the Retail Planning Guidelines 2012. (Refer to Map SD2)	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Swinford Settlement Plan Objectives	SDO 11	To use active land management measures, such as the vacant site levy and derelict site levy to ensure the delivery of the projected housing	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000	Refer to Chapter 9 of this NIR for mitigation measures for any plan

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		units for Swinford, as set out in the Core Strategy, on town centre consolidation sites and residential zoned lands (Map SD3).	network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	or project arising from this objective/ policy
Balla Settlement Plan Policies	BAP 1	Support community-led developments, where appropriate, including the implementation of Community Action Plan for Balla (2017-2022) or any amended or superseding Community Futures Plan for the town.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Balla Settlement Plan Objectives	BAO 1	Protect and enhance existing amenity areas/facilities in Balla, particularly the round tower, fair green, town park, and its sporting and recreational facilities.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Balla Settlement Plan Objectives	BAO 2	Protect the town park and encourage its enhancement through the Neighbourhood scheme (2017)	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
			conservation value, such as changes in water quality and quantity, and air quality.	
Balla Settlement Plan Objectives	BAO 3	Support and facilitate pedestrian mobility and safety in the town by introducing traffic calming measures and pedestrian crossings.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Balla Settlement Plan Objectives	BAO 4	To encourage and facilitate the re-use and regeneration of derelict land and buildings for retail and other town centre uses with due cognisance to the Sequential Approach prescribed in the Retail Planning Guidelines 2012. (Refer to Map BA2)	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Balla Settlement Plan Objectives	BAO 5	To support and facilitate the development of the two identified opportunity sites in Balla for residential development, as outlined in Section 12.9.10.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
Charlestown Settlement Plan Policies	CNP 1	Support community-led developments, where appropriate, including the implementation of Community Action Plan for Charlestown (2015-2020) or any amended or superseding Community Futures Plan for the town.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Charlestown Settlement Plan Policies	CNP 2	Promote and market Charlestown as an Enterprise Centre	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Charlestown Settlement Plan Policies	CNP 3	Support the reinstatement of the Western Rail Corridor, in particular actively seek and promote the re-opening of the Claremorris-Galway rail link, as well as a link to Sligo and to safeguard and protect these potential rail links from redevelopment for non-transport related purposes, in order not to preclude their future uses as an operational transportation network.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
Charlestown Settlement Plan Objectives	CNO 1	Revitalise the town centre and encourage the redevelopment of the backland areas to the rear of Church Street and Main Street where it can be demonstrated the development will not have adverse effects on the integrity of the River Moy SAC.	Policies or proposals which steer change in such as way as to protect international nature conservation sites from adverse effects	N/A
Charlestown Settlement Plan Objectives	CNO 2	Support and facilitate the construction of the N17 Knock to Collooney Atlantic Economic Corridor road project which includes the N17 Charlestown By-pass.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Charlestown Settlement Plan Objectives	CNO 3	Protect and enhance existing amenity facilities in Charlestown, particularly the fair green, bowling alley, football pitch and the river area.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Charlestown Settlement Plan Objectives	CNO 4	Protect and enhance areas considered unique and important in the town such as the church, the library, the Arts centre, the Town Hall, the railway line and railway	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		station, Westpoint Business Centre and the health centre.	species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	
Charlestown Settlement Plan Objectives	CNO 5	Secure sites for town centre parking.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Charlestown Settlement Plan Objectives	CNO 6	Protect the Mullaghanoë River as a salmonid status water course, and all water courses and surface waters within the area, their water quality, ecology and function as ecological corridors and as potential influences on the integrity of any Natura 2000 sites within an appropriate radial buffer zone.	Policies or proposals which steer change in such as way as to protect international nature conservation sites from adverse effects	N/A
Charlestown Settlement Plan Objectives	CNO 7	Continue to work and co-operate with Sligo County Council to ensure that a coherent and consistent strategy continues to be applied for the overall development of the Charlestown- Bellaghy Area.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
Charlestown Settlement Plan Objectives	CNO 8	Support local sports and community groups in delivering facilities, including the delivery of a new town park on a site identified adjacent to the Church.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Charlestown Settlement Plan Objectives	CNO 9	Improve and enhance the river walkway within the town and pedestrian linkages within the town.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Charlestown Settlement Plan Objectives	CNO 10	To encourage and facilitate the re-use and regeneration of derelict land and buildings for retail and other town centre uses with due cognisance to the Sequential Approach prescribed in the Retail Planning Guidelines 2012. (Refer to Map CN2)	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Charlestown Settlement Plan Objectives	CNO 11	To support and facilitate the development of the two identified opportunity sites in Charlestown for	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000	Refer to Chapter 9 of this NIR for mitigation measures for any plan

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective		Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red		Mitigation
		residential development, as outlined in Section 12.10.10.	network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	or project arising from this objective/ policy
Crossmolina Settlement Plan Policies	CAP 1	To investigate proposals for the enhancement of the public realm within the town core.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Crossmolina Settlement Plan Policies	CAP 2	Support community-led developments, where appropriate, including the implementation of Community Action Plan for Crossmolina (2015-2020) or any amended or superseding Community Futures Plan for the town.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Crossmolina Settlement Plan Objectives	CAO 1	To retain and enhance the distinctive character and layout of Crossmolina town, which reflects the historical street pattern of growth since the early nineteenth century configuration and to promote this as part of the town's unique identity.	Town enhancement works could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
Crossmolina Settlement Plan Objectives	CAO 2	To investigate the possible designation of built-up areas of special interest and importance in the town, such as an Architectural Conservation Areas (ACA).	General statement of policy/general aspiration which will not lead to development	N/A
Crossmolina Settlement Plan Objectives	CAO 3	To support the local community in exploring the creation of a Cultural and Heritage Centre in the town, as resources permit.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Crossmolina Settlement Plan Objectives	CAO 4	To support and facilitate the full implementation of the Office of Public Work's Flood Relief Scheme for Crossmolina.	Flood relief works could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Crossmolina Settlement Plan Objectives	CAO 5	To sustain, enhance and consolidate the retail and services offer within the core areas of Crossmolina and harness and develop the potential of heritage and tourism assets.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
Crossmolina Settlement Plan Objectives	CAO 6	To encourage and facilitate the re-use and regeneration of derelict land and buildings for retail and other town centre uses with due cognisance to the Sequential Approach prescribed in the Retail Planning Guidelines 2012. (Refer to Map CA2).	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Crossmolina Settlement Plan Objectives	CAO 7	To support and facilitate the development of the four identified opportunity sites in Crossmolina for residential development, as outlined in Section 12.11.10.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Foxford Settlement Plan Policies	FDP 1	To investigate proposals for the enhancement of the public realm within the town core.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Foxford Settlement Plan Policies	FDP 2	Support community-led developments, where appropriate, including the implementation of Community Action Plan for Foxford	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000	Refer to Chapter 9 of this NIR for mitigation measures for any plan

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		(2017-2022) or any amended or superseding Community Futures Plan for the town.	network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	or project arising from this objective/ policy
Foxford Settlement Plan Objectives	FDO 1	To retain and enhance the distinctive character and layout of Foxford town, which reflects the historical street pattern of growth since the early nineteenth century configuration and to promote this as part of the town's unique identity.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Foxford Settlement Plan Objectives	FDO 2	To support the local community in exploring the creation of a river walk in the town, as resources permit and in a manner that will not adversely affect the integrity of the River Moy candidate Special Area of Conservation.	Policies or proposals which steer change in such as way as to protect international nature conservation sites from adverse effects	N/A
Foxford Settlement Plan Objectives	FDO 3	To support the further investigation by the Office of Public Work's of the necessity of flood relief scheme for Foxford.	Flood relief works could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Foxford Settlement Plan Objectives	FDO 4	To sustain, enhance and consolidate the retail and services offer within	This objective could lead to increased development. Any development within the	Refer to Chapter 9 of this NIR for mitigation

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		the core areas of Foxford and harness and develop the potential of heritage and tourism assets.	county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	measures for any plan or project arising from this objective/ policy
Foxford Settlement Plan Objectives	FDO 5	To encourage and facilitate the re-use and regeneration of derelict land and buildings for retail and other town centre uses with due cognisance to the Sequential Approach prescribed in the Retail Planning Guidelines 2012. (Refer to Map FD2).	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Foxford Settlement Plan Objectives	FDO 6	To support and facilitate the development of the five identified opportunity sites in Foxford for residential development, as outlined in Section 12.12.10.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Killala Settlement Plan Policies	KAP 1	To support appropriate enterprise/employment uses that are sympathetic to and supportive of maintaining the architectural character and setting of Killala House (a Protected Structure) and	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		adjacent lands (zoned Enterprise & Employment).	species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	
Killala Settlement Plan Policies	KAP 2	To support the linkage of the Western Way with various local walking/cycling trails in the area including the Great Western Greenway.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Killala Settlement Plan Policies	KAP 3	Support community-led developments, where appropriate, including the implementation of Community Action Plan for Killala (2015-2020) or any amended or superseding Community Futures Plan for the town.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Killala Settlement Plan Objectives	KAO 1	To work with existing service providers and relevant interests to promote Killala as one of the key tourist locations within County Mayo, in accordance with the objectives contained within this Plan.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
Killala Settlement Plan Objectives	KAO 2	To protect and enhance the character of the pier area and to provide for the development of appropriate commercial, tourism, leisure uses and ancillary works such as car parking and navigation.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Killala Settlement Plan Objectives	KAO 3	To promote greater public accessibility to existing tourist attractions in the town.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Killala Settlement Plan Objectives	KAO 4	To investigate the provision of a bypass around the town and to improve traffic management through the provision of the inner relief road and the development of a town centre car park, subject to no significant adverse effects on the environment, including the integrity of the Natura 2000 network.	Policies or proposals which steer change in such as way as to protect international nature conservation sites from adverse effects	N/A
Killala Settlement Plan Objectives	KAO 5	To support and facilitate the provision of a wastewater treatment plant and associated works for the town.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000	Refer to Chapter 9 of this NIR for mitigation measures for any plan

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
			network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	or project arising from this objective/ policy
Killala Settlement Plan Objectives	KAO 6	To ensure that the historic architectural character of Killala is protected and enhanced.	General statement of policy/general aspiration which will not lead to development	N/A
Killala Settlement Plan Objectives	KAO 7	To designate within the lifetime of this plan an Architectural Conservation Area for Killala town centre including Market Street, Courthouse Street, Church Street, Church Lane, William Street, Georges Street, and Ballina Road (R314) and the coastline to the pier.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Killala Settlement Plan Objectives	KAO 8	To protect Killala Bay/Moy Estuary Special Area of Conservation (Site Code 000458) and Special Protection Area (Site Code 004036). Development proposals will be required to demonstrate that the development will not have an adverse effect on the integrity of the sites.	Policies or proposals which steer change in such as way as to protect international nature conservation sites from adverse effects	N/A
Killala Settlement Plan Objectives	KAO 9	To retain the tourist accommodation function of the holiday cottages at Steeple Hill.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
			habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	
Killala Settlement Plan Objectives	KAO 10	To encourage and facilitate the re-use and regeneration of derelict land and buildings for retail and other town centre uses with due cognisance to the Sequential Approach prescribed in the Retail Planning Guidelines 2012. (Refer to Map KA2).	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Killala Settlement Plan Objectives	KAO 11	To support and facilitate the development of the three identified opportunity sites in Killala for residential development, as outlined in Section 12.13.10.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Kiltimagh Settlement Plan Policies	KTP 1	Support the reinstatement of the Western Rail Corridor, in particular actively seek and promote the re-opening of the Claremorris-Galway rail link, as well as a link to Sligo and to safeguard and protect these potential rail links from redevelopment for non-transport related purposes, in order not to	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective		Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red		Mitigation
		preclude their future uses as an operational transportation network.		
Kiltimagh Settlement Plan Policies	KTP 2	Support the work of IRD Kiltimagh Ltd., in providing essential community services for the elderly and marginal groups of Kiltimagh.	General statement of policy/general aspiration which will not lead to development	N/A
Kiltimagh Settlement Plan Policies	KTP 3	Support the creation of an Arts residency programme with revolving art residency spaces.	Policy or proposal that could not have any conceivable effect on a site	N/A
Kiltimagh Settlement Plan Policies	KTP 4	Promote more frequent bus services to Kiltimagh allowing increased connectivity and accessibility for tourists and residents alike.	General statement of policy/general aspiration which will not lead to development	N/A
Kiltimagh Settlement Plan Policies	KTP 5	Support community-led developments, where appropriate, including the implementation of Community Action Plan for Kiltimagh (2019-2024) or any amended or superseding Community Futures Plan for the town.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Kiltimagh Settlement Plan Objectives	KTO 1	Protect and enhance existing amenity facilities in Kiltimagh, particularly the Pollagh River, the playground, amenity park, GAA pitch, wetland park and sculpture trail.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
Kiltimagh Settlement Plan Objectives	KTO 2	Protect existing public sculptures throughout the town and support the provision of additional sculptures at appropriate locations throughout the town.	Policy or proposal that could not have any conceivable effect on a site	N/A
Kiltimagh Settlement Plan Objectives	KTO 3	Support and facilitate pedestrian mobility and safety in the town by introducing traffic calming measures and pedestrian crossings.	General statement of policy/general aspiration which will not lead to development	N/A
Kiltimagh Settlement Plan Objectives	KTO 4	Maintain and further develop walking routes and linkages to the walking routes throughout the town.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Kiltimagh Settlement Plan Objectives	KTO 5	Implement effective vehicular linkages within and around the town, subject to further feasibility studies, detailed design and traffic impact assessment.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Kiltimagh Settlement Plan Objectives	KTO 6	To encourage and facilitate the re-use and regeneration of derelict land and buildings for retail and other town centre uses with due	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000	Refer to Chapter 9 of this NIR for mitigation measures for any plan

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		cognisance to the Sequential Approach prescribed in the Retail Planning Guidelines 2012. (Refer to Map KT2).	network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	or project arising from this objective/ policy
Kiltimagh Settlement Plan Objectives	KTO 7	To support and facilitate the development of the three identified opportunity sites in Kiltimagh for residential development, as outlined in Section 12.14.10.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Knock Settlement Plan Policies	KKP 1	To support and promote the development of Knock as a world-renowned religious tourist destination.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Knock Settlement Plan Policies	KKP 2	To support the provision of amenity walkways with dedicated pedestrian and cycle ways, planting and picnic areas in the town.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
			conservation value, such as changes in water quality and quantity, and air quality.	
Knock Settlement Plan Policies	KKP 3	To promote greater public accessibility to existing tourist attractions in the town.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Knock Settlement Plan Policies	KKP 4	Support community-led developments, where appropriate, including the implementation of Community Action Plan for Knock (2016-2021) or any amended or superseding Community Futures Plan for the town.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Knock Settlement Plan Objectives	KKO 1	To facilitate the development of the town's religious tourism products to the highest international standards, in a manner that respects, builds on, protects and enhances the cultural, built and natural heritage and local amenities of the town.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Knock Settlement Plan Objectives	KKO 2	To work with existing service providers and relevant interests to	This objective could lead to increased development. Any development within the	Refer to Chapter 9 of this NIR for mitigation

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		promote Knock as one of the key tourist locations within County Mayo, in accordance with the objectives contained in this Plan.	county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	measures for any plan or project arising from this objective/ policy
Knock Settlement Plan Objectives	KKO 3	To ensure that all new development makes a positive contribution to the built and natural environment of the Plan area, by ensuring that it is absorbed into the surrounding streetscape/landscape, so that it does not impinge in any significant way on the character, integrity or uniformity of the town and does not undermine or interfere with the iconic status of Knock Basilica.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Knock Settlement Plan Objectives	KKO 4	To safeguard and protect the Knock By-Pass from development for non-infrastructure related purposes.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Knock Settlement Plan Objectives	KKO 5	To encourage and facilitate the re-use and regeneration of derelict land and buildings for retail and other town centre uses with due	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000	Refer to Chapter 9 of this NIR for mitigation measures for any plan

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		cognisance to the Sequential Approach prescribed in the Retail Planning Guidelines 2012. (Refer to Map KK2).	network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	or project arising from this objective/ policy
Knock Settlement Plan Objectives	KKO 6	To support and facilitate the development of the three identified opportunity sites in Knock for residential development, as outlined in Section 12.15.10.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Louisburgh Settlement Plan Policies	LHP 1	Support the promotion of Louisburgh as a key tourist destination in the county and along the Wild Atlantic Way.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Louisburgh Settlement Plan Policies	LHP 2	Support community-led developments, where appropriate, including the implementation of Louisburgh Community Action Plan (2016-2021) or any amended or superseding Community Futures Plan for the town.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
			conservation value, such as changes in water quality and quantity, and air quality.	
Louisburgh Settlement Plan Objectives	LHO 1	To actively work with Irish Water to increase the capacity/upgrade Louisburgh wastewater treatment plant.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Louisburgh Settlement Plan Objectives	LHO 2	Re-establish a village green on Church Street.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Louisburgh Settlement Plan Objectives	LHO 3	Establish a pedestrian link between the town and Carrowmore Beach.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Louisburgh Settlement Plan Objectives	LHO 4	Encourage the development of a Bunowen River walkway and to	This objective could lead to increased development. Any development within the	Refer to Chapter 9 of this NIR for mitigation

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		safeguard the value of the river as an ecological “green corridor”. Riverside walkway provisions should be incorporated into development proposals bounding the river, where appropriate.	county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	measures for any plan or project arising from this objective/ policy
Louisburgh Settlement Plan Objectives	LHO 5	Protect the water quality and riparian zone of the Bunowen River. Any proposed developments adjacent to or close to watercourses shall be carefully assessed to ensure that there is no adverse impact to the water course, its riparian zone or to any waterbody into which it flows.	Policies or proposals which steer change in such as way as to protect international nature conservation sites from adverse effects	N/A
Louisburgh Settlement Plan Objectives	LHO 6	Support and, where possible, implement measures to create interpretative walking routes in and around the town, linking the town’s special features of built and natural heritage interest.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Louisburgh Settlement Plan Objectives	LHO 7	To encourage and facilitate the re-use and regeneration of derelict land and buildings for retail and other town centre uses with due cognisance to the Sequential Approach prescribed in the Retail	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		Planning Guidelines 2012. (Refer to Map LH2).	species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	
Louisburgh Settlement Plan Objectives	LHO 8	To support and facilitate the development of the three identified opportunity sites in Louisburgh for residential development, as outlined in Section 12.16.10.		
Newport Settlement Plan Policies	NTP 1	To have regard to the findings and provisions set out in the Newport Character Study.	General statement of policy/general aspiration which will not lead to development	N/A
Newport Settlement Plan Policies	NTP 2	To support the promotion of Newport as a key tourist destination in the county and along the Wild Atlantic Way, with emphasis on cultural heritage, sailing, angling, fishing and walking/cycling.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Newport Settlement Plan Policies	NTP 3	To promote more frequent bus services to Castlebar/Westport allowing increased connectivity and accessibility for tourists and residents alike.	This objective could lead to increased infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Newport Settlement Plan Policies	NTP 4	To support the favourable conservation status of the	Policies or proposals which steer change in such as way as to protect international	N/A

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		Freshwater Pearl Mussel (<i>Margaritifera margaritifera</i>) population located in Newport River, in accordance with the programme of measures contained in the Newport Sub-basin Management Plan. In this regard, development will only be permitted where it can be demonstrated that the proposed development will not have an adverse effect on the Freshwater Pearl Mussel population or its habitat, by virtue of their conservation objectives.	nature conservation sites from adverse effects therefore positive impacts are anticipated from the implementation of this objective/policy	
Newport Settlement Plan Objectives	NTO 1	To protect and enhance existing amenity areas/facilities in Newport, particularly the river/estuary, the playground, the tennis courts, the sporting pitches, Grace Kelly Amenity Park and the Great Western Greenway.	This objective could lead to increased infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Newport Settlement Plan Objectives	NTO 2	To support and facilitate pedestrian mobility and safety in the town, by introducing traffic calming measures and pedestrian crossings.	This objective could lead to increased infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
			conservation value, such as changes in water quality and quantity, and air quality.	
Newport Settlement Plan Objectives	NTO 3	To identify an appropriate site within the town for the development of a new community hall.	This objective could lead to increased infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Newport Settlement Plan Objectives	NTO 4	To protect the water quality and riparian zone of the Newport River and all water courses and surface waters within the area, their water quality, ecology and function as ecological corridors. In this regard, development will only be permitted where it can be demonstrated that the proposed development will not have an adverse effect on the integrity of the Natura 2000 sites including the Clew Bay Complex (Site Code 001482) and Newport River (Site Code 002144) candidate Special Areas of Conservation.	Protection or improvement to water quality will positively impact on Natura 2000 Sites	N/A
Newport Settlement Plan Objectives	NTO 5	To seek to make Tree Preservation Orders for individual and groups of trees in the town, particularly those located at Newport House, Grace	General statement of policy/general aspiration which will not lead to development	N/A

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		Kelly Amenity Park and along the Newport River.		
Newport Settlement Plan Objectives	NTO 6	To support and facilitate the provision of a wastewater treatment plant and associated works for the town.	This objective could lead to increased infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Newport Settlement Plan Objectives	NTO 7	To encourage and facilitate the re-use and regeneration of derelict land and buildings for retail and other town centre uses with due cognisance to the Sequential Approach prescribed in the Retail Planning Guidelines 2012. (Refer to Map NT2).	This objective could lead to increased infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Newport Settlement Plan Objectives	NTO 8	To support and facilitate the development of the identified opportunity site in Newport for residential development, as outlined in Section 12.17.10.	This objective could lead to increased infrastructure. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
Rural Settlement and Village Settlement Plan Policies	RSVP 1	To promote the development of rural settlements and villages to meet the needs of these established communities and to provide an alternative choice for those seeking to live in a more rural setting, while supporting existing local services and facilities.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Rural Settlement and Village Settlement Plan Policies	RSVP 2	To support the consolidation of Mayo's rural settlements and villages, by promoting proposals that contribute to the sustainable and sequential development of serviceable lands.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Rural Settlement and Village Settlement Plan Policies	RSVP 3	To encourage in-depth residential development in rural settlements and villages, of an appropriate scale, design and density, compatible with the intrinsic character and scale of those settlements/villages.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Rural Settlement and Village Settlement Plan Policies	RSVP 4	To support, promote and encourage the appropriate development of infilling, brownfield or the use of derelict or under-utilised land or	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000	Refer to Chapter 9 of this NIR for mitigation measures for any plan

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective		Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red		Mitigation
		premises, subject to siting, design, protection of residential amenities and normal planning considerations.	network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	or project arising from this objective/ policy
Rural Settlement and Village Settlement Plan Policies	RSVP 5	To encourage the re-use of existing vacant buildings for commercial or residential purposes and the development of infill sites to create compact, vibrant rural settlements and villages.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Rural Settlement and Village Settlement Plan Policies	RSVP 6	To support public realm enhancements in rural settlements and villages, including signage, public lighting (Dark Sky Friendly), public seating, hard and soft landscaping and improvements to the road and footpath network, where appropriate.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Rural Settlement and Village Settlement Plan Policies	RSVP 7	To support rural settlements and villages in their role as local rural service centres for their population and its rural hinterland.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
			conservation value, such as changes in water quality and quantity, and air quality.	
Rural Settlement and Village Settlement Plan Policies	RSVP 8	Support community-led developments in rural settlements/villages, where appropriate, including the implementation of Mayo Community Futures' Community Action Plans.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Rural Settlement and Village Settlement Plan Policies	RSVP 9	To support the development of a "New Homes in Small Towns and villages" initiative which would augment the delivery of actions by Local Authorities, Irish Water, communities and other stakeholders, in the provision of services and serviced sites to create "build your own home" opportunities, within the existing footprint of rural settlements and villages, in order to provide new homes to meet housing demand.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Rural Settlement and Village Settlement Plan Policies	RSVP 10	To liaise and work in conjunction with Irish Water in the delivery of an adequate level of water and wastewater services in rural settlements and villages, including pursuing wastewater treatment upgrades, where appropriate,	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective		Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red		Mitigation
		through Irish Water's Small Towns and Villages Growth Programme.	conservation value, such as changes in water quality and quantity, and air quality.	
Rural Settlement and Village Settlement Plan Policies	RSVP 11	Support the creation of cycling infrastructure within the rural villages and settlements, their hinterlands and at areas of interest and attractions.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Rural Settlement and Village Settlement Objectives	RSVO 1	To ensure that future housing occurs in rural settlements and villages within the settlement/village boundary (based on the sequential approach), where serviced lands are available.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Rural Settlement and Village Settlement Objectives	RSVO 2	To ensure that all rural settlements and villages develop in a self-sufficient manner, utilising existing physical and social infrastructure, where appropriate.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
Rural Settlement and Village Settlement Objectives	RSVO 3	To promote and facilitate residential development commensurate with the nature and scale of the particular rural village or settlement, utilising brownfield and infill opportunities in order to regenerate and consolidate the rural settlements and villages.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Rural Settlement and Village Settlement Objectives	RSVO 4	To support the development of appropriate housing in rural settlements and villages, in order to provide a choice for those who wish to live in a rural setting but not in the rural countryside, subject to a limited scope for individual small-scale multi-house developments of up to 12 houses only or 10% of the existing housing stock, unless it can be demonstrated to the satisfaction of the Planning Authority that local infrastructure, such as schools, community facilities and water services, are sufficiently developed to cater for a larger residential development.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Rural Settlement and Village Settlement Objectives	RSVO 5	To facilitate the expansion of and provision of new mixed-use and employment-generating development within rural settlements and villages at an	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
		appropriate size and scale, subject to normal planning requirements and the “good neighbour” principle.	habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	
Rural Settlement and Village Settlement Objectives	RSVO 6	To seek the improvement, consolidation and expansion of the public lighting and footpath network in rural settlements and village, including a footpath / cycle link, where appropriate and feasible.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Rural Settlement and Village Settlement Objectives	RSVO 7	To facilitate the expansion of the employment and service base in the village	General statement of policy/general aspiration which will not lead to development	N/A
Rural Settlement and Village Settlement Objectives	RSVO 8	To actively support the objectives of the ‘Rebuilding Ireland’ Strategy to address the shortage of housing.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Rural Settlement and Village Settlement Objectives	RSVO 9	To protect groundwater resources within Source Protection Zones.	Policies or proposals which steer change in such as way as to protect international nature conservation sites from adverse effects	N/A

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective			Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red	Mitigation
Rural Settlement and Village Settlement Objectives	RSVO 10	To improve recreational/community/social facilities in rural settlements and villages, where appropriate and as resources allow.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Rural Settlement and Village Settlement Objectives	RSVO 11	To facilitate additional community facilities and services within the rural settlement and village envelope, where possible.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Rural Settlement and Village Settlement Objectives	RSVO 12	To promote and facilitate development that is commensurate with the nature and extent of the existing settlement to support their role as local service centres.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Rural Settlement and Village Settlement Objectives	RSVO 13	To ensure new developments do not adversely impact on the setting and/or integrity of the built or	Policies or proposals which steer change in such as way as to protect international nature conservation sites from adverse effects	N/A

Settlements				
Potential Sources of Impact from the Draft Mayo County Development Plan 2020-2026 policy / objective		Potential Impacts for significant adverse effects on Natura 2000 site(s) No Impacts= Green, Likely impacts = Red		Mitigation
		natural heritage in or adjacent to rural settlements and villages.		
Rural Settlement and Village Settlement Objectives	RSVO 14	To facilitate the provision of gateway features and natural edges on the key approaches to rural settlements and villages.	General statement of policy/general aspiration which will not lead to development	N/A
Rural Settlement and Village Settlement Objectives	RSVO 15	To facilitate public realm improvements in rural settlements and villages, including signage, public seating, hard and soft landscaping and improvements to the road and footpath network, where appropriate and feasible.	This objective could lead to increased development. Any development within the county could potentially have direct and/or indirect impacts on the Natura 2000 network through fragmentation or loss of habitats, disturbance or fragmentation of species, or changes in key indicators of conservation value, such as changes in water quality and quantity, and air quality.	Refer to Chapter 9 of this NIR for mitigation measures for any plan or project arising from this objective/ policy
Rural Settlement and Village Settlement Objectives	RSVO 16	To consider proposals for small scale, clustered residential development in rural settlements and villages that are not serviced by a wastewater treatment plant. Subject to complying with the most up-to-date EPA Code of Practice for Wastewater Treatment and Disposal Systems Serving Single Houses and/or Small Communities Manuals.	Policies or proposals which steer change in such as way as to protect international nature conservation sites from adverse effects	N/A

B Qualifying Interests

Site Code	Site Name	Qualifying Interests
002268	Achill Head SAC	Mudflats and sandflats not covered by seawater at low tide [1140]
		Large shallow inlets and bays [1160]
		Reefs [1170]
000461	Ardkill Turlough SAC	Turloughs [3180]
000463	Balla Turlough SAC	Turloughs [3180]
002081	Ballinafad SAC	Rhinolophus hipposideros (Lesser Horseshoe Bat) [1303]
000474	Ballymaglancy Cave, Cong SAC	Rhinolophus hipposideros (Lesser Horseshoe Bat) [1303]
		Caves not open to the public [8310]
001922	Bellacorick Bog Complex SAC	Natural dystrophic lakes and ponds [3160]
		Northern Atlantic wet heaths with Erica tetralix [4010]
		Blanket bogs (* if active bog) [7130]
		Depressions on peat substrates of the Rhynchosporion [7150]
		Alkaline fens [7230]
		Vertigo geyeri (Geyer's Whorl Snail) [1013]
		Saxifraga hirculus (Marsh Saxifrage) [1528]
000466	Bellacorick Iron Flush SAC	Saxifraga hirculus (Marsh Saxifrage) [1528]
002005	Bellacragher Saltmarsh SAC	Atlantic salt meadows (Glaucopuccinellietalia maritimae) [1330]
		Mediterranean salt meadows (Juncetalia maritimi) [1410]
004177	Bills Rocks SPA	Storm Petrel (Hydrobates pelagicus) [A014]
		Puffin (Fratercula arctica) [A204]

Site Code	Site Name	Qualifying Interests
004037	Blacksod Bay/Broad Haven SPA	Red-throated Diver (<i>Gavia stellata</i>) [A001]
		Great Northern Diver (<i>Gavia immer</i>) [A003]
		Slavonian Grebe (<i>Podiceps auritus</i>) [A007]
		Light-bellied Brent Goose (<i>Branta bernicla hrota</i>) [A046]
		Common Scoter (<i>Melanitta nigra</i>) [A065]
		Red-breasted Merganser (<i>Mergus serrator</i>) [A069]
		Ringed Plover (<i>Charadrius hiaticula</i>) [A137]
		Sanderling (<i>Calidris alba</i>) [A144]
		Dunlin (<i>Calidris alpina</i>) [A149]
		Bar-tailed Godwit (<i>Limosa lapponica</i>) [A157]
		Curlew (<i>Numenius arquata</i>) [A160]
		Sandwich Tern (<i>Sterna sandvicensis</i>) [A191]
		Dunlin (<i>Calidris alpina schinzii</i>) [A466]
Wetland and Waterbirds [A999]		
000471	Brackloon Woods SAC	Old sessile oak woods with <i>Ilex</i> and <i>Blechnum</i> in the British Isles [91A0]
000472	Broadhaven Bay SAC	Mudflats and sandflats not covered by seawater at low tide [1140]
		Large shallow inlets and bays [1160]
		Reefs [1170]
		Atlantic salt meadows (<i>Glaucopuccinellietalia maritimae</i>) [1330]
		Submerged or partially submerged sea caves [8330]
000475	Carrowkeel Turlough SAC	Turloughs [3180]

Site Code	Site Name	Qualifying Interests
000476	Carrowmore Lake Complex SAC	Blanket bogs (* if active bog) [7130]
		Depressions on peat substrates of the Rhynchosporion [7150]
		Saxifraga hirculus (Marsh Saxifrage) [1528]
		Hamatocaulis vernicosus (Slender Green Feather-moss) [6216]
004052	Carrowmore Lake SPA	Sandwich Tern (<i>Sterna sandvicensis</i>) [A191]
002243	Clare Island Cliffs SAC	Vegetated sea cliffs of the Atlantic and Baltic coasts [1230]
		Calcareous rocky slopes with chasmophytic vegetation [8210]
		Siliceous rocky slopes with chasmophytic vegetation [8220]
004136	Clare Island SPA	Fulmar (<i>Fulmarus glacialis</i>) [A009]
		Shag (<i>Phalacrocorax aristotelis</i>) [A018]
		Common Gull (<i>Larus canus</i>) [A182]
		Kittiwake (<i>Rissa tridactyla</i>) [A188]
		Guillemot (<i>Uria aalge</i>) [A199]
		Razorbill (<i>Alca torda</i>) [A200]
		Chough (<i>Pyrhocorax pyrrhocorax</i>) [A346]
001482	Clew Bay Complex SAC	Mudflats and sandflats not covered by seawater at low tide [1140]
		Coastal lagoons [1150]
		Large shallow inlets and bays [1160]
		Annual vegetation of drift lines [1210]
		Perennial vegetation of stony banks [1220]

Site Code	Site Name	Qualifying Interests
		Atlantic salt meadows (<i>Glauco-Puccinellietalia maritimae</i>) [1330]
		Embryonic shifting dunes [2110]
		Shifting dunes along the shoreline with <i>Ammophila arenaria</i> (white dunes) [2120]
		Machairs (* in Ireland) [21A0]
		Old sessile oak woods with <i>Ilex</i> and <i>Blechnum</i> in the British Isles [91A0]
		<i>Vertigo geyeri</i> (Geyer's Whorl Snail) [1013]
		<i>Lutra lutra</i> (Otter) [1355]
		<i>Phoca vitulina</i> (Harbour Seal) [1365]
001899	Cloonakillina Lough SAC	Transition mires and quaking bogs [7140]
000479	Cloughmoyne SAC	Limestone pavements [8240]
000480	Clyard Kettle-Holes SAC	Turloughs [3180]
		Calcareous fens with <i>Cladium mariscus</i> and species of the <i>Caricion davallianae</i> [7210]
000485	Corraun Plateau SAC	Northern Atlantic wet heaths with <i>Erica tetralix</i> [4010]
		European dry heaths [4030]
		Alpine and Boreal heaths [4060]
		<i>Juniperus communis</i> formations on heaths or calcareous grasslands [5130]
		Siliceous scree of the montane to snow levels (<i>Androsacetalia alpinae</i> and <i>Galeopsietalia ladani</i>) [8110]
		Siliceous rocky slopes with chasmophytic vegetation [8220]
001955	Croaghaun/Slievemore SAC	Northern Atlantic wet heaths with <i>Erica tetralix</i> [4010]
		European dry heaths [4030]

Site Code	Site Name	Qualifying Interests
		Alpine and Boreal heaths [4060]
		Siliceous scree of the montane to snow levels (<i>Androsacetalia alpinae</i> and <i>Galeopsietalia ladani</i>) [8110]
		Siliceous rocky slopes with chasmophytic vegetation [8220]
000484	Cross Lough (Killadoon) SAC	Perennial vegetation of stony banks [1220]
004212	Cross Lough (Killadoon) SPA	Sandwich Tern (<i>Sterna sandvicensis</i>) [A191]
000492	Doocastle Turlough SAC	Turloughs [3180]
004235	Doogort Machair SPA	Dunlin (<i>Calidris alpina schinzii</i>) [A466]
001497	Doogort Machair/Lough Doo SAC	Machairs (* in Ireland) [21A0]
		Petalophyllum ralfsii (Petalwort) [1395]
000495	Duvillaun Islands SAC	Tursiops truncatus (Common Bottlenose Dolphin) [1349]
		Halichoerus grypus (Grey Seal) [1364]
004111	Duvillaun Islands SPA	Fulmar (<i>Fulmarus glacialis</i>) [A009]
		Storm Petrel (<i>Hydrobates pelagicus</i>) [A014]
		Barnacle Goose (<i>Branta leucopsis</i>) [A045]
001501	Erris Head SAC	Vegetated sea cliffs of the Atlantic and Baltic coasts [1230]
		Alpine and Boreal heaths [4060]
000497	Flughany Bog SAC	Active raised bogs [7110]
		Degraded raised bogs still capable of natural regeneration [7120]
		Depressions on peat substrates of the Rhynchosporion [7150]

Site Code	Site Name	Qualifying Interests
000500	Glenamoy Bog Complex SAC	Vegetated sea cliffs of the Atlantic and Baltic coasts [1230]
		Machairs (* in Ireland) [21A0]
		Natural dystrophic lakes and ponds [3160]
		Northern Atlantic wet heaths with <i>Erica tetralix</i> [4010]
		<i>Juniperus communis</i> formations on heaths or calcareous grasslands [5130]
		Blanket bogs (* if active bog) [7130]
		Transition mires and quaking bogs [7140]
		Depressions on peat substrates of the <i>Rhynchosporion</i> [7150]
		<i>Salmo salar</i> (Salmon) [1106]
		<i>Petalophyllum ralfsii</i> (Petalwort) [1395]
		<i>Saxifraga hirculus</i> (Marsh Saxifrage) [1528]
		<i>Hamatocaulis vernicosus</i> (Slender Green Feather-moss) [6216]
000503	Greaghans Turlough SAC	Turloughs [3180]
004074	Illanmaster SPA	Storm Petrel (<i>Hydrobates pelagicus</i>) [A014]
004084	Inishglora and Inishkeeragh SPA	Storm Petrel (<i>Hydrobates pelagicus</i>) [A014]
		Cormorant (<i>Phalacrocorax carbo</i>) [A017]
		Shag (<i>Phalacrocorax aristotelis</i>) [A018]
		Barnacle Goose (<i>Branta leucopsis</i>) [A045]
		Lesser Black-backed Gull (<i>Larus fuscus</i>) [A183]
		Herring Gull (<i>Larus argentatus</i>) [A184]

Site Code	Site Name	Qualifying Interests
		Arctic Tern (<i>Sterna paradisaea</i>) [A194]
000507	Inishkea Islands SAC	Machairs (* in Ireland) [21A0]
		Halichoerus grypus (Grey Seal) [1364]
		Petalophyllum ralfsii (Petalwort) [1395]
004004	Inishkea Islands SPA	Shag (<i>Phalacrocorax aristotelis</i>) [A018]
		Barnacle Goose (<i>Branta leucopsis</i>) [A045]
		Ringed Plover (<i>Charadrius hiaticula</i>) [A137]
		Sanderling (<i>Calidris alba</i>) [A144]
		Purple Sandpiper (<i>Calidris maritima</i>) [A148]
		Turnstone (<i>Arenaria interpres</i>) [A169]
		Common Gull (<i>Larus canus</i>) [A182]
		Herring Gull (<i>Larus argentatus</i>) [A184]
		Arctic Tern (<i>Sterna paradisaea</i>) [A194]
		Little Tern (<i>Sterna albifrons</i>) [A195]
		Dunlin (<i>Calidris alpina schinzii</i>) [A466]
001513	Keel Machair/Menaun Cliffs SAC	Perennial vegetation of stony banks [1220]
		Machairs (* in Ireland) [21A0]
		Alpine and Boreal heaths [4060]
		Petalophyllum ralfsii (Petalwort) [1395]
002320	Kildun Souterrain SAC	Rhinolophus hipposideros (Lesser Horseshoe Bat) [1303]
000504	Kilglassan/Cahevavoostia Turlough Complex SAC	Turloughs [3180]
000458	Killala Bay/Moy Estuary SAC	Estuaries [1130]

Site Code	Site Name	Qualifying Interests
		Mudflats and sandflats not covered by seawater at low tide [1140]
		Annual vegetation of drift lines [1210]
		Vegetated sea cliffs of the Atlantic and Baltic coasts [1230]
		Salicornia and other annuals colonising mud and sand [1310]
		Atlantic salt meadows (<i>Glaucopuccinellietalia maritimae</i>) [1330]
		Embryonic shifting dunes [2110]
		Shifting dunes along the shoreline with <i>Ammophila arenaria</i> (white dunes) [2120]
		Fixed coastal dunes with herbaceous vegetation (grey dunes) [2130]
		Humid dune slacks [2190]
		<i>Vertigo angustior</i> (Narrow-mouthed Whorl Snail) [1014]
		<i>Petromyzon marinus</i> (Sea Lamprey) [1095]
		<i>Phoca vitulina</i> (Harbour Seal) [1365]
004036	Killala Bay/Moy Estuary SPA	Ringed Plover (<i>Charadrius hiaticula</i>) [A137]
		Golden Plover (<i>Pluvialis apricaria</i>) [A140]
		Grey Plover (<i>Pluvialis squatarola</i>) [A141]
		Sanderling (<i>Calidris alba</i>) [A144]
		Dunlin (<i>Calidris alpina</i>) [A149]
		Bar-tailed Godwit (<i>Limosa lapponica</i>) [A157]
		Curlew (<i>Numenius arquata</i>) [A160]

Site Code	Site Name	Qualifying Interests
		Redshank (<i>Tringa totanus</i>) [A162]
		Wetland and Waterbirds [A999]
000516	Lackan Saltmarsh and Kilcummin Head SAC	Salicornia and other annuals colonising mud and sand [1310]
		Atlantic salt meadows (<i>Glaucopuccinellietalia maritima</i>) [1330]
		Mediterranean salt meadows (<i>Juncetalia maritimi</i>) [1410]
		Shifting dunes along the shoreline with <i>Ammophila arenaria</i> (white dunes) [2120]
		Fixed coastal dunes with herbaceous vegetation (grey dunes) [2130]
001529	Lough Cahasy, Lough Baun And Roonah Lough SAC	Coastal lagoons [1150]
		Perennial vegetation of stony banks [1220]
		Embryonic shifting dunes [2110]
		Shifting dunes along the shoreline with <i>Ammophila arenaria</i> (white dunes) [2120]
		Fixed coastal dunes with herbaceous vegetation (grey dunes) [2130]
		Machairs (* in Ireland) [21A0]
004051	Lough Carra SPA	Common Gull (<i>Larus canus</i>) [A182]
001774	Lough Carra/Mask Complex SAC	Oligotrophic waters containing very few minerals of sandy plains (<i>Littorelletalia uniflorae</i>) [3110]
		Oligotrophic to mesotrophic standing waters with vegetation of the <i>Littorelletea uniflorae</i> and/or <i>Isoeto-Nanojuncetea</i> [3130]

Site Code	Site Name	Qualifying Interests
		<p>Hard oligo-mesotrophic waters with benthic vegetation of Chara spp. [3140]</p> <p>European dry heaths [4030]</p> <p>Semi-natural dry grasslands and scrubland facies on calcareous substrates (Festuco-Brometalia) (* important orchid sites) [6210]</p> <p>Calcareous fens with Cladium mariscus and species of the Caricion davallianae [7210]</p> <p>Alkaline fens [7230]</p> <p>Limestone pavements [8240]</p> <p>Alluvial forests with Alnus glutinosa and Fraxinus excelsior (Alno-Padion, Alnion incanae, Salicion albae) [91E0]</p> <p>Rhinolophus hipposideros (Lesser Horseshoe Bat) [1303]</p> <p>Lutra lutra (Otter) [1355]</p> <p>Hamatocaulis vernicosus (Slender Green Feather-moss) [6216]</p>
004228	Lough Conn and Lough Cullin SPA	<p>Tufted Duck (Aythya fuligula) [A061]</p> <p>Common Scoter (Melanitta nigra) [A065]</p> <p>Common Gull (Larus canus) [A182]</p> <p>Greenland White-fronted Goose (Anser albifrons flavirostris) [A395]</p> <p>Wetland and Waterbirds [A999]</p>
000297	Lough Corrib SAC	<p>Oligotrophic waters containing very few minerals of sandy plains (Littorelletalia uniflorae) [3110]</p>

Site Code	Site Name	Qualifying Interests
		Oligotrophic to mesotrophic standing waters with vegetation of the Littorelletea uniflorae and/or Isoeto-Nanojuncetea [3130]
		Hard oligo-mesotrophic waters with benthic vegetation of Chara spp. [3140]
		Water courses of plain to montane levels with the Ranunculion fluitantis and Callitriche-Batrachion vegetation [3260]
		Semi-natural dry grasslands and scrubland facies on calcareous substrates (Festuco-Brometalia) (* important orchid sites) [6210]
		Molinia meadows on calcareous, peaty or clayey-silt-laden soils (Molinion caeruleae) [6410]
		Active raised bogs [7110]
		Degraded raised bogs still capable of natural regeneration [7120]
		Depressions on peat substrates of the Rhynchosporion [7150]
		Calcareous fens with Cladium mariscus and species of the Caricion davallianae [7210]
		Petrifying springs with tufa formation (Cratoneurion) [7220]
		Alkaline fens [7230]
		Limestone pavements [8240]
		Old sessile oak woods with Ilex and Blechnum in the British Isles [91A0]
		Bog woodland [91D0]

Site Code	Site Name	Qualifying Interests
		<p>Margaritifera margaritifera (Freshwater Pearl Mussel) [1029]</p> <p>Austropotamobius pallipes (White-clawed Crayfish) [1092]</p> <p>Petromyzon marinus (Sea Lamprey) [1095]</p> <p>Lampetra planeri (Brook Lamprey) [1096]</p> <p>Salmo salar (Salmon) [1106]</p> <p>Rhinolophus hipposideros (Lesser Horseshoe Bat) [1303]</p> <p>Lutra lutra (Otter) [1355]</p> <p>Najas flexilis (Slender Naiad) [1833]</p> <p>Hamatocaulis vernicosus (Slender Green Feather-moss) [6216]</p>
004042	Lough Corrib SPA	<p>Gadwall (Anas strepera) [A051]</p> <p>Shoveler (Anas clypeata) [A056]</p> <p>Pochard (Aythya ferina) [A059]</p> <p>Tufted Duck (Aythya fuligula) [A061]</p> <p>Common Scoter (Melanitta nigra) [A065]</p> <p>Hen Harrier (Circus cyaneus) [A082]</p> <p>Coot (Fulica atra) [A125]</p> <p>Golden Plover (Pluvialis apricaria) [A140]</p> <p>Black-headed Gull (Chroicocephalus ridibundus) [A179]</p> <p>Common Gull (Larus canus) [A182]</p> <p>Common Tern (Sterna hirundo) [A193]</p> <p>Arctic Tern (Sterna paradisaea) [A194]</p>

Site Code	Site Name	Qualifying Interests
		Greenland White-fronted Goose (<i>Anser albifrons flavirostris</i>) [A395]
		Wetland and Waterbirds [A999]
002177	Lough Dahybaun SAC	<i>Najas flexilis</i> (Slender Naiad) [1833]
000522	Lough Gall Bog SAC	Blanket bogs (* if active bog) [7130]
		Depressions on peat substrates of the <i>Rhynchosporion</i> [7150]
000633	Lough Hoe Bog SAC	Oligotrophic waters containing very few minerals of sandy plains (<i>Littorelletalia uniflorae</i>) [3110]
		Blanket bogs (* if active bog) [7130]
		<i>Vertigo geyeri</i> (Geyer's Whorl Snail) [1013]
		<i>Austropotamobius pallipes</i> (White-clawed Crayfish) [1092]
004062	Lough Mask SPA	Tufted Duck (<i>Aythya fuligula</i>) [A061]
		Black-headed Gull (<i>Chroicocephalus ridibundus</i>) [A179]
		Common Gull (<i>Larus canus</i>) [A182]
		Lesser Black-backed Gull (<i>Larus fuscus</i>) [A183]
		Common Tern (<i>Sterna hirundo</i>) [A193]
		Greenland White-fronted Goose (<i>Anser albifrons flavirostris</i>) [A395]
		Wetland and Waterbirds [A999]
001536	Mocorha Lough SAC	Calcareous fens with <i>Cladium mariscus</i> and species of the <i>Caricion davallianae</i> [7210]

Site Code	Site Name	Qualifying Interests
000527	Moore Hall (Lough Carra) SAC	Rhinolophus hipposideros (Lesser Horseshoe Bat) [1303]
004227	Mullet Peninsula SPA	Corncrake (Crex crex) [A122]
000470	Mullet/Blacksod Bay Complex SAC	Mudflats and sandflats not covered by seawater at low tide [1140]
		Large shallow inlets and bays [1160]
		Reefs [1170]
		Salicornia and other annuals colonising mud and sand [1310]
		Shifting dunes along the shoreline with Ammophila arenaria (white dunes) [2120]
		Fixed coastal dunes with herbaceous vegetation (grey dunes) [2130]
		Atlantic decalcified fixed dunes (Calluno-Ulicetea) [2150]
		Machairs (* in Ireland) [21A0]
		Natural eutrophic lakes with Magnopotamion or Hydrocharition - type vegetation [3150]
		Alkaline fens [7230]
		Lutra lutra (Otter) [1355]
Petalophyllum ralfsii (Petalwort) [1395]		
001932	Mweelrea/Sheeffry/Erriff Complex SAC	Coastal lagoons [1150]
		Annual vegetation of drift lines [1210]
		Atlantic salt meadows (Glaucopuccinellietalia maritimae) [1330]
		Mediterranean salt meadows (Juncetalia maritimi) [1410]
		Embryonic shifting dunes [2110]

Site Code	Site Name	Qualifying Interests
		Shifting dunes along the shoreline with <i>Ammophila arenaria</i> (white dunes) [2120]
		Atlantic decalcified fixed dunes (Calluno-Ulicetea) [2150]
		Dunes with <i>Salix repens</i> ssp. <i>argentea</i> (<i>Salicion arenariae</i>) [2170]
		Machairs (* in Ireland) [21A0]
		Oligotrophic waters containing very few minerals of sandy plains (<i>Littorelletalia uniflorae</i>) [3110]
		Oligotrophic to mesotrophic standing waters with vegetation of the <i>Littorelletea uniflorae</i> and/or <i>Isoeto-Nanojuncetea</i> [3130]
		Natural dystrophic lakes and ponds [3160]
		Water courses of plain to montane levels with the <i>Ranunculion fluitantis</i> and <i>Callitriche-Batrachion</i> vegetation [3260]
		Northern Atlantic wet heaths with <i>Erica tetralix</i> [4010]
		European dry heaths [4030]
		Alpine and Boreal heaths [4060]
		Juniperus communis formations on heaths or calcareous grasslands [5130]
		Hydrophilous tall herb fringe communities of plains and of the montane to alpine levels [6430]
		Blanket bogs (* if active bog) [7130]
		Transition mires and quaking bogs [7140]

Site Code	Site Name	Qualifying Interests
		Depressions on peat substrates of the Rhynchosporion [7150]
		Petrifying springs with tufa formation (Cratoneurion) [7220]
		Alkaline fens [7230]
		Siliceous scree of the montane to snow levels (Androsacetalia alpinae and Galeopsietalia ladani) [8110]
		Calcareous rocky slopes with chasmophytic vegetation [8210]
		Siliceous rocky slopes with chasmophytic vegetation [8220]
		Vertigo geyeri (Geyer's Whorl Snail) [1013]
		Vertigo angustior (Narrow-mouthed Whorl Snail) [1014]
		Margaritifera margaritifera (Freshwater Pearl Mussel) [1029]
		Salmo salar (Salmon) [1106]
		Lutra lutra (Otter) [1355]
		Petalophyllum ralfsii (Petalwort) [1395]
		Najas flexilis (Slender Naiad) [1833]
002144	Newport River SAC	Margaritifera margaritifera (Freshwater Pearl Mussel) [1029]
		Salmo salar (Salmon) [1106]
000532	Oldhead Wood SAC	European dry heaths [4030]
		Old sessile oak woods with Ilex and Blechnum in the British Isles [91A0]
000534	Owenduff/Nepin Complex SAC	Oligotrophic waters containing very few minerals of sandy plains (Littorelletalia uniflorae) [3110]

Site Code	Site Name	Qualifying Interests
		<p>Natural dystrophic lakes and ponds [3160]</p> <p>Water courses of plain to montane levels with the <i>Ranunculus fluitans</i> and <i>Callitriche-Batrachion</i> vegetation [3260]</p> <p>Northern Atlantic wet heaths with <i>Erica tetralix</i> [4010]</p> <p>Alpine and Boreal heaths [4060]</p> <p><i>Juniperus communis</i> formations on heaths or calcareous grasslands [5130]</p> <p>Blanket bogs (* if active bog) [7130]</p> <p>Transition mires and quaking bogs [7140]</p> <p><i>Salmo salar</i> (Salmon) [1106]</p> <p><i>Lutra lutra</i> (Otter) [1355]</p> <p><i>Saxifraga hirculus</i> (Marsh Saxifrage) [1528]</p> <p><i>Hamatocaulis vernicosus</i> (Slender Green Feather-moss) [6216]</p>
004098	Owenduff/Nephin Complex SPA	<p>Merlin (<i>Falco columbarius</i>) [A098]</p> <p>Golden Plover (<i>Pluvialis apricaria</i>) [A140]</p>
002006	Ox Mountains Bogs SAC	<p>Oligotrophic waters containing very few minerals of sandy plains (<i>Littorelletalia uniflorae</i>) [3110]</p> <p>Natural dystrophic lakes and ponds [3160]</p> <p>Northern Atlantic wet heaths with <i>Erica tetralix</i> [4010]</p> <p>European dry heaths [4030]</p> <p>Blanket bogs (* if active bog) [7130]</p> <p>Transition mires and quaking bogs [7140]</p>

Site Code	Site Name	Qualifying Interests
		Depressions on peat substrates of the Rhynchosporion [7150]
		Vertigo geyeri (Geyer's Whorl Snail) [1013]
		Saxifraga hirculus (Marsh Saxifrage) [1528]
002298	River Moy SAC	Active raised bogs [7110]
		Degraded raised bogs still capable of natural regeneration [7120]
		Depressions on peat substrates of the Rhynchosporion [7150]
		Alkaline fens [7230]
		Old sessile oak woods with Ilex and Blechnum in the British Isles [91A0]
		Alluvial forests with Alnus glutinosa and Fraxinus excelsior (Alno-Padion, Alnion incanae, Salicion albae) [91E0]
		Austropotamobius pallipes (White-clawed Crayfish) [1092]
		Petromyzon marinus (Sea Lamprey) [1095]
		Lampetra planeri (Brook Lamprey) [1096]
		Salmo salar (Salmon) [1106]
		Lutra lutra (Otter) [1355]
000525	Shrule Turlough SAC	Turloughs [3180]
000541	Skealaghan Turlough SAC	Turloughs [3180]
000542	Slieve Fyagh Bog SAC	Blanket bogs (* if active bog) [7130]
004072	Stags of Broad Haven SPA	Storm Petrel (Hydrobates pelagicus) [A014]
		Leach's Storm-petrel (Oceanodroma leucorhoa) [A015]
004093	Termoncarragh Lake and Annagh Machair SPA	Whooper Swan (Cygnus cygnus) [A038]

Site Code	Site Name	Qualifying Interests
		Barnacle Goose (<i>Branta leucopsis</i>) [A045]
		Corncrake (<i>Crex crex</i>) [A122]
		Lapwing (<i>Vanellus vanellus</i>) [A142]
		Chough (<i>Pyrhocorax pyrrhocorax</i>) [A346]
		Greenland White-fronted Goose (<i>Anser albifrons flavirostris</i>) [A395]
		Dunlin (<i>Calidris alpina schinzii</i>) [A466]
		Wetland and Waterbirds [A999]
002179	Towerhill House SAC	Rhinolophus hipposideros (Lesser Horseshoe Bat) [1303]
001571	Urlaur Lakes SAC	Hard oligo-mesotrophic waters with benthic vegetation of <i>Chara</i> spp. [3140]
002998	West Connacht Coast SAC	Tursiops truncatus (Common Bottlenose Dolphin) [1349]
000614	Cloonshanville Bog SAC	Active raised bogs [7110]
		Degraded raised bogs still capable of natural regeneration [7120]
		Depressions on peat substrates of the Rhynchosporion [7150]
		Bog woodland [91D0]
002034	Connemara Bog Complex SAC	Coastal lagoons [1150]
		Reefs [1170]
		Oligotrophic waters containing very few minerals of sandy plains (<i>Littorelletalia uniflorae</i>) [3110]
		Oligotrophic to mesotrophic standing waters with vegetation of the <i>Littorelletea uniflorae</i> and/or <i>Isoeto-Nanojuncetea</i> [3130]
		Natural dystrophic lakes and ponds [3160]

Site Code	Site Name	Qualifying Interests
		Water courses of plain to montane levels with the <i>Ranunculus fluitans</i> and <i>Callitriche-Batrachion</i> vegetation [3260]
		Northern Atlantic wet heaths with <i>Erica tetralix</i> [4010]
		European dry heaths [4030]
		<i>Molinia</i> meadows on calcareous, peaty or clayey-silt-laden soils (<i>Molinia caerulea</i>) [6410]
		Blanket bogs (* if active bog) [7130]
		Transition mires and quaking bogs [7140]
		Depressions on peat substrates of the <i>Rhynchosporion</i> [7150]
		Alkaline fens [7230]
		Old sessile oak woods with <i>Ilex</i> and <i>Blechnum</i> in the British Isles [91A0]
		<i>Euphydryas aurinia</i> (Marsh Fritillary) [1065]
		<i>Salmo salar</i> (Salmon) [1106]
		<i>Lutra lutra</i> (Otter) [1355]
		<i>Najas flexilis</i> (Slender Naiad) [1833]
004181	Connemara Bog Complex SPA	Cormorant (<i>Phalacrocorax carbo</i>) [A017]
		Merlin (<i>Falco columbarius</i>) [A098]
		Golden Plover (<i>Pluvialis apricaria</i>) [A140]
		Common Gull (<i>Larus canus</i>) [A182]
000296	Lisnageeragh Bog and Ballinastack Turlough SAC	Turloughs [3180]
		Active raised bogs [7110]
		Degraded raised bogs still capable of natural regeneration [7120]

Site Code	Site Name	Qualifying Interests
		Depressions on peat substrates of the Rhynchosporion [7150]
000285	Kilsallagh Bog SAC	Active raised bogs [7110]
		Degraded raised bogs still capable of natural regeneration [7120]
		Depressions on peat substrates of the Rhynchosporion [7150]
001669	Knockalongy and Knockachree Cliffs SAC	Trichomanes speciosum (Killarney Fern) [1421]
001898	Unshin River SAC	Water courses of plain to montane levels with the Ranunculion fluitantis and Callitriche-Batrachion vegetation [3260]
		Semi-natural dry grasslands and scrubland facies on calcareous substrates (Festuco-Brometalia) (* important orchid sites) [6210]
		Molinia meadows on calcareous, peaty or clayey-silt-laden soils (Molinion caeruleae) [6410]
		Alluvial forests with Alnus glutinosa and Fraxinus excelsior (Alno-Padion, Alnion incanae, Salicion albae) [91E0]
		Salmo salar (Salmon) [1106]
		Lutra lutra (Otter) [1355]
000592	Bellanagare Bog SAC	Active raised bogs [7110]
		Degraded raised bogs still capable of natural regeneration [7120]
		Depressions on peat substrates of the Rhynchosporion [7150]
004144	High Island, Inishshark and Davillaun SPA	Fulmar (Fulmarus glacialis) [A009]

Site Code	Site Name	Qualifying Interests
		Barnacle Goose (<i>Branta leucopsis</i>) [A045]
		Arctic Tern (<i>Sterna paradisaea</i>) [A194]
004105	Bellanagare Bog SPA	Greenland White-fronted Goose (<i>Anser albifrons flavirostris</i>) [A395]
001656	Bricklieve Mountains and Keishcorran SAC	Turloughs [3180]
		Semi-natural dry grasslands and scrubland facies on calcareous substrates (<i>Festuco-Brometalia</i>) (* important orchid sites) [6210]
		Lowland hay meadows (<i>Alopecurus pratensis</i> , <i>Sanguisorba officinalis</i>) [6510]
		Calcareous and calcshist screes of the montane to alpine levels (<i>Thlaspietea rotundifolii</i>) [8120]
		<i>Euphydryas aurinia</i> (Marsh Fritillary) [1065]
		<i>Austropotamobius pallipes</i> (White-clawed Crayfish) [1092]
004221	Illaunnaon SPA	Sandwich Tern (<i>Sterna sandvicensis</i>) [A191]
002110	Corliskea/Trien/Cloonfelliv Bog SAC	Active raised bogs [7110]
		Degraded raised bogs still capable of natural regeneration [7120]
		Depressions on peat substrates of the <i>Rhynchosporion</i> [7150]
		Bog woodland [91D0]
002130	Tully Lough SAC	Oligotrophic to mesotrophic standing waters with vegetation of the <i>Littorelletea uniflorae</i> and/or <i>Isoeto-Nanojuncetea</i> [3130]
		<i>Najas flexilis</i> (Slender Naiad) [1833]

Site Code	Site Name	Qualifying Interests
001311	Rusheenduff Lough SAC	Oligotrophic to mesotrophic standing waters with vegetation of the Littorelletea uniflorae and/or Isoeto-Nanojuncetea [3130]
		Najas flexilis (Slender Naiad) [1833]
000255	Croaghill Turlough SAC	Turloughs [3180]
000278	Inishbofin And Inishshark SAC	Coastal lagoons [1150]
		Oligotrophic waters containing very few minerals of sandy plains (Littorelletalia uniflorae) [3110]
		Northern Atlantic wet heaths with Erica tetralix [4010]
		European dry heaths [4030]
		Halichoerus grypus (Grey Seal) [1364]
002296	Williamstown Turloughs SAC	Turloughs [3180]
000595	Callow Bog SAC	Active raised bogs [7110]
		Degraded raised bogs still capable of natural regeneration [7120]
		Depressions on peat substrates of the Rhynchosporion [7150]
000218	Coolcam Turlough SAC	Turloughs [3180]
004048	Lough Gara SPA	Whooper Swan (Cygnus cygnus) [A038]
		Greenland White-fronted Goose (Anser albifrons flavirostris) [A395]
002354	Tullaghanrock Bog SAC	Active raised bogs [7110]
		Degraded raised bogs still capable of natural regeneration [7120]
		Depressions on peat substrates of the Rhynchosporion [7150]
000600	Cloonchambers Bog SAC	Active raised bogs [7110]
		Degraded raised bogs still capable of natural regeneration [7120]

Site Code	Site Name	Qualifying Interests
		Depressions on peat substrates of the Rhynchosporion [7150]
000636	Templehouse And Cloonacleigha Loughs SAC	Hard oligo-mesotrophic waters with benthic vegetation of Chara spp. [3140]
		Water courses of plain to montane levels with the Ranunculion fluitantis and Callitriche-Batrachion vegetation [3260]
002338	Drumalough Bog SAC	Active raised bogs [7110]
		Degraded raised bogs still capable of natural regeneration [7120]
		Depressions on peat substrates of the Rhynchosporion [7150]
000637	Turloughmore (Sligo) SAC	Turloughs [3180]
000634	Lough Nabrickkeagh Bog SAC	Blanket bogs (* if active bog) [7130]
000597	Carrowbehy/Caher Bog SAC	Active raised bogs [7110]
		Degraded raised bogs still capable of natural regeneration [7120]
		Depressions on peat substrates of the Rhynchosporion [7150]
002031	The Twelve Bens/Garraun Complex SAC	Oligotrophic waters containing very few minerals of sandy plains (Littorelletalia uniflorae) [3110]
		Oligotrophic to mesotrophic standing waters with vegetation of the Littorelletea uniflorae and/or Isoeto-Nanojuncetea [3130]
		Alpine and Boreal heaths [4060]
		Blanket bogs (* if active bog) [7130]

Site Code	Site Name	Qualifying Interests
		Depressions on peat substrates of the Rhynchosporion [7150]
		Siliceous scree of the montane to snow levels (<i>Androsacetalia alpinae</i> and <i>Galeopsietalia ladani</i>) [8110]
		Calcareous rocky slopes with chasmophytic vegetation [8210]
		Siliceous rocky slopes with chasmophytic vegetation [8220]
		Old sessile oak woods with <i>Ilex</i> and <i>Blechnum</i> in the British Isles [91A0]
		<i>Margaritifera margaritifera</i> (Freshwater Pearl Mussel) [1029]
		<i>Salmo salar</i> (Salmon) [1106]
		<i>Lutra lutra</i> (Otter) [1355]
		<i>Najas flexilis</i> (Slender Naiad) [1833]
000607	Errit Lough SAC	Hard oligo-mesotrophic waters with benthic vegetation of <i>Chara</i> spp. [3140]
002008	Maumturk Mountains SAC	Oligotrophic waters containing very few minerals of sandy plains (<i>Littorelletalia uniflorae</i>) [3110]
		Northern Atlantic wet heaths with <i>Erica tetralix</i> [4010]
		Alpine and Boreal heaths [4060]
		Blanket bogs (* if active bog) [7130]
		Depressions on peat substrates of the Rhynchosporion [7150]
		Siliceous rocky slopes with chasmophytic vegetation [8220]

Site Code	Site Name	Qualifying Interests
		Salmo salar (Salmon) [1106]
		Najas flexilis (Slender Naiad) [1833]
000604	Derrinea Bog SAC	Active raised bogs [7110]
		Degraded raised bogs still capable of natural regeneration [7120]
		Depressions on peat substrates of the Rhynchosporion [7150]

DRAFT

C Conservation Objectives

DRAFT

D Cumulative and In-combination Impacts

Plans and Policies	Level	Description	Potential cumulative and/or in-combination impacts	Mitigation measures required
Water Quality				
River Basin Management Plan for Ireland 2018-2021	National	Sets out the actions that Ireland will take to improve water quality and achieve 'good' ecological status in water bodies (rivers, lakes, estuaries and coastal waters) by 2021. Ireland is required to produce a river basin management plan under the Water Framework Directive (WFD). Water quality in Ireland has deteriorated over the past two decades. The Plan provides a more coordinated framework for improving the quality of our waters — to protect public health, the environment, water amenities and to sustain water-intensive industries, including agri-food and tourism, particularly in rural Ireland	Yes - positive	Project level assessments will be required to determine any potential mitigation measures, as these will be dependent on the relevant project and any potential impacts that may arise as a result of this
Water Services Strategic Plan 2015-2040 (Irish Water)	National	Sets out strategic objectives for the delivery of water services up to 2040. It details current and future challenges which affect the provision of water services and identifies the priorities to be tackled in the short and medium term	Yes - negative	Project level assessments will be required to determine any potential mitigation measures, as these will be dependent on the relevant project and any potential impacts that may arise as a result of this
Local Authorities Water Programme	Local	The LAWP is a shared service working with Local Authorities and State agencies to meet obligations under the EU Water Framework Directive for the development and implementation of River Basin Management Plans in Ireland	Yes - positive	Project level assessments will be required to determine any potential mitigation measures, as these will be dependent on the relevant project and any potential impacts that may arise as a result of this
Moy Water Management Unit Action Plan	Regional	Maps the river and lake water bodies of the Moy Water Management Unit, states the status, impacts, pressures and risk for these water bodies and defines action programmes and objectives	Yes - positive	Project level assessments will be required to determine any potential mitigation measures, as these will be dependent on the relevant project and any potential impacts that may arise as a result of this

Plans and Policies	Level	Description	Potential cumulative and/or in-combination impacts	Mitigation measures required
Plan for Forests & Freshwater Pearl Mussel in Ireland (DRAFT)	National	Aims to eliminate, reduce or mitigate diffuse and point sources of sediment and nutrients and the disruption of the natural hydrological regime, arising from forests and regulated forestry activities, to ensure that these do not threaten the achievement of the conservation objectives for Freshwater Pearl Mussel set for each of the SACs involved	Yes - positive & negative	Project level assessments will be required to determine any potential mitigation measures, as these will be dependent on the relevant project and any potential impacts that may arise as a result of this
The Pearl Mussel Project	National	A pilot agri-environment programme that seeks to improve the quality of watercourses to benefit the endangered Freshwater Pearl Mussel. Farmers will be rewarded for their environmental services whilst having the freedom and flexibility to farm. This will in turn improve the outcomes and long-term sustainability of agri-environment schemes for biodiversity, and for rural communities. The Programme is being run by the Pearl Mussel Project Team, based in counties Kerry and Mayo	Yes - positive & negative	Project level assessments will be required to determine any potential mitigation measures, as these will be dependent on the relevant project and any potential impacts that may arise as a result of this
Shellfish Water Action Programmes / Pollution Reduction Programmes for shellfish	Regional	Under the EU Shellfish Waters Directive Ireland is obliged to establish measures to protect shellfish waters against pollution and to safeguard certain shellfish populations from various harmful consequences, resulting from the discharge of pollutant substances into the sea	Yes - positive	Project level assessments will be required to determine any potential mitigation measures, as these will be dependent on the relevant project and any potential impacts that may arise as a result of this
Irish National Action Plan for the Sustainable Use of Pesticides	National	The EU Sustainable Use Directive requires each Member State to develop a National Action Plan for the sustainable use of pesticides, which shall be reviewed at least every 5 years. The Plan sets out a national strategy to achieve a sustainable use of pesticides and aims to achieve a balance between ensuring human and environmental safety while maintaining continued viability of the farming and amenity sectors	No	Project level assessments will be required to determine any potential mitigation measures, as these will be dependent on the relevant project and any potential impacts that may arise as a result of this
County Groundwater Protection Schemes	Regional	Provides guidelines for the planning and licensing authorities in carrying out their functions, and a framework to assist in decision-making on the location, nature and control of developments and activities in order to protect groundwater	Yes - positive & negative	Project level assessments will be required to determine any potential mitigation measures, as these will be dependent on the relevant project and any potential impacts that may arise as a result of this

Plans and Policies	Level	Description	Potential cumulative and/or in-combination impacts	Mitigation measures required
Ireland's Nitrates Action Programme	National	Designed to prevent pollution of surface waters and groundwater from agricultural sources and to protect and improve water quality. Under the national NAP, each Member State's NAP must include 1) a limit on the amount of livestock manure applied to the land each year, 2) set periods when land spreading is prohibited due to risk, and 3) set capacity levels for the storage of livestock manure	Yes - positive	Project level assessments will be required to determine any potential mitigation measures, as these will be dependent on the relevant project and any potential impacts that may arise as a result of this
Environmental River Enhancement Programme (EREP)	National	The Environmental River Enhancement Programme (EREP) is an OPW funded project that is being co-ordinated and managed by Inland Fisheries Ireland. The programme focuses on the enhancement of drained Salmonid rivers in Ireland	Yes - positive	Project level assessments will be required to determine any potential mitigation measures, as these will be dependent on the relevant project and any potential impacts that may arise as a result of this
Invasive Species				
Invasive Species Action Plans	Local	The National Biodiversity Action Plan requires Ireland to prepare strategies, in consultation with Northern Ireland, to control introduced species and to prevent, or minimise, future (accidental or deliberate) introduction of alien species, which might threaten biodiversity	Yes - positive	Project level assessments will be required to determine any potential mitigation measures, as these will be dependent on the relevant project and any potential impacts that may arise as a result of this
Climate change				
Climate Action Plan Ireland	National	Sets out an ambitious course of action over the coming years to address this issue. The Plan identifies the nature and scale of the challenge, outlines the current state of play across key sectors including Electricity, Transport, Built Environment, Industry and Agriculture and charts a course towards ambitious decarbonisation targets. Reflecting the central priority climate change will have in Ireland's political and administrative systems into the future, the Plan sets out governance arrangements including carbon-proofing policies, establishment of carbon budgets, a strengthened Climate Change Advisory Council and greater accountability to the Oireachtas	No	Project level assessments will be required to determine any potential mitigation measures, as these will be dependent on the relevant project and any potential impacts that may arise as a result of this

Plans and Policies	Level	Description	Potential cumulative and/or in-combination impacts	Mitigation measures required
Green, Low-Carbon Agri-Environment Scheme (GLAS)	National	The GLAS Scheme is part of the Rural Development Programme 2014-2020. It provides funding to farmers in return for delivering environmental management on their land. Farmers must commit to the scheme for a minimum period of 5 years	Yes - positive	Project level assessments will be required to determine any potential mitigation measures, as these will be dependent on the relevant project and any potential impacts that may arise as a result of this
Renewable Energy Strategy 2011-2022	Regional	Sets out a path to allow County Mayo to contribute to meeting the national legally-binding targets and sets out opportunities for individuals, communities and businesses to harness renewable energy in a sustainable manner and to assist in combating climate change. The Strategy also clarifies the approach Mayo County Council takes to renewable energy, and should assist direction and reduce uncertainty for the most regarding issues associated with renewable energy developments in Mayo. All major forms of renewable energy are considered in the Strategy, including micro renewables	Yes - positive & negative	Project level assessments will be required to determine any potential mitigation measures, as these will be dependent on the relevant project and any potential impacts that may arise as a result of this
Economics / Agri-food and fisheries				
Food Wise 2025	National	Sets out a ten year plan for the agri-food sector. It underlines the sector's unique and special position within the Irish economy, and it illustrates the potential which exists for this sector to grow even further. The Food Wise report was published in July 2015, with the Food Wise Implementation Plan and the final Environmental Analysis on the strategy published in December 2015	Yes - negative	Project level assessments will be required to determine any potential mitigation measures, as these will be dependent on the relevant project and any potential impacts that may arise as a result of this
Biodiversity				
National Biodiversity Action Plan 2017-2021	National	Captures the objectives, targets and actions for biodiversity that will be undertaken by a wide range of government, civil society and private sectors to achieve Ireland's Vision for Biodiversity. It provides a framework to track and assess progress towards Ireland's Vision for Biodiversity over a five-year timeframe from 2017 to 2021	Yes - positive	Project level assessments will be required to determine any potential mitigation measures, as these will be dependent on the relevant project and any potential impacts that may arise as a result of this

Plans and Policies	Level	Description	Potential cumulative and/or in-combination impacts	Mitigation measures required
County Mayo Biodiversity Action Plan 2010-2015	Regional	The CMBAP aims to raise awareness of and promote the conservation of the natural heritage and biodiversity of the County. The Plan provides a framework for the conservation of biodiversity at a local level and helps ensure that national & international targets for biodiversity conservation can be achieved. A new strategy for the protection, conservation and promotion of Mayo's biodiversity is currently in preparation. This strategy will form part of the new County Mayo Heritage Plan 2020 – 2025 currently being developed and mirrors the National Biodiversity Action Plan	Yes - positive	Project level assessments will be required to determine any potential mitigation measures, as these will be dependent on the relevant project and any potential impacts that may arise as a result of this
National Peatlands Strategy	National	Guides the Government's approach to peatlands management and conservation in the future, taking into account current and potential uses of this key resource	Yes - positive & negative	Project level assessments will be required to determine any potential mitigation measures, as these will be dependent on the relevant project and any potential impacts that may arise as a result of this
All-Ireland Pollinator Plan	Local	The All-Ireland Pollinator Plan (AIPP) is a strategy that works to support the decline of pollinators including Ireland's 98 bee species, one-third of which are threatened with extinction. Mayo County Council signed up as a partner to the All-Ireland Pollinator Plan in May 2019 and works to support the All-Ireland Pollinator Plan through working with communities and local authority staff and hosting workshops and talks on the Plan	Yes - positive	Project level assessments will be required to determine any potential mitigation measures, as these will be dependent on the relevant project and any potential impacts that may arise as a result of this
Tourism				
Destination Mayo - a strategy for the future development of tourism in County Mayo 2015-2020 (DRAFT)	Regional	Seeks to build on the work of a previous study of tourism in Mayo called A Tourism Plan for Mayo. Some of the outcomes of this 1991 Plan included Mayo 5000, Mayo Naturally, designation of Ballycroy National Park and facilitated the development of a number of visitor attractions throughout the County, including Céide Fields, Mayo Naturally, National Museum of Ireland - Country Life and the Lough Lannagh Holiday Village. The vision for Mayo's tourism industry in 2020 is to be a high quality, inclusive and authentic destination. Key to the successful delivery of this vision are the availability of a skilled and enthusiastic workforce and good access to the County by road, rail and air	Yes - negative	Project level assessments will be required to determine any potential mitigation measures, as these will be dependent on the relevant project and any potential impacts that may arise as a result of this

Plans and Policies	Level	Description	Potential cumulative and/or in-combination impacts	Mitigation measures required
Strategic planning				
Project Ireland 2040	National	The government's long-term overarching strategy to make Ireland a better country for all of its people. Alongside the development of physical infrastructure, Project Ireland 2040 supports business and communities across all of Ireland in realising their potential. The National Development Plan and the National Planning Framework combine to form Project Ireland 2040	Yes - negative	Project level assessments will be required to determine any potential mitigation measures, as these will be dependent on the relevant project and any potential impacts that may arise as a result of this
National Development Plan 2018-2027	National	Sets out the investment priorities that will underpin the successful implementation of the new National Planning Framework (NPF). This will guide national, regional and local planning and investment decisions in Ireland over the next two decades to cater for an expected population increase of over 1 million people	Yes - negative	Project level assessments will be required to determine any potential mitigation measures, as these will be dependent on the relevant project and any potential impacts that may arise as a result of this
National Planning Framework	National	Sets the vision and strategy for the development of Ireland to 2040. It is a framework to guide public and private investment, to create and promote opportunities for people, and to protect and enhance the environment	Yes - negative	Project level assessments will be required to determine any potential mitigation measures, as these will be dependent on the relevant project and any potential impacts that may arise as a result of this
The National Spatial Strategy 2002-2020	National	A coherent national planning framework for Ireland, which aims to achieve a better balance of social, economic and physical development across Ireland, supported by more effective planning. In order to drive development in the regions, the NSS proposes that areas of sufficient scale and critical mass will be built up through a network of gateways and hubs	Yes - negative	Project level assessments will be required to determine any potential mitigation measures, as these will be dependent on the relevant project and any potential impacts that may arise as a result of this


Plans and Policies	Level	Description	Potential cumulative and/or in-combination impacts	Mitigation measures required
Our Sustainable Future - Framework for Sustainable Development in Ireland	National	Identifies some 70 measures to be implemented across government and tasks a high-level inter-departmental group with ensuring that the vision set out in the policy document is translated into clear and effective action	Yes - negative	Project level assessments will be required to determine any potential mitigation measures, as these will be dependent on the relevant project and any potential impacts that may arise as a result of this
National Rural Development Programme 2014-2020	National	Part of the Common Agricultural Policy (CAP), a common set of objectives, principles and rules through which the European Union coordinates support for European agriculture. The CAP is structured around two complementary pillars: Pillar 1 deals with direct payments to farmers and market management measures; Pillar 2 covers multi-annual rural development measures which include those that have beneficial impacts on the environment and climate change	Yes - negative	Project level assessments will be required to determine any potential mitigation measures, as these will be dependent on the relevant project and any potential impacts that may arise as a result of this
Sligo County Development Plan 2017-2023	Regional	Sets out an overall strategy for the proper planning and sustainable development of County Sligo, in accordance with the Planning and Development Act 2000 (as amended)	Yes - negative	Project level assessments will be required to determine any potential mitigation measures, as these will be dependent on the relevant project and any potential impacts that may arise as a result of this
Galway County Development Plan 2015-2021	Regional	Sets out an overall strategy for the proper planning and sustainable development of County Galway	Yes - negative	Project level assessments will be required to determine any potential mitigation measures, as these will be dependent on the relevant project and any potential impacts that may arise as a result of this

Plans and Policies	Level	Description	Potential cumulative and/or in-combination impacts	Mitigation measures required
Roscommon County Development Plan 2014-2020	Regional	Sets out an overall strategy for the proper planning and sustainable development of County Roscommon. A new CDP for 2021-2027 is in preparation	Yes - negative	Project level assessments will be required to determine any potential mitigation measures, as these will be dependent on the relevant project and any potential impacts that may arise as a result of this
Mayo local economic and community plan 2015-2021	Regional	A Framework that identifies economic and local community issues in County Mayo and gives effect to the delivery of economic and community development in a manner that is consistent with higher level plans and strategies including the Regional Planning Guidelines for the West Region 2010-2022 and the The Mayo County Development Plan 2014-2020	Yes - negative	Project level assessments will be required to determine any potential mitigation measures, as these will be dependent on the relevant project and any potential impacts that may arise as a result of this
Forthcoming Castlebar, Ballina and Westport Local Area Plans	Local	Mayo County Council intends to prepare a Local Area Plans for the towns in the County which have a population exceeding 5000, these towns are Castlebar, Ballina & Westport. Castlebar and Ballina have been designated as Key Towns in the Regional Spatial and Economic Strategy (RSES), published by the Northern and Western Regional Assembly on 23 January 2020, whilst Westport was also identified as a regional driver. The formal preparation of the Local Area Plans for Castlebar, Ballina and Westport will commence after the Draft Mayo County Development Plan 2021-2027 has been published	Yes - negative	Project level assessments will be required to determine any potential mitigation measures, as these will be dependent on the relevant project and any potential impacts that may arise as a result of this
Northwest BAU (Business Area Unit) Strategic Plan 2016-2020	Regional	Sets out plans for the forest and non-forest business that will take place in the BAU during the plan period. In practicing sustainable forest management Coillte's aim is to develop its forests in a way that is environmentally sustainable, socially sustainable and economically sustainable	Yes - negative	Project level assessments will be required to determine any potential mitigation measures, as these will be dependent on the relevant project and any potential impacts that may arise as a result of this


Plans and Policies	Level	Description	Potential cumulative and/or in-combination impacts	Mitigation measures required
Sustainable Development Goals National Implementation Plan 2018 - 2020	National	The Sustainable Development Goals National Implementation Plan 2018 - 2020 is in direct response to the 2030 Agenda for Sustainable Development and provides a whole-of-government approach to implement the 17 Sustainable Development Goals (SDGs). The Plan identifies four strategic priorities to guide implementation: 1) Awareness: raise public awareness of the SDGs; 2) Participation: provide stakeholders opportunities to engage and contribute to follow-up and review processes, and further develop national implementation of the Goals; 3) Support: encourage and support efforts of communities and organisations to contribute towards meeting the SDGs, and foster public participation; and 4) Policy alignment: develop alignment of national policy with the SDGs and identify opportunities for policy coherence	No	Project level assessments will be required to determine any potential mitigation measures, as these will be dependent on the relevant project and any potential impacts that may arise as a result of this
Ireland Forestry Programme 2014-2020	National	Aims to develop a competitive and sustainable forest sector through the implementation of measures, in particular afforestation, woodland creation and the prevention and restoration of damage to forests	Yes - positive & negative	Project level assessments will be required to determine any potential mitigation measures, as these will be dependent on the relevant project and any potential impacts that may arise as a result of this

E Maps of Natura 2000 sites within close vicinity of each settlement


E.1 Ballinrobe




E.2 Ballyhaunis



E.3 Belmullet



E.4 Claremorris



Minogue & Associates
Planning & Environmental Consultants

JBA
consulting

Legend


- County Boundary
- Settlements
- ▨ SACs
- ▨ SPAs

Mayo County Development Plan 2021-2027: Natura Impact Report

Settlements (Claremorris) - Natura 2000 sites

Comhairle Contae Mhaigh Eo
Mayo County Council

E.5 Charlestown



Minogue & Associates
Planning & Consulting

JBA
consulting

Legend


- ▭ County Boundary
- Settlements
- ▨ SACs

Mayo County Development
Plan 2021-2027: Natura Impact
Report

Settlements (Charlestown) -
Natura 2000 sites

Comhairle Contae Mhaigh Eo
Mayo County Council

E.6 Foxford



Miragou & Associates
Environmental & Planning

JBA
consulting

Legend


- County boundary
- Settlements
- ▨ SACs
- ▨ SPAs

Mayo County Development
Plan 2021-2027: Natura Impact
Report

Settlements (Foxford) - Natura
2000 sites

 Comhairle Contae Mhaigh Eo
Mayo County Council

E.7 Crossmolina



Miragou & Associates
Environmental & Planning

JBA
consulting

Legend

- County Boundary
- Settlements
- ▨ SACs
- ▨ SPAs

Mayo County Development Plan 2021-2027: Natura Impact Report

Settlements (Crossmolina) - Natura 2000 sites

 Comhairle Contae Mhaigh Eo
Mayo County Council

E.8 Knock




- Legend**
- County boundary
 - Settlements
 - SACs

Mayo County Development
Plan 2021-2027:Natura Impact
Report

Settlements (Knock) - Natura
2000 sites



E.9 Kiltamagh



Minogue & Associates
Environmental Consulting

JBA
consulting

Legend


- County boundary
- Settlements
- ▨ SACs

Mayo County Development
Plan 2021-2027:Natura Impact
Report

Settlements (Kiltamagh) -
Natura 2000 sites

 Comhairle Contae Mhaigh Eo
Mayo County Council

E.10 Swinford



Minogue & Associates
Planning & Environmental Consultants

JBA
consulting

Legend


- ▬ County boundary
- Settlements
- ▨ SACs

Mayo County Development
Plan 2021-2027: Natura Impact
Report


Settlements (Swinford) - Natura
2000 sites

 Comhairle Contae Mhaigh Eo
Mayo County Council


E.11 Killala



E.12 Louisburgh



E.13 Newport



Minogue & Associates
Newport, Co. Mayo


JBA
consulting

Legend

- ▭ County boundary
- Settlements
- ▨ SACs

Mayo County Development
Plan 2021-2027: Natura Impact
Report

Settlements (Newport) - Natura
2000 sites

 Comhairle Contae Mhaigh Eo
Mayo County Council

References

- CIEEM, 2016. Guidelines for Ecological Impact Assessment in the UK and Ireland. Terrestrial, Freshwater and Coastal.
- Coillte, 2016. Northwest BAU (Business Area Unit) Strategic Plan 2016-2020.
- DoAFM, 2019. Irish National Action Plan for the Sustainable Use of Pesticides.
- DoAFM, 2018. Plan for Forests & Freshwater Pearl Mussel in Ireland (DRAFT).
- DoAFM, 2015. Food Wise 2025.
- DoAFM, 2014a. Green, Low-Carbon Agri-Environment Scheme (GLAS).
- DoAFM, 2014b. National Rural Development Programme 2014-2020.
- DoCCA, 2018. Sustainable Development Goals National Implementation Plan 2018 - 2020.
- DoCCA, 2012. 'Our Sustainable Future - Framework for Sustainable Development in Ireland.
- DoCHG, 2017. National Biodiversity Action Plan 2017-2021.
- DoEHLG, 2009. Appropriate Assessment of Plans and Projects in Ireland Guidance for Planning Authorities. Department of the Environment, Heritage and Local Government.
- DoHLGH, 2018. River Basin Management Plan for Ireland 2018-2021.
- DoHLGH, 2017. Ireland's Nitrates Action Programme.
- DoHLGH, 2002. The National Spatial Strategy 2002-2020.
- DoHPLG, 2018. River Basin Management Plan for Ireland 2018-2021.
- DoPER, 2018. Project Ireland 2040.
- EC, 2013. Interpretation Manual of European Union Habitats.
- European Commission, 2007. Guidance document on Article 6 (4) of the "Habitats Directive" 92/43/EEC - Clarification of the concepts of: alternative solutions, imperative reasons of overriding public interest, compensatory measures, overall coherence, opinion of the commission.
- European Commission (Ed.), 2000. Managing natura 2000 sites: the provisions of article 6 of the "Habitats" directive 92/43/EEC. Office for Official Publications of the European Communities, Luxembourg.
- European Commission. Directorate General for the Environment., Alterra., Eurosite., 2013. Guidelines on climate change and Natura 2000: dealing with the impact of climate change, on the management of the Natura 2000 network of areas of high biodiversity value. Publications Office, LU.
- European Commission, Directorate-General for the Environment, Oxford Brookes University, Impacts Assessment Unit (Eds.), 2002. Assessment of plans and projects significantly affecting Natura 2000 sites: methodological guidance on the provisions of Article 6(3) and (4) of the Habitats directive 92/43/EEC. EUR-OP, Luxembourg.
- Forest Service/DoAFM, 2015. Ireland Forestry Programme 2014-2020.
- Fossitt, J.A., 2000. A guide to habitats in Ireland, Heritage Council of Ireland series. Heritage Council/Chomhairle Oidhreachta, Kilkenny.
- Galway CoCo, 2015. Galway County Development Plan 2015-2021.
- Geological Survey Ireland, 2018. Mayo County Groundwater Protection Scheme.
- Government of Ireland, 2019. Climate Action Plan Ireland.
- Hanley, R., 2014. Screening Methodology - Stage 1: Appropriate Assessment Screening Methodology for the Maintenance of Arterial Drainage Schemes.
- IFI, 2008. The Environmental River Enhancement Programme (EREP).

- Irish Water, 2015. Water Services Strategic Plan 2015-2040.
- Mayo CoCo, 2019. Climate Ready Mayo - Mayo County Council Climate Adaptation Strategy.
- Mayo CoCo, 2016. Mayo's Management of Invasive Alien Plant Species (IAPS) for Japanese Knotweed.
- Mayo CoCo, 2011. Renewable Energy Strategy 2011-2022.
- Mayo CoCo, 2010. County Mayo Biodiversity Action Plan 2010-2015.
- Mayo CoCo Enterprise & Investment Unit, 2015. Destination Mayo - a strategy for the future development of tourism in County Mayo 2015-2020 (DRAFT).
- Mayo Local Community Development Committee, 2015. Mayo local economic and community plan 2015-2021.
- NBDC, 2015. National Biodiversity Data Series No.3: All-Ireland Pollinator Plan.
- Nolan, P., Flanagan, J., 2020. High-resolution Climate Projections for Ireland – A Multimodel Ensemble Approach (Research Report No. 339 Part1), EPA Research Programme 2014 - 2020. Environmental Protection Agency.
- NPWS, 2019. The Status of EU Protected Habitats and Species in Ireland.
- NPWS, 2015. National Peatlands Strategy.
- Ordnance Survey Ireland/Galway County Council, 2010. Moy Water Management Unit Action Plan.
- Roscommon CoCo, 2014. Roscommon County Development Plan 2014-2020.
- Sligo CoCo, 2014. Sligo County Development Plan 2017-2023.
- WFD, 2009. Western River Basin District - River Basin Management Plan (ERBDMP) 2009 - 2015.

JBA
consulting

Offices at

Dublin
Limerick

Registered Office
24 Grove Island
Corbally
Limerick
Ireland

+353(0)61 345463
info@jbaconsulting.ie
www.jbaconsulting.ie
Follow us:  

JBA Consulting Engineers and
Scientists Limited

Registration number 444752

JBA Group Ltd is certified to:
ISO 9001:2015
ISO 14001:2015
OHSAS 18001:2007



APPENDIX C

National Parks and Wildlife Service

Conservation Objectives Series

Coolcam Turlough SAC 000218



An Roinn
Cultúir, Oidhreachta agus Gaeltachta

Department of
Culture, Heritage and the Gaeltacht



**National Parks and Wildlife Service,
Department of Culture, Heritage and the Gaeltacht,
7 Ely Place, Dublin 2, Ireland.
Web: www.npws.ie
E-mail: nature.conservation@ahg.gov.ie**

Citation:

NPWS (2017) Conservation Objectives: Coolcam Turlough SAC 000218. Version 1. National Parks and Wildlife Service, Department of Culture, Heritage and the Gaeltacht.

**Series Editor: Rebecca Jeffrey
ISSN 2009-4086**

Introduction

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

** indicates a priority habitat under the Habitats Directive*

000218 Coolcam Turlough SAC

3180 TurloughsE

Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

NPWS Documents

Year :	1992
Title :	Turloughs over 10ha - Vegetation survey and evaluation
Author :	Goodwillie, R.N.
Series :	Unpublished report to NPWS
<hr/>	
Year :	2015
Title :	Turlough hydrology, ecology and conservation (Part 1)
Author :	Waldren, S. (ed.)
Series :	Unpublished report to NPWS
<hr/>	
Year :	2015
Title :	Turlough hydrology, ecology and conservation (Part 2)
Author :	Waldren, S. (ed.)
Series :	Unpublished report to NPWS
<hr/>	
Year :	2017
Title :	Conservation objectives supporting document: Turloughs* and Rivers with muddy banks with <i>Chenopodium rubri</i> p.p. and <i>Bidention</i> p.p. vegetation
Author :	O Connor, Á.
Series :	Conservation objectives supporting document

Other References

Year :	1986
Title :	A study of the geology, hydrology and geomorphology of turloughs
Author :	Coxon, C.
Series :	Unpublished Ph.D. Thesis, Trinity College Dublin

Spatial data sources

Year : 2015

Title : Goodwillie (1992) Turloughs over 10 hectares: vegetation survey and evaluation

GIS Operations : Goodwillie map scanned and georectified. Turlough as outlined on map digitised. New turlough dataset clipped to SAC boundary. Expert opinion used as necessary to resolve any issues arising

Used For : 3180 (map 2)


Conservation Objectives for : Coolcam Turlough SAC [000218]

3180 Turloughs


To restore the favourable conservation condition of Turloughs* in Coolcam Turlough SAC, which is defined by the following list of attributes and targets:


Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable at c.56ha or increasing, subject to natural processes	Based on the approximate area of 56ha for Coolcam turlough from Waldren (2015), of which 55.5ha lies within the boundary of Coolcam Turlough SAC. Coolcam was one of 22 Trinity College Dublin (TCD) turlough project study sites (Waldren, 2015). It was also studied by Goodwillie (1992). Neither of these studies surveyed the north-west of the SAC, so the figure of 56ha may underestimate the area of the habitat. See map 2 for the recorded extent (based on Waldren, 2015). Coolcam turlough was assessed as being in inadequate conservation condition (Waldren, 2015). See also Coxon (1986) for information on Coolcam turlough. See O Connor (2017) for information on all attributes and targets
Habitat distribution	Occurrence	No decline, subject to natural processes	See map 2
Hydrological regime	Various	Maintain appropriate natural hydrological regime necessary to support the natural structure and functioning of the habitat	Hydrological regime is sub-divided into more detailed attributes (groundwater distribution, flood duration, frequency, area and depth, and permanently flooded/wet areas) and targets in O Connor (2017). Coolcam turlough has basins separated by a narrow esker; the smaller Coolcam Lough dries out in summer, the larger seems to retain water throughout the year (Waldren, 2015). The hydrological data suggest that this turlough experiences one significant flooding event per annum and has become wetter over time (Waldren, 2015). Coolcam turlough is hydrologically linked with Croaghill turlough (SAC 000255) (Waldren, 2015)
Soil type	Hectares	Maintain variety, area and extent of soil types necessary to support turlough vegetation and other biota	Coolcam soils are moderately alkaline and mineral, with an alluvial mineral soil type covering almost 95% of the turlough area (Waldren, 2015)
Soil nutrient status: nitrogen and phosphorus	N and P concentration in soil	Maintain/restore nutrient status appropriate to soil types and vegetation communities	Waldren (2015) found mean total nitrogen (TN) at Coolcam of 4,983mg/kg TN and total phosphorus (TP) of 245mg/kg TP
Physical structure: bare ground	Presence	Maintain sufficient wet bare ground, as appropriate	See O Connor (2017) for further details on this and all attributes
Chemical processes: calcium carbonate deposition and concentration	Calcium carbonate deposition rate/soil concentration	Maintain appropriate calcium carbonate deposition rate and concentration in soil	Soils were moderately alkaline at Coolcam, with a percentage calcium carbonate content of 4.78% (Waldren, 2015)
Water quality	Various	Restore appropriate water quality to support the natural structure and functioning of the habitat	Water quality is sub-divided into more detailed attributes (nutrients, colour, phytoplankton and epiphyton biomass) and targets in O Connor (2017). Waldren (2015) recorded mean TP of 34µg/l and mean chlorophyll <i>a</i> of 18µg/l. Owing to the dominance of mineral alluvial soils at Coolcam, ≤20µg/l TP may be sufficient to restore favourable condition. It is likely, however, that Coolcam naturally had 10µg/l TP, so further nutrient reductions may be required to restore more oligotrophic communities, e.g. marl lake/pond
Active peat formation	Flood duration	Maintain active peat formation, where appropriate	No peat soils were recorded at Coolcam (Waldren, 2015)

Vegetation composition: area of vegetation communities	Hectares	Maintain area of sensitive and high conservation value vegetation communities/units	See Goodwillie (1992) and Waldren (2015) for information on vegetation communities at Coolcam
Vegetation composition: vegetation zonation	Distribution	Maintain vegetation zonation/mosaic characteristic of the site	See Goodwillie (1992) and Waldren (2015) for information on vegetation at Coolcam
Vegetation structure: sward height	Centimetres	Maintain sward heights appropriate to the vegetation unit, and a variety of sward heights across the turlough	See Goodwillie (1992) and Waldren (2015) for information on vegetation at Coolcam
Typical species	Presence	Maintain typical species within and across the turlough	Typical species is sub-divided into more detailed attributes (terrestrial, wetland and aquatic plants, invertebrates and birds) and targets in O Connor (2017). See Goodwillie (1992) and Waldren (2015) for information on plant and aquatic invertebrate species at Coolcam
Fringing habitats: area	Hectares	Maintain marginal fringing habitats that support turlough vegetation, invertebrate, mammal and/or bird populations	See O Connor (2017) for further details on this and all attributes
Vegetation structure: turlough woodland	Species diversity and woodland structure	Maintain appropriate turlough woodland diversity and structure	See O Connor (2017) for further details on this and all attributes



Legend

 Coolcam Turlough SAC 000218



An Roinn
Cultúir, Oidhreachta agus Gaeltachta
Department of
Culture, Heritage and the Gaeltacht

**MAP 1:
COOLCAM TURLOUGH SAC
CONSERVATION OBJECTIVES
SAC DESIGNATION**

Map to be read in conjunction with the NPWS Conservation Objectives Document.

**SITE CODE:
SAC 000218; version 3.01.
CO. ROSCOMMON / CO. GALWAY**

0 0.1 0.2 0.3 0.4 0.5 km


The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
Ordnance Survey of Ireland Licence No EN 0059216. © Ordnance Survey of Ireland Government of Ireland

Níl sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbheithnithe a déanamh ar theorainneacha na gceantar comharthaíthe. Suirbhéarachta Ordonáis na hÉireann Ceadúnas Uimh EN 0059216. © Suirbhéarachta Ordonáis na hÉireann Rialtas na hÉireann

N



**Map Version 1
Date: Aug 2017**



Legend

- 3180 Turlough
- Coolcam Turlough SAC 000218
- OSi Discovery Series County Boundary

National Parks and Wildlife Service

Conservation Objectives Series

Croaghill Turlough SAC 000255



An Roinn
Cultúir, Oidhreachta agus Gaeltachta
Department of
Culture, Heritage and the Gaeltacht



**National Parks and Wildlife Service,
Department of Culture, Heritage and the Gaeltacht,
7 Ely Place, Dublin 2, Ireland.
Web: www.npws.ie
E-mail: nature.conservation@ahg.gov.ie**

Citation:

NPWS (2017) Conservation Objectives: Croaghil Turlough SAC 000255. Version 1. National Parks and Wildlife Service, Department of Culture, Heritage and the Gaeltacht.

**Series Editor: Rebecca Jeffrey
ISSN 2009-4086**

Introduction

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

** indicates a priority habitat under the Habitats Directive*

000255 Croaghill Turlough SAC

3180 TurloughsE

Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

NPWS Documents

Year :	1992
Title :	Turloughs over 10ha - Vegetation survey and evaluation
Author :	Goodwillie, R.N.
Series :	Unpublished report to NPWS
<hr/>	
Year :	2015
Title :	Turlough hydrology, ecology and conservation (Part 1)
Author :	Waldren, S. (ed.)
Series :	Unpublished report to NPWS
<hr/>	
Year :	2015
Title :	Turlough hydrology, ecology and conservation (Part 2)
Author :	Waldren, S. (ed.)
Series :	Unpublished report to NPWS
<hr/>	
Year :	2017
Title :	Conservation objectives supporting document: Turloughs* and Rivers with muddy banks with <i>Chenopodium rubri</i> p.p. and <i>Bidenton</i> p.p. vegetation
Author :	O Connor, Á.
Series :	Conservation objectives supporting document

Other References

Year :	1986
Title :	A study of the geology, hydrology and geomorphology of turloughs
Author :	Coxon, C.
Series :	Unpublished Ph.D. Thesis, Trinity College Dublin
<hr/>	
Year :	1995
Title :	Additions to the Irish range of <i>Rorippa islandica</i> (Oeder ex Murray) Borbas
Author :	Goodwillie, R.
Series :	Irish Naturalists' Journal, 25(2): 57-59

Spatial data sources

Year : 2015

Title : Goodwillie (1992) Turloughs over 10 hectares: vegetation survey and evaluation

GIS Operations : Goodwillie map scanned and georectified. Turlough as outlined on map digitised. New turlough dataset clipped to SAC boundary. Expert opinion used as necessary to resolve any issues arising

Used For : 3180 (map 2)


Conservation Objectives for : Croaghill Turlough SAC [000255]

3180 Turloughs


To restore the favourable conservation condition of Turloughs* in Croaghill Turlough SAC, which is defined by the following list of attributes and targets:


Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable at c.37ha or increasing, subject to natural processes	Based on the approximate area of 38.6ha for Croaghill turlough from Waldren (2015), of which 36.67ha lies within the boundary of Croaghill Turlough SAC. Croaghill was one of 22 Trinity College Dublin (TCD) turlough project study sites (Waldren, 2015). It was also studied by Goodwillie (1992). See map 2 for the recorded extent (based on Waldren, 2015). Croaghill turlough was assessed as being in inadequate conservation condition (Waldren, 2015). See also Coxon (1986) and Goodwillie (1995) and for information on Croaghill turlough. See O Connor (2017) for information on all attributes and targets
Habitat distribution	Occurrence	No decline, subject to natural processes	See map 2
Hydrological regime	Various	Maintain appropriate natural hydrological regime necessary to support the natural structure and functioning of the habitat	Hydrological regime is sub-divided into more detailed attributes (groundwater contribution, flood duration, frequency, area and depth, and permanently flooded/wet areas) and targets in O Connor (2017). The hydrological data suggest that Croaghill turlough experiences a single significant flooding event per annum (Waldren, 2015). Croaghill turlough is hydrologically linked to Coolcam turlough (SAC 000218) (Waldren, 2015)
Soil type	Hectares	Maintain variety, area and extent of soil types necessary to support turlough vegetation and other biota	Croaghill soils are moderately acidic and peaty, with low amounts of calcium carbonate (Waldren, 2015)
Soil nutrient status: nitrogen and phosphorus	N and P concentration in soil	Maintain/restore nutrient status appropriate to soil types and vegetation communities	Waldren (2015) found mean total nitrogen (TN) at Croaghill was towards the high end of the range for turloughs (and very variable among samples) with a mean of 15,883mg/kg TN, and total phosphorus was moderate at 896mg/kg TP
Physical structure: bare ground	Presence	Maintain sufficient wet bare ground, as appropriate	See O Connor (2017) for further details on this and all attributes
Chemical processes: calcium carbonate deposition and concentration	Calcium carbonate deposition rate/soil concentration	Maintain appropriate calcium carbonate deposition rate and concentration in soil	Percentage calcium carbonate content was low at Croaghill (3.8%) (Waldren, 2015)
Water quality	Various	Restore appropriate water quality to support the natural structure and functioning of the habitat	Water quality is sub-divided into more detailed attributes (nutrients, colour, phytoplankton and epiphyton biomass) and targets in O Connor (2017). Waldren (2015) recorded average TP of 25µg/l. Owing to the dominance of peat soils at Croaghill, ≤10µg/l TP is likely to be required to restore favourable condition
Active peat formation	Flood duration	Maintain active peat formation, where appropriate	More than 90% of the Croaghill turlough area is fen peat (Waldren, 2015)
Vegetation composition: area of vegetation communities	Hectares	Maintain area of sensitive and high conservation value vegetation communities/units	See Goodwillie (1992) and Waldren (2015) for information on vegetation communities at Croaghill
Vegetation composition: vegetation zonation	Distribution	Maintain vegetation zonation/mosaic characteristic of the site	See Goodwillie (1992) and Waldren (2015) for information on vegetation at Croaghill

Vegetation structure: sward height	Centimetres	Maintain sward heights appropriate to the vegetation unit, and a variety of sward heights across the turlough	See Goodwillie (1992) and Waldren (2015) for information on vegetation at Croaghill
Typical species	Presence	Maintain typical species within and across the turlough	Typical species is sub-divided into more detailed attributes (terrestrial, wetland and aquatic plants, invertebrates and birds) and targets in O Connor (2017). See Goodwillie (1992) and Waldren (2015) for information on plant and aquatic invertebrate species at Croaghill. Northern yellow-cress (<i>Rorippa islandica</i>) occurs at Croaghill (Goodwillie, 1992, 1995)
Fringing habitats: area	Hectares	Maintain marginal fringing habitats that support turlough vegetation, invertebrate, mammal and/or bird populations	See O Connor (2017) for further details on this and all attributes
Vegetation structure: turlough woodland	Species diversity and woodland structure	Maintain appropriate turlough woodland diversity and structure	See O Connor (2017) for further details on this and all attributes



Legend

 Croaghill Turlough SAC 000255



An Roinn
Cultúir, Oidhreachta agus Gaeltachta
Department of
Culture, Heritage and the Gaeltacht

MAP 1:
CROAGHILL TURLOUGH SAC
CONSERVATION OBJECTIVES
SAC DESIGNATION

Map to be read in conjunction with the NPWS Conservation Objectives Document.

SITE CODE:
SAC 000255; version 3.01. CO. GALWAY

0 50 100 150 200 250 Meters


The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
Ordnance Survey of Ireland Licence No EN 0059216. © Ordnance Survey of Ireland Government of Ireland

Níl sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbhreithnithe a déanamh ar theorainneacha na gceantar comharthaíthe. Suirbhéarachta Ordonáis na hÉireann Ceadúnas Uimh EN 0059216. © Suirbhéarachta Ordonáis na hÉireann Rialtas na hÉireann




Map Version 1
Date: Aug 2017

© Ordnance Survey Ireland



Legend

- 3180 Turlough
- Croaghill Turlough SAC 000255
- OSi Discovery Series County Boundary




An Roinn
Cultúir, Oidhreachta agus Gaeltachta
Department of
Culture, Heritage and the Gaeltacht

**MAP 2:
CROAGHILL TURLOUGH SAC
CONSERVATION OBJECTIVES
TURLOUGH**

Map to be read in conjunction with the NPWS Conservation Objectives Document.

**SITE CODE:
SAC 000255; version 3.01. CO. GALWAY**

0 50 100 150 200 250 Meters



The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
Ordnance Survey of Ireland Licence No EN 0059216. © Ordnance Survey of Ireland Government of Ireland

Níl sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbhreithnithe a déanamh ar theorainneacha na gceantar comharthaithe. Suirbhéarachta Ordonáis na hÉireann Ceadúnas Uimh EN 0059216. © Suirbhéarachta Ordonáis na hÉireann Rialtas na hÉireann



**Map Version 1
Date: Aug 2017**

National Parks and Wildlife Service

Conservation Objectives Series

Inishbofin and Inishshark SAC 000278



An Roinn
Ealaíon, Oidhreachta agus Gaeltachta

Department of
Arts, Heritage and the Gaeltacht



**National Parks and Wildlife Service,
Department of Arts, Heritage and the Gaeltacht,**

7 Ely Place, Dublin 2, Ireland.

Web: www.npws.ie

E-mail: nature.conservation@ahg.gov.ie

Citation:

**NPWS (201) Conservation Objectives: Inishbofin and Inishshark SAC 000278.
Version 1. National Parks and Wildlife Service, Department of Arts, Heritage
and the Gaeltacht.**

Series Editor: Rebecca Jeffrey

ISSN 2009-4086

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

* indicates a priority habitat under the Habitats Directive

000278	Inishbofin and Inishshark SAC
1150	Coastal lagoonsE
1364	Grey Seal <i>Halichoerus grypus</i>
3110	Oligotrophic waters containing very few minerals of sandy plains (<i>Littorelletalia uniflorae</i>)
4010	Northern Atlantic wet heaths with <i>Calluna vulgaris</i> <i>Calluna vulgaris</i>
4030	European dry heaths

Please note that this SAC overlaps with High Island, Inishshark and Davillaun SPA (004144) and Inishbofin, Omey Island and Turbot Island SPA (004231). See map 2. The conservation objectives for this site should be used in conjunction with those for the overlapping sites as appropriate.

Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

NPWS Documents

Year :	1984
Title :	The vegetation of Irish lakes
Author :	Heuff, H.
Series :	Unpublished report to NPWS
Year :	2003
Title :	Grey seal population status at islands in the Inishkea group, as determined from breeding ground surveys in 2002
Author :	Ó Cadhla, O.; Strong, D.
Series :	Unpublished report to NPWS
Year :	2004
Title :	Harbour seal population assessment in the Republic of Ireland: August 2003
Author :	Cronin, M.; Duck, C.; O Cadhla, O.; Nairn, R.; Strong, D.; O'Keefe, C.
Series :	Irish Wildlife Manual No. 11
Year :	2004
Title :	Summary of National Parks and Wildlife Service surveys for common (harbour) seals (<i>Phoca vitulina</i>) and grey seals (<i>Halichoerus grypus</i>), 1978 to 2003
Author :	Lyons, D.O.
Series :	Irish Wildlife Manual No. 13
Year :	2007
Title :	Grey seal moult population survey in the Republic of Ireland, 2007
Author :	Ó Cadhla, O.; Strong, D.
Series :	Unpublished report to NPWS
Year :	2007
Title :	Inventory of Irish coastal lagoons (version 2)
Author :	Oliver, G.
Series :	Unpublished report to NPWS
Year :	2008
Title :	An assessment of the breeding population of grey seals in the Republic of Ireland, 2005
Author :	O Cadhla, O.; Strong, D.; O'Keefe, C.; Coleman, M.; Cronin, M.; Duck, C.; Murray, T.; Dower, P.; Nairn, R.; Murphy, P.; Smiddy, P.; Saich, C.; Lyons, D.O.; Hiby, L.
Series :	Irish Wildlife Manual No. 34
Year :	2013
Title :	A survey of the benthic macrophytes of three hard-water lakes: Lough Bunny, Lough Carra and Lough Owel
Author :	Roden, C.; Murphy, P.
Series :	Irish Wildlife Manual No. 70
Year :	2013
Title :	The status of EU protected habitats and species in Ireland. Volume 2. Habitats assessments
Author :	NPWS
Series :	Conservation assessments
Year :	2014
Title :	Guidelines for a national survey and conservation assessment of upland vegetation and habitats in Ireland, Version 2.0
Author :	Perrin, P.M.; Barron, S.J.; Roche, J.R.; O'Hanrahan, B.
Series :	Irish Wildlife Manual No. 79

Year : 2015
Title : Inishbofin and Inishshark SAC (site code:278) Conservation objectives supporting document-coastal lagoons V1
Author : NPWS
Series : Conservation objectives supporting document

Year : 2015
Title : Inishbofin and Inishshark SAC (site code: 278) Conservation objectives supporting document-marine species V1
Author : NPWS
Series : Conservation objectives supporting document

Other References

Year : 1982
Title : Eutrophication of waters. Monitoring assessment and control
Author : OECD
Series : OECD, Paris

Year : 1983
Title : The grey seal (*Halichoerus grypus*) in Ireland
Author : Summers, C.F.
Series : Unpublished Report to the Minister for Fisheries, Forestry and Wildlife

Year : 1988
Title : The Irish red data book 1. Vascular plants
Author : Curtis, T.G.F; McGough, H.N.
Series : Wildlife Service, Dublin

Year : 1998
Title : Population biology of grey seals (*Halichoerus grypus*, Fabricius 1791) in western Ireland
Author : Kiely, O.R.M.
Series : Unpublished PhD thesis, National University of Ireland, University College Cork

Year : 1998
Title : Grey seal (*Halichoerus grypus*) pup production at the Inishkea island group, Co. Mayo and the Blasket Islands, Co. Kerry
Author : Kiely, O.; Myers, A.A.
Series : Biology and Environment: Proc. Royal Ir. Acad. 98B (2): 113-122

Year : 2000
Title : Colour in Irish lakes
Author : Free, G.; Allott, N.; Mills, P.; Kennelly, C.; Day, S.
Series : Verhandlungen Internationale Vereinigung für theoretische und angewandte Limnologie. 27: 2620-2623

Year : 2001
Title : Grey seal interactions with fisheries in Irish coastal waters
Author : BIM
Series : Report to the European Commission DG XIV. Study 95/40

Year : 2001
Title : Aquatic plants in Britain and Ireland
Author : Preston, C.D.; Croft, J.M.
Series : Harley Books, Colchester

Year :	2002
Title :	Deterioration of Atlantic soft water macrophyte communities by acidification, eutrophication and alkalisation
Author :	Arts, G.H.P.
Series :	Aquatic Botany, 73: 373-393
Year :	2006
Title :	A reference-based typology and ecological assessment system for Irish lakes. Preliminary investigations. Final report. Project 2000-FS-1-M1 Ecological assessment of lakes pilot study to establish monitoring methodologies EU (WFD)
Author :	Free, G.; Little, R.; Tierney, D.; Donnelly, K.; Coroni, R.
Series :	EPA, Wexford
Year :	2008
Title :	Water Quality in Ireland 2004-2006
Author :	Clabby, K.J.; Bradley, C.; Craig, M.; Daly, D.; Lucey, J.; McGarrigle, M.; O'Boyle, S.; Tierney, D.; Bowman, J.
Series :	EPA, Wexford
Year :	2009
Title :	The identification, characterization and conservation value of isoetid lakes in Ireland
Author :	Free G.; Bowman, J.; McGarrigle, M.; Little, R.; Coroni, R.; Donnelly, K.; Tierney, D.; Trodd, W.
Series :	Aquatic Conservation: Marine and Freshwater Ecosystems 19 (3): 264–273
Year :	2010
Title :	Water quality in Ireland 2007-2009
Author :	McGarrigle, M., Lucey, J.; Ó Cinnéide, M.
Series :	EPA, Wexford
Year :	2013
Title :	Monitoring and assessment of Irish lagoons for the purposes of the EU Water Framework Directive, 2009-2011. Parts 1 and 2
Author :	Roden, C.M; Oliver, G.A.
Series :	Unpublished report to the Environmental Protection Agency
Year :	in prep.
Title :	Habitats Directive Annex I lake habitats: a working interpretation for the purposes of site-specific conservation objectives and Article 17 reporting
Author :	O Connor, A.
Series :	Unpublished report by NPWS
Year :	in prep.
Title :	Monitoring of hard-water lakes in Ireland using charophytes and other macrophytes
Author :	Roden, C.; Murphy, P.
Series :	Unpublished report to NPWS

Spatial data sources

Year :	Revision 2011
Title :	Inventory of Irish Coastal Lagoons. Version 3
GIS Operations :	Clipped to SAC boundary
Used For :	1150 (map 3)
<hr/>	
Year :	2008
Title :	OSi 1:5000 IG vector dataset
GIS Operations :	WaterPolygons feature class clipped to the SAC boundary. Expert opinion used to identify Annex I habitat and to resolve any issues arising
Used For :	3110 (map 4)
<hr/>	
Year :	2012
Title :	NPWS rare and threatened species database
GIS Operations :	Dataset created from spatial references in database records. Expert opinion used as necessary to resolve any issues arising
Used For :	1364 (map 5)
<hr/>	
Year :	2005
Title :	OSi Discovery series vector data
GIS Operations :	High Water Mark (HWM) polyline feature class converted into polygon feature class; clipped to SAC boundary. Expert opinion used as necessary to resolve any issues arising
Used For :	1364 (map 5)
<hr/>	

Conservation Objectives for : Inishbofin and Inishshark SAC [000278]

1150 Coastal lagoons

To restore the favourable conservation condition of Coastal lagoons in Inishbofin and Inishshark SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable, subject to slight natural variation. Favourable reference area 8.0ha. See map 3	Areas calculated from spatial data derived from Oliver (2007). Site code IL070 (Lough Bofin). See lagoons supporting document for further details
Habitat distribution	Occurrence	No decline, subject to natural processes. See map 3 for mapped lagoon	Site IL070 in Oliver (2007). See lagoons supporting document for further details
Salinity regime	Practical salinity units (psu)	Median annual salinity and temporal variation within natural ranges	Lough Bofin apparently undergoes extreme variations in salinity. See lagoons supporting document for further details
Hydrological regime	Metres	Annual water level fluctuations and minima within natural ranges	Lough Bofin is shallow (less than 1.5m in depth). See lagoons supporting document for further details
Barrier: connectivity between lagoon and sea	Permeability	Appropriate hydrological connections between lagoons and sea, including where necessary, appropriate management	Lough Bofin has a cobble barrier. See lagoons supporting document for further details
Water quality: Chlorophyll <i>a</i>	µg/L	Annual median chlorophyll <i>a</i> within natural ranges and less than 5µg/L	Target based on Roden and Oliver (2013). See lagoons supporting document for further details
Water quality: Molybdate Reactive Phosphorus (MRP)	mg/L	Annual median MRP within natural ranges and less than 0.1mg/L	Target based on Roden and Oliver (2013). See lagoons supporting document for further details
Water quality: Dissolved Inorganic Nitrogen (DIN)	mg/L	Annual median DIN within natural ranges and less than 0.15mg/L	Target based on Roden and Oliver (2013). See lagoons supporting document for further details
Depth of macrophyte colonisation	Metres	Macrophyte colonisation to maximum depth of lagoon	Increased depth of colonisation increases both the extent and diversity of submergent macrophytes. As Lough Bofin is less than 2m deep, it is expected that macrophytes would extend to its full depth
Typical plant species	Number and m ²	Maintain number and extent of listed lagoonal specialists, subject to natural variation	Species listed in Oliver (2007). See lagoons supporting document for further details
Typical animal species	Number	Maintain listed lagoon specialists, subject to natural variation	Species listed in Oliver (2007). See lagoons supporting document for further details
Negative indicator species	Number and % cover	Negative indicator species absent or under control	Low salinity, shallow water and elevated nutrient levels increase the threat of unnatural encroachment by reedbeds. See lagoons supporting document for further details

Conservation Objectives for : Inishbofin and Inishshark SAC [000278]

3110 Oligotrophic waters containing very few minerals of sandy plains (Littorelletalia uniflorae)

To maintain the favourable conservation condition of Oligotrophic waters containing very few minerals of sandy plains (Littorelletalia uniflorae) in Inishbofin and Inishshark SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	The selection of the SAC for habitat 3110 was based on information on Lough Gowlangower, Inishbofin, which has <i>Eriocaulon aquaticum</i> and <i>Lobelia dortmanna</i> , however the lakes within the SAC have not been comprehensively surveyed. There is a large number of small lakes and ponds on Inishbofin, many of which are likely to contain lake habitat 3110 (see map 4), but require field confirmation. Two measures of extent should be used: 1. the area of the lake itself and; 2. the extent of the vegetation communities/zones that typify the habitat. For further information on this and other attributes see the lake habitats supporting document for the purposes of site-specific conservation objectives and Article 17 reporting (O Connor, in prep.)
Habitat distribution	Occurrence	No decline, subject to natural processes	As noted above, it is likely that the habitat is widespread in the site (see map 4), however detailed field survey is required to confirm this potential distribution
Typical species	Occurrence	Typical species present, in good condition, and demonstrating typical abundances and distribution	For lists of typical plant species, see Article 17 habitat assessment for 3110 (NPWS, 2013) and the lake habitats supporting document for the purposes of site-specific conservation objectives and Article 17 reporting (O Connor, in prep.). Survey work is required to identify the typical and other species that characterise the lakes in the SAC
Vegetation composition: characteristic zonation	Occurrence	All characteristic zones should be present, correctly distributed and in good condition	The characteristic zonation of lake habitat 3140 has been described (Roden and Murphy, 2013; in prep.), however significant further work is necessary to describe the characteristic zonation and other spatial patterns in the other lake habitats
Vegetation distribution: maximum depth	Metres	Maintain maximum depth of vegetation, subject to natural processes	The maximum depth of vegetation is likely to be specific to the lake shoreline in question. An indicative target of >6m has been developed for hard water lakes (3140) (see Roden and Murphy, 2013; in prep.). Indicative targets will be developed for the other lake habitats with time
Hydrological regime: water level fluctuations	Metres	Maintain appropriate natural hydrological regime necessary to support the habitat	Fluctuations in lake water level are typical in Ireland, but can be amplified by activities such as abstraction and drainage. Increased water level fluctuations can increase wave action, up-root vegetation, increase turbidity, alter the substratum and lead to release of nutrients from the sediment. The hydrological regime of the lakes must be maintained so that the area, distribution and depth of the lake habitat and its constituent/characteristic vegetation zones and communities are not reduced
Lake substratum quality	Various	Maintain appropriate substratum type, extent and chemistry to support the vegetation	Research is required to further characterise the substratum types (particle size and origin) and substratum quality (notably pH, calcium, iron and nutrient concentrations) favoured by each of the five Annex I lake habitats in Ireland. It is likely that the oligotrophic soft water habitat is associated with a range of nutrient-poor substrates, from stones, cobble and gravel, through sands, silt, clay and peat. Substratum particle size is likely to vary with depth and along the shoreline within a single lake

Water quality: transparency	Metres	Maintain appropriate Secchi transparency. There should be no decline in Secchi depth/transparency	Transparency relates to light penetration and, hence, to the depth of colonisation of vegetation. It can be affected by phytoplankton blooms, water colour and turbidity. A target has been set for hard water lakes (3140), however targets have yet to be established for the remaining lake habitats. Habitat 3110 is associated with very clear water. The OECD fixed boundary system set transparency targets for oligotrophic lakes of $\geq 6\text{m}$ annual mean Secchi disk depth, and $\geq 3\text{m}$ annual minimum Secchi disk depth. Free et al. (2009) found high isoetid abundance in lakes with Secchi depths of more than 3m
Water quality: nutrients	$\mu\text{g/l P}$; mg/l N	Maintain the concentration of nutrients in the water column to sufficiently low levels to support the habitat and its typical species	As a nutrient poor habitat, oligotrophic and Water Framework Directive (WFD) 'high' status targets apply. Where a lake has nutrient concentrations that are lower than these targets, there should be no decline within class, i.e. no upward trend in nutrient concentrations. For the oligotrophic soft water lake habitat, annual average TP concentration should be $\leq 10\mu\text{g/l TP}$, average annual total ammonia concentration should be $\leq 0.040\text{ mg/l N}$ and annual 95th percentile for total ammonia should be $\leq 0.090\text{mg/l N}$. For further information see the European Communities Environmental Objectives (Surface Waters) Regulations 2009
Water quality: phytoplankton biomass	$\mu\text{g/l Chlorophyll } a$	Maintain appropriate water quality to support the habitat, including high chlorophyll <i>a</i> status	Oligotrophic and WFD 'high' status targets apply to the oligotrophic soft water habitat (3110). Where a lake has a chlorophyll <i>a</i> concentration that is lower than this target, there should be no decline within class, i.e. no upward trend in phytoplankton biomass. The average growing season (March-October) chlorophyll <i>a</i> concentration must be $< 5.8\mu\text{g/l}$. The annual average chlorophyll <i>a</i> concentration should be $< 2.5\mu\text{g/l}$ and the annual peak chlorophyll <i>a</i> concentration should be $\leq 8.0\mu\text{g/l}$. For further information see the European Communities Environmental Objectives (Surface Waters) Regulations 2009
Water quality: phytoplankton composition	EPA phytoplankton composition metric	Maintain appropriate water quality to support the habitat, including high phytoplankton composition status	The EPA has developed a phytoplankton composition metric for nutrient enrichment of Irish lakes. As for other water quality indicators, habitat 3110 requires WFD high status
Water quality: attached algal biomass	Algal cover and EPA phytobenthos metric	Maintain trace/ absent attached algal biomass ($< 5\%$ cover) and high phytobenthos status	Nutrient enrichment can favour epiphytic and epipelagic algae that can out-compete the submerged vegetation. The cover abundance of attached algae in the oligotrophic soft water habitat should, therefore, be trace/ absent ($< 5\%$ cover). EPA phytobenthos can be used as an indicator of changes in attached algal biomass. As for other water quality indicators, habitat 3110 requires high phytobenthos status
Water quality: macrophyte status	EPA macrophyte metric (The Free Index)	Maintain high macrophyte status	Nutrient enrichment can favour more competitive submerged macrophyte species that out-compete the typical and characteristic species for the lake habitat. The EPA monitors macrophyte status for WFD purposes using the 'Free Index'. The target for the oligotrophic soft water lake habitat is high status or an Ecological Quality Ratio (EQR) for lake macrophytes of ≥ 0.90 , as defined in Schedule Five of the European Communities Environmental Objectives (Surface Waters) Regulations 2009

Acidification status	pH units, mg/l	Maintain appropriate water and sediment pH, alkalinity and cation concentrations to support the habitat, subject to natural processes	Acidification can impact on species abundance and composition in soft water lake habitats. In Europe, acidification of isoetid lakes can lead to loss of isoetids and dominance by submerged <i>Sphagnum</i> mosses and <i>Juncus bulbosus</i> (Arts, 2002). The specific requirements of lake habitat 3110, in terms of water and sediment pH, alkalinity and cation concentration, have not been determined. For oligotrophic soft water lakes (3110), and adopting a precautionary approach based on Arts (2002), minimum pH should not be <5.5 pH units. Maximum pH should be <9.0 pH units, in line with the surface water standards established for soft waters (where water hardness is ≤100mg/l CaCO ₃). See Schedule Five of the European Communities Environmental Objectives (Surface Waters) Regulations 2009)
Water colour	mg/l PtCo	Maintain appropriate water colour to support the habitat	Increased water colour and turbidity decrease light penetration and can reduce the area of available habitat for lake macrophytes, particularly at the lower euphotic depths. The primary source of increased water colour in Ireland is disturbance to peatland. No habitat-specific or national standards for water colour currently exist. Studies have shown median colour concentrations in Irish lakes of 38mg/l PtCo (Free, et al., 2000) and 33mg/l PtCo (Free et al., 2006). It is likely that the water colour in all Irish lake habitats would naturally be <50 mg/l PtCo. Water colour can be very low (<20mg/l PtCo or even <0mg/l PtCo) in oligotrophic soft water lakes (3110), where the peatland in the lake's catchment is intact
Dissolved organic carbon (DOC)	mg/l	Maintain appropriate organic carbon levels to support the habitat	Dissolved (and particulate) organic carbon (OC) in the water column is linked to water colour and acidification (organic acids). Increasing DOC in water has been documented across the Northern Hemisphere, including afforested peatland catchments in Ireland. Damage and degradation of peatland, leading to decomposition of peat is likely to be the predominant source of OC in Ireland. OC in water promotes decomposition by fungi and bacteria that, in turn, releases dissolved nutrients. The increased biomass of decomposers can also impact directly on the characteristic lake communities through shading, competition, etc.
Turbidity	nephelometric turbidity units/ mg/l SS/ other appropriate units	Maintain appropriate turbidity to support the habitat	Turbidity can significantly affect the quantity and quality of light reaching rooted and attached vegetation and can, therefore, impact on lake habitats. The settlement of higher loads of inorganic or organic material on lake vegetation communities may also have impacts on sensitive, delicate species. Turbidity can increase as a result of re-suspension of material within the lake, higher loads entering the lake, or eutrophication. Turbidity measurement and interpretation is challenging. As a result, it is likely to be difficult to set habitat-specific targets for turbidity in lakes
Fringing habitat: area and condition	Hectares	Maintain the area and condition of fringing habitats necessary to support the natural structure and functioning of habitat 3110	Most lake shorelines have fringing habitats of reedswamp, other swamp, fen, marsh or wet woodland that intergrade with and support the structure and functions of the lake habitat. Equally, fringing habitats are dependent on the lake, particularly its water levels, and support wetland communities and species of conservation concern. Many of the fringing wetland habitats support higher invertebrate and plant species richness than the lake habitats themselves

Conservation Objectives for : Inishbofin and Inishshark SAC [000278]

4010 Northern Atlantic wet heaths with *Erica tetralix*

To restore the favourable conservation condition of Northern Atlantic wet heaths with *Erica tetralix* in Inishbofin and Inishshark SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Total area of this habitat has not been calculated although it is known to be distributed throughout the SAC, usually occurring in mosaic with other habitats such as other heath types (including European dry heaths (4030)), exposed rock, blanket bog and grasslands (NPWS internal files; Commonage Framework Plan (GA03))
Habitat distribution	Occurrence	No decline from current distribution, subject to natural processes	See note above
Ecosystem function: soil nutrient status	Soil pH and nutrient levels at a representative number of monitoring stops	Maintain soil nutrient status within natural range	Changes to soil nutrient status can occur from high stock densities or supplementary feeding above appropriate levels
Vegetation composition: cross-leaved heath	Occurrence in vicinity of a representative number of monitoring stops	Cross-leaved heath (<i>Erica tetralix</i>) present	Attribute and target based on Perrin et al. (2014)
Vegetation composition: positive indicator species	Percentage cover at a representative number of monitoring stops	Cover of positive indicator species, as listed in Perrin et al. (2014) at least 50%	Attribute and target based on Perrin et al. (2014). Ling (<i>Calluna vulgaris</i>), cross-leaved heath (<i>Erica tetralix</i>), sedges including dioecious sedge (<i>Carex dioica</i>) and bog-sedge (<i>C. limosa</i>) are listed for the heath in this SAC (NPWS internal files)
Vegetation composition: lichens and bryophytes	Percentage cover at a representative number of monitoring stops	Total cover of <i>Cladonia</i> and <i>Sphagnum</i> species, <i>Racomitrium lanuginosum</i> and pleurocarpous mosses at least 10%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: ericoid species	Percentage cover at a representative number of monitoring stops	Cover of ericoid species at least 15%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: rare/scarce species	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat	This includes species listed in the Flora (Protection) Order 1999 and/or the red data book (Curtis and McGough, 1988). Species that have been recorded in the past include marsh clubmoss (<i>Lycopodiella inundata</i>) and wood small-reed (<i>Calamagrostis epigejos</i>) (NPWS internal files)
Vegetation composition: dwarf-shrub species	Percentage cover at a representative number of monitoring stops	Cover of dwarf shrub species collectively less than 75%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: negative indicator species	Percentage cover at a representative number of monitoring stops	Cover of negative indicator species collectively less than 1%	Attribute and target based on Perrin et al. (2014), where negative indicator species are also listed
Vegetation composition: non-native species	Percentage cover at a representative number of monitoring stops and in local vicinity	Cover of non-native species less than 1%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: native trees and shrubs	Percentage cover in local vicinity	Cover of scattered native trees and shrubs less than 20%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: bracken	Percentage cover in local vicinity	Cover of bracken (<i>Pteridium aquilinum</i>) less than 10%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: soft rush	Percentage cover in local vicinity	Cover of soft rush (<i>Juncus effusus</i>) less than 10%	Attribute and target based on Perrin et al. (2014). Dense areas of soft rush can indicate disturbance

Vegetation structure: <i>Sphagnum</i> condition	Percentage at a representative number of monitoring stops	Less than 10% of <i>Sphagnum</i> cover is crushed, broken and/or pulled up	Attribute and target based on Perrin et al. (2014)
Vegetation structure: signs of browsing	Percentage at a representative number of monitoring stops	Last complete growing season's shoots of ericoids showing signs of browsing collectively less than 33%	Attribute and target based on Perrin et al. (2014)
Vegetation structure: burning	Occurrence in local vicinity of a representative number of monitoring stops	No signs of burning inside sensitive areas	Attribute and target based on Perrin et al. (2014), where sensitive areas are also defined
Physical structure: drainage	Percentage cover in local vicinity of a representative number of monitoring stops	Area showing signs of drainage from heavy trampling, tracking or ditches less than 10%	Attribute and target based on Perrin et al. (2014)
Physical structure: disturbed bare ground	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of disturbed bare ground less than 10%	Attribute and target based on Perrin et al. (2014)

Conservation Objectives for : Inishbofin and Inishshark SAC [000278]

4030 European dry heaths



To restore the favourable conservation condition of European dry heaths in Inishbofin and Inishshark SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Total area of this habitat has not been calculated although it is known to be distributed throughout the SAC, usually occurring in mosaic with other habitats such as other heath types (including Northern Atlantic wet heaths with <i>Erica tetralix</i> (4010)), exposed rock, blanket bog and grasslands (NPWS internal files; Commonage Framework Plan (GA03))
Habitat distribution	Occurrence	No decline, subject to natural processes	See note above
Ecosystem function: soil nutrient status	Soil pH and nutrient levels at a representative number of monitoring stops	Maintain soil nutrient status within natural range	Changes to soil nutrient status can occur from high stock densities or supplementary feeding above appropriate levels
Vegetation composition: positive indicator species	Number and percentage cover at a representative number of monitoring stops	At least two positive indicator species, as listed in Perrin et al. (2014), with combined cover of at least 50%	Attribute and target based on Perrin et al. (2014). Bell heather (<i>Erica cinerea</i>), ling (<i>Calluna vulgaris</i>) and Western gorse (<i>Ulex gallii</i>) are listed for the heath in this SAC (NPWS internal files)
Vegetation composition: bryophyte and non-crustose lichen species	Number at a representative number of monitoring stops	At least three bryophyte or non-crustose lichen species present, excluding <i>Campylopus</i> and <i>Polytrichum</i> moss species	Attribute and target based on Perrin et al. (2014)
Vegetation composition: rare/scarce species	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat	This includes species listed in the Flora (Protection) Order 1999 and/or the red data book (Curtis and McGough, 1988). Spotted rock-rose (<i>Tuberaria guttata</i>), a species listed in Curtis and McGough (1988) has been recorded on shallow peat on Inishbofin (NPWS internal files)
Vegetation structure: dwarf shrub species	Percentage cover at a representative number of monitoring stops	Cover of bog myrtle (<i>Myrica gale</i>), creeping willow (<i>Salix repens</i>) and Western gorse (<i>Ulex gallii</i>) collectively less than 50%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: negative indicator species	Percentage cover at a representative number of monitoring stops	Cover of negative indicator species collectively less than 1%	Attribute and target based on Perrin et al. (2014), where negative indicator species are also listed
Vegetation composition: non-native species	Percentage cover at a representative number of monitoring stops and in local vicinity	Cover of non-native species less than 1%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: native trees and shrubs	Percentage cover in local vicinity	Cover of scattered native trees and shrubs less than 20%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: bracken	Percentage cover in local vicinity	Cover of bracken (<i>Pteridium aquilinum</i>) less than 10%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: soft rush	Percentage cover in local vicinity	Cover of soft rush (<i>Juncus effusus</i>) less than 10%	Attribute and target based on Perrin et al. (2014). Dense areas of soft rush can indicate disturbance
Vegetation structure: senescent ling	Percentage cover at a representative number of monitoring stops	Senescent proportion of ling (<i>Calluna vulgaris</i>) cover less than 50%	Attribute and target based on Perrin et al. (2014)


Vegetation structure: growth phases of ling	Percentage cover in local vicinity	Outside boundaries of sensitive areas, all growth phases of ling (<i>Calluna vulgaris</i>) should occur throughout, with at least 10% of cover in mature phase	Attribute and target based on Perrin et al. (2014), where sensitive areas and growth phases are defined
Vegetation structure: signs of browsing	Percentage cover at a representative number of monitoring stops	Last complete growing season's shoots of ericoids showing signs of browsing collectively less than 33%	Attribute and target based on Perrin et al. (2014)
Vegetation structure: burning	Occurrence in local vicinity	No signs of burning inside sensitive areas	Attribute and target based on Perrin et al. (2014), where sensitive areas are defined
Physical structure: disturbed bare ground	Percentage cover at a representative number of monitoring stops and in local vicinity	Cover of disturbed bare ground less than 10%	Attribute and target based on Perrin et al. (2014)


Conservation Objectives for : Inishbofin and Inishshark SAC [000278]**1364 Grey Seal *Halichoerus grypus*****To maintain the favourable conservation condition of Grey Seal in Inishbofin and Inishshark SAC, which is defined by the following list of attributes and targets:**


Attribute	Measure	Target	Notes
Access to suitable habitat	Number of artificial barriers	Species range within the SAC should not be restricted by artificial barriers to site use. See map 5	See marine supporting document for further details
Breeding behaviour	Breeding sites	Conserve the breeding sites in a natural condition. See map 5	Attribute and target based on background knowledge of Irish breeding populations, comprehensive breeding surveys in 1995 (Kiely, 1998; Kiely and Myers, 1998), 1998 and 1999 (BIM, 2001), 2002 (Ó Cadhla and Strong, 2003) and 2005 (Ó Cadhla et al, 2008) and unpublished NPWS records, including those reported by Lyons (2004). See marine supporting document for further details
Moulting behaviour	Moult haul-out sites	Conserve the moult haul-out sites in a natural condition. See map 5	Attribute and target based on background knowledge of Irish populations, on review of data from Kiely (1998) and Lyons (2004), a national moult survey (Ó Cadhla and Strong, 2007) and unpublished NPWS records. See marine supporting document for further details
Resting behaviour	Resting haul-out sites	Conserve the resting haul-out sites in a natural condition. See map 5	Attribute and target based on review of data from Kiely (1998), BIM (2001), Lyons (2004), Cronin et al., (2004), Ó Cadhla et al, (2008) and unpublished NPWS records. See marine supporting document for further details
Disturbance	Level of impact	Human activities should occur at levels that do not adversely affect the grey seal population at the site	See marine supporting document for further details






Legend


 Inishbofin and Inishshark SAC 000278






Legend


-  Inishbofin and Inishshark SAC 000278
-  1150 *Coastal lagoons
-  OSi Discovery Series County Boundary






Legend

- Inishbofin and Inishshark SAC 000278
- Potential 3110 Oligotrophic waters containing very few minerals of sandy plains (*Littorelletalia unifloraræ*)
- OSi Discovery Series County Boundary





Legend

- Inishbofin and Inishshark SAC 000278
- ★ 1364 Grey Seal - *Halichoerus grypus* breeding sites
- ★ 1364 Grey Seal - *Halichoerus grypus* moulting sites
- ★ 1364 Grey Seal - *Halichoerus grypus* resting sites
- 1364 Grey Seal - *Halichoerus grypus* habitat
- OSi Discovery Series County Boundary

National Parks and Wildlife Service

Conservation Objectives Series

Kilsallagh Bog SAC 000285



An Roinn
Ealaíon, Oidhreachta agus Gaeltachta

Department of
Arts, Heritage and the Gaeltacht



**National Parks and Wildlife Service,
Department of Arts, Heritage and the Gaeltacht,
7 Ely Place, Dublin 2, Ireland.
Web: www.npws.ie
E-mail: nature.conservation@ahg.gov.ie**

Citation:

**NPWS (201) Conservation Objectives: Kilsallagh Bog SAC 000285. Version 1.
National Parks and Wildlife Service, Department of Arts, Heritage and the
Gaeltacht.**

**Series Editor: Rebecca Jeffrey
ISSN 2009-4086**

Introduction

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

** indicates a priority habitat under the Habitats Directive*

000285	Kilsallagh Bog SAC
7110	Active raised bogsE
7120	Degraded raised bogs still capable of natural regeneration
7150	Depressions on peat substrates of the Rhynchosporion

Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

NPWS Documents

Year :	2014
Title :	Raised Bog Monitoring and Assessment Survey 2013
Author :	Fernandez, F.; Connolly K.; Crowley W.; Denyer J.; Duff K.; Smith G.
Series :	Irish Wildlife Manual No. 81
<hr/>	
Year :	2014
Title :	Kilsallagh Bog (SAC 000285), Co. Galway, Site Report
Author :	Fernandez, F.; Connolly, K.; Crowley, W.; Denyer J.; Duff K.; Smith G.
Series :	Raised bog monitoring and assessment survey 2013
<hr/>	
Year :	2014
Title :	National raised bog SAC management plan
Author :	Department of Arts, Heritage and the Gaeltacht
Series :	Draft for consultation. 15 January 2014
<hr/>	
Year :	2015
Title :	Kilsallagh Bog SAC (site code: 285) Conservation objectives supporting document- raised bog habitats V1
Author :	NPWS
Series :	Conservation objectives supporting document

Other References

Year :	2011
Title :	Review and revision of empirical critical loads and dose-response relationships. Proceedings of an expert workshop, Noordwijkerhout, 23-25 June 2010
Author :	Bobbink, R.; Hettelingh, J.P.
Series :	RIVM report 680359002, Coordination Centre for Effects, National Institute for Public Health and the Environment (RIVM)
<hr/>	
Year :	2014
Title :	Nitrogen deposition and exceedance of critical loads for nutrient nitrogen in Irish grasslands
Author :	Henry, J.; Aherne, J.
Series :	Science of the Total Environment 470–471: 216–223

Spatial data sources

Year :	2014
Title :	Scientific Basis for Raised Bog Conservation in Ireland
GIS Operations :	RBSB13_SACs_ARB_DRB dataset, RBSB13_SACs_2012_HB dataset, RBSB13_SACs_DrainagePatterns_5k dataset and RBSB13_SAC_LIDAR_DTMs dataset clipped to SAC boundary. Expert opinion used as necessary to resolve any issues arising
Used For :	potential 7110; digital elevation model; drainage patterns (maps 2 and 4)
<hr/>	
Year :	2013
Title :	Raised Bog Monitoring and Assessment Survey 2013
GIS Operations :	RBMA13_ecotope_map dataset clipped to SAC boundary. Appropriate ecotopes selected and exported to new dataset. Expert opinion used as necessary to resolve any issues arising
Used For :	7110 ecotopes (map 3)
<hr/>	

Conservation Objectives for : Kilsallagh Bog SAC [000285]

7110 Active raised bogs

To restore the favourable conservation condition of Active raised bogs in Kilsallagh Bog SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Restore area of active raised bog to 35.6ha, subject to natural processes	Active Raised Bog (ARB) habitat was mapped at 11.5ha by Fernandez et al. (2014). Area of Degraded Raised Bog (DRB) on the High Bog (HB) has been modelled as 18.1ha. See map 2. It is estimated that all of this area is potentially restorable to ARB by drain blocking. The total potential ARB on the HB is therefore estimated to be 29.6ha. Eco-hydrological assessments of the cutover estimates that an additional 6.0ha of bog forming habitats could be restored. The long term target for ARB is therefore 35.6ha. See raised bog supporting document for further details on this and following attributes
Habitat distribution	Occurrence	Restore the distribution and variability of active raised bog across the SAC. See map 3 for distribution in 2012	ARB habitat at Kilsallagh comprises central and sub-central ecotopes that currently occurs on both the northern and southern parts of the bog. DRB also occurs on both parts of the bog, which will require restoration measures. There is also potential for ARB restoration on cutover areas of the bog (see area target above)
High bog area	Hectares	No decline in extent of high bog necessary to support the development and maintenance of active raised bog. See map 2	The area of high bog within Kilsallagh Bog SAC in 2012 (latest figure available) was 182.1ha (DAHG 2014)
Hydrological regime: water levels	Centimetres	Restore appropriate water levels throughout the site	For ARB, mean water level needs to be near or above the surface of the bog lawns for most of the year. Seasonal fluctuations should not exceed 20cm, and should only be 10cm below the surface, except for very short periods of time. Open water is often characteristic of soak systems
Hydrological regime: flow patterns	Flow direction; slope	Restore, where possible, appropriate high bog topography, flow directions and slopes. See map 4 for current situation	ARB depends on mean water levels being near or above the surface of bog lawns for most of the year. Long and gentle slopes are the most favourable to achieve these conditions. Changes to flow directions due to subsidence of bogs can radically change water regimes and cause drying out of high quality ARB areas and soak systems
Transitional areas between high bog and adjacent mineral soils (including cutover areas)	Hectares; distribution	Restore adequate transitional areas to support/protect active raised bog and the services it provides	No natural marginal habitats exist on the margins of this bog. Eco-hydrological assessments have evaluated the potential for ARB restoration on cutover areas (see note for habitat area attribute above)
Vegetation quality: central ecotope, active flush, soaks, bog woodland	Hectares	Restore 17.8ha of central ecotope/active flush/soaks/bog woodland as appropriate	At least 50% of ARB habitat should be high quality (i.e. central ecotope, active flush, soaks, bog woodland). Target area of active raised bog for the site has been set at 35.6ha (see area target above)
Vegetation quality: microtopographical features	Hectares	Restore adequate cover of high quality microtopographical features	High quality microtopography (hummocks, hollows and pools) is well developed on Kilsallagh Bog
Vegetation quality: bog moss (<i>Sphagnum</i>) species	Percentage cover	Restore adequate cover of bog moss (<i>Sphagnum</i>) species to ensure peat-forming capacity	<i>Sphagnum</i> cover varies naturally across Ireland with relatively high cover in the east to lower cover in the west. Hummock forming species such as <i>Sphagnum austini</i> are particularly good peat formers. <i>Sphagnum</i> cover and distribution also varies naturally across a site

Typical ARB species: flora	Occurrence	Restore, where appropriate, typical active raised bog flora	Typical flora species include widespread species, as well as those with more restricted distributions but typical of the habitat's subtypes or geographical range
Typical ARB species: fauna	Occurrence	Restore, where appropriate, typical active raised bog fauna	Typical fauna species include widespread species, as well as those with more restricted distributions but typical of the habitat's subtypes or geographical range
Elements of local distinctiveness	Occurrence	Maintain features of local distinctiveness, subject to natural processes	Kilsallagh Bog is noted for the presence of two flushes within the high bog. The flush in the south of the SAC contains several swallowholes. Notable bird species that have been recently recorded include breeding red grouse (<i>Lagopus lagopus</i>) and breeding curlew (<i>Numenius arquata</i>)
Negative physical indicators	Percentage cover	Negative physical features absent or insignificant	Negative physical indicators include: bare peat, algae dominated pools and hollows, marginal cracks, tear patterns, subsidence features such as dry mineral mounds /ridges emerging or expanding, and evidence of burning
Vegetation composition: native negative indicator species	Percentage cover	Native negative indicator species at insignificant levels	Disturbance indicators include species indicative of conditions drying out such as abundant bog asphodel (<i>Narthecium ossifragum</i>), deergrass (<i>Trichophorum germanicum</i>) and haretail cotton-grass (<i>Eriophorum vaginatum</i>) forming tussocks; abundant magellanic bog-moss (<i>Sphagnum magellanicum</i>) in pools previously dominated by <i>Sphagnum</i> species typical of very wet conditions (e.g. feathery bog-moss (<i>S. cuspidatum</i>)); and indicators of frequent burning events such as abundant <i>Cladonia floerkeana</i> and high cover of carnation sedge (<i>Carex panicea</i>) (particularly in true midlands raised bogs)
Vegetation composition: non-native invasive species	Percentage cover	Non-native invasive species at insignificant levels and not more than 1% cover	Common non-native invasive species include lodgepole pine (<i>Pinus contorta</i>), rhododendron (<i>Rhododendron ponticum</i>) and pitcherplant (<i>Sarracenia purpurea</i>)
Air quality: nitrogen deposition	kg N/ha/year	Air quality surrounding bog close to natural reference conditions. The total N deposition should not exceed 5kg N/ha/yr	Change in air quality can result from fertiliser drift; adjacent quarry activities; or other atmospheric inputs. The critical load range for ombrotrophic bogs has been set as between 5 and 10kg N/ha/yr (Bobbink and Hettelingh, 2011). The latest N deposition figures for the area around Kilsallagh Bog suggests that the current level is approximately 11.3kg N/ha/yr (Henry and Aherne, 2014)
Water quality	Hydrochemical measures	Water quality on the high bog and in transitional areas close to natural reference conditions	Water chemistry within raised bogs is influenced by atmospheric inputs (rainwater). However, within soak systems, water chemistry is influenced by other inputs such as focused flow or interaction with underlying substrates. Water chemistry in areas surrounding the high bog varies due to influences of different water types (bog water, regional groundwater, and run-off from surrounding mineral lands)

Conservation Objectives for : Kilsallagh Bog SAC [000285]

7120 Degraded raised bogs still capable of natural regeneration

The long-term aim for Degraded raised bogs still capable of natural regeneration is that its peat-forming capability is re-established; therefore, the conservation objective for this habitat is inherently linked to that of Active raised bogs (7110) and a separate conservation objective has not been set in Kilsallagh Bog SAC


Attribute	Measure	Target	Notes

Conservation Objectives for : Kilsallagh Bog SAC [000285]

7150 Depressions on peat substrates of the Rhynchosporion


Depressions on peat substrates of the Rhynchosporion is an integral part of good quality Active raised bogs (7110) and thus a separate conservation objective has not been set for the habitat in Kilsallagh Bog SAC


Attribute	Measure	Target	Notes
-----------	---------	--------	-------



Legend


 Kilsallagh Bog SAC 000285










Legend

- Kilsallagh Bog SAC 000285
- High Bog Boundary
- Potential 7110 *Active Raised Bog






Legend

-  Kilsallagh Bog SAC 000285
-  High Bog Boundary
- Active Raised Bog Ecotopes**
-  Central ecotope
-  Sub-central ecotope







Legend

- Kilsallagh Bog SAC 000285
- High Bog Boundary
- Drainage Patterns
- Contours

Elevation



National Parks and Wildlife Service

Conservation Objectives Series

Lisnageeragh Bog and Ballinastack Turlough SAC
000296



*An Roinn
Ealaíon, Oidhreachta agus Gaeltachta*

*Department of
Arts, Heritage and the Gaeltacht*



**National Parks and Wildlife Service,
Department of Arts, Heritage and the Gaeltacht,**

7 Ely Place, Dublin 2, Ireland.

Web: www.npws.ie

E-mail: nature.conservation@ahg.gov.ie

Citation:

**NPWS (2016) Conservation Objectives: Lisnageeragh Bog and Ballinastack
Turlough SAC 000296. Version 1. National Parks and Wildlife Service,
Department of Arts, Heritage and the Gaeltacht.**

Series Editor: Rebecca Jeffrey

ISSN 2009-4086

Introduction

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

** indicates a priority habitat under the Habitats Directive*

000296 Lisnageeragh Bog and Ballinastack Turlough SAC

3180 TurloughsE

7110 Active raised bogsE

7120 Degraded raised bogs still capable of natural regeneration

7150 Depressions on peat substrates of the Rhynchosporion

Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

NPWS Documents

Year :	2014
Title :	Raised Bog Monitoring and Assessment Survey 2013
Author :	Fernandez, F.; Connolly K.; Crowley W.; Denyer J.; Duff K.; Smith G.
Series :	Irish Wildlife Manual No. 81
<hr/>	
Year :	2014
Title :	Lisnageeragh Bog (SAC 000296), Co. Galway, Site Report
Author :	Fernandez, F.; Connolly, K.; Crowley, W.; Denyer J.; Duff K.; Smith G.
Series :	Raised bog monitoring and assessment survey 2013
<hr/>	
Year :	2014
Title :	National raised bog SAC management plan
Author :	Department of Arts, Heritage and the Gaeltacht
Series :	Draft for consultation. 15 January 2014
<hr/>	
Year :	2016
Title :	Lisnageeragh Bog and Ballinastack Turlough SAC (site code: 296) Conservation objectives supporting document- raised bog habitats V1
Author :	NPWS
Series :	Conservation objectives supporting document
<hr/>	
Year :	2016
Title :	Lisnageeragh Bog and Ballinastack Turlough SAC (site code: 296) Conservation objectives supporting document- turloughs V1
Author :	NPWS
Series :	Conservation objectives supporting document

Other References

Year :	2011
Title :	Review and revision of empirical critical loads and dose-response relationships. Proceedings of an expert workshop, Noordwijkerhout, 23-25 June 2010
Author :	Bobbink, R.; Hettelingh, J.P.
Series :	RIVM report 680359002, Coordination Centre for Effects, National Institute for Public Health and the Environment (RIVM)
<hr/>	
Year :	2014
Title :	Nitrogen deposition and exceedance of critical loads for nutrient nitrogen in Irish grasslands
Author :	Henry, J.; Aherne, J.
Series :	Science of the Total Environment 470-471: 216-223

Spatial data sources

Year :	2016
Title :	Goodwillie (1992) Turloughs over 10 hectares: vegetation survey and evaluation
GIS Operations :	Goodwillie map scanned and georectified. Turlough as outlined on map digitised. New Turlough dataset clipped to SAC Boundary. Expert opinion used as necessary to resolve any issues arising
Used For :	3180 (map 2)
<hr/>	
Year :	2014
Title :	Scientific Basis for Raised Bog Conservation in Ireland
GIS Operations :	RBSB13_SACs_ARB_DRB dataset, RBSB13_SACs_2012_HB dataset, RBSB13_SACs_DrainagePatterns_5k dataset and RBSB13_SAC_LIDAR_DTMs dataset clipped to SAC boundary. Expert opinion used as necessary to resolve any issues arising
Used For :	potential 7110; digital elevation model; drainage patterns (maps 3 and 5)
<hr/>	
Year :	2013
Title :	Raised Bog Monitoring and Assessment Survey 2013
GIS Operations :	RBMA13_ecotope_map dataset clipped to SAC boundary. Appropriate ecotopes selected and exported to new dataset. Expert opinion used as necessary to resolve any issues arising
Used For :	7110 ecotopes (map 4)
<hr/>	

3180 Turloughs

To maintain the favourable conservation condition of Turloughs in Lisnageeragh Bog and Ballinastack Turlough SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Current mapped area of Ballinastack Turlough is 23.7ha. See map 2. See turloughs supporting document for further details
Habitat distribution	Occurrence	No decline, subject to natural processes	The location of Ballinastack Turlough is shown in map 2. See turloughs supporting document for further details
Hydrological regime: flood duration, frequency, area, depth; permanently flooded area	Various	Maintain/restore appropriate natural hydrological regime necessary to support the natural structure and functioning of the habitat	Hydrological regime is sub-divided into more detailed attributes in the turloughs supporting document
Soil type: area	Hectares	Maintain variety, area and extent of soil types necessary to support turlough vegetation and other biota	See turloughs supporting document for further details
Soil nutrient status: nitrogen and phosphorous	N and P concentration in soil	Maintain/restore nutrient status appropriate to soil types	See turloughs supporting document for further details
Physical structure: bare ground	Presence	Maintain sufficient wet bare ground, as appropriate	See turloughs supporting document for further details
Chemical processes: calcium carbonate deposition and concentration	CaCO ₃ deposition rate/soil concentration	Maintain CaCO ₃ deposition rate/soil concentration	See turloughs supporting document for further details
Water quality: nutrients; colour; phytoplankton; epiphyton	Various	Maintain appropriate water quality to support the natural structure and functioning of the habitat	Water quality is sub-divided into more detailed attributes in the turloughs supporting document
Active peat formation	Flood duration	Restore active peat formation, where appropriate	See turloughs supporting document for further details
Vegetation composition: area of vegetation communities	Hectares	Maintain area of sensitive and high conservation value vegetation communities/units	See turloughs supporting document for further details
Vegetation composition: vegetation zonation	Distribution	Maintain vegetation zonation/mosaic characteristic of the site	See turloughs supporting document for further details
Vegetation structure: sward height	Centimetres	Maintain sward heights appropriate to the vegetation unit, and a variety of sward heights across the turlough	See turloughs supporting document for further details
Typical species: terrestrial, wetland and aquatic plants, invertebrates and birds	Presence	Maintain typical species	Typical species is sub-divided into more detailed attributes in the turloughs supporting document
Fringing habitats: area	Hectares	Maintain marginal fringing habitats that support turlough vegetation, invertebrate, mammal and/or bird populations	See turloughs supporting document for further details. See also the conservation objective for Active raised bogs (7110)

Vegetation structure:
turlough woodland

Species diversity and woodland structure

Maintain appropriate turlough woodland diversity and structure

See turloughs supporting document for further details

7110 Active raised bogs

To restore the favourable conservation condition of Active raised bogs in Lisnageeragh Bog and Ballinastack Turlough SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Restore area of active raised bog to 58.8ha, subject to natural processes	Active Raised Bog (ARB) habitat was mapped at 29.6ha by Fernandez et al. (2014). Area of Degraded Raised Bog (DRB) on the High Bog (HB) has been modelled as 38.0ha. See map 3. However, it is estimated that only 26.6ha is potentially restorable to ARB by drain blocking. The total potential ARB on the HB is therefore estimated to be 56.2ha. Eco-hydrological assessments of the cutover estimates that an additional 2.6ha of bog forming habitats could be restored. The long term target for ARB is therefore 58.8ha. See raised bog supporting document for further details on this and following attributes
Habitat distribution	Occurrence	Restore the distribution and variability of active raised bog across the SAC. See map 4 for distribution in 2012	ARB habitat at Lisnageeragh Bog is central and sub-central ecotopes and active flush. ARB currently occurs mainly in the south-central part of the bog. DRB occurs on both the south-central and northern parts of the bog, which will require restoration measures. There is also potential for ARB restoration on cutover areas of the bog (see area target above)
High bog area	Hectares	No decline in extent of high bog necessary to support the development and maintenance of active raised bog. See map 3	The area of high bog within Lisnageeragh Bog SAC in 2012 (latest figure available) was 269.5ha (DAHG, 2014)
Hydrological regime: water levels	Centimetres	Restore appropriate water levels throughout the site	For ARB, mean water level needs to be near or above the surface of the bog lawns for most of the year. Seasonal fluctuations should not exceed 20cm, and should only be 10cm below the surface, except for very short periods of time. Open water is often characteristic of soak systems
Hydrological regime: flow patterns	Flow direction; slope	Restore, where possible, appropriate high bog topography, flow directions and slopes. See map 5 for current situation	ARB depends on mean water levels being near or above the surface of bog lawns for most of the year. Long and gentle slopes are the most favourable to achieve these conditions. Changes to flow directions due to subsidence of bogs can radically change water regimes and cause drying out of high quality ARB areas and soak systems
Transitional areas between high bog and adjacent mineral soils (including cutover areas)	Hectares; distribution	Restore adequate transitional areas to support/protect active raised bog and the services it provides	Studies suggest that the ARB is threatened due to water loss from past drainage and peat-cutting at Lisnageeragh Bog. Only remnant semi-natural margins occur between the bog and the turlough in the north of the SAC. See also the conservation objective for Turloughs (3180). Eco-hydrological assessments have evaluated the potential for ARB restoration on cutover areas (see note for habitat area attribute above)
Vegetation quality: central ecotope, active flush, soaks, bog woodland	Hectares	Restore 29.4ha of central ecotope/active flush/soaks/bog woodland as appropriate	At least 50% of ARB habitat should be high quality (i.e. central ecotope, active flush, soaks, bog woodland). Target area of active raised bog for the site has been set at 58.8ha (see area target above)
Vegetation quality: microtopographical features	Hectares	Restore adequate cover of high quality microtopographical features	High quality microtopography (hummocks, hollows and pools) is well developed in the southern part of Lisnageeragh Bog

Vegetation quality: bog moss (<i>Sphagnum</i>) species	Percentage cover	Restore adequate cover of bog moss (<i>Sphagnum</i>) species to ensure peat-forming capacity	<i>Sphagnum</i> cover varies naturally across Ireland with relatively high cover in the east to lower cover in the west. Hummock forming species such as <i>Sphagnum austinii</i> are particularly good peat formers. <i>Sphagnum</i> cover and distribution also varies naturally across a site
Typical ARB species: flora	Occurrence	Restore, where appropriate, typical active raised bog flora	Typical flora species include widespread species, as well as those with more restricted distributions but typical of the habitat's subtypes or geographical range
Typical ARB species: fauna	Occurrence	Restore, where appropriate, typical active raised bog fauna	Typical fauna species include widespread species, as well as those with more restricted distributions but typical of the habitat's subtypes or geographical range
Elements of local distinctiveness	Occurrence	Maintain features of local distinctiveness, subject to natural processes	Lisnageeragh Bog is noted for the presence of a number of flushes on the high bog and the transition to turlough in the northern part of the SAC
Negative physical indicators	Percentage cover	Negative physical features absent or insignificant	Negative physical indicators include: bare peat, algae dominated pools and hollows, marginal cracks, tear patterns, subsidence features such as dry mineral mounds /ridges emerging or expanding and evidence of burning
Vegetation composition: native negative indicator species	Percentage cover	Native negative indicator species at insignificant levels	Disturbance indicators include species indicative of conditions drying out such as abundant bog asphodel (<i>Narthecium ossifragum</i>), deergrass (<i>Trichophorum germanicum</i>) and harestail cotton-grass (<i>Eriophorum vaginatum</i>) forming tussocks; abundant magellanic bog-moss (<i>Sphagnum magellanicum</i>) in pools previously dominated by <i>Sphagnum</i> species typical of very wet conditions (e.g. feathery bog-moss (<i>S. cuspidatum</i>)); and indicators of frequent burning events such as abundant <i>Cladonia floerkeana</i> and high cover of carnation sedge (<i>Carex panicea</i>) (particularly in true midlands raised bogs)
Vegetation composition: non-native invasive species	Percentage cover	Non-native invasive species at insignificant levels and not more than 1% cover	Most common non-native invasive species include lodgepole pine (<i>Pinus contorta</i>), rhododendron (<i>Rhododendron ponticum</i>) and pitcherplant (<i>Sarracenia purpurea</i>)
Air quality: nitrogen deposition	kg N/ha/year	Air quality surrounding bog close to natural reference conditions. The total N deposition should not exceed 5kg N/ha/yr	Change in air quality can result from fertiliser drift; adjacent quarry activities; or other atmospheric inputs. The critical load range for ombrotrophic bogs has been set as between 5 and 10kg N/ha/yr (Bobbink and Hettelingh, 2011). The latest N deposition figures for the area around Lisnageeragh Bog suggests that the current level is approximately 11.4kg N/ha/yr (Henry and Aherne, 2014)
Water quality	Hydrochemical measures	Water quality on the high bog and in transitional areas close to natural reference conditions	Water chemistry within raised bogs is influenced by atmospheric inputs (rainwater). However, within soak systems, water chemistry is influenced by other inputs such as focused flow or interaction with underlying substrates. Water chemistry in areas surrounding the high bog varies due to influences of different water types (bog water, regional groundwater and run-off from surrounding mineral lands)

Conservation Objectives for : Lisnageeragh Bog and Ballinastack Turlough SAC [000296]

7120 Degraded raised bogs still capable of natural regeneration

The long-term aim for Degraded raised bogs still capable of natural regeneration is that its peat-forming capability is re-established; therefore, the conservation objective for this habitat is inherently linked to that of Active raised bogs (7110) and a separate conservation objective has not been set in Lisnageeragh Bog and Ballinastack Turlough SAC


Attribute	Measure	Target	Notes
------------------	----------------	---------------	--------------


Conservation Objectives for : Lisnageeragh Bog and Ballinastack Turlough SAC [000296]


7150 Depressions on peat substrates of the Rhynchosporion

Depressions on peat substrates of the Rhynchosporion is an integral part of good quality Active raised bogs (7110) and thus a separate conservation objective has not been set for the habitat in Lisnageeragh Bog and Ballinastack Turlough SAC

Attribute	Measure	Target	Notes
-----------	---------	--------	-------



Legend
 Lisnageeragh Bog and Ballinastack Turlough SAC 000296

 *An Roinn Ealaíon, Oidhreachta agus Gaeltachta*
 Department of Arts, Heritage and the Gaeltacht

MAP 1:
LISNAGEERAGH BOG AND BALLINASTACK TURLOUGH SAC CONSERVATION OBJECTIVES SAC DESIGNATION
 Map to be read in conjunction with the NPWS Conservation Objectives Document.


SITE CODE:
SAC 000296; version 3. Co. Galway

0 250 500 750 1,000 m

The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
 Ordnance Survey of Ireland Licence No EN 0059216. © Ordnance Survey of Ireland Government of Ireland.


Níl sna teorainneacha ar na léarscáileanna ach nod garshuimhach ginearálta. Féadfar athbheithnihe a déanamh ar theorainneacha na gceantar comharthaite. Suirbhéarachta Ordonáis na hÉireann Ceadúnas Uimh EN 0059216. © Suirbhéarachta Ordonáis na hÉireann Rialtas na hÉireann.



Map Version 1
Date: March 2016



Legend


- Lisnageeragh Bog and Ballinastack Turlough SAC 000296
- 3180 *Turloughs






Legend

- Lisnageeragh Bog and Ballinastack Turlough SAC 000296
- High Bog Boundary
- Potential 7110 *Active Raised Bogs




Legend


 Lisnageeragh Bog and Ballinastack Turlough SAC 000296


 High Bog Boundary

Active Raised Bogs Ecotopes


 Central ecotope

 Soaks / active flush

 Sub-central ecotope



Legend



 Lisnageeragh Bog and Ballinastack Turlough SAC 000296

 High Bog Boundary

 Drainage Patterns

 Contours

Elevation



National Parks and Wildlife Service

Conservation Objectives Series

Lough Corrib SAC 000297



An Roinn Ealaíon, Oidhreachta,
Gnóthaí Réigiúnacha, Tuaithe agus Gaeltachta

Department of Arts, Heritage,
Regional, Rural and Gaeltacht Affairs



**National Parks and Wildlife Service,
Department of Arts, Heritage, Regional, Rural and Gaeltacht Affairs,
7 Ely Place, Dublin 2, Ireland.
Web: www.npws.ie
E-mail: nature.conservation@ahg.gov.ie**

Citation:

**NPWS (2017) Conservation Objectives: Lough Corrib SAC 000297. Version 1.
National Parks and Wildlife Service, Department of Arts, Heritage, Regional,
Rural and Gaeltacht Affairs.**

**Series Editor: Rebecca Jeffrey
ISSN 2009-4086**

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

* indicates a priority habitat under the Habitats Directive

000297	Lough Corrib SAC
1029	Freshwater Pearl Mussel <i>Margaritifera margaritifera</i>
1092	White-clawed Crayfish <i>Austropotamobius pallipes</i>
1095	Sea Lamprey <i>Petromyzon marinus</i>
1096	Brook Lamprey <i>Lampetra planeri</i>
1106	Salmon <i>Salmo salar</i>
1303	Lesser Horseshoe Bat <i>Rhinolophus hipposideros</i>
1355	Otter <i>Lutra lutra</i>
1393	Slender Green Feather-moss <i>Drepanocladus vernicosus</i>
1833	Slender Naiad <i>Najas flexilis</i>
3110	Oligotrophic waters containing very few minerals of sandy plains (Littorelletalia uniflorae)
3130	Oligotrophic to mesotrophic standing waters with vegetation of the Littorelletea uniflorae and/or Isoeto-Nanojuncetea
3140	Hard oligo-mesotrophic waters with benthic vegetation of Chara spp.
3260	Water courses of plain to montane levels with the Ranunculion fluitantis and Callitriche-Batrachion vegetation
6210	Semi-natural dry grasslands and scrubland facies on calcareous substrates (Festuco-Brometalia) (* important orchid sites)
6410	Terrestrial meadows on calcareous, peaty or clayey-silt-laden soils (Molinion caeruleae)
7110	Active raised bogsE
7120	Degraded raised bogs still capable of natural regeneration
7150	Depressions on peat substrates of the Rhynchosporion
7210	Calcareous fens with <i>Cladonia</i> and species of the Caricion davallianaeE
7220	Petrifying springs with tufa formation (Cratoneurion)E
7230	Alkaline fens
8240	Limestone pavementsE
91A0	Old sessile oak woods with <i>Quercus</i> and <i>Ulmus</i> in the British Isles
91D0	Bog woodlandE

Please note that this SAC overlaps with Lough Corrib SPA (004042). It is adjacent to Galway Bay Complex SAC (000268), Maumturk Mountains SAC (002008), Connemara Bog Complex SAC (002034) and Monivea Bog SAC (002352). See map 2. The conservation objectives for this site should be used in conjunction with those for overlapping and adjacent sites as appropriate.

Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

NPWS Documents

Year :	1984
Title :	The vegetation of Irish lakes
Author :	Heuff, H.
Series :	Unpublished report to NPWS
Year :	1995
Title :	Mapping of proposed SAC rivers for <i>Margaritifera margaritifera</i> . A report for the National Parks and Wildlife Service on work carried out from August to October 1995
Author :	Moorkens, E.
Series :	Unpublished report to NPWS
Year :	1998
Title :	Conservation management of the white-clawed crayfish, <i>Austropotamobius pallipes</i>
Author :	Reynolds, J.D.
Series :	Irish Wildlife Manual No. 1
Year :	2002
Title :	<i>Najas flexilis</i> in Donegal
Author :	Roden, C.M.
Series :	Unpublished report to NPWS
Year :	2004
Title :	Pilot project for monitoring populations of the freshwater pearl mussel <i>Margaritifera margaritifera</i> . Baseline survey of the Owenriff River SAC, County Galway
Author :	Moorkens, E.
Series :	Unpublished report to NPWS
Year :	2004
Title :	The distribution of <i>Najas flexilis</i> in Ireland 2002-2004
Author :	Roden, C.M.
Series :	Unpublished report to NPWS
Year :	2005
Title :	Monitoring populations of the freshwater pearl mussel <i>Margaritifera margaritifera</i> . Repeat survey of the Owenriff River cSAC, County Galway
Author :	Moorkens, E.
Series :	Unpublished report to NPWS
Year :	2006
Title :	Otter survey of Ireland 2004/2005
Author :	Bailey, M.; Rochford, J.
Series :	Irish Wildlife Manual No. 23
Year :	2006
Title :	Bat mitigation guidelines for Ireland
Author :	Kelleher, C.; Marnell, F.
Series :	Irish Wildlife Manual No. 25
Year :	2007
Title :	A survey of juvenile lamprey populations in the Corrib and Suir catchments
Author :	O'Connor, W.
Series :	Irish Wildlife Manual No. 26

Year :	2007
Title :	Supporting documentation for the Habitats Directive Conservation Status Assessment - backing documents. Article 17 forms and supporting maps
Author :	NPWS
Series :	Unpublished report to NPWS
Year :	2008
Title :	Monitoring populations of freshwater pearl mussel <i>Margaritifera margaritifera</i> . 2008 repeat survey of the Owenriff River SAC, County Galway
Author :	Moorkens, E.; Killeen, I.
Series :	Unpublished report to NPWS
Year :	2008
Title :	National survey of native woodlands 2003-2008
Author :	Perrin, P.M.; Martin, J.; Barron, S.; O'Neill, F.H.; McNutt, K.E.; Delaney, A.
Series :	Unpublished report to NPWS
Year :	2009
Title :	NS II freshwater pearl mussel sub-basin management plans: monitoring of the freshwater pearl mussel in the Owenriff
Author :	Moorkens, E.
Series :	Unpublished report to NPWS
Year :	2009
Title :	NS II freshwater pearl mussel sub-basin management plans: Report on biological monitoring of surface water quality in the Owenriff catchment (Corrib sub-catchment), Co. Galway
Author :	Williams, L.
Series :	Unpublished report to NPWS
Year :	2009
Title :	NS II freshwater pearl mussel sub-basin management plans: fisheries survey. Stage 1 report
Author :	Paul Johnston Associates
Series :	Unpublished report to NPWS
Year :	2010
Title :	A provisional inventory of ancient and long-established woodland in Ireland
Author :	Perrin, P.M.; Daly, O.H.
Series :	Irish Wildlife Manual No. 46
Year :	2010
Title :	A technical manual for monitoring white-clawed crayfish (<i>Austropotamobius pallipes</i>) in Irish lakes
Author :	Reynolds, J., O'Connor, W., O'Keeffe, C.; Lynn, D.
Series :	Irish Wildlife Manual No.45
Year :	2010
Title :	Second Draft Owenriff Freshwater Pearl Mussel Sub-basin Management Plan (2009-2015). March 2010
Author :	NPWS
Series :	Unpublished document to Department of the Environment, Heritage and Local Government
Year :	2011
Title :	Monitoring populations of the freshwater pearl mussel <i>Margaritifera margaritifera</i> – 2011 condition assessment survey of the freshwater pearl mussel in the Owenriff River, County Galway
Author :	Moorkens, E.
Series :	Unpublished report to NPWS

Year :	2012
Title :	Ireland Red List No. 8: Bryophytes
Author :	Lockhart, N.; Hodgetts, N.; Holyoak, D.
Series :	Ireland Red List series, NPWS
Year :	2013
Title :	Conservation status assessment for petrifying springs
Author :	Lyons, M.D.; Kelly, D.L.
Series :	Unpublished report to NPWS
Year :	2013
Title :	National otter survey of Ireland 2010/12
Author :	Reid, N.; Hayden, B.; Lundy, M.G.; Pietravalle, S.; McDonald, R.A.; Montgomery, W.I.
Series :	Irish Wildlife Manual No. 76
Year :	2013
Title :	Irish semi-natural grasslands survey 2007-2012
Author :	O'Neill, F.H.; Martin, J.R.; Devaney, F.M.; Perrin, P.M.
Series :	Irish Wildlife Manual No. 78
Year :	2013
Title :	National survey of limestone pavement and associated habitats in Ireland
Author :	Wilson, S.; Fernandez, F.
Series :	Irish Wildlife Manual No. 73
Year :	2013
Title :	Results of a monitoring survey of bog woodland
Author :	Cross, J.; Lynn, D.
Series :	Irish Wildlife Manual No. 69
Year :	2013
Title :	A survey of the benthic macrophytes of three hard-water lakes: Lough Bunny, Lough Carra and Lough Owel
Author :	Roden, C.; Murphy, P.
Series :	Irish Wildlife Manual No. 70
Year :	2013
Title :	The status of EU protected habitats and species in Ireland. Volume 2. Habitats assessments
Author :	NPWS
Series :	Conservation assessments
Year :	2013
Title :	The status of EU protected habitats and species in Ireland. Volume 3. Species assessments
Author :	NPWS
Series :	Conservation assessments
Year :	2014
Title :	Guidelines for a national survey and conservation assessment of upland vegetation and habitats in Ireland, Version 2.0
Author :	Perrin, P.M.; Barron, S.J.; Roche, J.R.; O'Hanrahan, B.
Series :	Irish Wildlife Manual No. 79
Year :	2014
Title :	Raised Bog Monitoring and Assessment Survey 2013
Author :	Fernandez, F.; Connolly, K.; Crowley, W.; Denyer, J.; Duff, K.; Smith, G.
Series :	Irish Wildlife Manual No. 81

Year :	2014
Title :	Addergoole Bog (SAC 000297), Co.Galway, Site Report
Author :	Fernandez, F.; Connolly, K.; Crowley, W.; Denyer, J.; Duff, K.; Smith, G.
Series :	Raised bog monitoring and assessment survey 2013
Year :	2014
Title :	National raised bog SAC management plan
Author :	Department of Arts, Heritage and the Gaeltacht
Series :	Draft for consultation. 15 January 2014
Year :	2015
Title :	Habitats Directive Annex I lake habitats: a working interpretation for the purposes of site-specific conservation objectives and Article 17 reporting
Author :	O Connor, Á.
Series :	Unpublished document by NPWS
Year :	2015
Title :	Monitoring methods for <i>Hamatocaulis vernicosus</i> (Mitt.) Hedenäs (Slender green feather-moss) in the Republic of Ireland
Author :	Campbell, C.; Hodgetts, N.; Lockhart, N.
Series :	Irish Wildlife Manual No. 91
Year :	2015
Title :	Monitoring populations of the freshwater pearl mussel <i>Margaritifera margaritifera</i> – 2014 Monitoring survey of the Owenriff River, County Galway. March 2015
Author :	Moorkens, E.A.
Series :	Unpublished report to NPWS
Year :	2015
Title :	Monitoring populations of the freshwater pearl mussel <i>Margaritifera margaritifera</i> – 2015 Monitoring Survey of the Owenriff River, County Galway. November 2015
Author :	Moorkens, E.A.
Series :	Unpublished report to NPWS
Year :	2016
Title :	Monitoring guidelines for the assessment of petrifying springs in Ireland
Author :	Lyons, M.D.; Kelly, D.L.
Series :	Irish Wildlife Manual No. 94
Year :	2016
Title :	Ireland Red List No. 10: Vascular Plants
Author :	Wyse Jackson, M.; FitzPatrick, Ú.; Cole, E.; Jebb, M.; McFerran, D.; Sheehy Skeffington, M.; Wright, M.
Series :	Ireland Red Lists series, NPWS
Year :	2017
Title :	2016 Survey and condition assessment of the population of the freshwater mussel <i>Margaritifera margaritifera</i> in the Owenriff River, County Galway
Author :	Moorkens, E.
Series :	Unpublished report to NPWS
Year :	2017
Title :	Lough Corrib SAC (site code: 297) Conservation objectives supporting document- <i>Najas flexilis</i> V1
Author :	NPWS
Series :	Conservation objectives supporting document

Year : 2017
Title : Lough Corrib SAC (site code: 297) Conservation objectives supporting document- raised bog habitats V1
Author : NPWS
Series : Conservation objectives supporting document

Other References

Year : 1982
Title : Otter survey of Ireland
Author : Chapman, P.J.; Chapman, L.L.
Series : Unpublished report to Vincent Wildlife Trust

Year : 1982
Title : Eutrophication of waters. Monitoring assessment and control
Author : OECD
Series : OECD, Paris

Year : 1984
Title : Studies on the biology of freshwater mussels (Lamellibranchia: Unionacea) in Ireland
Author : Ross, E.D.
Series : Unpublished MSc Thesis. National University of Ireland, Galway

Year : 1988
Title : The reproductive biology of freshwater mussels in Ireland, with observations on their distribution and demography
Author : Ross, E.D.
Series : Unpublished Ph.D. Thesis, National University of Ireland, Galway

Year : 1989
Title : The genera *Scorpidium* and *Hamatocaulis*, gen. nov., in northern Europe
Author : Hedenäs, L.
Series : Lindbergia, 15: 8-36

Year : 1990
Title : The phytosociology and ecology of the aquatic and wetland communities of the lower Lough Corrib basin, County Galway
Author : Mooney, E.P.; O'Connell, M.
Series : Proceedings of the Royal Irish Academy. Section B: Biological, Geological, and Chemical Science, 90B(5): 58-97

Year : 1991
Title : The spatial organization of otters (*Lutra lutra*) in Shetland
Author : Kruuk, H.; Moorhouse, A.
Series : Journal of Zoology, 224: 41-57

Year : 1994
Title : The ecological status of Lough Corrib, Ireland, as indicated by physiographic factors, water chemistry and macrophytic flora
Author : Krause, W.; King, J.J.
Series : Vegetatio, 110: 149-161

Year : 1996
Title : Studies on the biology and ecology of *Margaritifera* in Ireland
Author : Moorkens, E.
Series : Unpublished Ph.D. thesis, University of Dublin, Trinity College.

Year : 2000
Title : Colour in Irish lakes
Author : Free, G.; Allott, N.; Mills, P.; Kennelly, C.; Day, S.
Series : Verhandlungen Internationale Vereinigung für theoretische und angewandte Limnologie, 27: 2620-2623

Year : 2001
Title : Aquatic plants in Britain and Ireland
Author : Preston, C.D.; Croft, J.M.
Series : Harley Books, Colchester

Year : 2002
Title : Reversing the habitat fragmentation of British woodlands
Author : Peterken, G.
Series : WWF-UK, London

Year : 2002
Title : A survey of the white-clawed crayfish (*Austropotamobius pallipes*) Lereboullet and of water quality in two catchments of eastern Ireland
Author : Demers, A.; Reynolds, J.D.
Series : Bulletin Francais de la Peche et de la Pisciculture, 367: 729-740

Year : 2002
Title : Deterioration of Atlantic soft water macrophyte communities by acidification, eutrophication and alkalinisation
Author : Arts, G.H.P.
Series : Aquatic Botany, 73: 373-393

Year : 2003
Title : Monitoring the river, sea and brook lamprey, *Lampetra fluviatilis*, *L. planeri* and *Petromyzon marinus*
Author : Harvey, J.; Cowx, I.
Series : Conserving Natura 2000 Rivers Monitoring Series No. 5. English Nature, Peterborough

Year : 2003
Title : Ecology of watercourses characterised by *Ranunculus fluitans* and *Callitriche-Batrachion* Vegetation
Author : Hatton-Ellis, T.W.; Grieve, N.
Series : Conserving Natura 2000 Rivers Ecology Series No. 11. English Nature, Peterborough

Year : 2003
Title : Identifying lamprey. A field key for sea, river and brook lamprey
Author : Gardiner, R.
Series : Conserving Natura 2000 rivers, Conservation techniques No. 4. English Nature, Peterborough

Year : 2004
Title : The ecology of *Najas flexilis*
Author : Wingfield, R.A.; Murphy, K.J.; Hollingsworth, P.; Gaywood, M.J.
Series : Scottish Natural Heritage Commissioned Report No. 017 (ROAME No. F98PA02)

Year : 2005
Title : Common standards monitoring guidance for freshwater fauna
Author : JNCC
Series : Joint Nature Conservation Committee, Peterborough

Year : 2006
Title : Otters - ecology, behaviour and conservation
Author : Kruuk, H.
Series : Oxford University Press

Year :	2006
Title :	The status of host fish populations and fish species richness in European freshwater pearl mussel (<i>Margaritifera margaritifera</i>) streams
Author :	Geist, J.; Porkka, M.; Kuehn, R.
Series :	Aquatic Conservation: Marine and Freshwater Ecosystems, 16: 251-266
Year :	2006
Title :	A reference-based typology and ecological assessment system for Irish lakes. Preliminary investigations. Final report. Project 2000-FS-1-M1 Ecological assessment of lakes pilot study to establish monitoring methodologies EU (WFD)
Author :	Free, G.; Little, R.; Tierney, D.; Donnelly, K.; Coroni, R.
Series :	EPA, Wexford
Year :	2007
Title :	Evolutionary history of lamprey paired species <i>Lampetra fluviatilis</i> L. and <i>Lampetra planeri</i> Bloch as inferred from mitochondrial DNA variation
Author :	Espanhol, R.; Almeida, P.R.; Alves, M.J.
Series :	Molecular Ecology, 16: 1909-1924
Year :	2008
Title :	The lesser horseshoe bat conservation handbook
Author :	Schofield, H.W.
Series :	The Vincent Wildlife Trust
Year :	2008
Title :	Water Quality in Ireland 2004-2006
Author :	Clabby, K.J.; Bradley, C.; Craig, M.; Daly, D.; Lucey, J.; McGarrigle, M.; O'Boyle, S.; Tierney, D.; Bowman, J.
Series :	EPA, Wexford
Year :	2009
Title :	The identification, characterization and conservation value of isoetid lakes in Ireland
Author :	Free, G.; Bowman, J.; McGarrigle, M.; Little, R.; Coroni, R.; Donnelly, K.; Tierney, D.; Trodd, W.
Series :	Aquatic Conservation: Marine and Freshwater Ecosystems, 19 (3): 264–273
Year :	2010
Title :	Otter tracking study of Roaringwater Bay
Author :	De Jongh, A.; O'Neill, L.
Series :	Unpublished draft report to NPWS
Year :	2010
Title :	Addressing the conservation and rehabilitation of <i>Margaritifera margaritifera</i> populations in the Republic of Ireland within the framework of the habitats and species directive
Author :	Moorkens, E.
Series :	Journal of Conchology, 40: 339
Year :	2010
Title :	Water quality in Ireland 2007-2009
Author :	McGarrigle, M.; Lucey, J.; Ó Cinnéide, M.
Series :	EPA, Wexford
Year :	2011
Title :	Comparison of field- and GIS-based assessments of barriers to Atlantic salmon migration: a case study in the Nore Catchment, Republic of Ireland
Author :	Gargan, P.G.; Roche, W.K.; Keane, S.; King, J.J.; Cullagh, A.; Mills, P.; O'Keeffe, J.
Series :	Journal of Applied Ichthyology, 27 (Suppl. 3): 66-72

Year :	2011
Title :	Review and revision of empirical critical loads and dose-response relationships. Proceedings of an expert workshop, Noordwijkerhout, 23-25 June 2010
Author :	Bobbink, R.; Hettelingh, J.P.
Series :	RIVM report 680359002, Coordination Centre for Effects, National Institute for Public Health and the Environment (RIVM)
Year :	2012
Title :	A report on the sub-littoral environment around selected navigation markers in the north west sector of Lough Corrib
Author :	Roden, C.
Series :	Unpublished report to RPS Group
Year :	2013
Title :	Aspects of brook lamprey (<i>Lampetra planeri</i> Bloch) spawning in Irish waters
Author :	Rooney, S.M.; O'Gorman, N.M.; Green, F.; King, J.J.
Series :	Biology and Environment: Proceedings of the Royal Irish Academy, 113B(1): 13-25
Year :	2013
Title :	Conservation of selected legally protected and Red Listed bryophytes in Ireland
Author :	Campbell, C.
Series :	Unpublished Ph.D. Thesis, Trinity College Dublin
Year :	2014
Title :	Nitrogen deposition and exceedance of critical loads for nutrient nitrogen in Irish grasslands
Author :	Henry, J.; Aherne, J.
Series :	Science of the Total Environment, 470–471: 216–223
Year :	2014
Title :	Assessing near-bed velocity in a recruiting population of the endangered freshwater pearl mussel (<i>Margaritifera margaritifera</i>) in Ireland
Author :	Moorkens, E.; Killeen, I.
Series :	Aquatic Conservation: Marine and Freshwater Ecosystems, 24(6): 853-862
Year :	2015
Title :	Behaviour of sea lamprey (<i>Petromyzon marinus</i> L.) at man-made obstacles during upriver spawning migration: use of telemetry to assess efficacy of weir modifications for improved passage
Author :	Rooney, S.M.; Wightman, G.D.; O Conchuir, R.; King, J.J.
Series :	Biology and Environment: Proceedings of the Royal Irish Academy, 115B: 1-12
Year :	2015
Title :	River engineering works and lamprey ammocoetes; impacts, recovery, mitigation
Author :	King, J.J.; Wightman, G.D.; Hanna, G.; Gilligan, N.
Series :	Water and Environment Journal, 29: 482-488
Year :	2015
Title :	The flora and conservation status of petrifying springs in Ireland
Author :	Lyons, M.D.
Series :	Unpublished Ph.D. Thesis, Trinity College Dublin
Year :	2016
Title :	The Status of Irish Salmon Stocks in 2015 with Precautionary Catch Advice for 2016
Author :	SSCS (Standing Scientific Committee on Salmon)
Series :	Independent Scientific Report to Inland Fisheries Ireland

Spatial data sources

Year :	2008
Title :	OSi 1:5000 IG vector dataset
GIS Operations :	WaterPolygons feature class clipped to the SAC boundary. Expert opinion used to identify Annex I habitats and to resolve any issues arising
Used For :	3110, 3130, 3140 (map 3)
Year :	2014
Title :	Scientific Basis for Raised Bog Conservation in Ireland
GIS Operations :	RBSB13_SACs_ARB_DRB dataset, RBSB13_SACs_2012_HB dataset and RBSB13_SACs_DrainagePatterns_5k dataset clipped to SAC boundary. Expert opinion used as necessary to resolve any issues arising
Used For :	potential 7110; digital elevation model; drainage patterns (maps 4 and 6)
Year :	2003
Title :	Turf Cutting Impact Assesment Project
GIS Operations :	Ecotope dataset clipped to SAC boundary. Appropriate ecotopes selected and exported to new dataset. Expert opinion used as necessary to resolve any issues arising
Used For :	7110 ecotopes (map 5)
Year :	2013
Title :	Raised Bog Monitoring and Assessment Survey 2013
GIS Operations :	RBMA13_ecotope_map dataset clipped to SAC boundary. Appropriate ecotopes selected and exported to new dataset. Expert opinion used as necessary to resolve any issues arising
Used For :	7110 ecotopes, 91D0 (maps 5 and 8)
Year :	2013
Title :	National Survey of Limestone Pavement and Associated Habitats in Ireland distribution data
GIS Operations :	Dataset clipped to the SAC boundary. Expert opinion used as necessary to resolve any issues arising
Used For :	8240 (map 7)
Year :	Revision 2010
Title :	National Survey of Native Woodlands 2003-2008. Version 1
GIS Operations :	QIs selected; clipped to SAC boundary. Expert opinion used as necessary to resolve any issues arising
Used For :	91A0 (map 8)
Year :	Revision 2012
Title :	Margaritifera Sensitive Areas data
GIS Operations :	Relevant catchment boundary identified. Expert opinion used as necessary to resolve any issues arising
Used For :	1029 (map 9)
Year :	2017
Title :	NPWS rare and threatened species database
GIS Operations :	Dataset created from spatial references in database records. Expert opinion used as necessary to resolve any issues arising
Used For :	1029, 1092, 1393 (maps 9 and 10)
Year :	2012
Title :	NPWS lesser horseshoe bat database
GIS Operations :	Relevant roost identified by clipping to SAC boundary. Expert opinion used as necessary to resolve any issues arising
Used For :	1303 (map 11)
Year :	2007
Title :	Forest Inventory and Planning System (FIPS)
GIS Operations :	Dataset clipped to 2.5km buffer centred on roost location
Used For :	1303 (map 11)

Year :	2010
Title :	OSi 1:5000 IG vector dataset
GIS Operations :	Creation of 80m buffer on aquatic side of lake data; creation of 10m buffer on terrestrial side of lake data. These datasets combined with derived OSi Discovery Series river and canal datasets. Overlapping regions investigated and resolved; resulting dataset clipped to SAC boundary. Expert opinion used as necessary to resolve any issues arising. Creation of 250m buffer on aquatic side of lake boundary to highlight potential commuting points
Used For :	1355 (map 12)
Year :	2005
Title :	OSi Discovery series vector data
GIS Operations :	Creation of 10m buffer on terrestrial side of river banks data; creation of 20m buffer applied to canal centreline data. Creation of 20m buffer applied to river and stream centreline data; These datasets combined with derived OSi 1:5000 vector lake buffer data. Overlapping regions investigated and resolved; resulting dataset clipped to SAC boundary. Expert opinion used as necessary to resolve any issues arising
Used For :	1355 (no map)
Year :	2013
Title :	<i>Najas flexilis</i> data
GIS Operations :	Lake habitat for species clipped to SAC boundary. Expert opinion used as necessary to resolve any issues arising
Used For :	1833 (map 13)

3110 Oligotrophic waters containing very few minerals of sandy plains (Littorelletalia uniflorae)

To restore the favourable conservation condition of Oligotrophic waters containing very few minerals of sandy plains (*Littorelletalia uniflorae*) in Lough Corrib SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	The distribution of lake habitat 3110 in Lough Corrib SAC has not been fully surveyed. Krause and King (1994) recorded it in the "western arm proper". Within Lough Corrib, it is likely to be restricted to this 'western arm' (the north-western bay). It may, however, occur elsewhere along the northern or western shoreline of Lough Corrib, in Ballydoo Lough (N. of Corrib) and in small lakes in the Owenriff catchment. Two measures of extent should be used: 1. the area of the lake itself; 2. the extent of the vegetation communities/zones that typify the habitat. Further information on this and all other attributes is provided in the lake habitats supporting document for the purposes of site-specific conservation objectives and Article 17 reporting (O Connor, 2015)
Habitat distribution	Occurrence	No decline, subject to natural processes. See map 3 for indicative lake habitat distribution	The selection of the SAC for lake habitat 3110 was based on data on the macrophyte flora of Lough Corrib, particularly the north-western bay, from Krause and King (1994). Despite the occurrence of slender naiad (<i>Najas flexilis</i>) in this bay, the vegetation appears to be typical of lake habitat 3110. It is possible that habitat 3110 occurs elsewhere along the northern and western shores of Lough Corrib, on acid geology, as well as within some smaller lakes in the SAC, e.g. Lough Ateean in the Owenriff catchment
Typical species	Occurrence	Typical species present, in good condition, and demonstrating typical abundances and distribution	For lists of typical plant species, see the Article 17 habitat assessment for 3110 (NPWS, 2013) and O Connor (2015)
Vegetation composition: characteristic zonation	Occurrence	All characteristic zones should be present, correctly distributed and in good condition	Further work is necessary to describe the characteristic zonation and other spatial patterns in lake habitat 3110 (see O Connor, 2015)
Vegetation distribution: maximum depth	Metres	Restore maximum depth of vegetation, subject to natural processes	The maximum depth of vegetation is likely to be specific to the lake shoreline in question. Further work is necessary to develop indicative targets for lake habitat 3110. Recent investigations have indicated a significant decline in depth of colonisation throughout Lough Corrib (Roden, 2012; Roden and Murphy, in prep.)
Hydrological regime: water level fluctuations	Metres	Maintain appropriate natural hydrological regime necessary to support the habitat	Fluctuations in lake water level are typical in Ireland, but can be amplified by activities such as abstraction and drainage. Increased water level fluctuations can increase wave action, up-root vegetation, increase turbidity, alter the substratum and lead to release of nutrients from the sediment. The hydrological regime of the lakes must be maintained so that the area, distribution and depth of the lake habitat and its constituent/characteristic vegetation zones and communities are not reduced

Lake substratum quality	Various	Restore appropriate substratum type, extent and chemistry to support the vegetation	Research is required to further characterise the substratum types (particle size and origin) and substratum quality (notably pH, calcium, iron and nutrient concentrations) favoured by each of the five Annex I lake habitats in Ireland. It is likely that the oligotrophic soft water habitat is associated with a range of nutrient-poor substrates, from stones, cobble and gravel, through sands, silt, clay and peat. Substratum particle size is likely to vary with depth and along the shoreline within a single lake. Lake substratum quality in Lough Corrib is impaired as a result of peat deposition and possibly also nutrient enrichment (Roden, 2012; Roden and Murphy, in prep.)
Water quality: transparency	Metres	Restore appropriate Secchi transparency. There should be no decline in Secchi depth/transparency	Transparency relates to light penetration and, hence, to the depth of colonisation of vegetation. It can be affected by phytoplankton blooms, water colour and turbidity. Specific targets have yet to be established for lake habitat 3110. Habitat 3110 is associated with very clear water. The OECD fixed boundary system set transparency targets for oligotrophic lakes of $\geq 6\text{m}$ annual mean Secchi disk depth, and $\geq 3\text{m}$ annual minimum Secchi disk depth. Free et al. (2009) found high isoetid abundance in lakes with Secchi depths of more than 3m. Secchi depth did not exceed 2m in the 'western arm' of Lough Corrib and approximately 5m in the northern basin in 2012 (Roden, 2012)
Water quality: nutrients	$\mu\text{g/l P}$; mg/l N	Restore the concentration of nutrients in the water column to sufficiently low levels to support the habitat and its typical species	As a nutrient-poor habitat, oligotrophic and Water Framework Directive (WFD) 'high' status targets apply. Where a lake has nutrient concentrations that are lower than these targets, there should be no decline within class, i.e. no upward trend in nutrient concentrations. For lake habitat 3110, annual average Total Phosphorus (TP) concentration should be $\leq 10\mu\text{g/l TP}$, average annual total ammonia concentration should be $\leq 0.040\text{ mg/l N}$ and annual 95th percentile for total ammonia should be $\leq 0.090\text{ mg/l N}$. See also the European Communities Environmental Objectives (Surface Waters) Regulations 2009. TP has exceeded the $10\mu\text{g/l}$ target in Upper Corrib in a number of recent years (Clabby et al., 2008; McGarrigle et al., 2010)
Water quality: phytoplankton biomass	$\mu\text{g/l Chlorophyll } a$	Restore appropriate water quality to support the habitat, including high chlorophyll <i>a</i> status	Oligotrophic and WFD 'high' status targets apply to lake habitat 3110. Where a lake has a chlorophyll <i>a</i> concentration that is lower than this target, there should be no decline within class, i.e. no upward trend in phytoplankton biomass. The average growing season (March-October) chlorophyll <i>a</i> concentration must be $< 5.8\mu\text{g/l}$. The annual average chlorophyll <i>a</i> concentration should be $< 2.5\mu\text{g/l}$ and the annual peak chlorophyll <i>a</i> concentration should be $\leq 8.0\mu\text{g/l}$. See also the European Communities Environmental Objectives (Surface Waters) Regulations 2009. Maximum chlorophyll <i>a</i> in Upper Corrib has exceeded the OECD target of $\leq 8.0\mu\text{g/l}$ in five of the nine monitoring cycles from 1976-2009 (McGarrigle et al., 2010)
Water quality: phytoplankton composition	EPA phytoplankton composition metric	Maintain appropriate water quality to support the habitat, including high phytoplankton composition status	The Environmental Protection Agency (EPA) has developed a phytoplankton composition metric for nutrient enrichment of Irish lakes. As for other water quality indicators, lake habitat 3110 requires WFD high status
Water quality: attached algal biomass	Algal cover and EPA phytobenthos metric	Restore/maintain trace/absent attached algal biomass ($< 5\%$ cover) and high phytobenthos status	Nutrient enrichment can favour epiphytic and epipelic algae that can out-compete the submerged vegetation. The cover abundance of attached algae in lake habitat 3110 should, therefore, be trace/absent ($< 5\%$ cover). EPA phytobenthos can be used as an indicator of changes in attached algal biomass. As for other water quality indicators, lake habitat 3110 requires high phytobenthos status

Water quality: macrophyte status	EPA macrophyte metric (The Free Index)	Maintain high macrophyte status	Nutrient enrichment can favour more competitive submerged macrophyte species that out-compete the typical and characteristic species for the lake habitat. The EPA monitors macrophyte status for WFD purposes using the 'Free Index'. The target for lake habitat 3110 is high status or an Ecological Quality Ratio (EQR) for lake macrophytes of ≥ 0.90 , as defined in Schedule Five of the European Communities Environmental Objectives (Surface Waters) Regulations 2009
Acidification status	pH units; mg/l	Maintain appropriate water and sediment pH, alkalinity and cation concentrations to support the habitat, subject to natural processes	Acidification can impact on species abundance and composition in soft water lake habitats. In Europe, acidification of isoetid lakes can lead to loss of isoetids and dominance by submerged <i>Sphagnum</i> mosses and <i>Juncus bulbosus</i> (Arts, 2002). The specific requirements of lake habitat 3110, in terms of water and sediment pH, alkalinity and cation concentration, have not been determined. For lake habitat 3110, and adopting a precautionary approach based on Arts (2002), minimum pH should not be < 5.5 pH units. Maximum pH should be < 9.0 pH units, in line with the surface water standards established for soft waters (where water hardness is ≤ 100 mg/l calcium carbonate). See Schedule Five of the European Communities Environmental Objectives (Surface Water) Regulations (S.I. 272 of 2009)
Water colour	mg/l PtCo	Restore/maintain appropriate water colour to support the habitat	Increased water colour and turbidity decrease light penetration and can reduce the area of available habitat for lake macrophytes, particularly at the lower euphotic depths. The primary source of increased water colour in Ireland is disturbance to peatland. No habitat-specific or national standards for water colour currently exist. Studies have shown median colour concentrations in Irish lakes of 38mg/l PtCo (Free et al., 2000) and 33mg/l PtCo (Free et al., 2006). It is likely that the water colour in all Irish lake habitats would naturally be < 50 mg/l PtCo. Water colour can be very low (20mg/l PtCo or even < 10 mg/l PtCo) in oligotrophic soft water lakes (3110), where the peatland in the lake's catchment is intact. Roden (2012) recorded very dark water and poor visibility in Upper Corrib and the western arm
Dissolved organic carbon (DOC)	mg/l	Restore/maintain appropriate organic carbon levels to support the habitat	Dissolved (and particulate) organic carbon (OC) in the water column is linked to water colour and acidification (organic acids). Increasing DOC in water has been documented across the Northern Hemisphere, including afforested peatland catchments in Ireland. Damage and degradation of peatland, leading to decomposition of peat is likely to be the predominant source of OC in Ireland. OC in water promotes decomposition by fungi and bacteria that, in turn, releases dissolved nutrients. The increased biomass of decomposers can also impact directly on the characteristic lake communities through shading, competition, etc.
Turbidity	Nephelometric turbidity units/ mg/l SS/ other appropriate units	Restore/maintain appropriate turbidity to support the habitat	Turbidity can significantly affect the quantity and quality of light reaching rooted and attached vegetation and can, therefore, impact on lake habitats. The settlement of higher loads of inorganic or organic material on lake vegetation communities may also have impacts on sensitive, delicate species. Turbidity can increase as a result of re-suspension of material within the lake, higher loads entering the lake, or eutrophication. Turbidity measurement and interpretation is challenging. As a result, it is likely to be difficult to set habitat-specific targets for turbidity in lakes. Roden recorded low transparency, very hard water, poor visibility and peat deposition in Lough Corrib, all of which suggest increased turbidity (Roden, 2012)

Fringing habitat : Hectares
area and condition

Maintain the area and
condition of fringing
habitats necessary to
support the natural
structure and functioning
of habitat 3110

Most lake shorelines have fringing habitats of reedswamp, other swamp, fen, marsh or wet woodland that intergrade with and support the structure and functions of the lake habitat. Equally, fringing habitats are dependent on the lake, particularly its water levels, and support wetland communities and species of conservation concern. Many of the fringing wetland habitats support higher invertebrate and plant species-richness than the lake habitats themselves. Lough Corrib has important areas of fen, limestone pavement, marsh, reedswamp and some woodland along its shoreline. See also the conservation objectives for associated Annex I habitats and Annex II species within the SAC

Conservation Objectives for : Lough Corrib SAC [000297]

3130 Oligotrophic to mesotrophic standing waters with vegetation of the *Littorelletea uniflorae* and/or *Isoeto-Nanojuncetea*

To restore the favourable conservation condition of Oligotrophic to mesotrophic standing waters with vegetation of the *Littorelletea uniflorae* and/or *Isoeto-Nanojuncetea* in Lough Corrib SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	The full distribution and characteristics of lake habitat 3130 in Lough Corrib SAC have not been mapped. While the characteristic species slender naiad (<i>Najas flexilis</i>) was recorded in the western arm of Lough Corrib, that area appears to be dominated by lake habitat 3110, with lake habitat 3130 found towards the northern basin proper. The division between lake habitats 3130 and 3140 may be difficult to determine, and both habitats may occur throughout the lake. Habitat 3130 is thought likely to dominate Ballycurke Lake. Two measures of extent should be used: 1. the area of the lake itself; 2. the extent of the vegetation communities/zones that typify the habitat. For additional information see Krause and King (1994). Further information on this and all other attributes is contained in the lake habitats supporting document for the purposes of site-specific conservation objectives and Article 17 reporting (O Connor, 2015) and the <i>Najas flexilis</i> supporting document
Habitat distribution	Occurrence	No decline, subject to natural processes. See map 3 for indicative lake habitat distribution	The characteristics and distribution of lake habitat 3130 in Ireland are not yet fully understood. The Annex II macrophyte slender naiad (<i>Najas flexilis</i>) is considered to be characteristic of the habitat and occurs in Lough Corrib. Based on Environmental Protection Agency (EPA) macrophyte data, lake habitat 3130 is also likely to occur in Ballycurke Lough (Loch Bhaile Ui Choirc)
Typical species	Occurrence	Typical species present, in good condition, and demonstrating typical abundances and distribution	For lists of typical plant species, see the Article 17 habitat assessment for lake habitat 3130 (NPWS, 2013) and O Connor (2015)
Vegetation composition: characteristic zonation	Occurrence	All characteristic zones should be present, correctly distributed and in good condition	Further work is necessary to describe the characteristic zonation and other spatial patterns in lake habitat 3130 (see O Connor, 2015)
Vegetation distribution: maximum depth	Metres	Restore maximum depth of vegetation, subject to natural processes	The maximum depth of vegetation is likely to be specific to the lake shoreline in question. Further work is necessary to develop indicative targets for lake habitat 3130. Recent investigations have indicated a significant decline in depth of colonisation throughout Lough Corrib (Roden, 2012; Roden and Murphy, in prep.)
Hydrological regime: water level fluctuations	Metres	Maintain appropriate natural hydrological regime necessary to support the habitat	Fluctuations in lake water level are typical in Ireland, but can be amplified by activities such as abstraction and drainage. Increased water level fluctuations can increase wave action, up-root vegetation, increase turbidity, alter the substratum and lead to release of nutrients from the sediment. The hydrological regime of the lakes must be maintained so that the area, distribution and depth of the lake habitat and its constituent/characteristic vegetation zones and communities are not reduced

Lake substratum quality	Various	Restore appropriate substratum type, extent and chemistry to support the vegetation	Research is required to further characterise the substratum types (particle size and origin) and substratum quality (notably pH, calcium, iron and nutrient concentrations) favoured by each of the five Annex I lake habitats in Ireland. It is likely that lake habitat 3130 is associated with a range of substrate types that are more productive/base-rich relative to the substratum of lake habitat 3110. Substratum particle size is likely to vary with depth and along the shoreline within a single lake, however it should be noted that <i>Najas flexilis</i> is typically found on soft substrata of mud, silt or fine sand (Preston and Croft, 2001; Roden, 2002, 2004). Lake substratum quality in Lough Corrib is impaired as a result of peat deposition and possibly also nutrient enrichment (Roden, 2012; Roden and Murphy, in prep.)
Water quality: transparency	Metres	Restore appropriate Secchi transparency. There should be no decline in Secchi depth/transparency	Transparency relates to light penetration and, hence, to the depth of colonisation of vegetation. It can be affected by phytoplankton blooms, water colour and turbidity. Specific targets have yet to be established for lake habitat 3130 (O Connor, 2015). Habitat 3130 is associated with clear water, as evidenced by the growth of the character species <i>Najas flexilis</i> at depths of up to 10m. The OECD fixed boundary system set transparency targets for oligotrophic lakes of $\geq 6\text{m}$ annual mean Secchi disk depth, and $\geq 3\text{m}$ annual minimum Secchi disk depth. There is likely to be some variation across lakes with habitat 3130 in Secchi depth and site-specific conditions should also be considered. Secchi depth did not exceed 2m in the 'western arm' of Lough Corrib and approximately 5m in the northern basin in 2012 (Roden, 2012)
Water quality: nutrients	$\mu\text{g/l P}$; mg/l N	Restore the concentration of nutrients in the water column to sufficiently low levels to support the habitat and its typical species	Lake habitat 3130 is associated with high water quality, with low dissolved nutrients. It is naturally more productive than 3110, probably reflecting higher concentrations of nutrients such as calcium, rather than P alone. 3130 may reach favourable condition slightly above the oligotrophic boundary for nutrients, but in the absence of habitat-specific targets, the targets are Water Framework Directive (WFD) 'High Status' or oligotrophic (OECD, 1982). The "good-moderate" boundary is too enriched to support the habitat. Annual average Total Phosphorus (TP) concentration should be $\leq 10\mu\text{g/l}$ TP, average annual total ammonia should be $\leq 0.04\text{mg/l N}$ and annual 95th percentile for total ammonia should be $\leq 0.09\text{mg/l N}$. Where nutrient concentrations are lower, there should be no upward trend. See also the European Communities Environmental Objectives (Surface Waters) Regulations 2009. TP has exceeded the $10\mu\text{g/l}$ target in Upper Corrib in recent years (Clabby et al., 2008; McGarrigle et al., 2010)
Water quality: phytoplankton biomass	$\mu\text{g/l Chlorophyll } a$	Restore appropriate water quality to support the habitat, including high chlorophyll <i>a</i> status	Lake habitat 3130 is associated with high water quality, and naturally low algal growth. As for nutrients, the targets are WFD 'High Status' or oligotrophic (OECD, 1982). The "good-moderate" boundary is too enriched to support the habitat. The average growing season (March-October) chlorophyll <i>a</i> concentration must be $< 5.8\mu\text{g/l}$. The annual average chlorophyll <i>a</i> should be $< 2.5\mu\text{g/l}$ and the annual peak chlorophyll <i>a</i> should be $< 8.0\mu\text{g/l}$. Where a lake has a chlorophyll <i>a</i> concentration that is lower than this target, there should be no decline within class, i.e. no upward trend in phytoplankton biomass. See also the European Communities Environmental Objectives (Surface Waters) Regulations 2009. Maximum chlorophyll <i>a</i> in Upper Corrib has exceeded the OECD target of $\leq 8.0\mu\text{g/l}$ in five of the nine monitoring cycles from 1976-2009 (McGarrigle et al., 2010)

Water quality: phytoplankton composition	EPA phytoplankton composition metric	Maintain appropriate water quality to support the habitat, including high phytoplankton composition status	The EPA has developed a phytoplankton composition metric for nutrient enrichment of Irish lakes. As for other water quality indicators, lake habitat 3130 requires WFD high status
Water quality: attached algal biomass	Algal cover and EPA phytobenthos metric	Restore/maintain trace/absent attached algal biomass (<5% cover) and high phytobenthos status	Nutrient enrichment can favour epiphytic and epipelic algae that can out-compete the submerged vegetation. The cover abundance of attached algae in lake habitat 3130 should, therefore, be trace/absent (<5% cover). EPA phytobenthos can be used as an indicator of changes in attached algal biomass. As for other water quality indicators, lake habitat 3130 requires high phytobenthos status
Water quality: macrophyte status	EPA macrophyte metric (The Free Index)	Maintain high macrophyte status	Nutrient enrichment can favour more competitive submerged macrophyte species that out-compete the typical and characteristic species for the soft water lake habitat with base-rich influences (3130). The EPA monitors macrophyte status for WFD purposes using the 'Free Index'. The target for lake habitat 3130 is high status or an Ecological Quality Ratio (EQR) for lake macrophytes of ≥ 0.90 , as defined in Schedule Five of the European Communities Environmental Objectives (Surface Waters) Regulations 2009
Acidification status	pH units; mg/l	Maintain appropriate water and sediment pH, alkalinity and cation concentrations to support the habitat, subject to natural processes	Acidification can impact on species abundance and composition in habitat 3130. Acidification reduces the abundance and reproductive capacity of <i>Najas flexilis</i> (Wingfield et al., 2004). The specific requirements of lake habitat 3130, in terms of water and sediment pH, alkalinity and cation concentration, have not been determined. In line with targets for <i>Najas flexilis</i> , median pH values should be greater than 7 pH units. Water and sediment alkalinity and concentrations of cations (notably calcium) should be appropriate to the habitat. The target for WFD Acidification/Alkalisiation status is high. Maximum pH should be <9.0 pH units, in line with the surface water standards. See the European Communities Environmental Objectives (Surface Waters) Regulations 2009
Water colour	mg/l PtCo	Restore/maintain appropriate water colour to support the habitat	Increased water colour and turbidity decrease light penetration and can reduce the area of available habitat for lake macrophytes, particularly at the lower euphotic depths. The primary source of increased water colour in Ireland is disturbance to peatland. No habitat-specific or national standards for water colour currently exist. Studies have shown median colour concentrations in Irish lakes of 38mg/l PtCo (Free et al., 2000) and 33mg/l PtCo (Free et al., 2006). It is likely that the water colour in all Irish lake habitats would naturally be <50mg/l PtCo. Water colour is generally <30mg/l PtCo or, more naturally, <20mg/l PtCo in lake habitat 3130, where the peatland in the lake's catchment is intact. Roden (2012) recorded very dark water and poor visibility in Upper Corrib and the western arm
Dissolved organic carbon (DOC)	mg/l	Restore/maintain appropriate organic carbon levels to support the habitat	Dissolved (and particulate) organic carbon (OC) in the water column is linked to water colour and acidification (organic acids). Increasing DOC in water has been documented across the Northern Hemisphere, including afforested peatland catchments in Ireland. Damage and degradation of peatland, leading to decomposition of peat is likely to be the predominant source of OC in Ireland. OC in water promotes decomposition by fungi and bacteria that, in turn, releases dissolved nutrients. The increased biomass of decomposers can also impact directly on the characteristic lake communities through shading, competition, etc.

Turbidity	Nephelometric turbidity units/ mg/l SS/ other appropriate units	Restore/maintain appropriate turbidity to support the habitat	Turbidity can significantly affect the quantity and quality of light reaching rooted and attached vegetation and can, therefore, impact on lake habitats. The settlement of higher loads of inorganic or organic material on lake vegetation communities may also have impacts on sensitive, delicate species. Turbidity can increase as a result of re-suspension of material within the lake, higher loads entering the lake, or eutrophication. Turbidity measurement and interpretation is challenging. As a result, it is likely to be difficult to set habitat-specific targets for turbidity in lakes. Roden recorded low transparency, very hard water, poor visibility and peat deposition in Lough Corrib, all of which suggest increased turbidity (Roden, 2012)
Fringing habitat: area and condition	Hectares	Maintain the area and condition of fringing habitats necessary to support the natural structure and functioning of habitat 3130	Most lake shorelines have fringing habitats of reedswamp, other swamp, fen, marsh or wet woodland that intergrade with and support the structure and functions of the lake habitat. Equally, fringing habitats are dependent on the lake, particularly its water levels, and support wetland communities and species of conservation concern. Many of the fringing wetland habitats support higher invertebrate and plant species-richness than the lake habitats themselves. Lough Corrib has important areas of fen, limestone pavement, marsh, reedswamp and some woodland along its shoreline. See also the conservation objectives for associated Annex I habitats and Annex II species within the SAC

Conservation Objectives for : Lough Corrib SAC [000297]

3140 Hard oligo-mesotrophic waters with benthic vegetation of *Chara* spp.

To restore the favourable conservation condition of Hard oligo-mesotrophic waters with benthic vegetation of *Chara* spp. in Lough Corrib SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	The hard water lake habitat (3140) is found in Lough Corrib, notably the southern basin. Its exact distribution and area has not been mapped however, and it is likely to also extend along the eastern side of the northern basin. For additional information see Heuff (1984), Mooney and O'Connell (1990), Krause and King (1994) and Roden and Murphy (in prep.). Further information on this and all other attributes is provided in the lake habitats supporting document for the purposes of site-specific conservation objectives and Article 17 reporting (O Connor, 2015)
Habitat distribution	Occurrence	No decline, subject to natural processes. See map 3 for indicative lake habitat distribution	The distribution of the hard water lake habitat (3140) in Lough Corrib is determined, in the main, by geology and water chemistry. The hydrological regime (water circulation) in the Lough Corrib basins influences water mixing and, therefore, water chemistry. As a result, lake habitat 3140 is considered to dominate the southern basin, but is also likely to be widespread in the northern basin
Typical species	Occurrence	Typical species present, in good condition, and demonstrating typical abundances and distribution	For lists of typical species (cyanobacteria, algae, higher plants and water beetles), see the Article 17 habitat assessment for lake habitat 3140 (NPWS, 2013) and O Connor (2015). Lough Corrib is, naturally, a 'typical' marl lake as described by Roden and Murphy (2013; in prep.). Monitoring in 2012 demonstrated that the hard water vegetation in Lough Corrib is in bad condition, with shallow euphotic depth, some dead/dying charophytes and absent and decaying krustenstein (Roden and Murphy, in prep.)
Vegetation composition: characteristic zonation	Occurrence	All characteristic zones should be present, correctly distributed and in good condition	The characteristic zonation of lake habitat 3140 has been described (Roden and Murphy, 2013; in prep.). Charophyte zonation was reasonably intact in Lough Corrib in 2012, with <i>Chara curta</i> , <i>C. rudis</i> , <i>C. virgata</i> and <i>C. denudata</i> present, however the zones had become compressed since surveyed in 2004 and <i>C. contraria</i> and <i>Nitella flexilis</i> zones were not recorded (Roden and Murphy, in prep.). The krustenstein zone was either absent or decaying (Roden and Murphy, in prep.)
Vegetation distribution: maximum depth	Metres	Restore maximum depth of vegetation, subject to natural processes	The maximum depth of vegetation is likely to be specific to the lake shoreline in question, but is typically expected to be deep in clear, hard water lakes. An indicative target of more than 6m has been developed for hard water lakes (3140) (see Roden and Murphy, 2013; in prep.). Extremely clear marl lakes can have charophyte vegetation to far greater depths, such as Lough Rea (charophytes to 10-11m), or Coolorta (>9m) (Roden and Murphy, in prep.). In 2004, the euphotic zone of Lough Corrib extended to 6m, but in 2012 the maximum depth of vegetation was 4.2m (Roden and Murphy, in prep.). Decaying charophytes were recorded at 3m (Roden and Murphy, in prep.)

Hydrological regime: water level fluctuations	Metres	Maintain appropriate natural hydrological regime necessary to support the habitat	The hydrological regime of lakes with habitat 3140 is driven by groundwater flows. Groundwater can discharge directly to the lake, via springs or seepages, or to in-flowing rivers. Fluctuations in lake water level are typical in Ireland, but can be amplified by activities such as abstraction and drainage. Increased water level fluctuations can increase wave action, up-root vegetation, increase turbidity, alter the substratum and lead to release of nutrients from the sediment. The hydrological regime, particularly the groundwater contribution, must be maintained so that the area, distribution and depth of the lake habitat and its constituent/characteristic vegetation zones and communities are not reduced
Lake substratum quality	Various	Restore appropriate substratum type, extent and chemistry to support the vegetation	The hard water lake habitat is associated with a range of base-rich substratum types, from marl and limestone bedrock, through rocks, cobbles, gravel, muds and even peat. Lake substratum quality in Lough Corrib is impaired as a result of peat deposition, likely also nutrient enrichment, krustenstein decay and colonisation by zebra mussels (<i>Dreissena polymorpha</i>) (Roden, 2012; Roden and Murphy, in prep.). Further research into substratum quality (notably calcium, iron and nutrient concentrations) in the hard water lake habitat would be beneficial
Water quality: transparency	Metres	Restore appropriate Secchi transparency. There should be no decline in Secchi depth/transparency	Transparency relates to light penetration and, hence, to the depth of colonisation of vegetation. It can be affected by phytoplankton blooms, water colour and turbidity. A target has been set for hard water lakes (3140) of >6m (Roden and Murphy, in prep.). The OECD fixed boundary system set transparency targets for oligotrophic lakes of ≥6m annual mean Secchi disk depth and ≥3m annual minimum Secchi disk depth. Hard water lakes typically have high transparency, particularly in the very clear and typical marl forms (Roden and Murphy, in prep.). Secchi depth was 4m in lake habitat 3140 in 2012 (Roden and Murphy, in prep.)
Water quality: nutrients	µg/l P; mg/l N	Restore the concentration of nutrients in the water column to sufficiently low levels to support the habitat and its typical species	Lake habitat 3140 is typically associated with high water quality, including low dissolved nutrients. Some forms of the habitat appear to be naturally more productive than others, e.g. the machair form may be naturally more nutrient-rich. The default target for typical marl lakes is Water Framework Directive (WFD) 'High Status' or oligotrophic (OECD, 1982). Annual average Total Phosphorus (TP) concentration should be ≤10µg/l TP, average annual total ammonia should be ≤0.040mg/l N and annual 95th percentile for total ammonia should be ≤0.090mg/l N. Where nutrient concentrations are lower than the targets, there should be no upward trend in nutrient concentrations. For further information see the European Communities Environmental Objectives (Surface Waters) Regulations 2009. TP has exceeded the 10µg/l target in Lower Corrib in a number of recent years (Clabby et al., 2008; McGarrigle et al., 2010)

Water quality: phytoplankton biomass	µg/l Chlorophyll <i>a</i>	Maintain appropriate water quality to support the habitat, including high chlorophyll <i>a</i> status	Lake habitat 3140 is associated with high water quality, as demonstrated by naturally low algal growth. As for nutrients, the default target is WFD 'High Status' or oligotrophic (OECD, 1982). Average growing season (March-October) chlorophyll <i>a</i> concentration must be <5.8µg/l. Annual average chlorophyll <i>a</i> concentration should be <2.5µg/l and the annual peak should be <8.0µg/l. Where chlorophyll <i>a</i> concentrations are lower than the targets, there should be no upward trend in phytoplankton biomass. See the European Communities Environmental Objectives (Surface Waters) Regulations 2009. Maximum chlorophyll <i>a</i> in Lower Corrib exceeded the OECD target of ≤8.0 µg/l in five of the six monitoring cycles from 1976-2000 (McGarrigle et al., 2010). In recent years, however maximum concentrations have dropped to c.8µg/l, probably as a result of zebra mussel (<i>Dreissena polymorpha</i>) activity
Water quality: phytoplankton composition	EPA phytoplankton composition metric	Maintain appropriate water quality to support the habitat, including high phytoplankton composition status	The Environmental Protection Agency (EPA) has developed a phytoplankton composition metric for nutrient enrichment of Irish lakes. As for other water quality indicators, the default target for lake habitat 3140 is WFD high status
Water quality: attached algal biomass	Algal cover and EPA phytobenthos metric	Restore/maintain trace/absent attached algal biomass (<5% cover) and high phytobenthos status	Nutrient enrichment can favour epiphytic and epipelic algae that can out-compete the submerged vegetation. The cover abundance of attached algae in hard water lakes (3140) should, therefore, be trace/absent (<5% cover). EPA phytobenthos can be used as an indicator of changes in attached algal biomass. As for other water quality indicators, the default target for lake habitat 3140 is high phytobenthos status
Water quality: macrophyte status	EPA macrophyte metric (The Free Index)	Restore high macrophyte status	Nutrient enrichment can favour more competitive submerged macrophyte species that out-compete the typical and characteristic species for hard water lakes (3140). The EPA monitors macrophyte status for WFD purposes using the 'Free Index'. The target for lake habitat 3140 is high status or an Ecological Quality Ratio (EQR) for lake macrophytes of ≥0.90, as defined in Schedule Five of the European Communities Environmental Objectives (Surface Water) Regulations 2009
Acidification status	pH units; mg/l	Maintain appropriate water and sediment pH, alkalinity and cation concentrations to support the habitat, subject to natural processes	The specific requirements of lake habitat 3140, in terms of water and sediment pH, alkalinity and cation concentration, have not been fully determined. Acidification is not considered a threat to habitat 3140, however eutrophication can lead to at least temporary increases in pH to toxic levels (>9/9.5 pH units). Maximum pH should be <9.0 pH units, in line with the surface water standards. See the European Communities Environmental Objectives (Surface Waters) Regulations 2009
Water colour	mg/l PtCo	Restore/maintain appropriate water colour to support the habitat	Increased colour decreases light penetration and reduces the area of macrophyte habitat, particularly at the lower euphotic depths. Higher colour also appears to favour angiosperms over charophytes in hard water lakes (Roden and Murphy, in prep.). The primary source of increased colour in Ireland is peatland disturbance. No habitat-specific or national standards for water colour exist. Studies have shown median colour concentrations in Irish lakes of 38mg/l PtCo (Free et al., 2000) and 33mg/l PtCo (Free et al., 2006). Lake habitat 3140 is typically associated with very clear waters and expected colour would be <10 or, more likely, <5mg/l PtCo. Higher colour is found in some hard water lakes with significant areas of peatland in their catchment, but it is not clear whether this is natural or the result of peatland degradation. Roden and Murphy (in prep.) recorded dark water in Lough Corrib

Dissolved organic carbon (DOC)	mg/l	Restore/maintain appropriate organic carbon levels to support the habitat	Dissolved (and particulate) organic carbon (OC) in the water column is linked to water colour and acidification (organic acids). Increasing DOC in water has been documented across the Northern Hemisphere, including afforested peatland catchments in Ireland. Damage and degradation of peatland, leading to decomposition of peat is likely to be the predominant source of OC in Ireland. OC in water promotes decomposition by fungi and bacteria that, in turn, releases dissolved nutrients. The increased biomass of decomposers can also impact directly on the characteristic lake communities through shading, competition, etc.
Turbidity	Nephelometric turbidity units/ mg/l SS/ other appropriate unit	Restore/maintain appropriate turbidity to support the habitat	Turbidity can significantly affect the quantity and quality of light reaching rooted and attached vegetation and can, therefore, impact on lake habitats. The settlement of higher loads of inorganic or organic material on lake vegetation communities may also have impacts on sensitive, delicate species. Turbidity can increase as a result of re-suspension of material within the lake, higher loads entering the lake, or eutrophication. Turbidity measurement and interpretation is challenging. As a result, it is likely to be difficult to set habitat-specific targets for turbidity in lakes
Fringing habitat: area and condition	Hectares	Maintain the area and condition of fringing habitats necessary to support the natural structure and functioning of habitat 3140	Most lake shorelines have fringing habitats of reedswamp, other swamp, fen, marsh or wet woodland that intergrade with and support the structure and functions of the lake habitat. Equally, fringing habitats are dependent on the lake, particularly its water levels, and support wetland communities and species of conservation concern. Many of the fringing wetland habitats support higher invertebrate and plant species-richness than the lake habitats themselves. Fringing fen habitats can be particularly important around hard water lakes, notably the Annex I habitats alkaline fen, <i>Cladium</i> fen and petrifying springs (habitat codes 7230, 7210 and 7220). Lough Corrib has important areas of fen, limestone pavement, marsh, reedswamp and some woodland along its shoreline. See also the conservation objectives for associated Annex I habitats and Annex II species within the SAC

Conservation Objectives for : Lough Corrib SAC [000297]

3260 Water courses of plain to montane levels with the *Ranunculus fluitantis* and *Callitriche-Batrachion* vegetation

To maintain the favourable conservation condition of Water courses of plain to montane levels with the *Ranunculus fluitantis* and *Callitriche-Batrachion* vegetation in Lough Corrib SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Kilometres	Area stable or increasing, subject to natural processes	The description of 3260 is broad, from upland bryophyte/macroalgal dominated stretches, to lowland depositing rivers with pondweeds and starworts. Selection of SACs for the habitat used this broad interpretation. Site-specific objectives for the habitat concentrate upon high conservation value sub-types. Little is known, however, about the characteristics or sub-types in Lough Corrib SAC. Many of the rivers were included in the SAC for Atlantic salmon (<i>Salmo salar</i>). Most of the rivers are in arterial drainage schemes that have altered aquatic plant distribution and species composition. Note: rooted macrophytes should be absent or trace (<5% cover) in freshwater pearl mussel (<i>Margaritifera margaritifera</i>) habitat. The freshwater pearl mussel (1029) conservation objective takes precedence over this objective for habitat 3260 in the Owenriff River within this SAC, because the mussel requires environmental conditions closer to natural background levels
Habitat distribution	Occurrence	No decline, subject to natural processes	As noted above, little is known about the distribution of the habitat and its sub-types in this SAC. The vegetation of the River Corrib was documented in Mooney and O'Connell (1990). Macrophyte vegetation has expanded in the Owenriff River resulting in a decline in the condition of this priority freshwater pearl mussel (<i>Margaritifera margaritifera</i>) population and its habitat (NPWS, 2010). The Cornamonna, Owennaraha, Owenakilla and other rivers flowing into the north-western part of Lough Corrib are worthy of further investigation
Hydrological regime: river flow	Metres per second	Maintain appropriate hydrological regimes	Any high conservation value sub-types in the SAC will be associated with natural, fast and highly variable flows. Owing to regular disturbance (through variations in flow), river macrophytes rarely reach a climax condition but frequently occur as transient communities. A natural flow regime is required for both plant communities and channel geomorphology to be in favourable condition, exhibiting typical dynamics for the river type (Hatton-Ellis and Grieve, 2003). For many of the sub-types of this habitat, high flows are required to maintain the substratum necessary for the characteristic species. Flow variation can be particularly important, with high and flood flows being critical to the hydromorphology
Hydrological regime: groundwater discharge	Metres per second	Maintain appropriate hydrological regimes	It is likely that rivers over limestone bedrock have a significant groundwater contribution. There may be tufa formation associated with such groundwater springs and seepages. Such petrifying springs are an in-stream form of the Habitats Directive priority Annex I habitat "Petrifying springs with tufa formation (Cratoneurion)" (7220) and of high conservation value

Substratum composition: particle size range	Millimetres	Maintain appropriate substratum particle size range, quantity and quality, subject to natural process	Although many of the high conservation value sub-types are dominated by coarse substrata, for certain sub-types, notably tidal forms, fine substrata are required. The size and distribution of particles is largely determined by the river flow. The chemical composition (particularly minerals and nutrients) of the substratum is also important. The quality of finer sediment particles is a notable driver for rooted plant communities
Water quality	Various	Maintain appropriate water quality to support the natural structure and functioning of the habitat	The specific targets may vary among sub-types. Bryophyte-rich and tufaceous streams and rivers are considered highly sensitive to nutrient enrichment and are likely to require high status waters. Water quality for other sub-types should reach a minimum of Water Framework Directive (WFD) good status, in terms of nutrient and oxygenation standards, and EQRs (Ecological Quality Ratios) for macroinvertebrates and phytobenthos
Vegetation composition: typical species	Occurrence	Typical species of the relevant habitat sub-type should be present and in good condition	The sub-types of this habitat are poorly understood and their typical species have not yet been fully defined. The typical species may include higher plants, bryophytes, macroalgae and microalgae, and invertebrates
Floodplain connectivity: area	Hectares	The area of active floodplain at and upstream of the habitat should be maintained	River connectivity with the floodplain is important for the functioning of this habitat. Channels with a naturally functioning floodplain are better able to maintain habitat and water quality (Hatton-Ellis and Grieve, 2003). Floodplain connectivity is particularly important in terms of sediment sorting and nutrient deposition. High conservation value rivers are intimately connected to floodplain habitats and function as important wildlife corridors, connecting otherwise isolated or fragmented habitats in the wider countryside (Hatton-Ellis and Grieve, 2003)
Riparian habitat: area	Hectares	Maintain the area and condition of fringing habitats necessary to support the habitat and its sub-types	Riparian habitats, including those along lake fringes, particularly natural/semi-natural woodlands and wetlands, even where they do not form part of a natural floodplain, are an integral part of the structure and functioning of river systems. Fringing habitats can contribute to the aquatic food web (e.g. allochthonous matter such as leaf fall), provide habitat (refuge and resources) for certain life-stages of fish, birds and aquatic invertebrates, assist in the settlement of fine suspended material, protect banks from erosion and contribute to nutrient cycling. Shade may also be important in suppressing algal growth in enriched rivers and moderating temperatures. Equally, fringing habitats are dependent on rivers/lakes, particularly their water levels, and support wetland communities and species of conservation concern. See also the conservation objectives for associated Annex I habitats and Annex II species within the SAC

Conservation Objectives for : Lough Corrib SAC [000297]

6210 Semi-natural dry grasslands and scrubland facies on calcareous substrates (Festuco-Brometalia) (* important orchid sites)

To maintain the favourable conservation condition of Semi-natural dry grasslands and scrubland facies on calcareous substrates (Festuco-Brometalia) (* important orchid sites) in Lough Corrib SAC in owing list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Semi-natural dry grasslands and scrubland facies on calcareous substrates (Festuco-Brometalia) occurs mainly as small areas and in intimate association with other habitats in this SAC including other grassland types, fens and limestone pavements and is therefore difficult to map separately. O'Neill et al. (2013) surveyed and mapped some grassland sites within the SAC in detail and the surveys of limestone pavement sites carried out by Wilson and Fernandez (2013) included associated grassland habitats; however, as all areas of this habitat within the SAC have not been identified, the total area is unknown
Habitat distribution	Occurrence	No decline, subject to natural processes	See notes for area above
Vegetation composition: typical species	Number at a representative number of monitoring stops	At least seven positive indicator species present, including two "high quality" species	Attribute and target based on O'Neill et al. (2013), where the list of positive indicator species, including high quality species, as identified by the Irish Semi-natural Grasslands Survey (ISGS) is presented. O'Neill et al. (2013) should be consulted for further details
Vegetation composition: negative indicator species	Percentage at a representative number of monitoring stops	Negative indicator species collectively not more than 20% cover, with cover by an individual species not more than 10%	Attribute and target based on O'Neill et al. (2013), where the list of negative indicator species as identified by the ISGS is presented
Vegetation composition: non-native species	Percentage at a representative number of monitoring stops	Cover of non-native species not more than 1%	Attribute and target based on O'Neill et al. (2013)
Vegetation composition: woody species and bracken	Percentage at a representative number of monitoring stops	Cover of woody species (except certain listed species) and bracken (<i>Pteridium aquilinum</i>) not more than 5% cover	Attribute and target based on O'Neill et al. (2013). Woody species that can occur above 5% cover are juniper (<i>Juniperus communis</i>), burnet rose (<i>Rosa spinosissima</i>), mountain avens (<i>Dryas octopetala</i>) and hoary rock-rose (<i>Helianthemum oelandicum</i>)
Vegetation structure: broadleaf herb: grass ratio	Percentage at a representative number of monitoring stops	Broadleaf herb component of vegetation between 40% and 90%	Attribute and target based on O'Neill et al. (2013)
Vegetation structure: sward height	Percentage at a representative number of monitoring stops	At least 30% of sward between 5cm and 40cm tall	Attribute and target based on O'Neill et al. (2013)
Vegetation structure: litter	Percentage at a representative number of monitoring stops	Litter cover not more than 25%	Attribute and target based on O'Neill et al. (2013)
Physical structure: bare soil	Percentage at a representative number of monitoring stops	Not more than 10% bare soil	Attribute and target based on O'Neill et al. (2013)
Physical structure: disturbance	Square metres	Area showing signs of serious grazing or other disturbance less than 20m ²	Attribute and target based on O'Neill et al. (2013)

Conservation Objectives for : Lough Corrib SAC [000297]

6410 *Molinia* meadows on calcareous, peaty or clayey-silt-laden soils (*Molinion caeruleae*)

To maintain the favourable conservation condition of *Molinia* meadows on calcareous, peaty or clayey-silt-laden soils (*Molinion caeruleae*) in Lough Corrib SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	<i>Molinia</i> meadows on calcareous, peaty or clayey-silt-laden soils (<i>Molinion caeruleae</i>) occurs mainly as small areas and in intimate association with other habitats in this SAC such as other grassland types and fens and is therefore difficult to map separately. O'Neill et al. (2013) surveyed and mapped some grassland sites within Lough Corrib SAC. However, the full extent of this habitat in this SAC is currently unknown
Habitat distribution	Occurrence	No decline, subject to natural processes	See notes for area above
Vegetation composition: typical species	Number at a representative number of monitoring stops	At least seven positive indicator species present, including one "high quality" species as listed in O'Neill et al. (2013)	Attribute and target based on O'Neill et al. (2013), where the list of positive indicator species, including high quality species, as identified by the Irish Semi-natural Grasslands Survey (ISGS) is presented. O'Neill et al. (2013) should be consulted for further details
Vegetation composition: negative indicator species	Percentage at a representative number of monitoring stops	Negative indicator species collectively not more than 20% cover, with cover by an individual species not more than 10%	Attribute and target based on O'Neill et al. (2013), where the list of negative indicator species as identified by the ISGS is presented
Vegetation composition: non-native species	Percentage at a representative number of monitoring stops	Cover of non-native species not more than 1%	Attribute and target based on O'Neill et al. (2013)
Vegetation composition: moss species	Percentage at a representative number of monitoring stops	Hair mosses (<i>Polytrichum</i> spp.) not more than 25% cover	Attribute and target based on O'Neill et al. (2013)
Vegetation structure: woody species and bracken	Percentage at a representative number of monitoring stops	Cover of woody species and bracken (<i>Pteridium aquilinum</i>) not more than 5%	Attribute and target based on O'Neill et al. (2013)
Vegetation structure: broadleaf herb: grass ratio	Percentage at a representative number of monitoring stops	Broadleaf herb component of vegetation between 40% and 90%	Attribute and target based on O'Neill et al. (2013)
Vegetation structure: sward height	Percentage at a representative number of monitoring stops	At least 30% of sward between 10cm and 80cm tall	Attribute and target based on O'Neill et al. (2013)
Vegetation structure: litter	Percentage at a representative number of monitoring stops	Litter cover not more than 25%	Attribute and target based on O'Neill et al. (2013)
Physical structure: bare soil	Percentage at a representative number of monitoring stops	Not more than 10% bare soil	Attribute and target based on O'Neill et al. (2013)
Physical structure: disturbance	Square metres	Area showing signs of serious grazing or other disturbance less than 20m ²	Attribute and target based on O'Neill et al. (2013)

Conservation Objectives for : Lough Corrib SAC [000297]

7110 Active raised bogs

To restore the favourable conservation condition of Active raised bogs* in Lough Corrib SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Restore the area of active raised bog to 78.8ha, subject to natural processes	There are two raised bogs for which Active Raised Bog (ARB) has been selected in Lough Corrib SAC: Addergoole Bog and Lough Tee Bog. The total area of ARB habitat for these two bogs was mapped as 45.2ha. The area of Degraded Raised Bog (DRB) on the High Bog (HB) has been modelled as 72.0ha. See map 4. However, it is estimated that only 24.1ha is potentially restorable to ARB by drain blocking. The total potential ARB on the HB is therefore estimated to be 69.3ha. Eco-hydrological assessments of cutover bog estimate that an additional 9.5ha of peat-forming habitats could be restored. The long-term target for ARB is therefore 78.8ha. See the Lough Corrib SAC conservation objectives supporting document for raised bog habitats for further details on this and the following attributes
Habitat distribution	Occurrence	Restore the distribution and variability of active raised bog across the SAC. See map 5 for distribution in 2012	ARB currently occurs on the north-eastern and southern parts of Addergoole Bog and the north-western part of Lough Tee Bog. DRB occurs over much of the remainder of the high bog at Addergoole and also occurs at Lough Tee Bog. Such areas will require restoration measures. See also the conservation objective for Bog woodland (91D0)
High bog area	Hectares	No decline in extent of high bog subject to the conservation requirements of the SAC. See map 4 for mapped extent	The area of HB within Addergoole Bog in 2012 (latest figure available) is 157.4ha and 79.6ha at Lough Tee Bog (DAHG, 2014)
Hydrological regime: water levels	Centimetres	Restore appropriate water levels throughout each site	For ARB, mean water level needs to be near or above the surface of the bog lawns for most of the year. Seasonal fluctuations should not exceed 20cm and water levels should not be more than 10cm below the bog surface, except for very short periods of time. Open water is often characteristic of soak systems
Hydrological regime: flow patterns	Flow direction; slope	Restore, where possible, appropriate high bog topography, flow directions and slopes. See map 6 for current situation	ARB depends on mean water levels being near or above the surface of bog lawns for most of the year. Long and gentle slopes are the most favourable to achieve these conditions. Changes to flow directions due to subsidence of bogs can radically change water regimes and cause drying out of high quality ARB areas and soak systems
Transitional areas between high bog and adjacent mineral soils (including cutover areas)	Hectares; distribution	Restore adequate transitional areas to support/protect the raised bog ecosystem and the services it provides	Cutting has continued until recently around the eastern and southern margins of Addergoole Bog. Remnant semi-natural margins occur elsewhere (locally along the north-east and east margins of the site) and there are interesting transitional wetland habitats to the east of the high bog. Transitional areas between Lough Tee Bog and the adjacent mineral soils mostly corresponds with cutover bog at different stages of re-colonisation by vegetation. Eco-hydrological assessments are evaluating the potential for ARB on cutover areas (see note for habitat area attribute above)
Vegetation quality: central ecotope, active flush, soaks, bog woodland	Hectares	Restore 39.4ha of central ecotope/active flush/soaks/bog woodland as appropriate	At least 50% of ARB habitat should be high quality (i.e. central ecotope, active flush, soaks, bog woodland). The target area of active raised bog for the site has been set at 78.8ha (see habitat area target above)

Vegetation quality: microtopographical features	Hectares	Restore adequate cover of high quality microtopographical features	High quality microtopography (hummocks, hollows and pools) occurs at both Addergoole and Lough Tee bogs; however, both have been negatively affected by burning
Vegetation quality: bog moss (<i>Sphagnum</i>) species	Percentage cover	Restore adequate cover of bog moss (<i>Sphagnum</i>) species to ensure peat-forming capacity	<i>Sphagnum</i> cover varies naturally across Ireland with relatively high cover in the east to lower cover in the west. Hummock forming species such as <i>Sphagnum austinii</i> are particularly good peat formers
Typical ARB species: flora	Occurrence	Restore, where appropriate, typical active raised bog flora	Typical flora species include widespread species, as well as those with more restricted distributions, but typical of the habitat's subtypes or geographical range
Typical ARB species: fauna	Occurrence	Restore, where appropriate, typical active raised bog fauna	Typical fauna species include widespread species, as well as those with more restricted distributions, but typical of the habitat's subtypes or geographical range
Elements of local distinctiveness	Occurrence	Maintain features of local distinctiveness, subject to natural processes	Addergoole Bog is noted for the presence of a soak system, a rare feature of Irish raised bogs, which occurs in the centre of the bog near two lakes. This soak includes an area of Bog woodland (91D0) which is a priority Annex I habitat and a qualifying interest for the SAC (see also the conservation objective for 91D0)
Negative physical indicators	Percentage cover	Negative physical features absent or insignificant	Negative physical indicators include: bare peat, algae dominated pools and hollows, marginal cracks, tear patterns, subsidence features such as dry mineral mounds/ridges emerging or expanding, and evidence of burning
Vegetation composition: native negative indicator species	Percentage cover	Native negative indicator species at insignificant levels	Disturbance indicators include species indicative of conditions drying out such as abundant bog asphodel (<i>Nartheicum ossifragum</i>), deergrass (<i>Trichophorum germanicum</i>) and hare's-tail cotton-grass (<i>Eriophorum vaginatum</i>) forming tussocks; abundant magellanic bog-moss (<i>Sphagnum magellanicum</i>) in pools previously dominated by <i>Sphagnum</i> species typical of very wet conditions (e.g. feathery bog-moss (<i>S. cuspidatum</i>)); and indicators of frequent burning events such as abundant <i>Cladonia floerkeana</i> and high cover of carnation sedge (<i>Carex panicea</i>) (particularly in true midlands raised bogs)
Vegetation composition: non-native invasive species	Percentage cover	Non-native invasive species at insignificant levels and not more than 1% cover	The most common non-native invasive species include lodgepole pine (<i>Pinus contorta</i>), rhododendron (<i>Rhododendron ponticum</i>) and pitcherplant (<i>Sarracenia purpurea</i>)
Air quality: nitrogen deposition	kg N/ha/yr	Air quality surrounding the bogs close to natural reference conditions. The total nitrogen deposition should not exceed 5kg N/ha/yr	Change in air quality can result from fertiliser drift; adjacent quarry activities; or other atmospheric inputs. The critical load range for ombrotrophic bogs has been set as between 5 and 10kg N/ha/yr (Bobbink and Hettelingh, 2011). The latest nitrogen deposition figures for the area around the bogs in Lough Corrib SAC suggests that the current level is approximately 8.5kg N/ha/yr (Henry and Aherne, 2014)
Water quality	Hydrochemical measures	Water quality on the high bog and in transitional areas close to natural reference conditions	Water chemistry within raised bogs is influenced by atmospheric inputs (rainwater). However, within soak systems, water chemistry is influenced by other inputs such as focused flow or interaction with underlying substrates. Water chemistry in marginal areas surrounding the high bog varies due to influences of different water types (bog water, regional groundwater and run-off from surrounding mineral lands)

Conservation Objectives for : Lough Corrib SAC [000297]

7120 Degraded raised bogs still capable of natural regeneration

The long-term aim for Degraded raised bogs still capable of natural regeneration is that its peat-forming capability is re-established; therefore, the conservation objective for this habitat is inherently linked to that of Active raised bogs (7110) and a separate conservation objective has not been set in Lough Corrib SAC

Attribute	Measure	Target	Notes
-----------	---------	--------	-------

Conservation Objectives for : Lough Corrib SAC [000297]

7150 Depressions on peat substrates of the Rhynchosporion

Depressions on peat substrates of the Rhynchosporion is an integral part of good quality Active raised bogs (7110) and thus a separate conservation objective has not been set for the habitat in Lough Corrib SAC

Attribute	Measure	Target	Notes
------------------	----------------	---------------	--------------

Conservation Objectives for : Lough Corrib SAC [000297]

7210 Calcareous fens with *Cladium mariscus* and species of the Caricion davalliana

To maintain the favourable conservation condition of Calcareous fens with *Cladium mariscus* and species of the Caricion davalliana in Lough Corrib SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Calcareous fens with <i>Cladium mariscus</i> and species of the Caricion davalliana have not been mapped in detail for Lough Corrib SAC and thus total area of the qualifying habitat is unknown. While the full extent of Annex I fen habitats (both this habitat and Alkaline fens (7230)) within the SAC is currently unknown, their area is extensive and they often occur in association with and transitional to other habitats including <i>Molinia</i> meadows on calcareous, peaty or clayey-silt-laden soils (Molinion caeruleae) (6410), Active raised bogs (7110), Petrifying springs with tufa formation (Cratoneurion) (7220) and Limestone pavements (8240) (NPWS internal files). The conservation objectives for all these habitats in the SAC should be used in conjunction with each other as appropriate
Habitat distribution	Occurrence	No decline, subject to natural processes	The full distribution of this habitat in this SAC is currently unknown - see notes for area above
Ecosystem function: hydrology	Metres	Maintain appropriate natural hydrological regimes necessary to support the natural structure and functioning of the habitat	Maintenance of groundwater, surface water flows and water table levels within natural ranges is essential for this wetland habitat
Ecosystem function: peat formation	Flood duration	Maintain active peat formation, where appropriate	In order for peat to form, water levels need to be slightly below or above the soil surface for c.90% of the time (Jim Ryan, pers. comm.)
Ecosystem function: water quality	Water chemistry measures	Maintain appropriate water quality, particularly nutrient levels, to support the natural structure and functioning of the habitat	Fens receive natural levels of nutrients (e.g. iron, magnesium and calcium) from water sources. However, they are generally poor in nitrogen and phosphorus with the latter tending to be the limiting nutrient
Vegetation structure: typical species	Presence	Maintain vegetation cover of typical species including brown mosses and vascular plants	Mosses listed for fen habitat in this SAC includes <i>Campylopus stellatum</i> , while vascular plants include saw sedge (<i>Cladium mariscus</i>), slender sedge (<i>C. lasiocarpa</i>), long-stalked yellow sedge (<i>C. lepidocarpa</i>), black bog-rush (<i>Schoenus nigricans</i>), water mint (<i>Mentha aquatica</i>), wild angelica (<i>Angelica sylvestris</i>) and meadow thistle (<i>Cirsium dissectum</i>). Slender cottongrass (<i>Eriophorum gracile</i>) is known from fen habitat in this SAC (NPWS internal files)
Vegetation composition: non-native species	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of non-native species less than 1%	Attribute and target based on Perrin et al. (2014). Non-native species can be invasive and have deleterious effects on native vegetation. A low target is set as non-native species can spread rapidly and are most easily dealt with when still at lower abundances
Vegetation composition: trees and shrubs	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of scattered native trees and shrubs less than 10%	Attribute and target based on Perrin et al. (2014). Scrub and trees will tend to invade if fen conditions become drier
Physical structure: disturbed bare ground	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of disturbed bare ground not more than 10%. Where tufa is present, disturbed bare ground not more than 1%	Attribute and target based on Perrin et al. (2014). While grazing may be appropriate in this habitat, excessive areas of disturbed bare ground may develop due to unsuitable grazing regimes

Physical structure: drainage	Percentage area in local vicinity of a representative number of monitoring stops	Areas showing signs of drainage as a result of drainage ditches or heavy trampling not more than 10%	Attribute and target based Perrin et al. (2014). Drainage can result in loss of characteristic species and transition to drier habitats
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat	This includes species on the Flora (Protection) Order, 2015 and/or the red data lists (Lockhart et al., 2012; Wyse Jackson et al., 2016)

Conservation Objectives for : Lough Corrib SAC [000297]

7220 Petrifying springs with tufa formation (Cratoneurion)

To maintain the favourable conservation condition of Petrifying springs with tufa formation (Cratoneurion)* in Lough Corrib SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Square metres	Area stable or increasing, subject to natural processes	Petrifying springs with tufa formation (Cratoneurion) have not been mapped within Lough Corrib SAC and thus the total area of the qualifying habitat in the SAC is unknown. However, the necessary ecological conditions required for this habitat occur around Lough Corrib
Habitat distribution	Occurrence	No decline, subject to natural processes	As mentioned above, this habitat has not been mapped within the SAC. It is often associated with other habitats including Calcareous fens with <i>Cladium mariscus</i> and species of the Caricion davallianae (7210), Alkaline fens (7230) and Limestone pavements (8240). The conservation objectives for all these habitats in the SAC should be used in conjunction with each other as appropriate. Lyons and Kelly (2016) describe eight plant communities of Irish petrifying springs based on relevé data. Further information on the vegetation communities associated with this habitat is presented in Lyons and Kelly (2016)
Hydrological regime: height of water table; water flow	Metres; metres per second	Maintain appropriate hydrological regimes	Petrifying springs rely on permanent irrigation, usually from upwelling groundwater sources or seepage sources (Lyons and Kelly, 2013). In karst areas, water tends to flow away rapidly over bare rock surfaces, even on fairly flat ground (Lyons and Kelly, 2013). Water flow should not be altered anthropogenically. See Lyons and Kelly (2016) for further details
Water quality - nitrate level	mg/l	No increase from baseline nitrate level and less than 10mg/l	Target based on data from McGarrigle et al. (2010). See Lyons and Kelly (2016) for further details
Water quality - phosphate level	µg/l	No increase from baseline phosphate level and less than 15µg/l	Based on data from Lyons (2015). See Lyons and Kelly (2016) for further details
Vegetation composition: positive indicator species	Number per spring	At least three positive/high quality indicator species as listed in Lyons and Kelly (2016) and no loss from baseline number	Based on Lyons and Kelly (2016), where the lists of positive and high quality indicator species are presented
Vegetation composition: negative indicator species	Cover (DAFOR scale)	Potentially negative indicator species should not be Dominant or Abundant; invasive species should be absent	Based on Lyons and Kelly (2016), where the lists of potentially negative herbaceous, bryophyte (and alga) and woody species are presented. See Lyons and Kelly (2016) also for details on potentially invasive species, including sycamore (<i>Acer pseudoplatanus</i>) which is invasive in non-wooded springs and a negative indicator species in wooded springs. If two or more potentially negative bryophyte species are present, and if at least two are Frequent, or at least one is Abundant, then the habitat fails for this attribute. See Lyons and Kelly (2016) for further details
Vegetation structure: sward height	Centimetres	Field layer height between 10cm and 50cm (except for bryophyte-dominated ground <10cm)	See Lyons and Kelly (2016) for further details
Physical structure: trampling/dung	Cover (DAFOR scale)	Cover should not be Dominant or Abundant	See Lyons and Kelly (2016) for further details

Conservation Objectives for : Lough Corrib SAC [000297]

7230

Alkaline fens

To maintain the favourable conservation condition of Alkaline fens in Lough Corrib SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Alkaline fens have not been mapped in detail for Lough Corrib SAC and thus total area of the qualifying habitat is unknown. While the full extent of Annex I fen habitats (both this habitat and Calcareous fens with <i>Cladium mariscus</i> and species of the Caricion davallianae (7210)) within the SAC is currently unknown, their area is extensive and they often occur in association with and transitional to other habitats including <i>Molinia</i> meadows on calcareous, peaty or clayey-silt-laden soils (Molinion caeruleae) (6410), Active raised bogs (7110), Petrifying springs with tufa formation (Cratoneurion) (7220) and Limestone pavements (8240) (NPWS internal files). The conservation objectives for all these habitats in the SAC should be used in conjunction with each other as appropriate
Habitat distribution	Occurrence	No decline, subject to natural processes	The full distribution of this habitat in this SAC is currently unknown - see notes for area above
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil nutrient status within natural range	Relevant nutrients and their natural ranges are yet to be defined. However, nitrogen deposition is noted as being relevant to this habitat in NPWS (2013)
Ecosystem function: peat formation	Flood duration	Maintain active peat formation, where appropriate	In order for peat to form, water levels need to be slightly below or above the soil surface for c.90% of the time (Jim Ryan, pers. comm.)
Ecosystem function: hydrology	Metres	Maintain appropriate natural hydrological regimes necessary to support the natural structure and functioning of the habitat	Maintenance of groundwater, surface water flows and water table levels within natural ranges is essential for this wetland habitat
Ecosystem function: water quality	Water chemistry measures	Maintain appropriate water quality, particularly nutrient levels, to support the natural structure and functioning of the habitat	Fens receive natural levels of nutrients (e.g. iron, magnesium and calcium) from water sources. However, they are generally poor in nitrogen and phosphorus, with the latter tending to be the limiting nutrient
Community diversity	Abundance of variety of vegetation communities	Maintain variety of vegetation communities, subject to natural processes	Further information on the vegetation communities associated with alkaline fens in the uplands is presented in Perrin et al. (2014)
Vegetation composition: number of positive indicator species (brown mosses)	Number of species at a representative number of 2m x 2m monitoring stops	Number of brown moss species present at each monitoring stop is at least one	Attribute and target based on Perrin et al. (2014), where the list of positive indicator species for this habitat is also presented. Mosses listed for fen in this SAC includes <i>Campyllum stellatum</i> (NPWS internal files)
Vegetation composition: number of positive indicator species (vascular plants)	Number of species at a representative number of 2m x 2m monitoring stops	Number of positive vascular plant indicator species present at each monitoring stop is at least two for small-sedge flushes and at least three for black bog-rush (<i>Schoenus nigricans</i>) flush and bottle sedge (<i>Carex rostrata</i>) fen	Attribute and target based on Perrin et al. (2014), where the list of positive indicator species for this habitat is also presented. Vascular plants listed for fen in this SAC include saw sedge (<i>Cladium mariscus</i>), slender sedge (<i>C. lasiocarpa</i>), long-stalked yellow sedge (<i>C. lepidocarpa</i>), black bog-rush (<i>Schoenus nigricans</i>), water mint (<i>Mentha aquatica</i>), wild angelica (<i>Angelica sylvestris</i>) and meadow thistle (<i>Cirsium dissectum</i>). Slender cottongrass (<i>Eriophorum gracile</i>) is known from fen habitat in this SAC (NPWS internal files)

Vegetation composition: cover of positive indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of brown moss species and positive vascular plant indicator species at least 20% for small-sedge flushes and at least 75% cover for black bog-rush (<i>Schoenus nigricans</i>) flush and bottle sedge (<i>Carex rostrata</i>) fen	Attribute and target based on Perrin et al. (2014), where the list of positive indicator species for this habitat is also presented
Vegetation composition: negative indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of negative indicator species less than 1%	Attribute and target based on Perrin et al. (2014), where the list of negative indicator species for this habitat is also presented
Vegetation composition: non-native species	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of non-native species less than 1%	Attribute and target Perrin et al. (2014). Non-native species can be invasive and have deleterious effects on native vegetation. A low target is set as non-native species can spread rapidly and are most easily dealt with when still at lower abundances
Vegetation composition: native trees and shrubs	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of scattered native trees and shrubs less than 10%	Attribute and target based Perrin et al. (2014). Scrub and trees will tend to invade if fen conditions become drier
Vegetation composition: soft rush and common reed cover	Percentage cover in local vicinity of a representative number of monitoring stops	Total cover of soft rush (<i>Juncus effusus</i>) and common reed (<i>Phragmites australis</i>) less than 10%	Attribute and target based on Perrin et al. (2014)
Vegetation structure: height	Percentage of leaves/shoots at a representative number of 2m x 2m monitoring stops	Proportion of live leaves and/or flowering shoots of vascular plants that are more than 5cm above the ground surface should be at least 50%	Attribute and target based on Perrin et al. (2014). Vegetation heights lower than these would indicate undesirable levels of grazing
Physical structure: disturbed bare ground	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of disturbed bare ground less than 10%	Attribute and target based on Perrin et al. (2014). While grazing may be appropriate in this habitat, excessive areas of disturbed bare ground may develop due to unsuitable grazing regimes
Physical structure: drainage	Percentage area in local vicinity of a representative number of monitoring stops	Area showing signs of drainage as a result of drainage ditches or heavy trampling less than 10%	Attribute and target based Perrin et al. (2014). Drainage can result in loss of characteristic species and transition to drier habitats
Physical structure: tufa formations	Percentage cover in local vicinity of a representative number of 2m x 2m monitoring stops	Disturbed proportion of vegetation cover where tufa is present is less than 1%	Attribute and target based on Perrin et al. (2014)
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat	This includes species on the Flora (Protection) Order, 2015 and/or the red data lists (Lockhart et al., 2012; Wyse Jackson et al., 2016)

Conservation Objectives for : Lough Corrib SAC [000297]

8240 Limestone pavements

To maintain the favourable conservation condition of Limestone pavements* in Lough Corrib SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Limestone pavements often occur in intimate association with other habitats including the Annex I habitats Semi-natural dry grasslands and scrubland facies on calcareous substrates (Festuco-Brometalia) (6210), Petrifying springs with tufa formation (Cratoneurion) (7220) and Alkaline fens (7230). Therefore, these habitats cannot easily be mapped or considered separately. Conservation objectives for all these habitats should be used in conjunction with each other as appropriate. In Lough Corrib SAC, limestone pavements occur along the southern and eastern margins of Lough Corrib and more extensively in the southern part of the SAC. However, the total area of the Annex I habitat in the SAC is unknown. Wilson and Fernandez (2013) mapped the indicative area in the southern part, including mosaics with other habitats (map 7)
Distribution	Occurrence	No decline, subject to natural processes. Map 7 shows the indicative distribution in the southern part of the SAC, including mosaics with other habitats	See notes for area above. This habitat can be split into exposed pavement and wooded pavement (Wilson and Fernandez, 2013)
Vegetation composition: typical species	Number at a representative number of monitoring stops	At least seven positive indicator species present	Attribute and target based on Wilson and Fernandez (2013), where the positive indicator species for exposed and wooded pavement are listed
Vegetation composition: bryophyte layer	Percentage at a representative number of monitoring stops	Bryophyte cover at least 50% on wooded pavement	Attribute and target based on Wilson and Fernandez (2013)
Vegetation composition: negative indicator species	Percentage at a representative number of monitoring stops	Collective cover of negative indicator species on exposed pavement not more than 1%	Attribute and target based on Wilson and Fernandez (2013), where the negative indicator species are listed. Negative indicator species for wooded pavement overlap with non-native species (below)
Vegetation composition: non-native species	Percentage at a representative number of monitoring stops	Cover of non-native species not more than 1% on exposed pavement; on wooded pavement not more than 10% with no regeneration	Attribute and target based on Wilson and Fernandez (2013)
Vegetation composition: scrub	Percentage at a representative number of monitoring stops	Scrub cover no more than 25% of exposed pavement	Attribute and target based on Wilson and Fernandez (2013)
Vegetation composition: bracken cover	Percentage at a representative number of monitoring stops	Bracken (<i>Pteridium aquilinum</i>) cover no more than 10% on exposed pavement	Attribute and target based on Wilson and Fernandez (2013)
Vegetation structure: woodland canopy	Percentage at a representative number of monitoring stops	Canopy cover on wooded pavement at least 30%	Attribute and target based on Wilson and Fernandez (2013). Wooded limestone pavement is usually low-growing hazel (<i>Coryllus avellana</i>) woodland. Atlantic hazel woodland is an internationally rare woodland type. Despite its low stature it is nonetheless an important habitat for woodland species
Vegetation structure: dead wood	Occurrence in a representative number of monitoring stops	Sufficient quantity of dead wood on wooded pavement to provide habitat for saproxylic organisms	Dead wood is a valuable resource and an integral part of a healthy, functioning woodland ecosystem
Physical structure: disturbance	Occurrence in a representative number of monitoring stops	No evidence of grazing pressure on wooded pavement	Attribute and target based on Wilson and Fernandez (2013)

Indicators of local Occurrence
distinctiveness

Indicators of local
distinctiveness are
maintained

This includes species on the Flora (Protection) Order, 2015 and/or the red data lists (Lockhart et al., 2012; Wyse Jackson et al., 2016), and other rare or localised species, as well as archaeological and geological features, which often support distinctive species

Conservation Objectives for : Lough Corrib SAC [000297]

91A0 Old sessile oak woods with Ilex and Blechnum in the British Isles

To maintain the favourable conservation condition of Old sessile oak woods with *Ilex* and *Blechnum* in the British Isles in Lough Corrib SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Most of the woodland in Lough Corrib SAC occurs as narrow belts of low woodland around the lake, with occasional larger stands. Old sessile oakwoods are likely to occur as mosaics with other woodland types and the total extent within the SAC is unknown. As part of the National Survey of Native Woodlands (NSNW), Perrin et al. (2008) surveyed a number of sites in the SAC; one (Annaghwood; NSNW site no. 1624), contained this habitat (see map 8). It is important to note that further unsurveyed areas of old oak woodland are likely to be present within the SAC, including at the Hill of Doon and on some of the islands in the lake (NPWS internal files)
Habitat distribution	Occurrence	No decline. Surveyed location shown on map 8	Distribution shown based on Perrin et al. (2008). It is important to note that further unsurveyed areas are likely to be present within the SAC. See note on area above
Woodland size	Hectares	Area stable or increasing. Where topographically possible, "large"; woods at least 25ha in size and "small" woods at least 3ha in size.	The sizes of at least some of the existing woodlands need to be increased in order to reduce habitat fragmentation and benefit those species requiring 'deep' woodland conditions (Peterken, 2002). Topographical and land ownership constraints may restrict expansion
Woodland structure: cover and height	Percentage and metres	Diverse structure with a relatively closed canopy containing mature trees; subcanopy layer with semi-mature trees and shrubs; and well-developed herb layer	Described in Perrin et al (2008) and NPWS internal files
Woodland structure: community diversity and extent	Hectares	Maintain diversity and extent of community types	Based on data from Perrin et al. (2008)
Woodland structure: natural regeneration	Seedling: sapling: pole ratio	Seedlings, saplings and pole age-classes occur in adequate proportions to ensure survival of woodland canopy	Oak (<i>Quercus petraea</i>) regenerates poorly. In suitable sites, ash (<i>Fraxinus excelsior</i>) can regenerate in large numbers although few seedlings reach pole size
Woodland structure: dead wood	m ³ per hectare; number per hectare	At least 30m ³ /ha of fallen timber greater than 10cm diameter; 30 snags/ha; both categories should include stems greater than 40cm diameter	Dead wood is a valuable resource and an integral part of a healthy, functioning woodland ecosystem
Woodland structure: veteran trees	Number per hectare	No decline	Mature and veteran trees are important habitats for bryophytes, lichens, saproxylic organisms and some bird species. Their retention is important to ensure continuity of habitats/niches and propagule sources
Woodland structure: indicators of local distinctiveness	Occurrence	No decline	Includes ancient or long-established woodlands, archaeological and geological features as well as red-data and other rare or localised species. Perrin and Daly (2010) identified Annaghwood (NSNW site code 1624) as "possible ancient woodland"
Vegetation composition: native tree cover	Percentage	No decline. Native tree cover not less than 95%	Species reported in Perrin et al. (2008) and NPWS internal files

Vegetation composition: typical species	Occurrence	A variety of typical native species present, depending on woodland type, including oak (<i>Quercus petraea</i>) and birch (<i>Betula pubescens</i>)	Species reported in Perrin et al. (2008) and NPWS internal files
Vegetation composition: negative indicator species	Occurrence	Negative indicator species, particularly non-native invasive species, absent or under control	The following are the most common invasive species in this woodland type: rhododendron (<i>Rhododendron ponticum</i>), cherry laurel (<i>Prunus laurocerasus</i>), sycamore (<i>Acer pseudoplatanus</i>) and beech (<i>Fagus sylvatica</i>)

Conservation Objectives for : Lough Corrib SAC [000297]

91D0 Bog woodland

To maintain the favourable conservation condition of Bog woodland* in Lough Corrib SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes. At least 1.22ha. See map 8	Bog woodland occurs on Addergoole Bog in Lough Corrib SAC and is regarded as a component of the Active raised bogs habitat (7110) of that bog. Thus, the conservation objective and supporting document for active raised bog are also relevant to this habitat and common attributes have not been repeated here. The latest survey for bog woodland on Addergoole was carried out by the Raised Bog Monitoring Project (RBMP) in 2012 (Fernandez et al., 2014)
Habitat distribution	Occurrence	No decline, subject to natural processes. See map 8	Bog woodland occurs at one known location in this SAC, on Addergoole Bog. This is the most westerly known location for this habitat in Ireland
Vegetation composition: positive indicator species	Number in a representative number of monitoring stops	Birch (<i>Betula pubescens</i>), bog moss (<i>Sphagnum</i>) species and at least five other indicator species present	Bog woodland is typically species-poor but with a characteristic and distinctive flora. Positive indicator species are listed in Cross and Lynn (2013). Fernandez et al. (2014) note that Scot's pine (<i>Pinus sylvestris</i>) could be considered to be a positive indicator species; however, strong regeneration of the species can indicate that the site is drying out
Vegetation composition: negative indicator species	Percentage cover at a representative number of monitoring stops	Both native and non-native invasive species absent or under control. Total cover should be less than 10%	Negative indicator species include bracken (<i>Pteridium aquilinum</i>) and bramble (<i>Rubus fruticosus</i>), which can become invasive if the site begins drying out
Woodland structure: cover and height of birch	Percentage cover and metres at a representative number of monitoring stops	A minimum 30% cover of birch (<i>Betula pubescens</i>) with a median canopy height of 4m	Attribute and target based on bog woodland monitoring survey (Cross and Lynn, 2013)
Woodland structure: dwarf shrub cover	Percentage cover at a representative number of monitoring stops	Dwarf shrub cover not more than 50%	Attribute and target based on bog woodland monitoring survey (Cross and Lynn, 2013)
Woodland structure: ling cover	Percentage cover at a representative number of monitoring stops	Ling (<i>Calluna vulgaris</i>) cover not more than 40%	Attribute and target based on bog woodland monitoring survey (Cross and Lynn, 2013)
Woodland structure: bryophyte cover	Percentage cover at a representative number of monitoring stops	Bryophyte cover at least 50%, with bog moss (<i>Sphagnum</i> spp.) cover at least 25%	Attribute and target based on bog woodland monitoring survey (Cross and Lynn, 2013)
Woodland structure: tree size classes	Occurrence	Each size class present	Size classes are defined in Cross and Lynn (2013). The presence of all size classes suggests that a woodland has good structural variety with trees of varying ages
Woodland structure: senescent and dead wood	Occurrence	Senescent or dead wood present	Mature and veteran trees and dead wood are important for bryophytes, lichens, saproxylic organisms and some bird species. Their retention within a woodland is important to ensure continuity of habitats/niches and propagule sources over time. However, as birch (<i>Betula pubescens</i>) trees seldom exceed 30cm in diameter in this habitat and dead wood rots quickly and is engulfed by bog mosses (<i>Sphagnum</i> spp.), volume of dead wood may not be as high as in other woodland types

Conservation Objectives for : Lough Corrib SAC [000297]

1029 Freshwater Pearl Mussel *Margaritifera margaritifera*

To restore the favourable conservation condition of Freshwater Pearl Mussel in Lough Corrib SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Distribution	Kilometres	Maintain at 9.1km. See map 9	The conservation objective applies to the Owenriff freshwater pearl mussel (<i>Margaritifera margaritifera</i>) population in Lough Corrib SAC, which is of international importance and one of eight Irish populations prioritised for conservation action (Moorkens, 2010; NPWS, 2011). Its distribution is well-documented and full baseline monitoring took place in 2004 (Moorkens, 2004). The species is widespread in the Owenriff catchment, being found in the lower reaches of the Glengawbeg River, from Lough Agraftard to just upstream of the mouth of Lough Corrib in the Owenriff, and also in the Derrygauna tributary (Moorkens, 2004; NPWS, 2010). The Derrygauna River is in Connemara Bog Complex SAC (002034). The target length relates to the part of the distribution within Lough Corrib SAC. The objective is for the species to be sufficiently widespread to maintain itself on a long-term basis as a viable component of the Owenriff system. See NPWS (2010) for further information
Population size	Number of adult mussels	Restore Owenriff population to at least one million adult mussels	The Owenriff population was estimated as c.1 million in 2009 (NPWS, 2010). NPWS (2013) estimated that it had reduced to 940,000 in 2012, based on a 1% per year decline owing to insufficient recruitment. Declines were detected in 2011, 2014 and 2015. A large kill followed a drought in 2014 (Moorkens, 2015). The Owenriff population has been surveyed frequently since the 1990s, and monitored regularly since the baseline survey in 2004 (Moorkens, 2004, 2005, 2006, 2009, 2015, 2017; Moorkens and Killeen, 2008, 2014). Mussels are abundant (over 250 per 100m, often 150/m ²) from the hatchery at Canrawer East to upstream of Oughterard wastewater treatment plant (WWTP) discharge (Moorkens, 2004). The target is for the species to be sufficiently abundant to maintain itself on a long-term basis as a viable component of the Owenriff system
Population structure: recruitment	Percentage per size class	Restore to at least 20% of population no more than 65mm in length; and at least 5% of population no more than 30mm in length	Mussels ≤65mm are considered 'young mussels' and may be found buried in the substratum and/or beneath adult mussels. Mussels ≤30mm are 'juvenile mussels' and are always buried in the substratum. See the European Communities Environmental Objectives (Freshwater Pearl Mussel) Regulations 2009. The Owenriff had one of the best population profiles of any Irish population (Moorkens, 2004, 2009). NPWS (2010) summarises demographic work up to 2010. In 2013, it passed the targets, but failed both targets in 2011 (0% ≤65mm), 2014 (13.6% ≤65mm; 2% ≤30mm) and 2016 (10% ≤65mm; 0% ≤30mm) (Moorkens, 2011, 2015, 2017). In 2014, juveniles died owing to a combination of low flows and algal growth. In 2016, the smallest mussel was 11mm. The Owenriff population is unsustainable owing to lack of survival of juvenile mussels. The target is for sufficient juvenile recruitment to allow the species to maintain itself on a long-term basis as a viable component of the Owenriff system

Population structure: adult mortality	Percentage	No more than 5% decline from previous number of live adults counted; dead shells less than 1% of the adult population and scattered in distribution	5% is considered the cut-off between the combined errors associated with natural fluctuations and sampling methods and evidence of true population decline. 1% of dead shells is considered to be indicative of natural losses. Significant kills have been recorded on a number of occasions in the Owenriff, notably in 2004 and 2014. The Owenriff failed the target for dead shells in 2009, 2011, 2015 and 2016, but passed in 2014; and failed the 5% target in 2011, 2014 and 2015, but passed in 2016 (Moorkens, 2009, 2011, 2015, 2017; NPWS, 2010). The target is for sufficient survival of adults to allow the species to maintain itself on a long-term basis as a viable component of the Owenriff system
Suitable habitat: extent	Kilometres	Restore suitable habitat in more than 8.3km in the Owenriff and Glenawbeg rivers (see map 9) and any additional stretches necessary for salmonid spawning	The extent of the freshwater pearl mussel habitat in the Owenriff is well-documented; from the early surveys of Ross (1984, 1988) and Moorkens (1995, 1996) to full baseline monitoring in 2004 (Moorkens, 2004). Most of the available habitat in the Owenriff system is occupied by adult mussels, but below carrying capacity (Moorkens, 2004, 2005, 2006, 2009, 2011, 2015, 2017; Moorkens and Killeen, 2008, 2014). Annual monitoring has shown that episodes of poor habitat condition are causing loss of juveniles in the Owenriff, with very poor conditions recorded in 2011, 2014 and 2015, but near-favourable conditions in 2013 and 2016. Flow, macroalgal, macrophyte and sedimentation impacts are all concerns. Pollution appears to be chronic and episodes of good mussel habitat condition are found at higher flows. The target is for sufficient habitat in favourable condition to allow the species to maintain itself on a long-term basis as a viable component of the Owenriff system
Suitable habitat: condition	Kilometres	Restore condition of suitable habitat	The habitat is a combination of the area of 1) habitat adult and juvenile mussels can occupy; 2) spawning and nursery habitats host fish can occupy. Fish nursery and mussel habitat typically overlap. Fish spawning habitat is generally adjacent to mussel habitat, but may lie upstream of the generalised mussel distribution. Only spawning areas that regularly contribute juvenile fish to adult mussel habitat should be considered. Availability of mussel and fish habitat is determined by flow and substratum conditions. It is highly sensitive to hydromorphological changes, sedimentation and nutrient enrichment. Pressures throughout the catchment contribute to such impacts. The habitat in the Owenriff cannot support sufficient juvenile survival owing to nutrient and sediment pollution and hydrological impacts (Moorkens, 2017). The target is for sufficient habitat in favourable condition to allow the species to maintain itself on a long-term basis as a viable component of the Owenriff system
Water quality: macroinvertebrate and phytobenthos (diatoms)	Ecological quality ratio (EQR)	Restore water quality - macroinvertebrates: EQR greater than 0.90 (Q4-5 or Q5); phytobenthos: EQR greater than 0.93	These EQRs correspond to high ecological status for these two Water Framework Directive biological quality elements. They represent high water quality with very low nutrient concentrations (oligotrophic conditions). In 2009, the Owenriff River failed the macroinvertebrate standard at all sites and passed the phytobenthos standard (Ni Chatháin, 2009; Williams, 2009; NPWS, 2010). See also The European Communities Environmental Objectives (Surface Waters) Regulations 2009. The target is for sufficient habitat in favourable condition to allow the species to maintain itself on a long-term basis as a viable component of the Owenriff system

Substratum quality: filamentous algae (macroalgae); macrophytes (rooted higher plants)	Percentage	Restore substratum quality - filamentous algae: absent or trace (less than 5%); macrophytes: absent or trace (less than 5%)	The habitat in the Owenriff failed the macrophyte and the filamentous algal targets in 2009 (Moorkens, 2009; NPWS, 2010). Abundances of up to 100% filamentous algae and/or <i>Myriophyllum</i> were recorded in the best mussel habitat (Williams, 2009; NPWS, 2010). High cover abundance of both algae and macrophytes has been recorded regularly in the Owenriff from 2004-present, indicating on-going nutrient enrichment. In 2016, the algal target was exceeded in 57 of 65 quadrats and both filamentous algal and diatom biomass was elevated (Moorkens, 2017). Algae were abundant in 2015 and 2014 (Moorkens, 2015). Episodic poor habitat condition is leading to losses of juveniles. Good condition is linked to higher flows. Sufficient recruitment of juvenile mussels is being prevented by the poor condition of the river substratum. The target is for sufficient habitat in favourable condition to allow the species to maintain itself on a long-term basis as a viable component of the Owenriff system
Substratum quality: sediment	Occurrence	Restore substratum quality - stable cobble and gravel substrate with very little fine material; no artificially elevated levels of fine sediment	The habitat for the species in the Owenriff is unsuitable for the recruitment of juveniles owing to sedimentation and enrichment. The habitat failed this target in 2009 and significant sedimentation was recorded during all monitoring to 2009 (NPWS, 2010). Silt plumes and/or infiltration have been recorded 2010-2016 and the target was failed in 2011 and 2015 (Moorkens, 2011, 2015). Episodic poor habitat condition is leading to losses of juveniles. Good condition is linked to higher flows. Sufficient survival of juvenile mussels is being prevented by the poor condition of the river substratum. The target is for sufficient habitat in favourable condition to allow the species to maintain itself on a long-term basis as a viable component of the Owenriff system
Substratum quality: oxygen availability	Redox potential	Restore to no more than 20% decline from water column to 5cm depth in substrate	Differences in redox potential between the water column and the substrate correlate with differences in oxygen levels. Juvenile mussels require full oxygenation while buried in gravel. In suitable habitat, there should be very little loss of redox potential between the water column and underlying gravels. In 2009, the Owenriff failed the redox target, with averages at three locations of 18.9%, 24.2% and 24.5% at 5cm depth, giving an overall average of 22.5% (Moorkens, 2009; NPWS, 2010). The Owenriff passed the redox target in 2014 (average of 14.6%) and failed in 2011 (24%) and 2015 (22%) (Moorkens, 2011, 2015). In 2016, the average was 10.5%, and 100% of measurements had <20% loss (Moorkens, 2017). Episodic poor habitat condition is leading to losses of juveniles. Good condition is linked to higher flows. The target is for sufficient habitat in favourable condition to allow the species to maintain itself on a long-term basis as a viable component of the Owenriff system

Hydrological regime: flow variability	Metres per second	Restore appropriate hydrological regimes	The availability of suitable habitat is largely determined by flow (catchment geology being the other key factor). In order to restore the habitat for the species, flow variability over the annual cycle must be such that: 1) high flows can wash fine sediments from the substratum; 2) high flows are not artificially increased so as to cause excessive scour of mussel habitat; 3) low flows do not exacerbate the deposition of fine sediment or growth of algae/macrophytes and 4) low flows do not cause stress to mussels in terms of exposure, water temperatures, food availability or aspects of the reproductive cycle; see Moorikens and Killeen (2014). Groundwater inflow to the substratum also contributes to water-cycling and favourable habitat condition. Hydrological change and wetland loss in the Owenriff are significant concerns. The target is for sufficient habitat in favourable condition to allow the species to maintain itself on a long-term basis as a viable component of the Owenriff system
Host fish	Number	Maintain sufficient juvenile salmonids to host glochidial larvae	Salmonid fish are host to the larval stage of the freshwater pearl mussel and essential to completion of the life cycle. 0+ and 1+ fish are typically used, both because of habitat overlaps and the development of immunity with age in fish. Fish presence is sufficient, as higher fish density and biomass is indicative of enriched conditions in mussel rivers. Geist et al. (2006) found that higher densities of host fish coincided with eutrophication, poor substrate quality for mussels and a lack of mussel recruitment, while significantly lower densities and biomass of host fish were associated with high numbers of juvenile mussels. Fish movements must be such that 0+ fish remain in the mussel habitat until their 1+ summer. No fish stocking should occur within the mussel habitat, nor any works that may change the salmonid balance or residency time. No glochidia were found on fish in 2009, possibly owing to mussel stress (Johnston, 2009; NPWS, 2010). A fish host survey is planned for 2017/18
Fringing habitat: area and condition	Hectares	Maintain the area and condition of fringing habitats necessary to support the population	Riparian habitats, including those along lake fringes, even where not part of a natural floodplain, are integral to the structure and functioning of river systems. Fringing habitats aid in settlement of fine suspended material, protect banks from erosion, contribute to nutrient cycling and the aquatic food web (e.g. allochthonous matter such as leaf fall), and provide habitat (refuge and resources) for certain life-stages of fish, birds and aquatic invertebrates. Shade may be important in suppressing algal/macrophyte growth in enriched rivers and moderating temperatures. Fringing habitats are dependent on rivers/lakes, particularly water levels, and support wetland communities and species of conservation concern. Riparian wetlands and woodland are very important in the Owenriff and wetland damage is a significant concern. The target is for sufficient habitat in favourable condition to allow the species to maintain itself on a long-term basis as a viable component of the Owenriff system

Conservation Objectives for : Lough Corrib SAC [000297]

1092 White-clawed Crayfish *Austropotamobius pallipes*

To maintain the favourable conservation condition of White-clawed Crayfish in Lough Corrib SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Distribution: rivers	Occurrence	No reduction from baseline. See map 10	White-clawed crayfish (<i>Austropotamobius pallipes</i>) is recorded from the entire lengths of the four main tributaries of the River Clare. There are post-1996 records from the following tributaries: Abbert, Grange, Dalgan and Sinking Rivers. It is also present in some minor lower order streams within the Clare catchment
Distribution: Lough Corrib	Occurrence	No reduction from baseline. See map 10	The distribution of crayfish in Lough Corrib is uncertain. It certainly occurs in three 1km squares in the northern section of the lower basin (M2341, M2342, M2941) and is probably more widely distributed
Population structure: recruitment	Occurrence of juveniles and females with eggs	Juveniles and/or females with eggs in all occupied tributaries and occupied parts of Lough Corrib	See Reynolds et al. (2010) for further details
Negative indicator species	Occurrence	No alien crayfish species	Alien crayfish species are identified as a major direct threat to this species and as a disease vector. Ireland is currently free of non-native invasive crayfish species. See Reynolds (1998) for further details
Disease	Occurrence	No instances of disease	Disease is identified as a major threat and crayfish plague has occurred in Ireland even in the absence of alien vectors. Disease can, in some circumstances, be introduced through contaminated equipment and water in the absence of vector species. See Reynolds (1998) for further details
Water quality	EPA Q value	At least Q3-4 at all sites sampled by EPA	Target taken from Demers and Reynolds (2002). Q values based on triennial water quality surveys carried out by the Environmental Protection Agency (EPA)
Habitat quality: heterogeneity	Occurrence of positive habitat features	No decline in habitat heterogeneity or habitat quality	Crayfish need high habitat heterogeneity. Larger crayfish must have stones to hide under, or an earthen bank in which to burrow. Hatchlings shelter in vegetation, gravel and among fine tree-roots. Smaller crayfish are typically found among weed and debris in shallow water. Larger juveniles in particular may also be found among cobbles and detritus such as leaf litter. These conditions must be available throughout the occupied habitat

Conservation Objectives for : Lough Corrib SAC [000297]

1095 Sea Lamprey *Petromyzon marinus*

To restore the favourable conservation condition of Sea Lamprey in Lough Corrib SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Distribution: extent of anadromy	Percentage of river accessible	Greater than 75% of main stem length of rivers accessible from estuary	Sea lamprey (<i>Petromyzon marinus</i>) traditionally congregate and build spawning nests in the River Corrib in Galway city, both up- and downstream of the Salmon Weir Bridge. Their further upstream passage is impeded by the regulating weir immediately upstream. The combination of barriers to passage and low flows can impede further upstream passage in Irish catchments and prevent or reduce penetration and extensive colonisation (Gargan et al., 2011; Rooney et al., 2015). Sea lamprey have been recorded passing through the denil fish passage facility at the regulating weir. However, no quantitative assessment has been made, nor has any annual record been maintained. Sea lamprey have also been observed using their sucker mouths to project themselves up the damp concrete faces of the weir structure at low water levels (J. King, Inland Fisheries Ireland (IFI), pers. comm.)
Population structure of juveniles	Number of age/size groups	At least three age/size groups present	Attribute and target based on Harvey and Cowx (2003) and O'Connor (2007)
Juvenile density in fine sediment	Juveniles/m ²	Mean catchment juvenile density at least 1/m ²	Juveniles burrow in areas of fine sediment in still water. Attribute and target based on Harvey and Cowx (2003). No sites surveyed in 2006 (O'Connor, 2007) or 2013 (IFI, unpublished data) were positive for sea lamprey ammocoetes
Extent and distribution of spawning habitat	m ² and occurrence	No decline in extent and distribution of spawning beds	Attribute and target based on spawning bed habitat mapping by Inland Fisheries Ireland (IFI). Lampreys spawn in clean gravels. Artificial barriers can prevent lampreys from accessing suitable spawning habitat. As mentioned above, artificial barriers are currently preventing lamprey from accessing suitable spawning habitat above the regulating weir in the River Corrib
Availability of juvenile habitat	Number of positive sites in 3rd order channels (and greater), downstream of spawning areas	More than 50% of sample sites positive, with a minimum of four positive sites in a catchment, which are at least 5km apart	Artificial barriers can prevent juvenile lampreys from accessing the full extent of suitable habitat. Silting habitat is essential for larval lamprey and they can be severely impacted by sediment removal. Recovery can be rapid and newly-created habitat can be rapidly colonised (King et al., 2015). However, it is vital that such sedimenting habitats are retained

Conservation Objectives for : Lough Corrib SAC [000297]

1096 Brook Lamprey *Lampetra planeri*

To maintain the favourable conservation condition of Brook Lamprey in Lough Corrib SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Distribution	Percentage of river accessible	Access to all watercourses down to first order streams	Artificial barriers can block or cause difficulties to brook lampreys' migration both up- and downstream, thereby possibly limiting species to specific stretches, restricting access to spawning areas and creating genetically isolated populations (Espanhol et al., 2007)
Population structure of juveniles	Number of age/size groups	At least three age/size groups of brook/river lamprey present	Attribute and target based on data from Harvey and Cowx (2003) and JNCC (2005). It is impossible to distinguish between brook and river lamprey ammocoetes in the field (Gardiner, 2003), hence they are considered together in this target
Juvenile density in fine sediment	Ammocoetes/m ²	Mean catchment ammocoete density of brook/river lamprey at least 5/m ²	Ammocoetes burrow in areas of fine sediment in still water. Attribute target revised upward based on more recent proposals of JNCC (2005) and replacing initial proposals of Harvey and Cowx (2003). New criterion set at 5 ammocoetes/m ² on a catchment basis. The majority of sub-catchments in the SAC achieved this target in 2013 (IFI, unpublished data)
Extent and distribution of spawning habitat	m ² and occurrence	No decline in extent and distribution of spawning beds	Attribute and target based on spawning bed mapping by Inland Fisheries Ireland (IFI). Brook lamprey spawning habitat attributes compiled in Rooney et al. (2013) and the particle size required is considered to be available very widely in all river systems within the SAC, apart from very steep and torrential areas of boulder and bedrock. It is not considered that spawning habitat is a limiting feature for the conservation status of this species
Availability of juvenile habitat	Number of positive sites in 2nd order channels (and greater), downstream of spawning areas	More than 50% of sample sites positive	Target of 50% presence in suitable habitat based on Irish experience to date in catchment-wide surveys. 50% of surveyed sites in the catchment were positive in 2013 (IFI, unpublished data) compared with 49% in 2006 (O'Connor, 2007). Silting habitat is essential for larval lamprey and they can be severely impacted by sediment removal. Recovery can be rapid and newly-created habitat can be rapidly colonised (King et al., 2015). However, it is vital that such sedimenting habitats are retained

Conservation Objectives for : Lough Corrib SAC [000297]

1106 Salmon *Salmo salar*

To maintain the favourable conservation condition of Atlantic Salmon in Lough Corrib SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Distribution: extent of anadromy	Percentage of river accessible	100% of river channels down to second order accessible from estuary	There are no barriers to migration of salmon (<i>Salmo salar</i>) in Lough Corrib SAC. Salmon spawn in the headwaters of Lough Corrib tributaries. There is an artificial canal joining Lough Corrib and Lough Mask where salmon did not have access historically and does not constitute a limit on the distribution of salmon in Lough Corrib SAC
Adult spawning fish	Number	Conservation limit (CL) for each system consistently exceeded	A conservation limit (CL) is defined by the North Atlantic Salmon Conservation Organisation (NASCO) as "the spawning stock level that produces long-term average maximum sustainable yield as derived from the adult to adult stock and recruitment relationship". The target is based on the Standing Scientific Committee on Salmon (SSCS) annual model output of CL attainment levels. See SSCS (2016). Attainment of CL estimates are derived from direct counts of adults (rod catch, fish counter) or indirectly by fry abundance counts. The Corrib catchment is currently exceeding its CL
Salmon fry abundance	Number of fry/5 minutes electrofishing	Maintain or exceed 0+ fry mean catchment-wide abundance threshold value. Currently set at 17 salmon fry/5 minutes sampling	The target is the threshold value for rivers currently exceeding their conservation limit (CL)
Out-migrating smolt abundance	Number	No significant decline	Smolt abundance can be negatively affected by a number of impacts such as estuarine pollution, predation and sea lice (<i>Lepeophtheirus salmonis</i>)
Number and distribution of redds	Number and occurrence	No decline in number and distribution of spawning redds due to anthropogenic causes	Salmon spawn in clean gravels. The habitat for salmon is good and habitat rehabilitation programmes have been undertaken throughout the Corrib catchment to restore drained channels and repair habitat damaged by overgrazing
Water quality	EPA Q value	At least Q4 at all sites sampled by EPA	Q values based on triennial water quality surveys carried out by the Environmental Protection Agency (EPA)

Conservation Objectives for : Lough Corrib SAC [000297]

1303 Lesser Horseshoe Bat *Rhinolophus hipposideros*

To restore the favourable conservation condition of Lesser Horseshoe Bat in Lough Corrib SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Population per roost	Number	Minimum number of 100 bats for summer roost (roost id. 217 in NPWS database). See map 11	A figure of 100 bats for summer roosts was set as a minimum qualifying standard (MQS) when SACs were being selected for lesser horseshoe bats (<i>Rhinolophus hipposideros</i>). NPWS conduct annual counts at each qualifying roost. Qualified means from the 2006-2012 data have been calculated whereby the year with the highest maximum count and the year with the lowest maximum count were removed and the mean of the remaining years was calculated. This mean, or the MQS (i.e. 100 bats), whichever is higher, is set as the target figure for the roost
Summer roosts	Condition	No decline	Lough Corrib SAC has been selected for lesser horseshoe bats because of the presence of one important summer roost (roost id. 217 in NPWS database). Damage or disturbance to the roost or to the habitat immediately surrounding the roost will lead to a decline in its condition (Kelleher and Marnell, 2006)
Number of auxillary roosts	Number and condition	No decline	Lesser horseshoe bat populations will use a variety of roosts during the year besides the main summer maternity roost. Such additional roosts within the SAC may be important as night roosts/satellite roosts etc. A database of all known lesser horseshoe roosts is available on the National Biodiversity Data Centre website. NB further unrecorded roosts may also be present within this SAC
Extent of potential foraging habitat	Hectares	No significant decline	Lesser horseshoe bats normally forage in woodlands/scrub within 2.5km of their roosts (Schofield, 2008). See map 11 which shows a 2.5km zone around the above named roost and identifies potential foraging grounds
Linear features	Kilometres	No significant loss, within 2.5km of qualifying roosts. See map 11	This species follows commuting routes from its roost to its foraging grounds. Lesser horseshoe bats will not cross open ground. Consequently, linear features such as hedgerows, treelines and stone walls provide vital connectivity for this species, most importantly within 2.5km around each roost (Schofield, 2008)
Light pollution	Lux	No significant increase in artificial light intensity adjacent to named roost or along commuting routes within 2.5km of the roost. See map 11	Lesser horseshoes are very sensitive to light pollution and will avoid brightly lit areas. Inappropriate lighting around roosts may cause abandonment; lighting along commuting routes may cause preferred foraging areas to be abandoned, thus increasing the energetic cost for bats (Schofield, 2008)

Conservation Objectives for : Lough Corrib SAC [000297]

1355 Otter *Lutra lutra*

To maintain the favourable conservation condition of Otter in Lough Corrib SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Distribution	Percentage positive survey sites	No significant decline	Measure based on standard otter survey technique. Favourable Conservation Status (FCS) target, based on 1980/81 survey findings, is 88% in SACs. Current range is estimated at 93.6% (Reid et al., 2013)
Extent of terrestrial habitat	Hectares	No significant decline. Area mapped and calculated as 1,054ha along river banks/ lake shoreline/around ponds	No field survey. Areas mapped to include 10m terrestrial buffer along shoreline and river banks identified as critical for otters (NPWS, 2007)
Extent of freshwater (river) habitat	Kilometres	No significant decline. Length mapped and calculated as 314.2km	No field survey. River length calculated on the basis that otters will utilise freshwater habitats from estuary to headwaters (Chapman and Chapman, 1982)
Extent of freshwater (lake) habitat	Hectares	No significant decline. Area mapped and calculated as 4,178ha	No field survey. Area mapped based on evidence that otters tend to forage within 80m of the shoreline (NPWS, 2007)
Couching sites and holts	Number	No significant decline	Otters need lying up areas throughout their territory where they are secure from disturbance (Kruuk and Moorhouse, 1991; Kruuk, 2006)
Fish biomass available	Kilograms	No significant decline	Broad diet that varies locally and seasonally, but dominated by fish, in particular salmonids, eels and sticklebacks in freshwater (Bailey and Rochford, 2006; Reid et al., 2013)
Barriers to connectivity	Number	No significant increase. For guidance, see map 12	Otters will regularly commute across stretches of open water up to 500m e.g. between the mainland and an island; between two islands; across an estuary (De Jongh and O'Neill, 2010). It is important that such commuting routes are not obstructed

Conservation Objectives for : Lough Corrib SAC [000297]

1393 Slender Green Feather-moss *Drepanocladus vernicosus*

To maintain the favourable conservation condition of Slender Green Feather-moss (Shining Sickie-moss) in Lough Corrib SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Distribution of populations	Number and geographical spread of populations	No decline, subject to natural processes. See map 10 for known location at NW of Gortachalla Lough	(Please note that <i>Drepanocladus vernicosus</i> was reclassified as <i>Hamatocaulis vernicosus</i> by Hedenäs (1989)). The known population of slender green feather-moss (<i>Hamatocaulis vernicosus</i>) in Lough Corrib SAC occurs at NW of Gortachalla Lough in transition mire which is bounded to the west by acid bog. Data from NPWS surveys (NPWS internal files), Campbell (2013) and Campbell et al. (2015)
Population size	Number of individuals	No decline, subject to natural processes	The population at NW of Gortachalla Lough was estimated by Campbell (2013) to be 153,376,875 shoots (c.153,377,000 shoots). Counts of shoots were based on the mean of number of shoots in four 10cm x 10cm areas, extrapolated to 41,175 shoots/m ² in 3,725m ² (Campbell, 2013). See Campbell et al. (2015) for further details
Population cover	Percentage cover in a representative number of 2m x 2m monitoring plots	Mean percentage cover of slender green feather-moss (<i>Hamatocaulis vernicosus</i>) should be at least 45%	The mean percentage cover of slender green feather-moss (<i>Hamatocaulis vernicosus</i>) recorded in four 2m x 2m plots at NW of Gortachalla Lough was c.60% (Campbell, 2013). The target cover figure is a c.20% reduction of the recorded cover to allow for a margin of error and variability over monitoring seasons. See Campbell et al. (2015) for further details
Area of suitable habitat	Hectares	No decline, subject to natural processes	The area of occupancy at NW of Gortachalla Lough is estimated from mapping of GPS co-ordinates to be 6,209m ² . However, only c.60% of this is suitable habitat i.e. c.3,725m ² (0.373ha). See Campbell et al. (2015) for further details
Hydrological conditions: water table level	Metres	Maintain suitable hydrological conditions	Slender green feather-moss (<i>Hamatocaulis vernicosus</i>) is mostly confined to mesotrophic fens, a transitional habitat between acid bog and base-rich fen. This appears to occur in at least two forms in Ireland: upland transitional flushes, where the plants can occur in lawns that rise and fall with fluctuating water table levels; and wet lowland sedge meadows, where plants can be inundated in winter, but may be subject to some desiccation in the summer, such as at NW of Gortachalla Lough. Based on Campbell (2013) and Campbell et al. (2015)
Vegetation composition: tree cover	Percentage cover in a representative number of 2m x 2m monitoring plots	Mean percentage tree cover should be less than 15%	Slender green feather-moss (<i>Hamatocaulis vernicosus</i>) grows in moss-dominated, open communities, generally with a low cover of trees and shrubs. Campbell (2013) recorded 0% tree cover at NW of Gortachalla Lough. See also Campbell et al. (2015)
Vegetation composition: shrub cover	Percentage cover in a representative number of 2m x 2m monitoring plots	Mean percentage shrub cover should be less than 20%	Slender green feather-moss (<i>Hamatocaulis vernicosus</i>) grows in moss-dominated, open communities, generally with a low cover of trees and shrubs. Campbell (2013) recorded 0-1% shrub cover at NW of Gortachalla Lough. See also Campbell et al. (2015)
Vegetation composition: grass cover	Percentage cover in a representative number of 2m x 2m monitoring plots	Mean percentage grass species cover should be less than 25%	Slender green feather-moss (<i>Hamatocaulis vernicosus</i>) grows in moss-dominated, open communities, generally with a low cover of grasses, maintained by a low grazing intensity by rabbits (<i>Oryctolagus cuniculus</i>) at NW of Gortachalla Lough. Campbell (2013) recorded grass cover of 0-10% in four 2m x 2m plots at NW of Gortachalla Lough. See also Campbell et al. (2015)


Vegetation composition: bryophyte cover	Percentage cover in a representative number of 2m x 2m monitoring plots	Mean percentage bryophyte cover should be more than 50%	Campbell (2013) recorded bryophyte cover of 34-95% in four 2m x 2m plots at NW of Gortachalla Lough. See also Campbell et al. (2015)
Vegetation composition: cover of <i>Calliergonella cuspidata</i>	Percentage cover in a representative number of 2m x 2m monitoring plots	Mean percentage cover of <i>Calliergonella cuspidata</i> should be less than 15%	<i>Calliergonella cuspidata</i> , a moss species often associated with high nutrient conditions, is usually present, but with low cover and never dominant. Cover of <i>Calliergonella cuspidata</i> was 0-3% in four 2m x 2m plots recorded by Campbell (2013) at NW of Gortachalla Lough. See also Campbell et al. (2015)
Vegetation structure: vegetation height	Centimetres in a representative number of 2m x 2m monitoring plots	Mean vegetation height should not exceed 40cm	Campbell (2013) recorded a mean vegetation height of 40cm in four 2m x 2m plots at NW of Gortachalla Lough. See also Campbell et al. (2015)


Conservation Objectives for : Lough Corrib SAC [000297]

1833 Slender Naiad *Najas flexilis*

To restore the favourable conservation condition of Slender Naiad in Lough Corrib SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Population extent	Hectares; distribution	Restore the spatial extent of <i>Najas flexilis</i> within the lake, subject to natural processes. See map 13 for known locations	See the <i>Najas flexilis</i> supporting document for further details
Population depth	Metres	Restore the depth range of <i>Najas flexilis</i> within the lake, subject to natural processes	See the <i>Najas flexilis</i> supporting document for further details
Population viability	Plant traits	Restore plant fitness, subject to natural processes	See the <i>Najas flexilis</i> supporting document for further details
Population abundance	Square metres	Restore the cover abundance of <i>Najas flexilis</i> , subject to natural processes	See the <i>Najas flexilis</i> supporting document for further details
Species distribution	Occurrence	Restore to at least the north-western bay, subject to natural processes	See the <i>Najas flexilis</i> supporting document for further details
Habitat extent	Hectares	Restore, subject to natural processes	See the <i>Najas flexilis</i> supporting document for further details
Hydrological regime: water level fluctuations	Metres	Maintain appropriate natural hydrological regime necessary to support the habitat for the species	See the <i>Najas flexilis</i> supporting document for further details
Lake substratum quality	Various	Restore appropriate substratum type, extent and chemistry to support the population of the species	See the <i>Najas flexilis</i> supporting document for further details
Water quality	Various	Restore appropriate water quality to support the population of the species	See the <i>Najas flexilis</i> supporting document for further details
Acidification status	pH units; mg/l	Maintain appropriate water and sediment pH, alkalinity and cation concentrations to support the population of <i>Najas flexilis</i> , subject to natural processes	See the <i>Najas flexilis</i> supporting document for further details
Water colour	mg/l PtCo	Restore/maintain appropriate water colour to support the population of <i>Najas flexilis</i>	See the <i>Najas flexilis</i> supporting document for further details
Associated species	Species composition and abundance	Restore appropriate associated species and vegetation communities to support the population of <i>Najas flexilis</i>	See the <i>Najas flexilis</i> supporting document for further details
Fringing habitat: area and condition	Hectares	Maintain the area and condition of fringing habitats necessary to support the population of <i>Najas flexilis</i>	See the <i>Najas flexilis</i> supporting document for further details






 An Roinn Ealaíon, Oidhreacht, Gnóthaí Réigiúnacha, Tuaithe agus Gaeltachta
 Department of Arts, Heritage, Regional, Rural and Gaeltacht Affairs

MAP 1:
LOUGH CORRIB SAC
CONSERVATION OBJECTIVES
SAC DESIGNATION
 Map to be read in conjunction with the NPWS Conservation Objectives Document.

SITE CODE:
SAC 000297; version 3.09.
CO. GALWAY / MAYO / ROSCOMMON
 0 2 4 6 8 10 km


The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
 Ordnance Survey of Ireland Licence No EN 0059216. © Ordnance Survey of Ireland Government of Ireland
 Níl sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbhreithnithe a déanamh ar theorainneacha na gceantar comharthaíthe. Suirbhéarachta Ordonáis na hÉireann Ceadúnas Uimh EN 0059216. © Suirbhéarachta Ordonáis na hÉireann Rialtas na hÉireann


Map Version 1
Date: April 2017



Legend

- Lough Corrib SAC 000297
- Maumturk Mountains SAC 002008
- Connemara Bog Complex SAC 002034
- Monivea Bog SAC 002352
- Galway Bay Complex SAC 000268
- Lough Corrib SPA 004042
- OSI Discovery Series County Boundaries





Legend

- Lough Corrib SAC 000297
- OSI Discovery Series County Boundaries

Indicative Lake Habitats


- Dominant 3110, Dominant Oligotrophic waters containing very few minerals of sandy plains (*Littorelletalia uniflorarae*)
- Dominant 3130, Dominant Oligotrophic to mesotrophic standing waters with vegetation of the *Littorelletalia uniflorarae* and/or *Isoeto-Nanojuncetea*
- Dominant 3140, Dominant Hard oligo-mesotrophic waters with benthic vegetation of *Chara* spp.
- Potential 3110, Potential Oligotrophic waters containing very few minerals of sandy plains (*Littorelletalia uniflorarae*)
- Potential 3130, Potential Oligotrophic to mesotrophic standing waters with vegetation of the *Littorelletalia uniflorarae* and/or *Isoeto-Nanojuncetea*
- Potential 3110 / Potential 3130, Potential Oligotrophic waters containing very few minerals of sandy plains (*Littorelletalia uniflorarae*) / Potential Oligotrophic to mesotrophic standing waters with vegetation of the *Littorelletalia uniflorarae* and/or *Isoeto-Nanojuncetea*





Legend

- Lough Corrib SAC 000297
- High Bog Boundary
- Potential 7110 Active Raised Bogs
- OSI Discovery Series County Boundaries

 An Roinn Ealaíon, Oidhreacht, Gníothaí Réigiúnacha, Tuaithe agus Gaeltachta
 Department of Arts, Heritage, Regional, Rural and Gaeltacht Affairs

**MAP 4:
 LOUGH CORRIB SAC
 CONSERVATION OBJECTIVES
 EXTENT OF POTENTIAL
 ACTIVE RAISED BOGS**

Map to be read in conjunction with the NPWS Conservation Objectives Document.


SITE CODE:
 SAC 000297; version 3.09.
 CO. GALWAY / MAYO / ROSCOMMON

0 3 6 9 12 15 Kilometers

The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
 Ordnance Survey of Ireland Licence No EN 0059216. © Ordnance Survey of Ireland Government of Ireland


Níl sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbheithníthe a déanamh ar theorainneacha na gceantar comharthaithe. Suirbhéarachta Ordonáis na hÉireann Ceadúnas Uimh EN 0059216. © Suirbhéarachta Ordonáis na hÉireann Rialtas na hÉireann


Map Version 1
Date: April 2017



Legend

- Lough Corrib SAC 000297
- High Bog Boundary
- Active Raised Bog Ecotopes**
- Active Flush
- Bog Woodland
- Central
- Sub-central
- OSI Discovery Series County Boundaries



Legend

- Lough Corrib SAC 000297
- OSI Discovery Series County Boundaries

An Roinn Ealaíon, Oidhreacht, Gnóthaí Réigiúnacha, Tuairne agus Gaeltachta
Department of Arts, Heritage, Regional, Rural and Gaeltacht Affairs

**MAP 6:
LOUGH CORRIB SAC
CONSERVATION OBJECTIVES
DIGITAL ELEVATION MODEL
& DRAINAGE PATTERNS**

Map to be read in conjunction with the NPWS Conservation Objectives Document.

SITE CODE:
SAC 000297; version 3.09.
CO. GALWAY / MAYO / ROSCOMMON



0 3 6 9 12 15 Kilometers

The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
Ordnance Survey of Ireland Licence No EN 0059216. © Ordnance Survey of Ireland Government of Ireland

Nil sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbhreithnithe a déanamh ar theorainneacha na gceantar comharthaithe. Suirbhéarachta Ordonáis na hÉireann Ceadúnas Uimh EN 0059216. © Suirbhéarachta Ordonáis na hÉireann Rialtas na hÉireann






**Map Version 1
Date: April 2017**



Legend


- Lough Corrib SAC 000297
- Potential 8240 Limestone pavements including associated habitats






Legend

- Lough Corrib SAC 000297
- 1029 Freshwater Pearl Mussel - *Margaritifera margaritifera* Suitable habitat target
- 1029 Freshwater Pearl Mussel - *Margaritifera margaritifera* Distribution target
- 1029 Freshwater Pearl Mussel - *Margaritifera margaritifera* Catchment



Legend

- Lough Corrib SAC 000297
- ★ 1393 Slender Green Feather Moss - *Drepanocladus vernicosus*
- 1092 White-clawed Crayfish - *Austropotamobius pallipes*
- OSI Discovery Series County Boundaries



Legend


- Lough Corrib SAC 000297
- 1303 Lesser Horseshoe Bat - *Rhinolophus hipposideros* 2.5km buffer
- 1303 Lesser Horseshoe Bat - *Rhinolophus hipposideros* Potential Foraging Grounds

An Roinn Ealaíon, Oidhreacht, Gréthaí Réigiúnaíochta, Tuisithe agus Caistachta
 Department of Arts, Heritage, Regional, Rural and Gaeltacht Affairs

**MAP 11:
 LOUGH CORRIB SAC
 CONSERVATION OBJECTIVES
 LESSER HORSESHOE BAT**

Map to be read in conjunction with the NPWS Conservation Objectives Document.

SITE CODE:
 SAC 000297; version 3.09.
 CO. GALWAY / MAYO / ROSCOMMON




The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
 Ordnance Survey of Ireland Licence No EN 0059216. © Ordnance Survey of Ireland Government of Ireland

Nil sna teorainneacha ar na léarscáileanna ach nod garshuimhach ginearálta. Féadfar athbheirithne a déanamh ar theorainneacha na gceantar comharthaithe. Suirbhéarachta Ordonáis na hÉireann Ceadúnas Uimh EN 0059216. © Suirbhéarachta Ordonáis na hÉireann Rialtas na hÉireann




Map Version 1
 Date: April 2017



Legend

- Lough Corrib SAC 000297
- 1355 Otter - *Lutra lutra* Commuting 250m buffer



An Roinn Ealaíon, Oidhreachta, Gróthál Réigiúnacha, Tuaithe agus Gaeltachta
 Department of Arts, Heritage, Regional, Rural and Gaeltacht Affairs

MAP 12:
LOUGH CORRIB SAC
CONSERVATION OBJECTIVES
OTTER COMMUTING

Map to be read in conjunction with the NPWS Conservation Objectives Document.

SITE CODE:
 SAC 000297; version 3.04.
 CO. GALWAY / MAYO / ROSCOMMON


0 1 2 3 4 5 km

The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
 Ordnance Survey of Ireland Licence No EN 0059216. © Ordnance Survey of Ireland Government of Ireland

Níl sna teorainneacha ar na léarscáileanna ach nod garshuimhach ginearálta. Féadfar athbheirithne a déanamh ar theorainneacha na gceantar comharthaithe. Suirbhéarachta Ordonáis na hÉireann Ceadúnas Uimh EN 0059216. © Suirbhéarachta Ordonáis na hÉireann Rialtas na hÉireann

Map Version 1
Date: April 2017

© Ordnance Survey Ireland



Legend

- Lough Corrib SAC 000297
- 1833 Slender Naiad - *Najas flexilis* known habitat
- 1833 Slender Naiad - *Najas flexilis* possible habitat

An Foinn Ealaíon, Oidhreacht, Gnóthaí Réigiúnacha, Tuailíne agus Gaeltachta
 Department of Arts, Heritage, Regional, Rural and Gaeltacht Affairs

MAP 13:
LOUGH CORRIB SAC
CONSERVATION OBJECTIVES
SLENDER NAIAD - NAJAS FLEXILIS

Map to be read in conjunction with the NPWS Conservation Objectives Document.

SITE CODE:
 SAC 000297; version 3.09.
 CO. GALWAY / MAYO / ROSCOMMON

0 1.5 3 4.5 6 km

The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
 Ordnance Survey of Ireland Licence No EN 0059216. © Ordnance Survey of Ireland Government of Ireland

Nil sna teorainneacha ar na léarscáileanna ach nod garshuimhach ginearálta. Féadfar athbheithníthe a déanamh ar theorainneacha na gceantar comharthaithe. Suirbhéarachta Ordonáis na hÉireann Ceadúnas Uimh EN 0059216. © Suirbhéarachta Ordonáis na hÉireann Rialtas na hÉireann

N

Map Version 1
Date: April 2017

National Parks and Wildlife Service

Conservation Objectives Series

Killala Bay/Moy Estuary SAC 000458



An Roinn
Ealaíon, Oidhreachta agus Gaeltachta
Department of
Arts, Heritage and the Gaeltacht



**National Parks and Wildlife Service,
Department of Arts, Heritage and the Gaeltacht,
7 Ely Place, Dublin 2, Ireland.
Web: www.npws.ie
E-mail: nature.conservation@ahg.gov.ie**

Citation:

NPWS (2012) Conservation Objectives: Killala Bay/Moy Estuary SAC 000458. Version 1.0. National Parks and Wildlife Service, Department of Arts, Heritage and the Gaeltacht.

**Series Editors: Rebecca Jeffrey & Naomi Kingston
ISSN 2009-4086**

Introduction

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

* indicates a priority habitat under the Habitats Directive

000458	Killala Bay/Moy Estuary SAC
1014	Narrow-mouthed Whorl Snail <i>Vertigo angustior</i>
1095	Sea Lamprey <i>Petromyzon marinus</i>
1130	Estuaries
1140	Mudflats and sandflats not covered by seawater at low tide
1210	Annual vegetation of drift lines
1310	Salicornia and other annuals colonizing mud and sand
1330	Atlantic salt meadows (<i>Glauco-Puccinellietalia maritimae</i>)
1365	Harbour Seal <i>Phoca vitulina</i>
2110	Embryonic shifting dunes
2120	Shifting dunes along the shoreline with <i>Ammophila arenaria</i> ('white dunes')
2130	*Fixed coastal dunes with herbaceous vegetation ('grey dunes')
2190	Humid dune slacks

Please note that this SAC overlaps with Killala Bay/Moy Estuary SPA (004036) and is adjacent to River Moy SAC (002298). See map 2. The conservation objectives for this site should be used in conjunction with those for the overlapping and adjacent sites as appropriate.

Supporting documents, relevant reports & publications (listed by date)

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

Title: Harbour seal pilot monitoring project, 2011

Year: 2012

Author: NPWS

Series: Unpublished Report to NPWS

Title: Killala Bay/Moy Estuary SAC (000458). Conservation objectives supporting document - marine habitats and species. [Version 1]

Year: 2012

Author: NPWS

Series: Unpublished Report to NPWS

Title: Killala Bay/Moy Estuary SAC (000458). Conservation objectives supporting document - coastal habitats. [Version 1]

Year: 2012

Author: NPWS

Series: Unpublished Report to NPWS

Title: Subtidal Benthic Investigations in Killala Bay/Moy Estuary cSAC (Site Code: IE000458) Co. Sligo/Mayo

Year: 2011

Author: Aquafact

Series: Unpublished Report to NPWS & MI

Title: A survey of mudflats and sandflats in Ireland An intertidal soft sediment survey of Killala Bay

Year: 2011

Author: ASU

Series: Unpublished Report to NPWS & MI

Title: Monitoring and Condition Assessment of Populations of *Vertigo geyeri*, *Vertigo angustior* and *Vertigo moulinsiana* in Ireland

Year: 2011

Author: Moorkens, E.A.; Killeen, I.J.

Series: Irish Wildlife Manuals, No. 55

Title: Harbour seal pilot monitoring project, 2010

Year: 2011

Author: NPWS

Series: Unpublished Report to NPWS

Title: Harbour seal population monitoring 2009-2012: Report no. 1. Report on a pilot monitoring study carried out in southern and western Ireland, 2009

Year: 2010

Author: NPWS

Series: Unpublished Report to NPWS

Title: Saltmarsh Monitoring Report 2007-2008

Year: 2009

Author: McCorry, M.; Ryle, T.

Series: Unpublished Report to NPWS

Title:	Coastal Monitoring Project 2004-2006
Year:	2009
Author:	Ryle, T.; Murray, A.; Connolly, C.; Swann, M.
Series:	Unpublished Report to NPWS
Title:	The phytosociology and conservation value of Irish sand dunes
Year:	2008
Author:	Gaynor, K.
Series:	Unpublished PhD thesis, National University of Ireland, Dublin
Title:	Saltmarsh Monitoring Report 2006
Year:	2007
Author:	McCorry, M.
Series:	Unpublished Report to NPWS
Title:	A Survey of Juvenile Lamprey Populations in the Corrib and Suir Catchments
Year:	2007
Author:	O'Connor, W.
Series:	Irish Wildlife Manuals No. 26
Title:	Harbour seal population assessment in the Republic of Ireland: August 2003
Year:	2004
Author:	Cronin, M.; Duck, C.; Ó Cadhla, O.; Nairn, R.; Strong, D.; O'Keeffe, C.
Series:	Irish Wildlife Manuals No. 11
Title:	Summary of National Parks & Wildlife Service surveys for common (harbour) seals (<i>Phoca vitulina</i>) and grey seals (<i>Halichoerus grypus</i>), 1978 to 2003
Year:	2004
Author:	Lyons, D.O.
Series:	Irish Wildlife Manuals No. 13
Title:	A survey of juvenile lamprey populations in the Moy catchment
Year:	2004
Author:	O'Connor, W.
Series:	Irish Wildlife Manuals No. 15
Title:	Monitoring the river, sea and brook lamprey, <i>Lampetra fluviatilis</i> , <i>L. planeri</i> and <i>Petromyzon marinus</i>
Year:	2003
Author:	Harvey, J.; Cowx, I.
Series:	Conserving Natura 2000 Rivers Monitoring Series No. 5. English Nature, Peterborough
Title:	A survey of bottlenose dolphins (<i>Tursiops truncatus</i>) in the Shannon Estuary
Year:	2000
Author:	Rogan, E.; Ingram, S.; Holmes, B.; O'Flanagan, C.
Series:	Marine Institute Marine Resource Series No. 9
Title:	1989 survey of breeding herds of common seal <i>Phoca vitulina</i> with reference to previous surveys
Year:	1990
Author:	Harrington, R.
Series:	Unpublished Report to Wildlife Service

Title: An assessment of the status of the common seal *Phoca vitulina vitulina* in Ireland

Year: 1980

Author: Summers, C.F.; Warner, P.J; Nairn, R.G.W.; Curry, M.G.; Flynn, J.

Series: Biological Conservation 17: 115-123

Spatial data sources

Year:	2010
Title:	EPA WFD transitional waterbody data
GIS operations:	Clipped to SAC boundary. Expert opinion used as necessary to resolve any issues arising
Used for:	1130 (map 3)
Year:	Interpolated 2012
Title:	Mudflat and sandflat survey 2010; subtidal benthic survey 2010
GIS operations:	Polygon feature classes from marine community types base data sub-divided based on interpolation of marine survey data. Expert opinion used as necessary to resolve any issues arising
Used for:	Marine community types, 1140 (maps 4 and 5)
Year:	2005
Title:	OSi Discovery series vector data
GIS operations:	High water mark (HWM) and low water mark (LWM) polyline feature classes converted into polygon feature classes and combined; EU Annex I Saltmarsh and Coastal data erased out if present
Used for:	Marine community types base data (map 5)
Year:	Revision 2010
Title:	Saltmarsh Monitoring Project 2007-2008. Version 1
GIS operations:	QIs selected; clipped to SAC boundary; overlapping regions with Coastal CO data investigated and resolved with expert opinion used
Used for:	1310, 1330 (map 6)
Year:	2009
Title:	Coastal Monitoring Project 2004-2006. Version 1
GIS operations:	QIs selected; clipped to SAC boundary; overlapping regions with Saltmarsh CO data investigated and resolved with expert opinion used
Used for:	1210, 2110, 2120, 2130, 2190 (map 7)
Year:	2012
Title:	NPWS rare and threatened species database
GIS operations:	Dataset created from spatial references in database records. Expert opinion used as necessary to resolve any issues arising
Used for:	1014, 1365 (maps 8 and 9)
Year:	2005
Title:	OSi Discovery series vector data
GIS operations:	High Water Mark (HWM) polyline feature class converted into polygon feature class; clipped to SAC boundary. Expert opinion used as necessary to resolve any issues arising
Used for:	1365 (map 9)

1014 Narrow-mouthed Whorl Snail *Vertigo angustior*

To maintain the favourable conservation condition of Narrow-mouthed Whorl Snail in Killala Bay/Moy Estuary SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Distribution: occupied sites	Number	No decline. There is one known site for this species in this SAC. See map 8	From Moorkens and Killeen (2011)
Presence on transect	Occurrence	Adult or sub-adult snails are present in at least 3 places on the transect where optimal or sub-optimal habitat occurs (minimum 5 samples)	Transect established as part of condition assessment monitoring at this site (Moorkens and Killeen, 2011). See habitat area target below for definition of optimal and sub-optimal habitat
Abundance	Number per sample	At least 2 samples on the transect have more than 10 <i>V. angustior</i> individuals (minimum 5 samples)	From Moorkens and Killeen (2011)
Transect habitat quality	Metres	More than 50m of habitat along the transect is classed as optimal or sub-optimal	From Moorkens and Killeen (2011). See habitat area target below for definition of optimal and sub-optimal habitat
Transect optimal wetness	Metres	Soils, at time of sampling, are damp (optimal wetness) and covered with a layer of humid thatch for more than 50m along the transect	From Moorkens and Killeen (2011)
Habitat area	Hectares	1.465ha of potential habitat (optimal and sub-optimal); Optimal habitat is defined as marsh with transition of ecotone between red fescue (<i>Festuca rubra</i>) and silverweed (<i>Potentilla anserina</i>) wet grassland and waterlogged marsh dominated by yellow iris (<i>Iris pseudacorus</i>) and low growing herbs. Vegetation height 20-40cm. Habitat growing on wet to saturated soil covered with a deep layer of mosses and humid, open structured thatch. Sub-optimal habitat is defined as for optimal habitat, but either vegetation height is less than 20cm, or between 40 and 50cm; or the soil is dry, or covered with standing water	From Moorkens and Killeen (2011)

1095 Sea Lamprey *Petromyzon marinus*

To maintain the favourable conservation condition of Sea Lamprey in Killala Bay/Moy Estuary SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Distribution: extent of anadromy	% of estuary accessible	No barriers for migratory life stages of lamprey moving from freshwater to marine habitats and vice versa	This SAC only covers the estuarine portion of the River Moy. The adjacent River Moy SAC (site code: 2298) encompasses the freshwater elements of sea lamprey habitat. Artificial barriers can block or cause difficulties to lampreys' upstream migration, thereby limiting species to lower stretches and restricting access to spawning areas. See O'Connor (2004) for further information on artificial barriers in the Moy catchment
Population structure of juveniles	Number of age/size groups	At least three age/size groups present	Attribute and target based on data from Harvey and Cowx (2003) and O'Connor (2007). Important juvenile habitat identified immediately downstream of Ballina (see O'Connor, 2004)
Juvenile density in fine sediment	Juveniles/m ²	Juvenile density at least 1/m ²	Juveniles burrow in areas of fine sediment in still water. Attribute and target based on data from Harvey and Cowx (2003). Important juvenile habitat identified immediately downstream of Ballina (see O'Connor, 2004)

1130 Estuaries

To maintain the favourable conservation condition of Estuaries in Killala Bay/Moy Estuary SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	The permanent habitat area is stable or increasing, subject to natural processes. See map 3	Habitat area was estimated as 736ha using OSi data and the defined Transitional Water Body area under the Water Framework Directive
Community extent	Hectares	Maintain the extent of the <i>Zostera</i> -dominated community, subject to natural processes. See map 5	Estimated by EPA during 2011 intertidal survey. See marine supporting document for further details
Community structure: <i>Zostera</i> density	Shoots per m ²	Conserve the high quality of the <i>Zostera</i> -dominated community, subject to natural processes	Estimated by EPA during 2011 intertidal survey. See marine supporting document for further details
Community distribution	Hectares	Conserve the following community types in a natural condition: Muddy sand to fine sand dominated by <i>Hydrobia ulvae</i> , <i>Pygospio elegans</i> and <i>Tubificoides benedii</i> community complex; Estuarine muddy sand dominated by <i>Hediste diversicolor</i> and <i>Heterochaeta costata</i> community complex; and Fine sand dominated by <i>Nephtys cirrosa</i> community complex. See map 5	Habitat structure was elucidated from intertidal and subtidal surveys undertaken in 2010 (Aquafact, 2011; ASU, 2011). See marine supporting document for further details

1140 Mudflats and sandflats not covered by seawater at low tide

To maintain the favourable conservation condition of Mudflats and sandflats not covered by seawater at low tide in Killala Bay/Moy Estuary SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	The permanent habitat area is stable or increasing, subject to natural processes. See map 4	Habitat area was estimated as 1,332ha using OSi data
Community extent	Hectares	Maintain the extent of the <i>Zostera</i> -dominated community, subject to natural processes. See map 5	Estimated by EPA during 2011 intertidal survey. See marine supporting document for further details
Community structure: <i>Zostera</i> density	Shoots per m ²	Conserve the high quality of the <i>Zostera</i> -dominated community, subject to natural processes	Estimated by EPA during 2011 intertidal survey. See marine supporting document for further details
Community distribution	Hectares	Conserve the following community types in a natural condition: Muddy sand to fine sand dominated by <i>Hydrobia ulvae</i> , <i>Pygospio elegans</i> and <i>Tubificoides benedii</i> community complex; Estuarine muddy sand dominated by <i>Hediste diversicolor</i> and <i>Heterochaeta costata</i> community complex and Fine sand dominated by <i>Nephtys cirrosa</i> community complex. See map 5	Habitat structure was elucidated from intertidal survey undertaken in 2010 (ASU, 2011). See marine supporting document for further details

1210 Annual vegetation of drift lines

To maintain the favourable conservation condition of Annual vegetation of drift lines in Killala Bay/Moy Estuary SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes, including erosion and succession. For sub-site mapped: Bartragh Island- 0.58ha. See map 7	Based on data from the Coastal Monitoring Project (Ryle et al. 2009). Habitat is very difficult to measure in view of its dynamic nature which means that it can appear and disappear within a site from year to year. This habitat was only recorded from Bartragh Island. See coastal habitats supporting document for further details
Habitat distribution	Occurrence	No decline, or change in habitat distribution, subject to natural processes	Based on data from Ryle et al. (2009). Two separate narrow strips of strandline habitat were recorded on the northern side of Bartragh Island. See coastal habitats supporting document for further details
Physical structure: functionality and sediment supply	Presence/ absence of physical barriers	Maintain the natural circulation of sediment and organic matter, without any physical obstructions	Dunes are naturally dynamic systems that require continuous supply and circulation of sand. Accumulation of organic matter in tidal litter is essential for trapping sand and initiating dune formation. Sea defence/coastal protection works are present near the main access point to the beach at Inishcrone (Ryle et al. 2009). See coastal habitats supporting document for further details
Vegetation structure: zonation	Occurrence	Maintain the range of coastal habitats including transitional zones, subject to natural processes including erosion and succession	Based on data from Ryle et al. (2009). At Bartragh Island there are transitions from sand dunes into saltmarsh habitats. See coastal habitats supporting document for further details
Vegetation composition: typical species and sub-communities	Percentage cover at a representative sample of monitoring stops	Maintain the presence of species-poor communities with typical species: sea rocket (<i>Cakile maritima</i>), sea sandwort (<i>Honckenya peploides</i>), prickly saltwort (<i>Salsola kali</i>) and Orache (<i>Atriplex</i> spp.)	Based on data from Ryle et al. (2009). See coastal habitats supporting document for further details
Vegetation composition: negative indicator species	Percentage cover	Negative indicator species (including non-natives) to represent less than 5% cover	Negative indicators include non-native species, species indicative of changes in nutrient status and species not considered characteristic of the habitat. Based on data from Ryle et al. (2009). See coastal habitats supporting document for further details

1310 Salicornia and other annuals colonizing mud and sand

To maintain the favourable conservation condition of *Salicornia* and other annuals colonizing mud and sand in Killala Bay/Moy Estuary SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes, including erosion and succession. For sub-sites mapped: Bartragh Island- 0.26ha, Ross- 0.29ha. See map 6	Based on data from Saltmarsh Monitoring Project (SMP) (McCorry, 2007). Habitat mapped at two of the four sub-sites surveyed, giving a total estimated area of 0.55ha. NB further unsurveyed areas maybe present within the site. See coastal habitats supporting document for further details
Habitat distribution	Occurrence	No decline, or change in habitat distribution, subject to natural processes. See map 6 for known distribution	Based on data from McCorry (2007). <i>Salicornia</i> is an annual species, so its distribution can vary significantly from year to year. See coastal habitats supporting document for further details
Physical structure: sediment supply	Presence/ absence of physical barriers	Maintain natural circulation of sediments and organic matter, without any physical obstructions	Based on data from McCorry (2007). Sediment supply is particularly important for this pioneer saltmarsh community, as the distribution of this habitat depends on accretion rates. Accretion was noted at Ross and Bartragh Island. Old seawalls were recorded at Bartragh Island and some protection works were noted around buildings close to the shoreline at Ross. See coastal habitats backing document for further details
Physical structure: creeks and pans	Occurrence	Maintain creek and pan structure, subject to natural processes, including erosion and succession	Based on data from McCorry and Ryle (2009). Creeks deliver sediment throughout saltmarsh system. Creeks and pan structures are well developed at Ross. See coastal habitats supporting document for further details
Physical structure: flooding regime	Hectares flooded; frequency	Maintain natural tidal regime	This pioneer saltmarsh community requires regular tidal inundation. See coastal habitats supporting document for further details
Vegetation structure: zonation	Occurrence	Maintain the range of coastal habitats including transitional zones, subject to natural processes including erosion and succession	Based on data from McCorry (2007). Transitions to dune habitats are found at Bartragh Island and Ross. See coastal habitats supporting document for further details
Vegetation structure: vegetation height	Centimeters	Maintain structural variation within sward	Based on data from McCorry (2007). At Castleconor, grazing is absent. There are moderate levels of grazing at Rusheens, while grazing at Ross is heavy in places. Grazing intensity is low on Bartragh Island See coastal habitats supporting document for further details

1310 Salicornia and other annuals colonizing mud and sand

To maintain the favourable conservation condition of *Salicornia* and other annuals colonizing mud and sand in Killala Bay/Moy Estuary SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Vegetation structure: vegetation cover	Percentage cover at a representative sample of monitoring stops	Maintain more than 90% of the area outside of the creeks vegetated	Based on data from McCorry (2007). Castleconor and Rusheens are heavily poached in places. There are moderate levels of poaching at Bartragh Island and Ross. See coastal habitats supporting document for further details
Vegetation composition: typical species & sub-communities	Percentage cover	Maintain the presence of species-poor communities with typical species listed in the Saltmarsh Monitoring Project (McCorry and Ryle, 2009)	See coastal habitats supporting document for further details
Vegetation structure: negative indicator species- <i>Spartina anglica</i>	Hectares	No significant expansion of common cordgrass (<i>Spartina anglica</i>), with an annual spread of less than 1%	Based on data from McCorry (2007). See coastal habitats supporting document for further details

1330 Atlantic salt meadows (*Glauco-Puccinellietalia maritimae*)

To maintain the favourable conservation condition of Atlantic salt meadows (*Glauco-Puccinellietalia*) in Killala Bay/Moy Estuary SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes, including erosion and succession. For sub-sites mapped: Bartragh Island- 29.22ha, Ross- 14.95ha, Rusheens- 1.24ha, Castleconor - 1.61ha. See map 6	Based on data from the Saltmarsh Monitoring Project (SMP) (McCorry, 2007; McCorry and Ryle 2009). Four sub-sites that supported Atlantic salt meadow were mapped (47.02ha) and additional areas of potential ASM (3.34ha) were identified from an examination of aerial photographs, giving a total estimated area of 50.37ha. NB further unsurveyed areas maybe present within the site. See coastal habitats supporting document for further details
Habitat distribution	Occurrence	No decline, or change in habitat distribution, subject to natural processes. See map 6 for known distribution	Based on data from McCorry (2007). ASM is the dominant saltmarsh type with a wide distribution throughout the SAC. See coastal habitats supporting document for further details
Physical structure: sediment supply	Presence/ absence of physical barriers	Maintain natural circulation of sediments and organic matter, without any physical obstructions	Based on data from McCorry and Ryle (2009). The SMP noted accretion at Ross and Bartragh Island. Old seawalls were recorded at Bartragh Island and there are some protection works around buildings close to the shoreline at Ross. See coastal habitats supporting document for further details
Physical structure: creeks and pans	Occurrence	Maintain creek and pan structure/ allow to develop, subject to natural processes, including erosion and succession	Based on data from McCorry and Ryle (2009). Creeks and pan structures are well developed at Ross. See coastal habitats supporting document for further details
Physical structure: flooding regime	Hectares flooded; frequency	Maintain natural tidal regime	See coastal habitats supporting document for further details
Vegetation structure: zonation	Occurrence	Maintain the range of coastal habitats including transitional zones, subject to natural processes including erosion and succession	Based on data from McCorry (2007). Transitions to dune habitats are found at Bartragh Island and Ross. See coastal habitats supporting document for further details
Vegetation structure: vegetation height	Centimeters	Maintain structural variation within sward	Based on data from McCorry (2007). At Castleconor, grazing is absent. At Rusheens there are moderate levels of grazing. At Ross grazing is heavy in places. At Bartragh Island grazing intensity is low. See coastal habitats supporting document for further details

1330 Atlantic salt meadows (*Glauco-Puccinellietalia maritimae*)

To maintain the favourable conservation condition of Atlantic salt meadows (*Glauco-Puccinellietalia*) in Killala Bay/Moy Estuary SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Vegetation structure: vegetation cover	Percentage cover at a representative sample of monitoring stops	Maintain more than 90% of the area outside of the creeks vegetated	Based on data from McCorry (2007). Castleconor and Rusheens are heavily poached in places. There are moderate levels of poaching at Bartragh Island and Ross. See coastal habitats supporting document for further details
Vegetation composition: typical species and sub-communities	Percentage cover at a representative sample of monitoring stops	Maintain range of sub-communities with typical species listed in Saltmarsh Monitoring Project (McCorry and Ryle, 2009)	Based on data from McCorry and Ryle (2009). See coastal habitats supporting document for further details
Vegetation structure: negative indicator species- <i>Spartina anglica</i>	Hectares	No significant expansion of common cordgrass (<i>Spartina anglica</i>), with an annual spread of less than 1%	Based on data from McCorry (2007). See coastal habitats supporting document for further details

1365 Harbour Seal *Phoca vitulina*

To maintain the favourable conservation condition of Harbour Seal in Killala Bay/Moy Estuary SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Access to suitable habitat	Number of artificial barriers	Species range within the site should not be restricted by artificial barriers to site use. See map 9 for suitable habitat	See marine supporting document for further details
Breeding behaviour	Breeding sites	Conserve the breeding sites in a natural condition. See map 9	Attribute and target based on background knowledge of Irish breeding populations, review of data summarised by Summers et al. (1980), Harrington (1990), Lyons (2004) and unpublished National Parks and Wildlife Service records. See marine supporting document for further details
Moulting behaviour	Moult haul-out sites	Conserve the moult haul-out sites in a natural condition. See map 9	Attribute and target based on background knowledge of Irish populations, review of data from Lyons (2004), Cronin et al. (2004), NPWS (2010), NPWS (2011), NPWS (2012) and unpublished National Parks and Wildlife Service records. See marine supporting document for further details
Resting behaviour	Resting haul-out sites	Conserve the resting haul-out sites in a natural condition. See map 9	Attribute and target based on background knowledge of Irish populations, review of data from Lyons (2004), unpublished National Parks and Wildlife Service records and unpublished data collected by University College Cork/Inland Fisheries Ireland. See marine supporting document for further details
Disturbance	Level of impact	Human activities should occur at levels that do not adversely affect the harbour seal population at the site	See marine supporting document for further details

2110 Embryonic shifting dunes

To restore the favourable conservation condition of Embryonic shifting dunes in Killala Bay/Moy Estuary SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area increasing, subject to natural processes, including erosion and succession. For sub-site mapped: Ross- 0.81ha, Bartragh Island - 0.75ha. See map 7	Based on data from the Coastal Monitoring Project (Ryle et al., 2009). Habitat is very difficult to measure in view of its dynamic nature and was only recorded at Bartragh Island and Ross, giving a total estimated area of 1.56ha. Accretion was noted from the western end of Bartragh Island. Embryo dune habitat is restricted to a small area on the seaward edge at Ross. See coastal habitats supporting document for further details
Habitat distribution	Occurrence	No decline, or change in habitat distribution, subject to natural processes. See map 7 for known distribution	Based on data from Ryle et al. (2009). See coastal habitats supporting document for further details
Physical structure: functionality and sediment supply	Presence/ absence of physical barriers	Maintain the natural circulation of sediment and organic matter, without any physical obstructions	Dunes are naturally dynamic systems that require continuous supply and circulation of sand. Sea defence/coastal protection works are present near the main access point to the beach at Inishcrone (Ryle et al. 2009). See coastal habitats supporting document for further details
Vegetation structure: zonation	Occurrence	Maintain the range of coastal habitats including transitional zones, subject to natural processes including erosion and succession	Based on data from Gaynor (2008) and Ryle et al. (2009). At Bartragh Island and Ross there are transitions from sand dunes into saltmarsh habitats. See coastal habitats supporting document for further details
Vegetation composition: plant health of foredune grasses	Percentage cover	More than 95% of sand couch (<i>Elytrigia juncea</i>) and/or lyme-grass (<i>Leymus arenarius</i>) should be healthy (i.e. green plant parts above ground and flowering heads present)	Based on data from Ryle et al. (2009). See coastal habitats supporting document for further details
Vegetation composition: typical species and sub-communities	Percentage cover at a representative sample of monitoring stops	Maintain the presence of species-poor communities with typical species: sand couch (<i>Elytrigia juncea</i>) and/or lyme-grass (<i>Leymus arenarius</i>)	Based on data from Ryle et al. (2009). See coastal habitats supporting document for further details
Vegetation composition: negative indicator species	Percentage cover	Negative indicator species (including non-natives) to represent less than 5% cover	Based on data from Ryle et al. (2009). Negative indicators include non-native species, species indicative of changes in nutrient status and species not considered characteristic of the habitat. Sea-buckthorn (<i>Hippophae rhamnoides</i>) should be absent or effectively controlled. See coastal habitats supporting document for further details

2120 Shifting dunes along the shoreline with *Ammophila arenaria* ('white dunes')

To restore the favourable conservation condition of Shifting dunes along the shoreline with *Ammophila arenaria* (white dunes) in Killala Bay/Moy Estuary SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area increasing, subject to natural processes including erosion and succession. For sub-sites mapped: Ross- 1.58; Bartragh Island- 7.52ha ; Inishcrone- 3.65ha. See map 7	Habitat was mapped during the Coastal Monitoring Project (Ryle et al., 2009). Habitat was mapped at three sub-sites to give a total estimated area of 12.75ha. Habitat is very difficult to measure in view of its dynamic nature. See coastal habitats supporting document for further details
Habitat distribution	Occurrence	No decline, or change in habitat distribution, subject to natural processes. See map 7 for known distribution	Based on data from Ryle et al. (2009). Mobile dunes are well developed at Bartragh Island, while at Inishcrone they are patchy in distribution and eroded back to the fixed dune in places. See coastal habitats supporting document for further details
Physical structure: functionality and sediment supply	Presence/ absence of physical barriers	Maintain the natural circulation of sediment and organic matter, without any physical obstructions	Dunes are naturally dynamic systems that require continuous supply and circulation of sand. Marram (<i>Ammophila arenaria</i>) reproduces vegetatively and requires constant accretion of fresh sand to maintain active growth, thus encouraging further accretion. There are coastal protection works in place at Inishcrone. See coastal habitats supporting document for further details
Vegetation structure: zonation	Occurrence	Maintain the range of coastal habitats including transitional zones, subject to natural processes including erosion and succession	Based on data from Gaynor (2008) and Ryle et al. (2009). At both Bartragh Island and Ross there are transitions from sand dune to saltmarsh habitats. See coastal habitats supporting document for further details
Vegetation composition: plant health of dune grasses	Percentage cover	More than 95% of marram (<i>Ammophila arenaria</i>) and/or lyme-grass (<i>Leymus arenarius</i>) should be healthy (i.e. green plant parts above ground and flowering heads present)	Based on data from Ryle et al. (2009). See coastal habitats supporting document for further details
Vegetation composition: typical species and sub-communities	Percentage cover at a representative sample of monitoring stops	Maintain the presence of species-poor communities dominated by marram (<i>Ammophila arenaria</i>) and/or lyme-grass (<i>Leymus arenarius</i>)	Based on data from Ryle et al. (2009). Bartragh Island, Ross and Inishcrone all support a characteristic dune flora. See coastal habitats supporting document for further details

2120 Shifting dunes along the shoreline with *Ammophila arenaria* ('white dunes')

To restore the favourable conservation condition of Shifting dunes along the shoreline with *Ammophila arenaria* (white dunes) in Killala Bay/Moy Estuary SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Vegetation composition: negative indicator species	Percentage cover	Negative indicator species (including non-natives) to represent less than 5% cover	Based on data from Ryle et al. (2009). Negative indicators include non-native species, species indicative of changes in nutrient status and species not considered characteristic of the habitat. Sea-buckthorn (<i>Hippophae rhamnoides</i>) should be absent or effectively controlled. The mobile dune habitat at Ross has a high cover of creeping thistle (<i>Cirsium arvense</i>) and common ragwort (<i>Senecio jacobaea</i>). At Inishcrone and Bartragh Island, ragwort (<i>Senecio jacobaea</i>) is also common. See coastal habitats supporting document for further details

2130 *Fixed coastal dunes with herbaceous vegetation ('grey dunes')

To restore the favourable conservation condition of Fixed coastal dunes with herbaceous vegetation (grey dunes) in Killala Bay/Moy Estuary SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area increasing, subject to natural processes including erosion and succession. For sub-site mapped: Ross - 100.79ha; Bartragh Island - 120.13ha; Inishcrone - 38.53ha. See map 7	Based on data from the Coastal Monitoring Project (Ryle et al., 2009). Habitat mapped at three sub-sites to give a total estimated area of 259.46ha. See coastal habitats supporting document for further details
Habitat distribution	Occurrence	No decline, or change in habitat distribution, subject to natural processes. See map 7 for known distribution	Based on data from the Coastal Monitoring Project (Ryle et al., 2009). Fixed dune habitat is extensive at Bartragh Island. The extent of the fixed dune habitat is reduced at Inishcrone owing to presence of Enniscrone golf course. See coastal habitats supporting document for further details
Physical structure: functionality and sediment supply	Presence/ absence of physical barriers	Maintain the natural circulation of sediment and organic matter, without any physical obstructions.	Based on data from the Coastal Monitoring Project (Ryle et al., 2009). Physical barriers can lead to fossilisation or over-stabilisation of dunes, as well as beach starvation resulting in increased rates of erosion. There are coastal protection works at the main access to the beach at Inishcrone. See coastal habitats supporting document for further details
Vegetation structure: zonation	Occurrence	Maintain the range of coastal habitats including transitional zones, subject to natural processes including erosion and succession	Based on data from Ryle et al. (2009). At both Bartragh Island and Ross there are transitions from sand dune to saltmarsh habitats. See coastal habitats supporting document for further details
Vegetation structure: bare ground	Percentage cover	Bare ground should not exceed 10% of fixed dune habitat, subject to natural processes.	Based on data from Gaynor (2008) and Ryle et al. (2009). See coastal habitats supporting document for further details
Vegetation composition: sward height	Centimeters	Maintain structural variation within sward.	Based on data from Gaynor (2008) and Ryle et al. (2009). Vegetation is quite rank in places at Ross, Inishcrone and Bartragh Island due to undergrazing. See coastal habitats supporting document for further details
Vegetation composition: typical species and sub-communities	Percentage cover at a representative sample of monitoring stops	Maintain range of sub-communities with typical species listed in Ryle et al. (2009)	Based on data from Gaynor (2008) and Ryle et al. (2009). See coastal habitats supporting document for further details

2130 *Fixed coastal dunes with herbaceous vegetation ('grey dunes')

To restore the favourable conservation condition of Fixed coastal dunes with herbaceous vegetation (grey dunes) in Killala Bay/Moy Estuary SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Vegetation composition: negative indicator species (including <i>Hippophae rhamnoides</i>)	Percentage cover	Negative indicator species (including non-natives) to represent less than 5% cover	Based on data from Ryle et al. (2009). Negative indicators include non-native species, species indicative of changes in nutrient status and species not considered characteristic of the habitat. Sea-buckthorn (<i>Hippophae rhamnoides</i>) should be absent or effectively controlled. Bracken (<i>Pteridium aquilinum</i>) was recorded at Bartragh Island. At Inishcrone, common ragwort (<i>Senecio jacobaea</i>), creeping thistle (<i>Cirsium vulgare</i>) and bramble (<i>Rubus fruticosus</i>) occur. At Ross, creeping thistle (<i>Cirsium arvense</i>), common ragwort (<i>Senecio jacobaea</i>) and hogweed (<i>Heracleum sphondylium</i>) occur. See coastal habitats supporting document for further details
Vegetation composition: scrub/trees	Percentage cover	No more than 5% cover or under control	Based on data from Ryle et al. (2009). Scattered shrubs and stunted trees occur at Ross, while occasional scrub occurs at Bartragh Island. See coastal habitats supporting document for further details

2190 Humid dune slacks

To maintain the favourable conservation condition of Humid dune slacks in Killala Bay/Moy Estuary SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes including erosion and succession. For sub-sites mapped: Ross: 3.87ha; Bartragh Island: 1.22ha. See map 6	Based on data from the Coastal Monitoring Project (Ryle et al., 2009). Habitat was mapped at two sub-sites, giving a total estimated area of 5.09ha. See coastal habitats supporting document for further details
Habitat distribution	Occurrence	No decline or change in habitat distribution, subject to natural processes. See map 6 for known distribution	Based on data from Ryle et al. (2009). Dune slacks at Bartragh Island are narrow linear features. See coastal habitats supporting document for further details.
Physical structure: functionality and sediment supply	Presence/ absence of physical barriers	Maintain natural circulation of sediment and organic matter, without any physical obstructions	Physical barriers can lead to fossilisation or over-stabilisation of dunes, as well as beach starvation resulting in increased rates of erosion. See coastal habitats supporting document for further details
Physical structure: hydrological and flooding regime	Presence/ absence of water abstraction or drainage works	Maintain natural hydrological regime	Based on data from Gaynor (2008) and Ryle et al. (2009). See coastal habitats supporting document for further details
Vegetation structure: zonation	Occurrence	Maintain the range of coastal habitats including transitional zones, subject to natural processes including erosion and succession	Based on data from Ryle et al., (2009). At both Bartragh Island and Ross sub-sites there are transitions from sand dune to saltmarsh habitats. See coastal habitats supporting document for further details
Vegetation structure: bare ground	Percentage cover	Bare ground should not exceed 5% of dune slack habitat, with the exception of pioneer slacks which can have up to 20% bare ground.	Based on data from Gaynor (2008) and Ryle et al. (2009). At Ross, the dune slacks are poached by cattke in places. See coastal habitats supporting document for further details
Vegetation structure: vegetation height	Centimeters	Maintain structural variation within sward.	Based on data from Ryle et al. (2009). See coastal habitats supporting document for further details
Vegetation composition: typical species and sub-communities	Percentage cover at a representative sample of monitoring stops	Maintain range of sub-communities with typical species listed in Ryle et al. (2009)	Based on data from Gaynor (2008) and Ryle et al. (2009). See coastal habitats supporting document for further details
Vegetation composition: cover of <i>S. repens</i>	% cover; centimeters	Maintain more than 40% cover of creeping willow (<i>Salix repens</i>)	Based on data from Ryle et al. (2009). Cover of creeping willow (<i>Salix repens</i>) needs to be controlled (e.g. through an appropriate grazing regime) to prevent the development of a coarse, rank vegetation cover. <i>Salix repens</i> ssp. <i>argentea</i> was noted at Bartragh Island, but its cover was only 10% and it was not widespread. See coastal habitats supporting document for further details

2190 Humid dune slacks


To maintain the favourable conservation condition of Humid dune slacks in Killala Bay/Moy Estuary SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Vegetation composition: negative indicator species	Percentage cover	Negative indicator species (including non-natives) to represent less than 5% cover	Based on data from Ryle et al. (2009). Negative indicators include non-native species, species indicative of changes in nutrient status and species not considered characteristic of the habitat. Sea-buckthorn (<i>Hippophae rhamnoides</i>) should be absent or effectively controlled. See coastal habitats supporting document for further details
Vegetation composition: scrub/trees	Percentage cover	No more than 5% cover or under control	Based on data from Ryle et al. (2009). See coastal habitats supporting document for further details




Legend

SAC 000458




Legend

- SAC 000458 Killala Bay/Moy Estuary
- SAC 002298 River Moy
- SPA 004036 Killala Bay/Moy Estuary
- OSi Discovery Series County Boundaries




Legend

- SAC 000458
- 1130 Estuaries
- OSi Discovery Series County Boundaries




Legend

- SAC 000458
- 1140 Mudflats and sandflats not covered by seawater at low tide
- OSi Discovery Series County Boundaries



Legend

- SAC 000458
- OSi Discovery Series County Boundaries
- Marine Community Types**
- Estuarine muddy sand dominated by *Hediste diversicolor* and *Heterochaeta costata* community complex
- Fine sand dominated by *Nephtys cirrosa* community complex
- Muddy sand to fine sand dominated by *Hydrobia ulvae*, *Pygospio elegans* and *Tubificoides benedii* community complex
- River
- Zostera*-dominated community



Legend

- SAC 000458
- OSi Discovery Series County Boundaries
- SMP: 0026 Saltmarsh Monitoring Project Site Codes


Saltmarsh Habitats


Qualifying Interests

- 1310 *Salicornia* and other annuals colonising mud and sand
- 1330 Atlantic salt meadows (*Glauco-Puccinellietalia maritimae*)
- Potential 1330 Atlantic salt meadows (*Glauco-Puccinellietalia maritimae*)
- Potential 1330 / 1410 Atlantic salt meadows (*Glauco-Puccinellietalia maritimae*) / Mediterranean salt meadows (*Juncetalia maritimi*)

Non-Qualifying Interests

- 1410 Mediterranean salt meadows (*Juncetalia maritimi*)
- Potential 1410 Mediterranean salt meadows (*Juncetalia maritimi*)






Legend


- SAC 000458
- OSi Discovery Series County Boundaries

CMP: 131 Coastal Monitoring Project Site Codes

Qualifying Interests


- 1210 Annual vegetation of drift lines
- 2110 Embryonic shifting dunes
- 2120 Shifting dunes along the shoreline with *Ammophila arenaria* ('white dunes')
- 2130 *Fixed coastal dunes with herbaceous vegetation ('grey dunes')
- 2190 Humid dune slacks





Legend

- SAC 000458
- 1014 Narrow-Mouthed Whorl Snail - *Vertigo angustior*
- OSi Discovery Series County Boundaries



Legend

- SAC 000458
- ★ 1365 Harbour Seal - *Phoca vitulina* breeding sites
- ★ 1365 Harbour Seal - *Phoca vitulina* moulting sites
- ★ 1365 Harbour Seal - *Phoca vitulina* resting sites
- 1365 Harbour Seal - *Phoca vitulina* habitat
- OSi Discovery Series County Boundaries



Conservation objectives for Ardkill Turlough SAC [000461]

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Objective: To maintain or restore the favourable conservation condition of the Annex I habitat(s) and/or the Annex II species for which the SAC has been selected:

Code	Description
------	-------------

3180	Turloughs*
------	------------

* denotes a priority habitat



Citation: NPWS (2020) Conservation objectives for Ardkill Turlough SAC [000461]. Generic Version
7.0. Department of Culture, Heritage and the Gaeltacht.



Conservation objectives for Balla Turlough SAC [000463]

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Objective: To maintain or restore the favourable conservation condition of the Annex I habitat(s) and/or the Annex II species for which the SAC has been selected:

Code	Description
------	-------------

3180	Turloughs*
------	------------

* denotes a priority habitat



Citation: NPWS (2020) Conservation objectives for Balla Turlough SAC [000463]. Generic Version 7.0.
Department of Culture, Heritage and the Gaeltacht.

National Parks and Wildlife Service

Conservation Objectives Series

Bellacorick Iron Flush SAC 000466



An Roinn Cultúir,
Oidhreacht agus Gaeltachta
Department of Culture,
Heritage and the Gaeltacht

**National Parks and Wildlife Service,
Department of Culture, Heritage and the Gaeltacht,
90 King Street North, Dublin 7, D07 N7CV, Ireland.**

**Web: www.npws.ie
E-mail: nature.conservation@chg.gov.ie**

Citation:

**NPWS (2019) Conservation Objectives: Bellacorick Iron Flush SAC 000466.
Version 1. National Parks and Wildlife Service, Department of Culture, Heritage
and the Gaeltacht.**

**Series Editor: Rebecca Jeffrey
ISSN 2009-4086**

Introduction

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

** indicates a priority habitat under the Habitats Directive*

000466 Bellacorick Iron Flush SAC

1528 Marsh Saxifrage *Saxifraga hirculus*

Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

NPWS Documents

Year :	2015
Title :	Monitoring recommendations for Marsh Saxifrage (<i>Saxifraga hirculus</i> L.) in the Republic of Ireland
Author :	Muldoon, C.S.; Waldren, S.; Lynn, D.
Series :	Irish Wildlife Manuals, No. 88
Year :	2019
Title :	Results of a monitoring survey of the Annex II species <i>Saxifraga hirculus</i> (Marsh Saxifrage) in Ireland 2015–2018
Author :	O'Neill, F.H.; Hodd, R.L; Long, M.P.
Series :	Irish Wildlife Manuals, No. 112

Other References


Year :	1958
Title :	<i>Saxifraga hirculus</i> in Co. Mayo (Dooleeg)
Author :	Scannell, M.J.P.
Series :	Irish Naturalists' Journal, 12: 248
Year :	1960
Title :	Notes on the vegetation of a mineral flush in Co. Mayo
Author :	King, A.L.K.; Scannell, M.J.P.
Series :	Irish Naturalists' Journal, 13: 137-140
Year :	1988
Title :	International Mires Research Group Field Excursion To Ireland 1988
Author :	Fojt, W.
Series :	Nature Conservancy Council
Year :	1991
Title :	Phytosociological and ecological studies of lowland blanket bog flushes in West Galway and North Mayo
Author :	Lockhart, N.D.
Series :	Unpublished Ph.D. Thesis, National University of Ireland, Galway
Year :	2011
Title :	Conservation biology of <i>Saxifraga hirculus</i> L. in Ireland
Author :	Muldoon, C.S.
Series :	Unpublished Ph.D. Thesis, Trinity College Dublin
Year :	2012
Title :	Bellacorick Iron Flush vegetation study as part of hydro-geological investigations
Author :	BES (Biosphere Environmental Services)
Series :	Unpublished report for ESB International

1528 Marsh Saxifrage *Saxifraga hirculus*


To restore the favourable conservation condition of Marsh Saxifrage in Bellacorick Iron Flush SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Distribution	Number and geographical spread of populations	No loss in geographical spread and number of known populations, subject to natural processes	The known population of marsh saxifrage (<i>Saxifraga hirculus</i>) in Bellacorick Iron Flush SAC occurs in a rich flush. The SAC is surrounded by drains and extensive areas of mechanically-cut peat. The population was surveyed by O'Neill et al. (2019; site SH03 Bellacorick) in 2018; previous surveys include Muldoon (2011), BES (2012) and an NPWS commissioned rare plant survey in 1995 (NPWS internal files). Surveys of the flush also include Fojt (1988) and Lockhart (1991). See also Scannell (1958) and King and Scannell (1960)
Number of rosettes	Number	No decline in number of rosettes, subject to natural processes	See Muldoon et al. (2015) and O'Neill et al. (2019) for full details on methodology used to calculate number of rosettes. Muldoon (2011) recorded 700 rosettes at Bellacorick and a target of 560 rosettes was set, i.e. 80% of that recorded to allow a margin of error over monitoring seasons (Muldoon et al., 2015). O'Neill et al. (2019) recorded a total count of 23 rosettes. While a decline of 97% is suggested from these figures, differences in survey intensity and weather conditions may play a role. Previous survey numbers have fluctuated, e.g. a count of 27–30 flowering stalks plus many vegetative rosettes was recorded in 1995 (NPWS internal files), an estimate of 300 individuals was made in 1999 (NPWS internal files) and 40 flowering heads (no individual rosette counts made) were recorded in 2012 (BES, 2012). Overall, however, the population may have reached an equilibrium with the conditions prevailing in the SAC (Lockhart, pers. comm.). See O'Neill et al. (2019) for further details
Density of rosettes (rosettes/square metre)	Mean number of rosettes in a representative number of 1m x 1m monitoring stops	No decline in density of rosettes, subject to natural processes	See Muldoon et al. (2015) and O'Neill et al. (2019) for full details on methodology used to calculate density of rosettes. Density of rosettes calculation was not undertaken by O'Neill et al. (2019) at Bellacorick (site code SH03) as the species occurs as isolated clumps. However, a total count of rosettes was conducted (O'Neill et al., 2019)
Number of flowering heads	Number	No decline in number of flowering heads, subject to natural processes	Number of flowering heads is estimated as an order of magnitude (10s, 100s, 1,000s, or 10,000s). See Muldoon et al. (2015) and O'Neill et al. (2019) for full details on methodology used to calculate number of flowering heads
Area of occupancy	Square metres	No decline in the area occupied by the population, subject to natural processes	See Muldoon et al. (2015) and O'Neill et al. (2019) for full details on methodology used to estimate area of occupancy. An area of occupancy of 950m ² was recorded by Muldoon (2011) and a target of 855m ² was set, i.e. 90% of that recorded to allow a margin of error over monitoring seasons (Muldoon et al., 2015). An area of occupancy of c.156m ² was recorded by O'Neill et al. (2019) at Bellacorick in 2018. However, the core area of the population has remained largely unchanged since its discovery in the 1950s; see O'Neill et al. (2019) for further details

Hydrological conditions: wetness of substrate	Light pressure from hand onto substrate in a representative number of 1m x 1m monitoring stops	Water from the soil or peat substrate should cover the fingers of hand in at least 40% of monitoring stops	Marsh saxifrage (<i>Saxifraga hirculus</i>) requires the presence of groundwater close to the surface, but the species will not tolerate long periods of flooding and the water should be moving or flowing to some extent. Attribute and target based on O'Neill et al. (2019). At Bellacorick, this attribute failed when surveyed after a period of drought in August 2018 and no standing water or water visible with pressure was noted by O'Neill et al. (2019). While active drainage is not currently an issue at Bellacorick, significant drainage had occurred in the past (beginning in the 1950s for peat extraction) and water levels have not recovered (O'Neill et al., 2019)
Vegetation composition: positive indicator species	Occurrence in a representative number of 1m x 1m monitoring stops	Knotted pearlwort (<i>Sagina nodosa</i>) should be present in at least 40% of monitoring stops	Attribute and target based on O'Neill et al. (2019). Knotted pearlwort (<i>Sagina nodosa</i>) was absent from all monitoring stops at Bellacorick (O'Neill et al., 2019)
Vegetation composition: negative indicator species	Percentage cover in a representative number of 1m x 1m monitoring stops	Mean percentage cover of purple moor-grass (<i>Molinia caerulea</i>) should not exceed 5%	Attribute and target based on Muldoon et al. (2015) and O'Neill et al. (2019)
Vegetation composition: negative indicator species	Percentage cover in a representative number of 1m x 1m monitoring stops	Mean percentage cover of Yorkshire-fog (<i>Holcus lanatus</i>) should not exceed 15%	Attribute and target based on Muldoon et al. (2015) and O'Neill et al. (2019)
Vegetation structure: mean vegetation height	Centimetres in a representative number of 1m x 1m monitoring stops	Mean vegetation height should not exceed 20cm	See O'Neill et al. (2019) for full details on methodology used to calculate mean vegetation height (cm). At Bellacorick, O'Neill et al. (2019) recorded a mean vegetation height of 42cm and noted insufficient grazing, with tall, rank vegetation a feature of the habitat for the species
Vegetation structure: grazing level	Level in a representative number of 1m x 1m monitoring stops	Grazing should be at light to moderate levels (26-50%) to ensure an open vegetation structure and to allow flowering to occur	At each monitoring stop, grazing levels are assigned to one of four categories 0-25% (little/no grazing), 26-50%, 51-75% and 76-100% (heavy overgrazing). The median of each category is calculated and averaged among all stops to assign the grazing level to one of the four categories. See Muldoon et al. (2015) and O'Neill et al. (2019) for full details on methodology. At Bellacorick, grazing levels were found to be too low (0-25%) when surveyed by Muldoon (2011; see also Muldoon et al., 2015). In 2018, an insufficient level of grazing was recorded as a negative impact at Bellacorick (O'Neill et al., 2019). O'Neill et al. (2019) noted that it was probable that sheep grazing was being further discouraged by construction roads and traffic associated with a nearby wind farm development



Legend

 Bellacorick Iron Flush SAC 000466





An Roinn Cultúir,
Oidhreacht agus Gaeltachta
Department of Culture,
Heritage and the Gaeltacht

**MAP 1:
BELLACORICK IRON FLUSH SAC
CONSERVATION OBJECTIVES
SAC DESIGNATION**

Map to be read in conjunction with the NPWS Conservation Objectives Document.

**SITE CODE:
SAC 000466; version 3.01. CO. MAYO**

0 30 60 90 120 Meters



The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
Ordnance Survey of Ireland Licence No OSI-NMA-014. © Ordnance Survey of Ireland Government of Ireland

Níl sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbheithnithe a déanamh ar theorainneacha na gceantar comharthaithe. Suirbhéarachta Ordonáis na hÉireann Ceadúnas Uimh OSI-NMA-014. © Suirbhéarachta Ordonáis na hÉireann Rialtas na hÉireann



**Map Version 1
Date: May 2019**

National Parks and Wildlife Service

Conservation Objectives Series

Mullet/Blacksod Bay Complex SAC 000470



*An Roinn
Ealaíon, Oidhreachta agus Gaeltachta*

*Department of
Arts, Heritage and the Gaeltacht*



**National Parks and Wildlife Service,
Department of Arts, Heritage and the Gaeltacht,**

7 Ely Place, Dublin 2, Ireland.

Web: www.npws.ie

E-mail: nature.conservation@ahg.gov.ie

Citation:

**NPWS (201) Conservation Objectives: Mullet/Blacksod Bay Complex SAC
000470. Version 1. National Parks and Wildlife Service, Department of Arts,
Heritage and the Gaeltacht.**

Series Editor: Rebecca Jeffrey

ISSN 2009-4086

Introduction

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

* indicates a priority habitat under the Habitats Directive

000470	Mullet/Blacksod Bay Complex SAC
1140	Mudflats and sandflats not covered by seawater at low tide
1160	Large shallow inlets and bays
1170	Reefs
1310	Upland heath and other annuals colonising mud and sand
1355	Otter <i>Lutra lutra</i>
1395	Petalwort <i>Petalophyllum ralfsii</i>
2120	Shifting dunes along the shoreline with Cladonia <i>Cladonia</i> (white dunes)
2130	Fixed coastal dunes with herbaceous vegetation (grey dunes)E
2150	Atlantic decalcified fixed dunes (Calluno-Ulicetea)
21A0	Machairs (* in Ireland)
3150	Natural eutrophic lakes with Magnopotamion or Hydrocharition - type vegetation
7230	Alkaline fens

Please note that this SAC overlaps with Blacksod Bay/Broadhaven SPA (004037), Termoncarragh Lake and Annagh Machair SPA (004093) and Mullet Peninsula SPA (004227). It adjoins West Connaught Coast SAC (002998). See map 2. The conservation objectives for this site should be used in conjunction with those for the overlapping and adjacent sites as appropriate.

Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

NPWS Documents

Year :	1996
Title :	Biomar survey of Irish machair sites
Author :	Crawford, I.; Bleasdale, A.; Conaghan, J.
Series :	Irish Wildlife Manual No. 3
Year :	2007
Title :	Supporting documentation for the Habitats Directive Conservation Status Assessment - backing documents. Article 17 forms and supporting maps
Author :	NPWS
Series :	Unpublished report to NPWS
Year :	2008
Title :	Survey of sensitive subtidal benthic marine communities in Mullet/Blacksod Bay Complex SAC, Rutland Island and Sound SAC, Mulroy Bay SAC
Author :	MERC
Series :	Unpublished report to NPWS
Year :	2009
Title :	Coastal Monitoring Project 2004-2006
Author :	Ryle, T.; Murray, A.; Connolly, K.; Swann, M.
Series :	Unpublished report to NPWS
Year :	2009
Title :	Saltmarsh monitoring project 2007-2008
Author :	McCorry, M.; Ryle, T.
Series :	Unpublished report to NPWS
Year :	2013
Title :	National otter survey of Ireland 2010/12
Author :	Reid, N.; Hayden, B.; Lundy, M.G.; Pietravalle, S.; McDonald, R.A.; Montgomery, W.I.
Series :	Irish Wildlife Manual No. 76
Year :	2013
Title :	A survey of the benthic macrophytes of three hard-water lakes: Lough Bunny, Lough Carra and Lough Owel
Author :	Roden, C.; Murphy, P.
Series :	Irish Wildlife Manual No. 70
Year :	2013
Title :	Monitoring survey of Annex I sand dune habitats in Ireland
Author :	Delaney, A.; Devaney, F.M; Martin, J.M.; Barron, S.J.
Series :	Irish Wildlife Manual No. 75
Year :	2013
Title :	The status of EU protected habitats and species in Ireland. Volume 2. Habitats assessments
Author :	NPWS
Series :	Conservation assessments
Year :	2013
Title :	The status of EU protected habitats and species in Ireland. Volume 3. Species assessments
Author :	NPWS
Series :	Conservation assessments

Year : 2014
Title : Guidelines for a national survey and conservation assessment of upland vegetation and habitats in Ireland, Version 2.0
Author : Perrin, P.M.; Barron, S.J.; Roche, J.R.; O'Hanrahan, B.
Series : Irish Wildlife Manual No. 79

Year : 2014
Title : Mullet/Blacksod Bay Complex SAC (site code: 470) Conservation objectives supporting document- coastal habitats V1
Author : NPWS
Series : Conservation objectives supporting document

Year : 2014
Title : Mullet/Blacksod Bay Complex SAC (site code: 470) Conservation objectives supporting document- marine habitats V1
Author : NPWS
Series : Conservation objectives supporting document

Other References

Year : 1982
Title : Otter survey of Ireland
Author : Chapman, P.J.; Chapman, L.L.
Series : Unpublished report to Vincent Wildlife Trust

Year : 1982
Title : Eutrophication of waters. Monitoring assessment and control
Author : OECD
Series : OECD, Paris

Year : 1991
Title : The spatial organization of otters (*Lutra lutra*) in Shetland
Author : Kruuk, H.; Moorhouse, A.
Series : J. Zool, 224: 41-57

Year : 1997
Title : The BioMar biotope viewer: a guide to marine habitats, fauna and flora in Britain and Ireland
Author : Picton, B.E.; Costello, M.J.
Series : Environmental Science Unit, Trinity College Dublin

Year : 1999
Title : Diet of otters (*Lutra lutra*) on Inishmore, Aran Islands, west coast of Ireland
Author : Kingston, S.; O'Connell, M.; Fairley, J.S.
Series : Biol & Environ Proc R Ir Acad B 99B:173-182

Year : 2000
Title : Colour in Irish lakes
Author : Free, G.; Allott, N.; Mills, P.; Kennelly, C.; Day, S.
Series : Verhandlungen Internationale Vereinigung für theoretische und angewandte Limnologie. 27: 2620-2623

Year : 2006
Title : Otters - ecology, behaviour and conservation
Author : Kruuk, H.
Series : Oxford University Press

Year :	2006
Title :	The vegetation of Irish machair
Author :	Gaynor, K.
Series :	Biology and Environment: Proceedings of the Royal Irish Academy, vol 106B, No. 3: 311-321
Year :	2006
Title :	A reference-based typology and ecological assessment system for Irish lakes. Preliminary investigations. Final report. Project 2000-FS-1-M1 Ecological assessment of lakes pilot study to establish monitoring methodologies EU (WFD)
Author :	Free, G.; Little, R.; Tierney, D.; Donnelly, K.; Coroni, R.
Series :	EPA, Wexford
Year :	2008
Title :	The phytosociology and conservation value of Irish sand dunes
Author :	Gaynor, K.
Series :	Unpublished PhD thesis, National University of Ireland, Dublin
Year :	2010
Title :	Otter tracking study of Roaringwater Bay
Author :	De Jongh, A.; O'Neill, L.
Series :	Unpublished draft report to NPWS
Year :	2010
Title :	Subtidal benthic investigations in Mullet/Blacksod Bay Complex cSAC (cSAC code: IE000470) and Blacksod Bay/Broadhaven SPA (site code: IE004037)
Author :	Aquafact
Series :	Unpublished report to the Marine Institute and NPWS
Year :	2010
Title :	Reef investigations in Blacksod Bay cSAC (site code: IE000470) Co. Mayo
Author :	Aquafact
Series :	Unpublished report to the Marine Institute and NPWS
Year :	2013
Title :	Conservation of selected legally protected and Red Listed bryophytes in Ireland
Author :	Campbell, C.
Series :	Unpublished Ph.D. Thesis, Trinity College Dublin
Year :	2013
Title :	Benthic survey services framework. Blacksod Bay intertidal surveys 2009 & 2010
Author :	RPS
Series :	Unpublished report to the Marine Institute and NPWS
Year :	in prep.
Title :	Habitats Directive Annex I lake habitats: a working interpretation for the purposes of site-specific conservation objectives and Article 17 reporting
Author :	O Connor, A.
Series :	Unpublished report to NPWS
Year :	in prep.
Title :	Monitoring of hard-water lakes in Ireland using charophytes and other macrophytes
Author :	Roden, C.; Murphy, P.
Series :	Unpublished report to NPWS

Spatial data sources

Year :	Interpolated 2014
Title :	1994 BioMar Survey; 2008 sensitive species survey; 2009, 2010 intertidal and subtidal surveys
GIS Operations :	Polygon feature classes from marine community types base data sub-divided based on interpolation of marine survey data. Expert opinion used to resolve any issues arising
Used For :	1140, 1170, marine community types (maps 3, 5 and 6)
Year :	2005
Title :	OSi Discovery series vector data
GIS Operations :	High Water Mark (HWM) polyline feature class converted into polygon feature class; clipped to SAC boundary. EPA WFD transitional waterbody data erased from extent. Expert opinion used as necessary to resolve any issues arising
Used For :	1160 (map 4)
Year :	2005
Title :	OSi Discovery series vector data
GIS Operations :	High water mark (HWM) and low water mark (LWM) polyline feature classes converted into polygon feature classes and combined; EU Annex I Saltmarsh and Coastal data erased out if present
Used For :	Marine community types base data (map 6)
Year :	Revision 2010
Title :	Saltmarsh Monitoring Project 2007-2008. Version 1
GIS Operations :	QIs selected; clipped to SAC boundary; overlapping regions with Coastal CO data investigated and resolved with expert opinion used
Used For :	1310 (map 7)
Year :	2009
Title :	Coastal Monitoring Project 2004-2006. Version 1
GIS Operations :	QIs selected; clipped to SAC boundary; overlapping regions with Saltmarsh CO data investigated and resolved with expert opinion used
Used For :	2120, 2130, 2150, 21A0 (map 8)
Year :	2013
Title :	Sand Dune Monitoring Project 2011. Version 1
GIS Operations :	QIs selected; clipped to SAC boundary; overlapping regions with Saltmarsh CO data investigated and resolved with expert opinion used
Used For :	2120, 2130, 2150, 21A0 (map 8)
Year :	2008
Title :	OSi 1:5000 IG vector dataset
GIS Operations :	WaterPolygons feature class clipped to the SAC boundary. Expert opinion used to identify Annex I habitat and to resolve any issues arising
Used For :	3150 (map 9)
Year :	2005
Title :	OSi Discovery series vector data
GIS Operations :	Creation of an 80m buffer on marine side of high water mark (HWM); creation of a 10m buffer on terrestrial side of HWM; combination of 80m and 10m HWM buffer datasets; creation of a 10m buffer on terrestrial side of river banks data; creation of 20m buffer applied to canal centreline data. Datasets combined with the derived EPA WFD Waterbodies data. Overlapping regions investigated and resolved; resulting dataset clipped to SAC boundary. Expert opinion used as necessary to resolve any issues arising. Creation of 250m buffer on marine side of HWM to highlight potential commuting points
Used For :	1355 (map 10)
Year :	2014
Title :	NPWS rare and threatened species database
GIS Operations :	Dataset created from spatial references in database records. Expert opinion used as necessary to resolve any issues arising
Used For :	1395 (map 11)

Conservation Objectives for : Mullet/Blacksod Bay Complex SAC [000470]

1140 Mudflats and sandflats not covered by seawater at low tide

To maintain the favourable conservation condition of Mudflats and sandflats not covered by seawater at low tide in Mullet/Blacksod Bay Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	The permanent habitat area is stable or increasing, subject to natural processes. See map 3	Habitat area was estimated using OSI data as 1,428ha
Community distribution	Hectares	Conserve the following community types in a natural condition: Mobile sand with <i>Bathyporeia guilliamsoniana</i> community; Sand with <i>Angulus tenuis</i> and <i>Pygospio elegans</i> community complex. See map 6	Based on intertidal surveys undertaken in 2009 and 2010 (RPS, 2013). See marine supporting document for further information

Conservation Objectives for : Mullet/Blacksod Bay Complex SAC [000470]

1160 Large shallow inlets and bays

To maintain the favourable conservation condition of Large shallow inlets and bays in Mullet/Blacksod Bay Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	The permanent habitat area is stable or increasing, subject to natural processes. See map 4	Habitat area was estimated as 11,169ha using OSI data and the Transitional Water Body area as defined under the Water Framework Directive
Community extent	Hectares	Maintain the extent of the <i>Zostera</i> - and maërl-dominated communities and <i>Serpula vermicularis</i> -dominated community complex, subject to natural processes. See map 6	Based on dive surveys undertaken in 2008 (MERC, 2008). See marine supporting document for further details
Community structure: shoot density	Shoots per m ²	Conserve the high quality of <i>Zostera</i> -dominated community, subject to natural processes	Based on diver observation and underwater viewer (MERC, 2008). See marine supporting document for further details
Community structure	Biological composition	Conserve the high quality of the Maërl dominated community, subject to natural processes	Based on diver observation (MERC, 2008). See marine supporting document for further details
Community structure	Biological composition	Conserve the high quality of the <i>Serpula vermicularis</i> -dominated community complex, subject to natural processes	Based on diver observation (MERC, 2008). See marine supporting document for further details
Community distribution	Hectares	Conserve the following community types in a natural condition: Sand with <i>Angulus tenuis</i> and <i>Pygospio elegans</i> community complex; Sand with <i>Gastrosaccus spinifer</i> community complex; Fine sand with <i>Angulus fabula</i> community complex; Intertidal reef community complex; Sheltered subtidal reef community complex and <i>Laminaria</i> -dominated community complex. See map 6	Based on a BioMar survey in 1994 (Picton and Costello, 1997), a sensitive species survey in 2008 (MERC, 2008) and intertidal and subtidal surveys in 2009 and 2010 (Aquafact, 2010; RPS, 2013). See marine supporting document for further details

Conservation Objectives for : Mullet/Blacksod Bay Complex SAC [000470]

1170 Reefs

To maintain the favourable conservation condition of Reefs in Mullet/Blacksod Bay Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	The permanent habitat area is stable or increasing, subject to natural processes. See map 5	Habitat area estimated as 1,531ha from a 1994 BioMar survey (Picton and Costello, 1996), 2009 intertidal and subtidal reef surveys (Aquafact, 2010) and a walkover survey in 2013
Distribution	Occurrence	The distribution of reefs remains stable, subject to natural processes. See map 5 for mapped distribution	Based on information from a 1994 BioMar survey (Picton and Costello, 1996), 2009 intertidal and subtidal reef surveys (Aquafact, 2010) and a walkover survey in 2013
Community extent	Hectares	Maintain the extent of the <i>Serpula vermicularis</i> -dominated community complex, subject to natural processes. See map 6	Based on dive surveys undertaken in 2008 (MERC, 2008). See marine supporting document for further details
Community structure	Biological composition	Conserve the high quality of the <i>Serpula vermicularis</i> -dominated community complex, subject to natural processes	Based on diver observation (MERC, 2008). See marine supporting document for further details
Community structure	Biological composition	Conserve the following community types in a natural condition: Intertidal reef community complex; Sheltered subtidal reef community; <i>Laminaria</i> -dominated community complex. See map 6	Reef mapping based on information from a 1994 BioMar survey (Picton and Costello, 1996), 2009 intertidal and subtidal reef surveys (Aquafact, 2010) and a walkover survey in 2013. See marine supporting document for further details

Conservation Objectives for : Mullet/Blacksod Bay Complex SAC [000470]

1310 Salicornia and other annuals colonising mud and sand

To maintain the favourable conservation condition of *Salicornia* and other annuals colonising mud and sand in Mullet/Blacksod Bay Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes, including erosion and succession. For sub-site mapped: Elly Harbour - 0.02ha. See map 7	Based on data from Saltmarsh Monitoring Project (SMP) (McCorry and Ryle, 2009). Habitat recorded at one of the four sub-sites surveyed and mapped, giving a total estimated area of 0.02ha. NB further unsurveyed areas maybe present within the site. See coastal habitats supporting document for further details
Habitat distribution	Occurrence	No decline, or change in habitat distribution, subject to natural processes. See map 7 for known distribution	Based on data from McCorry and Ryle (2009). <i>Salicornia</i> is an annual species, so its distribution can vary significantly from year to year. See coastal habitats supporting document for further details
Physical structure: sediment supply	Presence/ absence of physical barriers	Maintain, or where necessary restore, natural circulation of sediments and organic matter, without any physical obstructions	Based on data from McCorry and Ryle (2009). Sediment supply is particularly important for this pioneer saltmarsh community, as the distribution of this habitat depends on accretion rates. See coastal habitats supporting document for further details
Physical structure: creeks and pans	Occurrence	Maintain creek and pan structure, subject to natural processes, including erosion and succession	Based on data from McCorry and Ryle (2009). Creeks deliver sediment throughout saltmarsh system. At Elly Harbour there are signs of modification of the former saltmarsh structure such as drainage channels and old peat-cutting banks. See coastal habitats supporting document for further details
Physical structure: flooding regime	Hectares flooded; frequency	Maintain natural tidal regime	This pioneer saltmarsh community requires regular tidal inundation. See coastal habitats supporting document for further details
Vegetation structure: zonation	Occurrence	Maintain the range of coastal habitats including transitional zones, subject to natural processes including erosion and succession	Based on data from McCorry and Ryle (2009). At Elly Harbour this habitat occurs in mosaic with other saltmarsh habitats. There are also notable successions to brackish and wet grassland vegetation communities around Leam Lough which increases the diversity of the site. See coastal habitats supporting document for further details
Vegetation structure: vegetation height	Centimetres	Maintain structural variation within sward	Based on data from McCorry and Ryle (2009). See coastal habitats supporting document for details
Vegetation structure: vegetation cover	Percentage cover at a representative sample of monitoring stops	Maintain more than 90% of area outside creeks vegetated	Based on data from McCorry and Ryle (2009). See coastal habitats supporting document for details
Vegetation composition: typical species and sub-communities	Percentage cover	Maintain the presence of species-poor communities with typical species including common glasswort (<i>Salicornia europaea</i>), common saltmarsh grass (<i>Puccinellia maritima</i>), sea aster (<i>Aster tripolium</i>) and annual sea-blite (<i>Suaeda maritima</i>)	Based on data from McCorry and Ryle (2009). Turf fucoids were recorded at all sub-sites. Saltmarsh flat-sedge (<i>Blysmus rufus</i>) was recorded at Doolough and Bunnahowen sub-sites and is also a species of local distinctiveness. See coastal habitats supporting document for further details
Vegetation structure: negative indicator species - <i>Spartina anglica</i>	Hectares	There is no record of common cordgrass (<i>Spartina anglica</i>) in the SAC and its establishment should be prevented	Based on data from McCorry and Ryle (2009). See coastal habitats supporting document for further details

Conservation Objectives for : Mullet/Blacksod Bay Complex SAC [000470]

2120 Shifting dunes along the shoreline with *Ammophila arenaria* (white dunes)

To restore the favourable conservation condition of Shifting dunes along the shoreline with *Ammophila arenaria* ('white dunes') in Mullet/Blacksod Bay Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area increasing, subject to natural processes including erosion and succession. Doo Lough - 3.56ha, Cross Lough - 2.61ha, Aghleam - 3.85ha, Dooyork -0.14ha , Srah South - 2.23ha, Srah North - 1.63ha, Leam Lough -2.36ha, Termoncarragh Lough - 2.58ha. See map 8	Based on data from the Coastal Monitoring Project (CMP) (Ryle et al., 2009) and the Sand Dunes Monitoring Project (SDM) (Delaney et al., 2013). Habitat was mapped at eight sub-sites, giving a total estimated area of 18.95ha. Habitat is very difficult to measure in view of its dynamic nature. See coastal habitats supporting document for further details
Habitat distribution	Occurrence	No decline, or change in habitat distribution, subject to natural processes. See map 8 for known distribution	Based on data from Ryle et al. (2009) and Delaney et al. (2013). This habitat was recorded at all eight subsites. The shifting dunes at Aghleam are undergoing natural erosion. At Cross Lough the establishment of an equestrian centre has reduced the extent of this habitat. See coastal habitats supporting document for further details
Physical structure: functionality and sediment supply	Presence/ absence of physical barriers	Maintain the natural circulation of sediment and organic matter, without any physical obstructions	Based on data from Ryle et al. (2009) and Delaney et al. (2013). Dunes are naturally dynamic systems that require continuous supply and circulation of sand. Marram grass (<i>Ammophila arenaria</i>) reproduces vegetatively and requires constant accretion of fresh sand to maintain active growth encouraging further accretion. At Srah North, coastal protection measures in the form of fencing (access control and sand-trapping), marram planting and dune reconstruction using gabions and geotextile fabrics have been put in place. Sand and shingle extraction is known to occur at a number of locations throughout the Mullet sand dune sites. See coastal habitats supporting document for further details
Vegetation structure: zonation	Occurrence	Maintain the range of coastal habitats including transitional zones, subject to natural processes including erosion and succession	Based on data from Gaynor (2008), Ryle et al. (2009) and Delaney et al. (2013). See coastal habitats supporting document for further details
Vegetation composition: plant health of dune grasses	Percentage cover	95% of marram grass (<i>Ammophila arenaria</i>) and/or lyme-grass (<i>Leymus arenarius</i>) should be healthy (i.e. green plant parts above ground and flowering heads present)	Based on data from Ryle et al. (2009) and Delaney et al. (2013). See coastal habitats supporting document for further details
Vegetation composition: typical species and sub-communities	Percentage cover at a representative number of monitoring stops	Maintain the presence of species-poor communities dominated by marram grass (<i>Ammophila arenaria</i>) and/or lyme-grass (<i>Leymus arenarius</i>)	Based on data from Ryle et al. (2009) and Delaney et al. (2013). The flora of mobile dunes in this SAC also includes sea bindweed (<i>Calystegia soldanella</i>), which has a scattered distribution along the north-west coast. See coastal habitats supporting document for further details
Vegetation composition: negative indicator species	Percentage cover	Negative indicator species (including non-natives) to represent less than 5% cover	Based on data from Ryle et al. (2009) and Delaney et al. (2013). Negative indicators include non-native species, species indicative of changes in nutrient status and species not considered characteristic of the habitat. Sea-buckthorn (<i>Hippophae rhamnoides</i>) should be absent or effectively controlled. Creeping thistle (<i>Cirsium arvense</i>) was recorded in mobile dune at Cross Lough. See coastal habitats supporting document for further details

Conservation Objectives for : Mullet/Blacksod Bay Complex SAC [000470]

2130 Fixed coastal dunes with herbaceous vegetation (grey dunes)

To restore the favourable conservation condition of Fixed coastal dunes with herbaceous vegetation ('grey dunes') in Mullet/Blacksod Bay Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes including erosion and succession. For sub-sites mapped: Doo Lough - 46.35ha, Cross Lough - 184.74ha, Aghleam - 292.43ha, Dooyork - 3.73ha, Srah South - 9.10ha, Srah North - 5.99ha, Leam Lough - 170.63ha, Termoncarragh Lough - 224.11ha. See map 8	Based on data from the Coastal Monitoring Project (CMP) (Ryle et al., 2009) and the Sand Dunes Monitoring Project (SDM) (Delaney et al., 2013). This habitat was recorded from all eight sub-sites, giving a total estimated area of 937.07ha. See coastal habitats supporting document for further details
Habitat distribution	Occurrence	No decline, or change in habitat distribution, subject to natural processes. See map 8 for known distribution	Based on data from Ryle et al. (2009) and Delaney et al. (2013). At Doo Lough there is some erosion of this habitat at the seaward edge and poaching by livestock has caused erosion at the fixed dune edge. At Aghleam the extent of the fixed dunes has been reduced due to overgrazing. At Cross Lough the establishment of an equestrian centre on the fixed dunes has led to a modification and reduction in extent of this habitat. See coastal habitats supporting document for further details
Physical structure: functionality and sediment supply	Presence/ absence of physical barriers	Maintain the natural circulation of sediment and organic matter, without any physical obstructions	Based on data from Ryle et al. (2009) and Delaney et al. (2013). Physical barriers can lead to fossilisation or over-stabilisation of dunes, as well as beach starvation resulting in increased rates of erosion. At Srah North, coastal protection measures in the form of fencing (access control and sand-trapping), marram planting and dune reconstruction using gabions and geotextile fabrics have been put in place. Sand and shingle extraction is known to occur a number of locations throughout the Mullet sand dune sites. See coastal habitats supporting document for further details
Vegetation structure: zonation	Occurrence	Maintain the range of coastal habitats including transitional zones, subject to natural processes including erosion and succession	Based on data from Ryle et al. (2009) and Delaney et al. (2013). See coastal habitats supporting document for further details
Vegetation structure: bare ground	Percentage cover	Bare ground should not exceed 10% of fixed dune habitat, subject to natural processes	Based on data from Gaynor (2008) Ryle et al. (2009) and Delaney et al. (2013). See coastal habitats supporting document for further details
Vegetation structure: sward height	Centimetres	Maintain structural variation within sward	Based on data from Ryle et al. (2009) and Delaney et al. (2013). All of the sub-sites are grazed to varying extents. At Doo Lough, sustainable grazing regimes have led to the creation and maintenance of a large area of species-rich turf. The positive impact of grazing is also seen in parts of Aghleam, Leam Lough and Cross Lough. Parts of Doo Lough are overgrazed as are Srah North, Aghleam, Leam Lough, Cross Lough and Termoncarragh. At some areas of Srah South, undergrazing is an issue. See coastal habitats supporting document for further details

Vegetation composition: typical species and sub-communities	Percentage cover at a representative sample of monitoring stops	Maintain range of sub-communities with typical species listed in Delaney et al. (2013)	Based on data from Ryle et al. (2009) and Delaney et al. (2013). See coastal habitats supporting document for further details.
Vegetation composition: negative indicator species (including <i>Hippophae rhamnoides</i>)	Percentage cover	Negative indicator species (including non-natives) to represent less than 5% cover	Based on data from Ryle et al. (2009) and Delaney et al. (2013). Negative indicators include non-native species, species indicative of changes in nutrient status and species not considered characteristic of the habitat. Sea-buckthorn (<i>Hippophae rhamnoides</i>) should be absent or effectively controlled. Common ragwort (<i>Senecio jacobaea</i>), creeping thistle (<i>Cirsium arvense</i>), perennial rye grass (<i>Lolium perenne</i>) and common nettle (<i>Urtica dioica</i>) were regularly recorded in the fixed dune habitat. New Zealand flax (<i>Phormium tenax</i>) was recorded at Leam Lough. See coastal habitats supporting document for further details
Vegetation composition: scrub/trees	Percentage cover	No more than 5% cover or under control	Based on data from Ryle et al. (2009) and Delaney et al. (2013). Within Mullet/Blacksod Complex SAC, the levels of grazing are such that prevents scrub from becoming dominant. See coastal habitats supporting document for further details

Conservation Objectives for : Mullet/Blacksod Bay Complex SAC [000470]

2150 Atlantic decalcified fixed dunes (*Calluno-Ulicetea*)

To maintain the favourable conservation condition of Atlantic decalcified fixed dunes (*Calluno-Ulicetea*) in Mullet/Blacksod Bay Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes including erosion and succession. For sub-sites mapped: Aghleam - 4.08ha, Termoncarragh Lough - 6.20ha. See map 8	Based on data from the Coastal Monitoring Project (CMP) (Ryle et al., 2009) and the Sand Dunes Monitoring Project (SDM) (Delaney et al., 2013). Habitat was mapped at two sub-sites, giving a total estimated area of 10.29ha. However, there is some doubt over the validity of the record of this habitat at Termoncarragh, based on a subsequent site visit by NPWS staff. See coastal habitats supporting document for further details
Habitat distribution	Occurrence	No decline or change in habitat distribution, subject to natural processes. See map 8 for known distribution	Based on data from Ryle et al. (2009) and Delaney et al. (2013). See coastal habitats supporting document for further details
Physical structure: functionality and sediment supply	Presence/ absence of physical barriers	Maintain the natural circulation of sediment and organic matter, without any physical obstructions	Based on data from Ryle et al. (2009) and Delaney et al. (2013). Physical barriers can lead to fossilisation or over-stabilisation of dunes, as well as beach starvation resulting in increased rates of erosion. See coastal habitats supporting document for further details
Vegetation structure: zonation	Occurrence	Maintain the range of coastal habitats including transitional zones, subject to natural processes including erosion and succession	Based on data from Ryle et al. (2009) and Delaney et al. (2013). See coastal habitats supporting document for further details
Vegetation structure: bare ground	Percentage cover	Bare ground should not exceed 10% of fixed dune habitat, subject to natural processes	Based on data from Gaynor (2008), Ryle et al. (2009) and Delaney et al. (2013). See coastal habitats supporting document for further details
Vegetation composition: sward height	Centimeters	Maintain structural variation within sward	Based on data from Gaynor (2008), Ryle et al. (2009) and Delaney et al. (2013). See coastal habitats supporting document for further details
Vegetation composition: typical species and sub-communities	Percentage cover at a representative number of monitoring stops	Maintain range of sub-communities with typical species listed in Delaney et al. (2013)	Based on data from Gaynor (2008), Ryle et al. (2009) and Delaney et al. (2013). Rarer species in this habitat include Autumn lady's tresses (<i>Spiranthes spiralis</i>), lesser butterfly orchid (<i>Platanthera bifolia</i>) and field gentian (<i>Gentianella campestris</i>). See coastal habitats supporting document for further details
Vegetation composition: negative indicator species	Percentage cover	Negative indicator species (including non-natives) to represent less than 5% cover	Based on data from Ryle et al. (2009) and Delaney et al. (2013). Negative indicators include non-native species, species indicative of changes in nutrient status and species not considered characteristic of the habitat. Sea-buckthorn (<i>Hippophae rhamnoides</i>) should be absent or effectively controlled. See coastal habitats supporting document for further details
Vegetation composition: scrub/trees	Percentage cover	No more than 5% cover or under control	Based on data from Ryle et al. (2009) and Delaney et al. (2013). See coastal habitats supporting document for further details

Conservation Objectives for : Mullet/Blacksod Bay Complex SAC [000470]

21A0 Machairs (* in Ireland)

To restore the favourable conservation condition of Machairs in Mullet/Blacksod Bay Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes including erosion and succession. For sub-sites mapped: Doo Lough - 59.52ha, Cross Lough - 60.73ha, Aghleam - 138.59ha, Dooyork - 31.55ha, Srah South - 15.50ha, Srah North - 21.44ha, Leam Lough - 45.55ha, Termoncarragh Lough - 222.76ha. See map 8	Based on data from the Coastal Monitoring Project (CMP) (Ryle et al., 2009) and the Sand Dunes Monitoring Project (SDM) (Delaney et al., 2013). Eight sub-sites were mapped, giving a total estimated area of 595.64ha. See coastal habitats supporting document for further details
Habitat distribution	Occurrence	No decline, subject to natural processes. See map 8 for known distribution	The largest machair site is at Termoncarragh Lough. See coastal habitats supporting document for further details
Physical structure: functionality and sediment supply	Presence/ absence of physical barriers	Maintain the natural circulation of sediment and organic matter, without any physical obstructions	Based on data from Ryle et al. (2009) and Delaney et al. (2013). Physical barriers can lead to fossilisation or over-stabilisation of dunes, as well as beach starvation resulting in increased rates of erosion. See coastal habitats supporting document for further details
Physical structure: hydrological and flooding regime	Presence/absence of water abstraction or drainage works	Maintain natural hydrological regime	Based on data from Ryle et al. (2009), Delaney et al. (2013), Crawford et al. (1996) and Gaynor (2006). See coastal habitats supporting document for further details
Vegetation structure: zonation	Occurrence	Maintain the range of coastal habitats including transitional zones, subject to natural processes including erosion and succession	Based on data from Ryle et al. (2009) and Delaney et al. (2013). See coastal habitats supporting document for further details
Vegetation structure: bare ground	Percentage cover	Bare ground should not exceed 20% of machair habitat, subject to natural processes	Based on data from Ryle et al. (2009) and Delaney et al. (2013). See coastal habitats supporting document for further details
Vegetation structure: sward height	Centimeters	Maintain structural variation within sward	Based on data from Ryle et al. (2009) and Delaney et al. (2013). See coastal habitats supporting document for further details
Vegetation composition: typical species and sub-communities	Percentage cover at a representative number of monitoring stops	Maintain range of sub-communities with typical species listed in Delaney et al. (2013)	Based on data from Gaynor (2008), Ryle et al. (2009) and Delaney et al. (2013). Notable species include the Annex II liverwort species petalwort (<i>Petalophyllum ralfsii</i>). See conservation objective for petalwort (1395) and coastal habitats supporting document for further details
Vegetation composition: negative indicator species	Percentage cover	Negative indicator species (including non-natives) to represent less than 5% cover	Based on data from Ryle et al. (2009) and Delaney et al. (2013). Negative indicators include non-native species, species indicative of changes in nutrient status and species not considered characteristic of the habitat. See coastal habitats supporting document for further details
Vegetation composition: bryophytes	Percentage cover	Should always be at least an occasional component of the vegetation	Based on data from Ryle et al. (2009) and Delaney et al. (2013). See coastal habitats supporting document for further details

Conservation Objectives for : Mullet/Blacksod Bay Complex SAC [000470]

3150 Natural eutrophic lakes with Magnopotamion or Hydrocharition - type vegetation

To maintain the favourable conservation condition of Natural eutrophic lakes with Magnopotamion or Hydrocharition-type vegetation in Mullet/Blacksod Bay Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes. See map 9	The selection of this SAC for habitat 3150 was based on data for Cross Lough. Little is known about the characteristics or ecology of this lake habitat type in Ireland. A working hypothesis of Irish lake habitats defines it as pondweed (<i>Potamogeton</i> spp.) dominated with circumneutral or higher pH, found in low-lying, large, naturally more productive catchments. It is likely that the coastal form inter-grades with or is related to lake habitats 3130 and 3140. Few, if any, Irish lakes are naturally "eutrophic" as defined using total phosphorus, chlorophyll <i>a</i> and water transparency (OECD, 1982). Lakes with habitat 3150 are considered to be associated with mesotrophic conditions as defined by OECD (1982). Two measures of extent should be used: 1. the area of the lake itself and 2. the extent of the vegetation communities/zones that typify the habitat. For further information see NPWS (2013) and the lake habitats supporting document (O Connor, in prep.)
Habitat distribution	Occurrence	No decline, subject to natural processes. See map 9	As noted above, the habitat is considered to occur in Cross Lough. Cross Lough appears to be influenced by the sea, through spray, wind-borne materials and/or subsurface flows, and has a high chloride content. It is likely that this maritime influence increases the lake's productivity
Vegetation composition: typical species	Occurrence	Typical species present, in good condition, and demonstrating typical abundances and distribution	For lists of typical plant species, see Article 17 habitat assessment for 3150 (NPWS, 2013) and the lake habitats supporting document for the purposes of site-specific conservation objectives and Article 17 reporting (O Connor, in prep.). The phytoplankton of Cross Lough is noted as being dominated by cyanobacteria. Pondweeds (<i>Potamogeton praelongus</i> , <i>P. filiformis</i> and <i>P. pectinatus</i>), <i>Littorella uniflora</i> and <i>Myriophyllum spicatum</i> have been recorded in Cross Lough
Vegetation composition: characteristic zonation	Occurrence	All characteristic zones should be present, correctly distributed and in good condition	Very little is known about the characteristic zonation of lake habitat 3150. Like the coastal forms of the hard water lake habitat (3140), coastal 3150 lakes may have compressed zonation (see Roden and Murphy, in prep.)
Vegetation distribution: maximum depth	Metres	Maintain maximum depth of vegetation, subject to natural processes	The maximum depth of vegetation is likely to be specific to the lake shoreline in question. An indicative target of >6m has been developed for hard water lakes (3140) (see Roden and Murphy, 2013; in prep.). Indicative targets will be developed for the other lake habitats with time. Coastal lakes with 3150 may have naturally shallow maximum vegetation depth, as for the coastal form of the hard water habitat. Colonisation tends to be shallower in the machair form of hard water lakes, owing to cloudier water and shallower lake depth (Roden and Murphy, in prep.). For further information see the lake habitats supporting document for the purposes of site-specific conservation objectives and Article 17 reporting (O Connor, in prep.)

Hydrological regime: water level fluctuations	Metres	Maintain appropriate natural hydrological regime necessary to support the habitat	Fluctuations in lake water level are typical in Ireland, but can be amplified by activities such as abstraction and drainage. Increased water level fluctuations can increase wave action, up-root vegetation, increase turbidity, alter the substratum and lead to release of nutrients from the sediment. The hydrological regime of the lakes must be maintained so that the area, distribution and depth of the lake habitat and its constituent/characteristic vegetation zones and communities are not reduced. For further information see the lake habitats supporting document for the purposes of site-specific conservation objectives and Article 17 reporting (O Connor, in prep.)
Lake substratum quality	Various	Maintain appropriate substratum type, extent and chemistry to support the vegetation	Research is required to further characterise the substratum types (particle size and origin) and substratum quality (notably pH, calcium, iron and nutrient concentrations) favoured by each of the five Annex I lake habitats in Ireland. It is likely that habitat 3150 is found on fine (silt and mud), neutral to alkaline, and more nutrient-rich substrates. Substratum particle size is likely to vary with depth and along the shoreline within a single lake. For further information see the lake habitats supporting document for the purposes of site-specific conservation objectives and Article 17 reporting (O Connor, in prep.)
Water quality: transparency	Metres	Maintain appropriate Secchi transparency. There should be no decline in Secchi depth/transparency	Transparency relates to light penetration and, hence, to the depth of colonisation of vegetation. It can be affected by phytoplankton blooms, water colour and turbidity. A target has been set for hard water lakes (3140); however, targets have yet to be established for the remaining lake habitats. Habitat 3150 is associated with clear water; however, coastal examples such as Cross Lough may have naturally lower transparency owing to the influence of the sea producing greater phytoplankton biomass. The OECD fixed boundary system set transparency targets for mesotrophic lakes of 3-6m annual mean Secchi disk depth, and 1.5-3m annual minimum Secchi disk depth. For further information see the lake habitats supporting document for the purposes of site-specific conservation objectives and Article 17 reporting (O Connor, in prep.)
Water quality: nutrients	µg/l phosphorus; mg/l nitrogen	Maintain the concentration of nutrients in the water column at sufficiently low levels to support the habitat and its typical species	As a relatively-productive habitat, mesotrophic and Water Framework Directive 'good' status targets apply. Where a lake has nutrient concentrations that are lower than these targets, there should be no decline within class, i.e. no upward trend in nutrient concentrations. For the lake habitat 3150, annual average TP concentration should be ≤ 20µg/l TP, average annual total ammonia concentration should be ≤ 0.065mg/l N and annual 95th percentile for total ammonia should be ≤ 0.140mg/l N. For further information see the lake habitats supporting document for the purposes of site-specific conservation objectives and Article 17 reporting (O Connor, in prep.). See also the European Communities Environmental Objectives (Surface Waters) Regulations 2009

Water quality: phytoplankton biomass	µg/l Chlorophyll <i>a</i>	Maintain appropriate water quality to support the habitat, including good chlorophyll <i>a</i> status	Mesotrophic and Water Framework Directive 'good' status targets apply to habitat 3150. Where a lake has a chlorophyll <i>a</i> concentration that is lower than this target, there should be no decline within class, i.e. no upward trend in phytoplankton biomass. The average growing season (March-October) chlorophyll <i>a</i> concentration must be <10µg/l. The annual average chlorophyll <i>a</i> concentration should be 2.5-8.0µg/l and the annual peak chlorophyll <i>a</i> concentration should be 8.0-25.0µg/l (OECD, 1982). For further information see the lake habitats supporting document for the purposes of site-specific conservation objectives and Article 17 reporting (O Connor, in prep.). See also The European Communities Environmental Objectives (Surface Waters) Regulations 2009
Water quality: phytoplankton composition	EPA phytoplankton composition metric	Maintain appropriate water quality to support the habitat, including good phytoplankton composition status	The EPA has developed a phytoplankton composition metric for nutrient enrichment of Irish lakes. As for other water quality indicators, habitat 3150 is considered to require WFD good status. For further information see the lake habitats supporting document for the purposes of site-specific conservation objectives and Article 17 reporting (O Connor, in prep.)
Water quality: attached algal biomass	Algal cover and EPA phytobenthos metric	Maintain trace/ absent attached algal biomass (<5% cover) and good phytobenthos status	Nutrient enrichment can favour epiphytic and epipelic algae that can out-compete the submerged vegetation. The cover abundance of attached algae in habitat 3150 should, ideally therefore, be trace/absent (<5% cover). EPA phytobenthos can be used as an indicator of changes in attached algal biomass. As for other water quality indicators, habitat 3150 is considered to require good phytobenthos status. For further information see the lake habitats supporting document for the purposes of site-specific conservation objectives and Article 17 reporting (O Connor, in prep)
Water quality: macrophyte status	EPA macrophyte metric (The Free Index)	Maintain good macrophyte status	Nutrient enrichment can favour more competitive submerged macrophyte species that out-compete the typical and characteristic species for the lake habitat. The EPA monitors macrophyte status for Water Framework Directive purposes using the 'Free Index'. The target for habitat 3150 is good status or an Ecological Quality Ratio (EQR) for lake macrophytes of ≥ 0.68 , as defined in Schedule Five of the European Communities Environmental Objectives (Surface Waters) Regulations. For further information see the lake habitats supporting document for the purposes of site-specific conservation objectives and Article 17 reporting (O Connor, in prep.)
Acidification status	pH units; mg/l-1	Maintain appropriate water and sediment pH, alkalinity and cation concentrations to support the habitat, subject to natural processes	The specific requirements of habitat 3150, in terms of water and sediment pH, alkalinity and cation concentration, have not been fully determined. Acidification is not considered a threat to habitat 3150, however eutrophication can lead to at least temporary increases in pH to toxic levels (>9/9.5 pH units). Maximum pH should be <9.0 pH units, in line with the surface water standards. See The European Communities Environmental Objectives (Surface Waters) Regulations 2009. For further information see the lake habitats supporting document for the purposes of site-specific conservation objectives and Article 17 reporting (O Connor, in prep.)

Water colour	mg/l PtCo	Maintain appropriate water colour to support the habitat	Increased water colour and turbidity decreases light penetration and can reduce the area of available habitat for lake macrophytes, particularly at the lower euphotic depths. The primary source of increased water colour in Ireland is disturbance to peatland. No habitat-specific or national standards for water colour currently exist. Studies have shown median colour concentrations in Irish lakes of 38mg/l PtCo (Free, et al., 2000) and 33mg/l PtCo (Free et al., 2006). It is likely that the water colour in all Irish lake habitats would naturally be <50mg/l PtCo. For further information see the lake habitats supporting document (O Connor, in prep.)
Dissolved organic carbon (DOC)	mg/l	Maintain appropriate organic carbon levels to support the habitat	Dissolved (and particulate) organic carbon (OC) in the water column is linked to water colour and acidification (organic acids). Increasing DOC in water has been documented across the Northern Hemisphere, including afforested peatland catchments in Ireland. Damage and degradation of peatland, leading to decomposition of peat is likely to be the predominant source of OC in Ireland. OC in water promotes decomposition by fungi and bacteria that, in turn, releases dissolved nutrients. The increased biomass of decomposers can also impact directly on the characteristic lake communities through shading, competition, etc. For further information see the lake habitats supporting document for the purposes of site-specific conservation objectives and Article 17 reporting (O Connor, in prep.)
Turbidity	Nephelometric turbidity units/ mg/l SS/ other appropriate units	Maintain appropriate turbidity to support the habitat	Turbidity can significantly affect the quantity and quality of light reaching rooted and attached vegetation and can, therefore, impact on lake habitats. Turbidity can increase as a result of re-suspension of material within the lake, higher loads entering the lake, or eutrophication. The settlement of higher loads of inorganic or organic material on lake vegetation communities may also have impacts on sensitive, delicate species. Turbidity measurement and interpretation is challenging. As a result, it is likely to be difficult to set habitat-specific targets for turbidity in lakes. For further information see the lake habitats supporting document for the purposes of site-specific conservation objectives and Article 17 reporting (O Connor, in prep.)
Fringing habitat : area	Hectares	Maintain the area and condition of fringing habitats necessary to support the natural structure and functioning of habitat 3150	3150 lakes typically have well-developed reedswamp, fen and/or marsh communities around much of their shoreline. Wet woodland would have surrounded much of their shoreline in the past and has survived or re-colonised patches of many 3150 lake shores. These fringing habitats intergrade with and support the structure and functions of the lake habitat. Equally, fringing habitats are dependent on the lake, particularly its water levels, and support wetland communities and species of conservation concern. Many of the fringing wetland habitats support higher invertebrate and plant species richness than the lake habitats themselves. For further information see the lake habitats supporting document for the purposes of site-specific conservation objectives and Article 17 reporting (O Connor, in prep.)

Conservation Objectives for : Mullet/Blacksod Bay Complex SAC [000470]

7230 Alkaline fens

To maintain the favourable conservation condition of Alkaline fens in Mullet/Blacksod Bay Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	The full extent of this habitat within the SAC is currently unknown. The main area described is to the southwest of Termoncarragh Lake; however, other areas are known to occur. This habitat is found in mosaic with, and transitional to, other habitat types including saltmarsh and sand dunes (Crawford et al., 1996; McCorry and Ryle, 2009; Ryle et al., 2009; Delaney et al., 2013; NPWS internal files)
Habitat distribution	Occurrence	No decline, subject to natural processes	Full distribution of this habitat in this SAC is currently unknown- see note above
Hydrological regime	Metres	Appropriate natural hydrological regimes necessary to support the natural structure and functioning of the habitat	Maintenance of groundwater, surface water flows and water table levels within natural ranges is essential for this wetland habitat
Peat formation	Flood duration	Active peat formation, where appropriate	In order for peat to form, water levels need to be slightly below or above the soil surface for c.90% of the time (Jim Ryan, pers. comm.)
Water quality: nutrients	Water chemistry measures	Appropriate water quality to support the natural structure and functioning of the habitat	Fens receive natural levels of nutrients (e.g. iron, magnesium and calcium) from water sources. However, they are generally poor in nitrogen and phosphorus with the latter tending to be the limiting nutrient
Vegetation composition: typical species	Percentage	Maintain vegetation cover of typical species including brown mosses and vascular plants	Species recorded at Termoncarragh include jointed rush (<i>Juncus articulatus</i>), glaucous sedge (<i>Carex flacca</i>), few-flowered spike-rush (<i>Eleocharis quinqueflora</i>), common spike-rush (<i>E. palustris</i>), common butterwort (<i>Pinguicula vulgaris</i>) and lesser clubmoss (Selaginella selaginoides). Marsh helloborine (<i>Epipactis palustris</i>) has also been recorded here. Saline influences are demonstrated by the presence of species such as sea-milkwort (<i>Glaux maritima</i>) and sea arrowgrass (<i>Triglochin maritimum</i>) (NPWS internal files)
Vegetation composition: trees and shrubs	Percentage cover in local vicinity	Cover of scattered native trees and shrubs less than 10%	Scrub and trees will tend to invade if fen conditions become drier. Attribute and target based on alkaline fen conservation assessment criteria in Perrin et al. (2014)
Physical structure: disturbed bare ground	Percentage cover at a representative number of monitoring stops and in local vicinity	Cover of disturbed bare ground less than 10%. Where tufa is present, disturbed bare ground less than 1%	While grazing may be appropriate in this habitat, excessive area of disturbed bare ground may develop due to unsuitable grazing regimes. Attribute and target based on alkaline fen conservation assessment criteria in Perrin et al. (2014)
Physical structure: drainage	Percentage cover in local vicinity	Area showing signs of drainage as a result of drainage ditches or heavy trampling less than 10%	Attribute and target based on alkaline fen conservation assessment criteria in Perrin et al. (2014)

Conservation Objectives for : Mullet/Blacksod Bay Complex SAC [000470]

1355 Otter *Lutra lutra*

To maintain the favourable conservation condition of Otter in Mullet/Blacksod Bay Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Distribution	Percentage positive survey sites	No significant decline	Measure based on standard otter survey technique. FCS target, based on 1980/81 survey findings, is 88% in SACs. Current range estimated at 93.6% (Reid et al., 2013)
Extent of terrestrial habitat	Hectares	No significant decline. Area mapped and calculated as 168.7ha	No field survey. Areas mapped to include 10m terrestrial buffer along shoreline (above HWM and along river banks) identified as critical for otters (NPWS, 2007)
Extent of marine habitat	Hectares	No significant decline. Area mapped and calculated as 929.6ha	No field survey. Area mapped based on evidence that otters tend to forage within 80m of the shoreline (HWM) (NPWS, 2007; Kruuk, 2006)
Extent of freshwater (river) habitat	Kilometres	No significant decline. Length mapped and calculated as 31.0km	No field survey. River length calculated on the basis that otters will utilise freshwater habitats from estuary to headwaters (Chapman and Chapman, 1982)
Extent of freshwater (lake/lagoon) habitat	Hectares	No significant decline. Area mapped and calculated as 87.7ha	No field survey. Area mapped based on evidence that otters tend to forage within 80m of the shoreline (NPWS, 2007)
Couching sites and holts	Number	No significant decline	Otters need lying up areas throughout their territory where they are secure from disturbance (Kruuk, 2006; Kruuk and Moorhouse, 1991)
Fish biomass available	Kilograms	No significant decline	Broad diet that varies locally and seasonally, but dominated by fish, in particular salmonids, eels and sticklebacks in freshwater (Bailey and Rochford, 2006; Reid et al., 2013) and wrasse and rockling in coastal waters (Kingston et al., 1999)
Barriers to connectivity	Number	No significant increase. For guidance, see map 10	Otters will regularly commute across stretches of open water up to 500m e.g. between the mainland and an island; between two islands; across an estuary (De Jongh and O'Neill, 2010). It is important that such commuting routes are not obstructed

Conservation Objectives for : Mullet/Blacksod Bay Complex SAC [000470]

1395 Petalwort *Petalophyllum ralfsii*


To maintain the favourable conservation condition of Petalwort in Mullet/Blacksod Bay Complex SAC, which is defined by the following list of attributes and targets:


Attribute	Measure	Target	Notes
Distribution of populations	Number and geographical spread of populations	No decline. See map 11 for the two recorded locations	There are currently two known sub-populations in the SAC: (8a) Doolough machair and (8b) Dooyork machair. The population at Dooyork machair has not been seen since 1998. Data from unpublished NPWS surveys
Population size	Number of individuals	No decline. Current known population at Doolough machair estimated c.33 thalli; at Dooyork machair estimated to be c.3 thalli. Total c.36 thalli	Counts of thalli: for Doolough machair- from mean number of thalli recorded by Lockhart in 1999 (20 thalli); Holyoak in 1999 (77 thalli) and Lockhart in 2006 (3 thalli) = 33.33 thalli; for Dooyork machair- from mean number of thalli recorded by Lockhart in 1998 (6 thalli) and Lockhart in 1999 (0 thalli) = 3 thalli. Total number of thalli = 36.33 (c.36 thalli)
Area of suitable habitat	Hectares	No decline. Area of suitable habitat at Doolough machair and Dooyork machair currently unknown, but thought to be very small. Total estimated to be c.0.0005ha	The extent of suitable habitat at Doolough machair and at Dooyork machair has not been accurately measured using GPS, but is known to be very small (c.0.5m ² and 4m ² respectively)
Hydrological conditions: soil moisture	Occurrence of damp soil conditions	Maintain hydrological conditions so that substrate is kept moist and damp throughout the year, but not subject to prolonged inundation by flooding in winter	<i>Petalophyllum ralfsii</i> grows in damp sand. Based on Campbell (2013)
Vegetation: open structure	Height and percentage cover of vegetation	Maintain and enhance open, low vegetation, with a high percentage cover of bryophytes (small acrocarps and liverwort turf) and bare ground	<i>Petalophyllum ralfsii</i> grows in compacted, sandy ground, maintained by rabbit (<i>Oryctolagus cuniculus</i>) and cattle grazing. Recorded at Doolough machair on tightly cropped turf on sides of low sandhills at the highest part of the plain, above a flushed calcareous slope to the south-east (by Holyoak in 1999; Lockhart in 2006); recorded at Dooyork machair by Lockhart in 1998 on damp flats between low sandhills, which is unusual for <i>P. ralfsii</i> ; not refound during a visit by Lockhart in 1999









Legend

 Mullet/Blacksod Bay Complex SAC 000470






Legend


-  Mullet/Blacksod Bay Complex SAC 000470
-  West Connacht Coast SAC 002998
-  Blacksod Bay/Broadhaven SPA 004037
-  Termoncarragh Lake and Annagh Machair SPA 004093
-  Mullet Peninsula SPA 004227
-  OSi Discovery Series County Boundaries





Legend

- Mullet/Blacksod Bay Complex SAC 000470
- 1140 Mudflats and sandflats not covered by sea water at low tide
- OSi Discovery Series County Boundaries



Legend


- Mullet/Blacksod Bay Complex SAC 000470
- 1160 Large shallow inlets and bays
- OSi Discovery Series County Boundaries

Map to be read in conjunction with the NPWS Conservation Objectives Document.



Legend

- Mullet/Blacksod Bay Complex SAC 000470
- 1170 Reefs
- OSi Discovery Series County Boundaries




Legend

- Mullet/Blacksod Bay Complex SAC 000470
- OSi Discovery Series County Boundaries

Marine Community Types

- Fine sand with *Angulus fabula* community complex
- Intertidal reef community complex
- Laminaria*-dominated community complex
- Maerl-dominated community
- Mobile sand with *Bathyporeia guilliamsoniana* community
- Sand with *Angulus tenuis* and *Pygospio elegans* community complex
- Sand with *Gastrosaccus spinifer* community complex
- Serpula vermicularis*-dominated community complex
- Sheltered subtidal reef community complex
- Shingle
- Zostera*-dominated community



Legend

- Mullet/Blacksod Bay Complex SAC 000470
- OSi Discovery Series County Boundaries
- SMP: 0113 Saltmarsh Monitoring Project Site Codes


Saltmarsh Habitats


Qualifying Interests

- Salicornia and other annuals colonising mud and sand, *Salicornia* and other annuals colonising mud and sand

Non Qualifying Interests


- 1330 Atlantic salt meadows (*Glauco-Puccinellietalia maritimae*)
- 1330 / 1410 Atlantic salt meadows (*Glauco-Puccinellietalia maritimae*) / Mediterranean salt meadows (*Juncetalia maritimi*)
- 1410 Mediterranean salt meadows (*Juncetalia maritimi*)
- Potential 1330 Atlantic salt meadows (*Glauco-Puccinellietalia maritimae*)
- Potential 1330 / 1410 Atlantic salt meadows (*Glauco-Puccinellietalia maritimae*) / Mediterranean salt meadows (*Juncetalia maritimi*)
- Potential 1410 Mediterranean salt meadows (*Juncetalia maritimi*)






Legend

- Mullet/Blacksod Bay Complex SAC 000470
- 3150 Natural eutrophic lakes with Magnopotamion or Hydrocharition-type vegetation
- OSi Discovery Series County Boundaries



Legend

- Mullet/Blacksod Bay Complex SAC 000470
- 1355 Otter - *Lutra lutra*
- OSi Discovery Series County Boundaries



Legend

- Mullet/Blacksod Bay Complex SAC 000470
- ★ 1395 Petalwort - *Petalophyllum ralfsii*
- OSi Discovery Series County Boundaries



Conservation objectives for Brackloon Woods SAC [000471]

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Objective: To maintain or restore the favourable conservation condition of the Annex I habitat(s) and/or the Annex II species for which the SAC has been selected:

Code	Description
------	-------------

91A0	Old sessile oak woods with <i>Ilex</i> and <i>Blechnum</i> in the British Isles
------	---

* denotes a priority habitat



Citation: NPWS (2020) Conservation objectives for Brackloon Woods SAC [000471]. Generic Version
7.0. Department of Culture, Heritage and the Gaeltacht.

National Parks and Wildlife Service

Conservation Objectives Series

Broadhaven Bay SAC 000472



An Roinn
Ealaíon, Oidhreachta agus Gaeltachta

Department of
Arts, Heritage and the Gaeltacht



**National Parks and Wildlife Service,
Department of Arts, Heritage and the Gaeltacht,
7 Ely Place, Dublin 2, Ireland.
Web: www.npws.ie
E-mail: nature.conservation@ahg.gov.ie**

Citation:

**NPWS (201) Conservation Objectives: Broadhaven Bay SAC 000472. Version 1.
National Parks and Wildlife Service, Department of Arts, Heritage and the
Gaeltacht.**

**Series Editor: Rebecca Jeffrey
ISSN 2009-4086**

Introduction

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

** indicates a priority habitat under the Habitats Directive*

000472	Broadhaven Bay SAC
--------	--------------------

1140	Mudflats and sandflats not covered by seawater at low tide
1160	Large shallow inlets and bays
1170	Reefs
1330	Atlantic salt meadows (<i>Glauco-Puccinellietalia maritimae</i>)
8330	Submerged or partially submerged sea caves

Please note that this SAC overlaps with Blacksod Bay/Broadhaven Bay SPA (004037). It is adjacent to Mullet/Blacksod Bay Complex SAC (000470), Glenamoy Bog Complex SAC (000500), Erris Head SAC (001501) and West Connaught Coast SAC (002998). See map 2. The conservation objectives for this site should be used in conjunction with those for overlapping and adjacent sites as appropriate.

Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

NPWS Documents

Year :	2007
Title :	Surveys of sensitive subtidal benthic communities
Author :	MERC
Series :	Unpublished report to NPWS
Year :	2009
Title :	Saltmarsh monitoring project 2007-2008
Author :	McCorry, M.; Ryle, T.
Series :	Unpublished report to NPWS
Year :	2014
Title :	Broadhaven Bay SAC (site code: 472) Conservation objectives supporting document- coastal habitats V1
Author :	NPWS
Series :	Conservation objectives supporting document
Year :	2014
Title :	Broadhaven Bay SAC (site code: 472) Conservation objectives supporting document- marine habitats V1
Author :	NPWS
Series :	Conservation objectives supporting document

Other References

Year :	1997
Title :	The BioMar biotope viewer: a guide to marine habitats, fauna and flora in Britain and Ireland
Author :	Picton, B.E.; Costello, M.J.
Series :	Environmental Science Unit, Trinity College Dublin
Year :	1998
Title :	The saltmarshes of Ireland: an inventory and account of their geographical variation
Author :	Curtis, T.G.F.; Sheehy Skeffington, M.J.
Series :	Biology and Environment, Proceedings of the Royal Irish Academy 98B: 87-104
Year :	2010
Title :	Subtidal benthic investigations in Broadhaven Bay cSAC (site code: IE000472) and Blacksod/Broadhaven SPA (site code: IE004037)
Author :	Aquafact
Series :	Unpublished report to the Marine Institute and NPWS
Year :	2010
Title :	Reef investigations in Broadhaven Bay cSAC (site code: IE000472) Co. Mayo
Author :	Aquafact
Series :	Unpublished report to the Marine Institute and NPWS
Year :	2013
Title :	Benthic survey services framework. Blacksod Bay intertidal surveys 2009 & 2010
Author :	RPS
Series :	Unpublished report to the Marine Institute and NPWS

Spatial data sources

Year :	Interpolated 2014
Title :	1994 BioMar Survey; 2009 and 2010 intertidal surveys; 2007 and 2010 subtidal surveys
GIS Operations :	Polygon feature classes from marine community types base data sub-divided based on interpolation of marine survey data. Expert opinion used as necessary to resolve any issues arising
Used For :	1140, 1170, marine community types (maps 3, 5 and 6)
Year :	2005
Title :	OSi Discovery series vector data
GIS Operations :	High Water Mark (HWM) polyline feature class converted into polygon feature class; clipped to SAC boundary. EPA WFD transitional waterbody data erased from extent. Expert opinion used as necessary to resolve any issues arising
Used For :	1160 (map 4)
Year :	Derived 2014
Title :	Coast of Ireland Oblique Imagery Survey 2003
GIS Operations :	Point dataset created from visual inspection of survey
Used For :	8330 (map 4)
Year :	2005
Title :	OSi Discovery series vector data
GIS Operations :	High water mark (HWM) and low water mark (LWM) polyline feature classes converted into polygon feature classes and combined; EU Annex I Saltmarsh and Coastal data erased out if present
Used For :	Marine community types base data (map 6)
Year :	Revision 2010
Title :	Saltmarsh Monitoring Project 2007-2008. Version 1
GIS Operations :	QIs selected; clipped to SAC boundary; overlapping regions with Coastal CO data investigated and resolved with expert opinion used
Used For :	1330 (map 7)

Conservation Objectives for : Broadhaven Bay SAC [000472]

1140 Mudflats and sandflats not covered by seawater at low tide

To maintain the favourable conservation condition of Mudflats and sandflats not covered by seawater at low tide in Broadhaven Bay SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	The permanent habitat area is stable or increasing, subject to natural processes. See map 3	Habitat area was estimated using OSi data as 495ha
Community distribution	Hectares	Conserve the following community types in a natural condition: Coarse sediment to sandy mud with <i>Pygospio elegans</i> community complex; Sand with <i>Angulus tenuis</i> community complex. See map 6	Based on intertidal surveys in 2009 and 2010 (RPS, 2013). See marine supporting document for further information

Conservation Objectives for : Broadhaven Bay SAC [000472]

1160 Large shallow inlets and bays

To maintain the favourable conservation condition of Large shallow inlets and bays in Broadhaven Bay SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	The permanent habitat area is stable or increasing, subject to natural processes. See map 4	Habitat area was estimated as 8,674ha using OSI data and the Transitional Water Body area as defined under the Water Framework Directive
Community extent	Hectares	Maintain the extent of the <i>Zostera</i> -dominated community, subject to natural processes. See map 6	Based on a dive survey undertaken in 2007 (MERC, 2007). See marine supporting document for further details
Community structure: <i>Zostera</i> density	Shoots per m ²	Conserve the high quality of <i>Zostera</i> -dominated community, subject to natural processes	Based on diver observation and underwater viewer (MERC, 2007). See marine supporting document for further details
Community distribution	Hectares	Conserve the following community types in a natural condition: Coarse sediment to sandy mud with <i>Pygospio elegans</i> community complex; Sand with <i>Angulus tenuis</i> community complex; Sand to coarse sediment with crustaceans and <i>Polyophthalmus pictus</i> community complex; Subtidal sand with polychaetes community complex; Furoid-dominated reef community complex; Subtidal reef community complex. See map 6	Based on a BioMar survey in 1994 (Picton and Costello, 1997), a sensitive species survey in 2007 (MERC, 2007), intertidal and subtidal surveys in 2009 and 2010 (Aquafact, 2010; RPS, 2013) and an intertidal walkover undertaken in 2013. See marine supporting document for further information

1170 Reefs

To maintain the favourable conservation condition of Reefs in Broadhaven Bay SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	The permanent habitat area is stable or increasing, subject to natural processes. See map 5	Habitat area estimated as 1,103ha from 1994 BioMar survey (Picton and Costello, 1997) and intertidal and subtidal reef surveys in 2009 (Aquafact, 2010)
Distribution	Occurrence	The distribution of reefs remains stable, subject to natural processes. See map 5 for mapped distribution	Based on information from 1994 BioMar survey (Picton and Costello, 1997) and intertidal and subtidal reef surveys in 2009 (Aquafact, 2010). See marine supporting document for further details
Community structure	Biological composition	Conserve the following community types in a natural condition: Furoid-dominated reef community complex; Subtidal reef community complex. See map 6	Reef mapping based on information from 1994 BioMar survey (Picton and Costello, 1997) and intertidal and subtidal reef surveys in 2009 (Aquafact, 2010) and an intertidal walkover undertaken in 2013. See marine supporting document for further details

Conservation Objectives for : Broadhaven Bay SAC [000472]

1330 Atlantic salt meadows (*Glauco-Puccinellietalia maritimae*)

To maintain the favourable conservation condition of Atlantic salt meadows (*Glauco-Puccinellietalia maritimae*) in Broadhaven Bay SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes, including erosion and succession. Potential area mapped: 13.46ha. See map 7	Based on data from Saltmarsh Monitoring Project (SMP) (McCorry and Ryle, 2009). Although the site was not physically surveyed, 13.46ha of potential habitat were identified through an examination of orthophotographs (OSi 2005 Series). NB further unsurveyed areas maybe present within the site. See coastal habitats supporting document for further details
Habitat distribution	Occurrence	No decline or change in habitat distribution, subject to natural processes. See map 7 for potential distribution	Based on data from Curtis and Sheehy Skeffington (1998) McCorry and Ryle (2009). Two saltmarsh sites are known to occur at Tallagh and Barnatra. Both sites are in sheltered locations in the inner part of the bay. See coastal habitats supporting document for further details
Physical structure: sediment supply	Presence/ absence of physical barriers	Maintain natural circulation of sediments and organic matter, without any physical obstructions	Based on data from NPWS internal files. The natural functioning of the saltmarsh at Tallagh has been affected by a long established road. Some reclamation works have also been carried out at this site in the past. See coastal habitats supporting document for further details
Physical structure: creeks and pans	Occurrence	Maintain/restore creek and pan structure, subject to natural processes, including erosion and succession	Pools and channels occur at the saltmarshes at Barnatra and Tallagh. See coastal habitats supporting document for further details
Physical structure: flooding regime	Hectares flooded; frequency	Maintain natural tidal regime	See coastal habitats supporting document for further details
Vegetation structure: zonation	Occurrence	Maintain range of coastal habitats including transitional zones, subject to natural processes including erosion and succession	Based on data from NPWS internal files. At both Barnatra and Tallagh, the saltmarshes merge into wet pasture and bog habitats. See coastal habitats supporting document for further details
Vegetation structure: vegetation height	Centimetres	Maintain structural variation within sward	Based on data from NPWS internal files. The structure of the saltmarshes at Barnatra and Tallagh has been affected by grazing, which is heavy in places. See coastal habitats supporting document for further details
Vegetation structure: vegetation cover	Percentage cover at a representative number of monitoring stops	Maintain more than 90% area outside creeks vegetated	See coastal habitats supporting document for further details
Vegetation composition: typical species and sub-communities	Percentage cover at a representative number of monitoring stops	Maintain range of sub-communities with typical species listed in SMP (McCorry and Ryle, 2009)	Based on data from NPWS internal files. Turf fucoids are present. See coastal habitats supporting document for further details
Vegetation structure: negative indicator species - <i>Spartina anglica</i>	Hectares	There is no record of common cordgrass (<i>Spartina anglica</i>) in this SAC. Prevent establishment of cordgrass	<i>Spartina</i> is thought to be absent from the SAC. See coastal habitats supporting document for further details

Conservation Objectives for : Broadhaven Bay SAC [000472]


8330 Submerged or partially submerged sea caves


To maintain the favourable conservation condition of Submerged or partially submerged sea caves in Broadhaven Bay SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Distribution	Occurrence	The distribution of sea caves remains stable, subject to natural processes. See map 4 for known caves	Sea cave distribution at this site was derived from an oblique aerial survey and therefore only detects the presence of sea caves visible intertidally in the flight path. NB other sea caves may occur within the SAC
Community structure	Biological composition	Human activities should occur at levels that do not adversely affect the ecology of sea caves in this SAC	




Legend


 Broadhaven Bay SAC 000472



Legend


- Broadhaven Bay SAC 000472
- Mullet/Blacksod Bay Complex SAC 000470
- Glenamoy Bog Complex SAC 000500
- Erris Head SAC 001501
- West Connacht Coast SAC 002998
- Blacksod Bay/Broadhaven SPA 004037
- OSi Discovery Series County Boundaries






Legend

- Broadhaven Bay SAC 000472
- 1140 Mudflats and sandflats not covered by sea water at low tide
- OSi Discovery Series County Boundaries




Legend

- Broadhaven Bay SAC 000472
- 1160 Large shallow inlets and bays
- 8330 Submerged or partially submerged sea caves
- OSi Discovery Series County Boundaries



Legend

- Broadhaven Bay SAC 000472
- 1170 Reefs
- OSi Discovery Series County Boundaries




Legend


- Broadhaven Bay SAC 000472
- OSi Discovery Series County Boundaries

Marine Community Types

- Coarse sediment to sandy mud with *Pygospio elegans* community complex
- Furoid-dominated reef community complex
- Sand to coarse sediment with crustaceans and *Polyophthalmus pictus* community complex
- Sand with *Angulus tenuis* community complex
- Shingle
- Subtidal reef community complex
- Subtidal sand with polychaetes community complex
- Zostera*-dominated community



Legend

 Broadhaven Bay SAC 000472

 OSi Discovery Series County Boundaries

Saltmarsh Habitats

 Potential 1330 / 1410 Atlantic salt meadows (*Glauco-Puccinellietalia maritima*) / Mediterranean salt meadows (*Juncetalia maritimi*)



National Parks and Wildlife Service

Conservation Objectives Series

Ballymaglancy Cave, Cong SAC 000474



An Roinn Cultúir,
Oidhreacht agus Gaeltachta
Department of Culture,
Heritage and the Gaeltacht

**National Parks and Wildlife Service,
Department of Culture, Heritage and the Gaeltacht,
90 King Street North, Dublin 7, D07 N7CV, Ireland.**

**Web: www.npws.ie
E-mail: nature.conservation@chg.gov.ie**

Citation:

**NPWS (2018) Conservation Objectives: Ballymaglancy Cave, Cong SAC 000474.
Version 1. National Parks and Wildlife Service, Department of Culture, Heritage
and the Gaeltacht.**

**Series Editor: Rebecca Jeffrey
ISSN 2009-4086**

Introduction

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

** indicates a priority habitat under the Habitats Directive*

000474	Ballymaglancy Cave, Cong SAC
1303	Lesser Horseshoe Bat <i>Rhinolophus hipposideros</i>
8310	Caves not open to the public

Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

NPWS Documents

Year : 2018
Title : Conservation objectives supporting document – lesser horseshoe bat (*Rhinolophus hipposideros*)
Author : NPWS
Series : Conservation objectives supporting document

Other References

Year : 2007
Title : Protecting and managing underground sites for bats
Author : Mitchell-Jones, A.J.; Bihari, Z.; Masing, M.; Rodrigues, L.
Series : EUROBATS Publication Series No. 2

Year : 2008
Title : The lesser horseshoe bat conservation handbook
Author : Schofield, H.W.
Series : The Vincent Wildlife Trust

Year : 2009
Title : Importance of night roosts for bat conservation: roosting behaviour of the lesser horseshoe bat *Rhinolophus hipposideros*
Author : Knight, T.; Jones, G.
Series : Endangered Species Research, 8: 79-86

Spatial data sources

Year : 2018
Title : NPWS lesser horseshoe bat database
GIS Operations : Roost identified, clipped to SAC boundary. Expert opinion used as necessary to resolve any issues arising
Used For : 8310, 1303 (map 2)

Year : 2007
Title : Forest Inventory and Planning System (FIPS)
GIS Operations : Dataset clipped to 2.5km buffer centred on roost location
Used For : 1303 (map 2)

Conservation Objectives for : Ballymaglancy Cave, Cong SAC [000474]

8310 Caves not open to the public

Caves not open to the public (8310) is integrally linked to lesser horseshoe bat (*Rhinolophus hipposideros*) (1303) as part of the habitat for the species; therefore, a separate conservation objective has not been set for the habitat in Ballymaglancy Cave, Cong SAC. See map 2. See the conservation objectives supporting document for lesser horseshoe bat (NPWS, 2018) for further details


Attribute	Measure	Target	Notes
------------------	----------------	---------------	--------------


Conservation Objectives for : Ballymaglancy Cave, Cong SAC [000474]

1303 Lesser Horseshoe Bat *Rhinolophus hipposideros*

To restore the favourable conservation condition of Lesser Horseshoe Bat in Ballymaglancy Cave, Cong SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Population per roost	Number	Minimum number of 50 bats for the winter roost (roost id. 245 in NPWS database). See map 2	A figure of 100 bats for summer roosts and 50 bats for winter roosts was set as a minimum qualifying standard (MQS) when SACs were being selected for lesser horseshoe bat (<i>Rhinolophus hipposideros</i>). NPWS conduct annual counts at each qualifying roost. Qualified means from the 2013-2017 winter data have been calculated whereby the year with the highest maximum count and the year with the lowest maximum count were removed and the mean of the remaining years was calculated. This mean is usually set as the target figure for the roost. However, in the case of the winter roost (roost id. 245 in NPWS database) in Ballymaglancy Cave, Cong SAC, where a mean of 28 bats was recorded (2013-2017), the target is instead set at the MQS of 50 bats. See the conservation objectives supporting document for lesser horseshoe bat (NPWS, 2018) for further information on all attributes and targets
Winter roosts	Condition	No decline	Ballymaglancy Cave, Cong SAC has been selected for lesser horseshoe bat because of the presence of one internationally important winter roost (roost id. 245 in NPWS database). Damage or disturbance to the roost or to the habitat immediately surrounding it will lead to a decline in its condition (Mitchell-Jones et al., 2007)
Auxiliary roosts	Number and condition	No decline	Lesser horseshoe bat populations will use a variety of roosts during the year besides the main summer maternity and winter hibernation roosts. Such additional roosts within the SAC may be important as night roosts, satellite roosts, etc. Night roosts are also considered an integral part of core foraging areas and require protection (Knight and Jones, 2009). In addition, in response to weather conditions for example, bats may use different seasonal roosts from year to year; this is particularly noticeable in winter. A database of all known lesser horseshoe bat roosts is available on the National Biodiversity Data Centre website. NB further unrecorded roosts may also be present within this SAC
Extent of potential foraging habitat	Hectares	No significant decline within 2.5km of qualifying roost	Lesser horseshoe bats normally forage in woodlands/scrub within 2.5km of their roosts (Schofield, 2008). See map 2 which shows a 2.5km zone around the above roost and identifies potential foraging grounds
Linear features	Kilometres	No significant loss within 2.5km of qualifying roost. See map 2	This species follows commuting routes from its roost to its foraging grounds. Lesser horseshoe bats will not cross open ground. Consequently, linear features such as hedgerows, treelines and stone walls provide vital connectivity for this species within 2.5km around each roost (Schofield, 2008)
Light pollution	Lux	No significant increase in artificial light intensity adjacent to named roost or along commuting routes within 2.5km of the roost. See map 2	Lesser horseshoe bats are very sensitive to light pollution and will avoid brightly lit areas. Inappropriate lighting around roosts may cause abandonment; lighting along commuting routes may cause preferred foraging areas to be abandoned, thus increasing energetic costs for bats (Schofield, 2008)



Legend
 Ballymaglancy Cave, Cong SAC 000474

 An Roinn Cultúir,
 Oidhreacht agus Gaeltachta
 Department of Culture,
 Heritage and the Gaeltacht

**MAP 1:
 BALLYMAGLANCY CAVE, CONG SAC
 CONSERVATION OBJECTIVES
 SAC DESIGNATION**
 Map to be read in conjunction with the NPWS Conservation Objectives Document.


**SITE CODE:
 SAC 000474; version 3.01. CO. GALWAY, CO.MAYO**

0 25 50 75 100 Meters

The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
 Ordnance Survey of Ireland Licence No EN 0059216. © Ordnance Survey of Ireland Government of Ireland

Níl sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbheithnithe a déanamh ar theorainneacha na gceantar comharthaithe. Suirbhéarachta Ordonáis na hÉireann Ceadúnas Uimh EN 0059216. © Suirbhéarachta Ordonáis na hÉireann Rialtas na hÉireann


**Map Version 1
 Date: June 2018**



Legend

- Ballymaglancy Cave, Cong SAC 000474
- OS Discovery Series County Boundary
- 8310 Caves not open to the public
- 1303 Lesser Horseshoe Bat *Rhinolophus hipposideros***
- Roost Location
- Roost ID 245 Foraging Range
- Potential Foraging Grounds



Conservation objectives for Carrowkeel Turlough SAC [000475]

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Objective: To maintain or restore the favourable conservation condition of the Annex I habitat(s) and/or the Annex II species for which the SAC has been selected:

Code	Description
------	-------------

3180	Turloughs*
------	------------

* denotes a priority habitat



Citation: NPWS (2020) Conservation objectives for Carrowkeel Turlough SAC [000475]. Generic Version 7.0. Department of Culture, Heritage and the Gaeltacht.

National Parks and Wildlife Service

Conservation Objectives Series

Carrowmore Lake Complex SAC 000476



An Roinn Ealaíon, Oidhreachta,
Gnóthaí Réigiúnacha, Tuaithe agus Gaeltachta

Department of Arts, Heritage,
Regional, Rural and Gaeltacht Affairs



**National Parks and Wildlife Service,
Department of Arts, Heritage, Regional, Rural and Gaeltacht Affairs,
7 Ely Place, Dublin 2, Ireland.
Web: www.npws.ie
E-mail: nature.conservation@ahg.gov.ie**

Citation:

**NPWS (2017) Conservation Objectives: Carrowmore Lake Complex SAC 000476.
Version 1. National Parks and Wildlife Service, Department of Arts, Heritage,
Regional, Rural and Gaeltacht Affairs.**

**Series Editor: Rebecca Jeffrey
ISSN 2009-4086**

Introduction

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

** indicates a priority habitat under the Habitats Directive*

000476	Carrowmore Lake Complex SAC
1393	Slender Green Feather-moss <i>Drepanocladus vernicosus</i>
1528	Marsh Saxifrage <i>Saxifraga hirculus</i>
7130	Blanket bogs (* if active bog)
7150	Depressions on peat substrates of the Rhynchosporion

Please note that this SAC overlaps with Carrowmore Lake SPA (004052) and adjoins Slieve Fyagh Bog SAC (000542). See map 2. The conservation objectives for this site should be used in conjunction with those for the overlapping and adjoining sites as appropriate.

Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

NPWS Documents

Year :	1987
Title :	A survey to locate blanket bogs of scientific interest in County Mayo. Part I
Author :	Foss, P.; McGee, E.
Series :	A report commissioned by the Wildlife Service
Year :	1989
Title :	Survey to locate blanket bogs of scientific interest in Mayo. Part II
Author :	Douglas, C.; Garvey, L.; Kelly, L.; O'Sullivan, A.; Van Doorsleer, L.
Series :	A report commissioned by the Wildlife Service
Year :	2012
Title :	Ireland Red List No. 8: Bryophytes
Author :	Lockhart, N.; Hodgetts, N.; Holyoak, D.
Series :	Ireland Red List series, NPWS
Year :	2014
Title :	Guidelines for a national survey and conservation assessment of upland vegetation and habitats in Ireland, Version 2.0
Author :	Perrin, P.M.; Barron, S.J.; Roche, J.R.; O'Hanrahan, B.
Series :	Irish Wildlife Manual No. 79
Year :	2015
Title :	Monitoring recommendations for Marsh Saxifrage (<i>Saxifraga hirculus</i> L.) in the Republic of Ireland
Author :	Muldoon, C.S.; Waldren, S.; Lynn, D.
Series :	Irish Wildlife Manual No. 88
Year :	2015
Title :	Monitoring methods for <i>Hamatocaulis vernicosus</i> (Mitt.) Hedenäs (Slender green feather-moss) in the Republic of Ireland
Author :	Campbell, C.; Hodgetts, N.; Lockhart, N.
Series :	Irish Wildlife Manual No. 91
Year :	2016
Title :	Ireland Red List No. 10: Vascular Plants
Author :	Wyse Jackson, M.; FitzPatrick, Ú.; Cole, E.; Jebb, M.; McFerran, D.; Sheehy Skeffington, M.; Wright, M.
Series :	Ireland Red Lists series, NPWS
Year :	2017
Title :	Carrowmore Lake Complex SAC (site code: 476) Conservation objectives supporting document- blanket bogs and associated habitats V1
Author :	NPWS
Series :	Conservation objectives supporting document

Other References

Year :	1989
Title :	The genera <i>Scorpidium</i> and <i>Hamatocaulis</i> , gen. nov., in northern Europe
Author :	Hedenäs, L.
Series :	Lindbergia, 15: 8-36

Year : 2011
Title : Conservation biology of *Saxifraga hirculus* L. in Ireland
Author : Muldoon, C.S.
Series : Unpublished Ph.D. Thesis, Trinity College Dublin

Year : 2013
Title : Conservation of selected legally protected and Red Listed bryophytes in Ireland
Author : Campbell, C.
Series : Unpublished Ph.D. Thesis, Trinity College Dublin

Spatial data sources

Year : 2017

Title : NPWS rare and threatened species database

GIS Operations : Datasets created from spatial references in database records. Expert opinion used as necessary to resolve any issues arising

Used For : 1393, 1528 (maps 3 and 4)

Conservation Objectives for : Carrowmore Lake Complex SAC [000476]

7130 Blanket bogs (* if active bog)

To restore the favourable conservation condition of Blanket bogs (* if active bog) in Carrowmore Lake Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Blanket bog has not been mapped in detail for Carrowmore Lake Complex SAC but from current available data the total area of the qualifying habitat is estimated to be approximately 2,285ha, covering 63% of the SAC (NPWS internal files). Further details on this and the following attributes can be found in the Carrowmore Lake Complex SAC conservation objectives supporting document for blanket bogs and associated habitats
Habitat distribution	Occurrence	No decline, subject to natural processes	Blanket bog habitat lies east of Carrowmore Lake. Three large areas of blanket bog are incorporated into the SAC at Glenturk, Carrowmore (or Glencullin) and Largan More. Further information can be found in Foss and McGee (1987), Douglas et al. (1989), NPWS internal files and the blanket bogs and associated habitats supporting document
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil nutrient status within natural range	See the blanket bogs and associated habitats supporting document for further details
Ecosystem function: peat formation	Active blanket bog as a proportion of the total area of Annex I blanket bog habitat	At least 99% of the total Annex I blanket bog area is active	See the blanket bogs and associated habitats supporting document for further details
Ecosystem function: hydrology	Flow direction, water levels, occurrence of drains and erosion gullies	Natural hydrology unaffected by drains and erosion	Further details and a brief discussion of restoration potential is presented in the blanket bogs and associated habitats supporting document
Community diversity	Abundance of variety of vegetation communities	Maintain variety of vegetation communities, subject to natural processes	A variety of blanket bog vegetation communities have been recorded in this SAC (Foss and McGee, 1987; Douglas et al., 1989; NPWS internal files), four of which correspond to communities recorded in the National Survey of Upland Habitats and listed in the provisional list of vegetation communities described in Perrin et al. (2014). Further information on vegetation communities associated with this habitat is presented in Perrin et al. (2014)
Vegetation composition: positive indicator species	Number of species at a representative number of 2m x 2m monitoring stops	Number of positive indicator species present at each monitoring stop is at least seven	Attribute and target based on Perrin et al. (2014), where the list of positive indicator species for this habitat is also presented
Vegetation composition: lichens and bryophytes	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of bryophytes or lichens, excluding <i>Sphagnum fallax</i> , at least 10%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: potential dominant species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of each of the potential dominant species less than 75%	Attribute and target based on Perrin et al. (2014). See the blanket bogs and associated habitats supporting document for the list of potential dominant species
Vegetation composition: negative indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of negative indicator species less than 1%	Attribute and target based on Perrin et al. (2014), where the list of negative indicator species for this habitat is also presented
Vegetation composition: non-native species	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of non-native species less than 1%	Attribute and target based on Perrin et al. (2014). Rhododendron (<i>Rhododendron ponticum</i>) was recorded from blanket bog within the SAC (NPWS internal files)

Vegetation composition: native trees and shrubs	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of scattered native trees and shrubs less than 10%	Attribute and target based on Perrin et al. (2014)
Vegetation structure: <i>Sphagnum</i> condition	Condition at a representative number of 2m x 2m monitoring stops	Less than 10% of the <i>Sphagnum</i> cover is crushed, broken and/or pulled up	Attribute and target based on Perrin et al. (2014)
Vegetation structure: signs of browsing	Percentage of shoots browsed at a representative number of 2m x 2m monitoring stops	Last complete growing season's shoots of ericoids, crowberry (<i>Empetrum nigrum</i>) and bog-myrtle (<i>Myrica gale</i>) showing signs of browsing collectively less than 33%	Attribute and target based on Perrin et al. (2014)
Vegetation structure: burning	Occurrence in local vicinity of a representative number of monitoring stops	No signs of burning in sensitive areas, into the moss, liverwort or lichen layer or exposure of peat surface due to burning	Attribute and target based on Perrin et al. (2014), where the list of sensitive areas for this habitat is also presented
Physical structure: disturbed bare ground	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of disturbed bare ground less than 10%	Attribute and target based on Perrin et al. (2014)
Physical structure: drainage	Percentage area in local vicinity of a representative number of monitoring stops	Area showing signs of drainage from heavy trampling, tracking or ditches less than 10%	Attribute and target based on Perrin et al. (2014)
Physical structure: erosion	Percentage area in local vicinity of a representative number of monitoring stops	Less than 5% of the greater bog mosaic comprises erosion gullies and eroded areas	Attribute and target based on Perrin et al. (2014)
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat	This includes species listed in the Flora (Protection) Order, 2015 (FPO) and/or the red data lists (Lockhart et al., 2012; Wyse Jackson et al., 2016). There are rare and threatened species in flushes within blanket bog in the SAC (NPWS internal files), including the Annex II and Annex IV listed, FPO listed and Near Threatened marsh saxifrage (<i>Saxifraga hirculus</i>) (Wyse Jackson et al., 2016) and the Annex II listed, FPO listed and Near Threatened slender green feather-moss (<i>Hamatocaulis (Drepanocladus) vernicosus</i>) (Lockhart et al., 2012). Although these species cannot be assigned specifically to blanket bog habitat, i.e. they are flush/fen species, they do occur in association with it (Muldoon et al., 2015; Campbell et al., 2015). See also the conservation objectives for marsh saxifrage (1528) and slender green feather-moss (1393)

Conservation Objectives for : Carrowmore Lake Complex SAC [000476]

7150 Depressions on peat substrates of the Rhynchosporion

To restore the favourable conservation condition of Depressions on peat substrates of the Rhynchosporion in Carrowmore Lake Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Depressions on peat substrates of the Rhynchosporion has not been mapped in detail for Carrowmore Lake Complex SAC and thus the total area of the qualifying habitat is unknown. Further details on this and the following attributes can be found in the Carrowmore Lake Complex SAC conservation objectives supporting document for blanket bogs and associated habitats
Habitat distribution	Occurrence	No decline, subject to natural processes	This habitat is scattered throughout the blanket bog in the SAC. It is best developed in areas of deep, quaking peat and around pools (Douglas et al., 1989; NPWS internal files). Further information can be found in Douglas et al. (1989), NPWS internal files and the blanket bogs and associated habitats supporting document
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil nutrient status within natural range	See the blanket bogs and associated habitats supporting document for further details
Vegetation composition: positive indicator species	Number of species at a representative number of 2m x 2m monitoring stops	Number of positive indicator species at each monitoring stop is at least five	Attribute and target based on Perrin et al. (2014), where the list of positive indicator species for this habitat is also presented
Vegetation composition: <i>Rhynchospora</i> spp.	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of white beaked sedge (<i>Rhynchospora alba</i>) and brown beaked sedge (<i>R. fusca</i>) at least 10%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: potential dominant species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of each of the potential dominant species individually less than 35%	Attribute and target based on Perrin et al. (2014). See the blanket bogs and associated habitats supporting document for the list of potential dominant species
Vegetation composition: negative indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of negative indicator species less than 1%	Attribute and target based on Perrin et al. (2014), where the list of negative indicator species for this habitat is also presented
Vegetation composition: non-native species	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of non-native species less than 1%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: native trees and shrubs	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of scattered native trees and shrubs less than 10%	Attribute and target based on Perrin et al. (2014)
Vegetation structure: <i>Sphagnum</i> condition	Condition at a representative number of 2m x 2m monitoring stops	Less than 10% of the <i>Sphagnum</i> cover is crushed, broken and/or pulled up	Attribute and target based on Perrin et al. (2014)
Vegetation structure: signs of browsing	Percentage of shoots browsed at a representative number of 2m x 2m monitoring stops	Last complete growing season's shoots of ericoids, crowberry (<i>Empetrum nigrum</i>) and bog-myrtle (<i>Myrica gale</i>) showing signs of browsing collectively less than 33%	Attribute and target based on Perrin et al. (2014)

Vegetation structure: burning	Occurrence in local vicinity of a representative number of monitoring stops	No signs of burning in sensitive areas, into the moss, liverwort or lichen layer or exposure of peat surface due to burning	Attribute and target based on Perrin et al. (2014), where the list of sensitive areas for this habitat is also presented
Physical structure: disturbed bare ground	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of disturbed bare ground less than 10%	Attribute and target based on Perrin et al. (2014)
Physical structure: drainage	Percentage area in local vicinity of a representative number of monitoring stops	Area showing signs of drainage from heavy trampling, tracking or ditches less than 10%	Attribute and target based on Perrin et al. (2014)
Physical structure: erosion	Percentage area in local vicinity of a representative number of monitoring stops	Less than 5% of the greater bog mosaic comprises erosion gullies and eroded areas	Attribute and target based on Perrin et al. (2014)
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat	This includes species listed in the Flora (Protection) Order, 2015 and/or the red data lists (Lockhart et al., 2012; Wyse Jackson et al., 2016)

Conservation Objectives for : Carrowmore Lake Complex SAC [000476]

1393 Slender Green Feather-moss *Drepanocladus vernicosus*

To maintain the favourable conservation condition of Slender Green Feather-moss (Shining Sickle-moss) in Carrowmore Lake Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Distribution of populations	Number and geographical spread of populations	No decline, subject to natural processes. See map 3 for known locations at Largan More	(Please note that <i>Drepanocladus vernicosus</i> was reclassified as <i>Hamatocaulis vernicosus</i> by Hedenäs (1989)). The known population of slender green feather-moss (<i>Hamatocaulis vernicosus</i>) in Carrowmore Lake Complex SAC occurs in flushes, spring heads and beside water tracks within the blanket bog at Largan More. Data from NPWS surveys (NPWS internal files), Campbell (2013) and Campbell et al. (2015)
Population size	Number of individuals	No decline, subject to natural processes	The population at Largan More was estimated by Campbell (2013) to be 3,979,350 shoots (c.3,979,000 shoots). Counts of shoots were based on the mean of number of shoots in four 10cm x 10cm areas, extrapolated to 8,325 per m ² in 478m ² (Campbell, 2013). See Campbell et al. (2015) for further details
Population cover	Percentage cover in a representative number of 2m x 2m monitoring plots	Mean percentage cover of slender green feather-moss (<i>Hamatocaulis vernicosus</i>) should be at least 24%	The mean percentage cover of slender green feather-moss (<i>Hamatocaulis vernicosus</i>) recorded in four 2m x 2m plots at Largan More was 29.5% (Campbell, 2013). The target cover figure is a c.20% reduction of the recorded cover to allow for a margin of error and variability over monitoring seasons. See Campbell et al. (2015) for further details
Area of suitable habitat	Hectares	No decline, subject to natural processes	The extent of occupancy for the species at Largan More is estimated to be c.1,593m ² ; however, only c.30% of this area is suitable habitat, i.e. c.478m ² (0.048ha). See Campbell (2013) and Campbell et al. (2015) for further details
Hydrological conditions: water table level	Metres	Maintain suitable hydrological conditions	Slender green feather-moss (<i>Hamatocaulis vernicosus</i>) is mostly confined to mesotrophic fens, a transitional habitat between acid bog and base-rich fen. This appears to occur in at least two forms in Ireland: transitional flushes, where the plants can occur in lawns that rise and fall with fluctuating water table levels, such as at Largan More; and wet lowland sedge meadows, where plants can be inundated in winter, but may be subject to some desiccation in the summer. Based on Campbell (2013) and Campbell et al. (2015)
Vegetation composition: tree cover	Percentage cover in a representative number of 2m x 2m monitoring plots	Mean percentage tree cover should be less than 15%	Slender green feather-moss (<i>Hamatocaulis vernicosus</i>) grows in moss-dominated, open communities, generally with a low cover of trees and shrubs. Campbell (2013) recorded 0% tree cover at Largan More. See also Campbell et al. (2015)
Vegetation composition: shrub cover	Percentage cover in a representative number of 2m x 2m monitoring plots	Mean percentage shrub cover should be less than 20%	Slender green feather-moss (<i>Hamatocaulis vernicosus</i>) grows in moss-dominated, open communities, generally with a low cover of trees and shrubs. Campbell (2013) recorded 0% shrub cover at Largan More. See also Campbell et al. (2015)
Vegetation composition: grass cover	Percentage cover in a representative number of 2m x 2m monitoring plots	Mean percentage grass species cover should be less than 25%	Slender green feather-moss (<i>Hamatocaulis vernicosus</i>) grows in moss-dominated, open communities, generally with a low cover of grasses, maintained by a low grazing intensity by sheep and cattle at Largan More. Campbell (2013) recorded grass cover of 1-10% in four 2m x 2m plots at Largan More. See also Campbell et al. (2015)
Vegetation composition: bryophyte cover	Percentage cover in a representative number of 2m x 2m monitoring plots	Mean percentage bryophyte cover should be more than 50%	Campbell (2013) recorded bryophyte cover of 34-90% in four 2m x 2m plots at Largan More. See also Campbell et al. (2015)



Vegetation composition: cover of <i>Calliergonella cuspidata</i>	Percentage cover in a representative number of 2m x 2m monitoring plots	Mean percentage cover of <i>Calliergonella cuspidata</i> should be less than 15%	<i>Calliergonella cuspidata</i> , a moss species often associated with high nutrient conditions, is usually present, but with low cover and never dominant. Cover of <i>Calliergonella cuspidata</i> was 0-2% in four 2m x 2m plots recorded by Campbell (2013) at Largan More. See also Campbell et al. (2015)
Vegetation structure: vegetation height	Centimetres in a representative number 2m x 2m monitoring plots	Mean vegetation height should not exceed 40cm	Campbell (2013) recorded a mean vegetation height of 13.7cm in four 2m x 2m plots at Largan More. See also Campbell et al. (2015)

Conservation Objectives for : Carrowmore Lake Complex SAC [000476]


1528 Marsh Saxifrage *Saxifraga hirculus*

To maintain the favourable conservation condition of Marsh Saxifrage in Carrowmore Lake Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Distribution of populations	Number and geographical spread of populations	No loss in geographical spread and number of populations, subject to natural processes. See map 4 for 1km grid square locations	Marsh saxifrage (<i>Saxifraga hirculus</i>) is known to occur in three flushes in Carrowmore Lake Complex SAC, at Largan Mor A, Largan Mor B and Largan Mor C. See Muldoon (2011) and Muldoon et al. (2015)
Population size: number of rosettes	Number	Maintain the size of each known population, subject to natural processes. The target numbers of rosettes are: at least 2,800 at Largan Mor A, at least 440 at Largan Mor B and at least 80 at Largan Mor C	The numbers of rosettes recorded were: 3,500 at Largan Mor A, 550 at Largan Mor B (Muldoon, 2011) and 100 at Largan Mor C (NPWS internal files). The target figures are a 20% reduction of the recorded numbers to allow for a margin of error and variability over monitoring seasons. See Muldoon et al. (2015) for further details
Population size: area of occupancy	Hectares	Maintain the area of occupancy of each known population, subject to natural processes. The target areas are: at least 0.0084ha at Largan Mor A, at least 0.00045ha at Largan Mor B and at least 0.00027ha at Largan Mor C	The areas of occupancy for the species estimated by Muldoon (2011) were: 93m ² (0.0093ha) at Largan Mor A and 5m ² (0.0005ha) at Largan Mor B. The area of occupancy at Largan Mor C has been estimated to be 3m ² (0.0003ha) (NPWS internal files). The target area figures are a 10% reduction of the recorded areas to allow for a margin of error. See Muldoon et al. (2015) for further details
Hydrological conditions: water level	Occurrence of high or fluctuating water levels	Maintain the appropriate natural hydrological regime necessary to support the habitat for the species	In Ireland, marsh saxifrage (<i>Saxifraga hirculus</i>) is now restricted to mineral flushes in blanket bog where rising groundwater forms small streams and seepage areas suitable for the species. Based on Muldoon (2011) and Muldoon et al. (2015)
Vegetation composition: positive indicator species	Occurrence in a number of 1m x 1m monitoring stops	Knotted pearlwort (<i>Sagina nodosa</i>) should be present in at least two of five 1m x 1m monitoring stops	See Muldoon (2011) and Muldoon et al. (2015) for further details
Vegetation composition: negative indicator species	Mean percentage cover in five 1m x 1m monitoring stops	Mean percentage cover of purple moor-grass (<i>Molinia caerulea</i>) should not exceed 5%; mean percentage cover of Yorkshire fog (<i>Holcus lanatus</i>) should not exceed 15%	Low cover of the negative indicator species purple moor-grass (<i>Molinia caerulea</i>) and Yorkshire fog (<i>Holcus lanatus</i>) should be maintained. See Muldoon (2011) and Muldoon et al. (2015) for further details
Vegetation structure: sward height	Centimetres	Maintain a mean vegetation height of less than 15cm	See Muldoon (2011) and Muldoon et al. (2015) for further details
Vegetation structure: grazing level	Evidence of grazing	Maintain grazing at light to moderate levels to ensure an open vegetation structure and to allow flowering to occur	Low grazing was identified as an issue at Largan Mor A and Largan Mor B; however, vegetation height was within the target range at both sites. See Muldoon (2011) and Muldoon et al. (2015) for further details



Legend

 Carrowmore Lake Complex SAC 000476

 An Roinn Ealaíon, Oidhreachta, Gnóthai Réigiúnacha, Tuaithe agus Gaeltachta
Department of Arts, Heritage, Regional, Rural and Gaeltacht Affairs

MAP 1:
CARROWMORE LAKE COMPLEX SAC
CONSERVATION OBJECTIVES
SAC DESIGNATION

Map to be read in conjunction with the NPWS Conservation Objectives Document.

SITE CODE:
SAC 000476; version 3
CO. MAYO


0 1 2 3 4 km

The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision. Ordnance Survey of Ireland Licence No EN 0059216. © Ordnance Survey of Ireland Government of Ireland.

Níl sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbheirniú a déanamh ar theorainneacha na gceantar comharthaithe. Suirbhéarachta Ordonáis na hÉireann Ceadúnas Uimh EN 0059216. © Suirbhéarachta Ordonáis na hÉireann Rialtas na hÉireann.



Map Version 1
Date: April 2017

© Ordnance Survey Ireland




Legend

- Carrowmore Lake Complex SAC 000476
- Slieve Fyagh Bog SAC 000542
- Carrowmore Lake SPA 004052





Legend

- ★ 1393 Slender Green Feather-moss *Drepanocladus vernicosus*
- Carrowmore Lake Complex SAC 000476



Legend

-  1528 Marsh Saxifrage - *Saxifraga hirculus*
-  Carrowmore Lake Complex SAC 000476

National Parks and Wildlife Service

Conservation Objectives Series

Cloughmoyne SAC 000479



An Roinn Cultúir,
Oidhreacht agus Gaeltachta
Department of Culture,
Heritage and the Gaeltacht

**National Parks and Wildlife Service,
Department of Culture, Heritage and the Gaeltacht,
90 King Street North, Dublin 7, D07 N7CV, Ireland.
Web: www.npws.ie
E-mail: nature.conservation@chg.gov.ie**

Citation:

**NPWS (2019) Conservation Objectives: Cloughmoyne SAC 000479. Version 1.
National Parks and Wildlife Service, Department of Culture, Heritage and the
Gaeltacht.**

**Series Editor: Rebecca Jeffrey
ISSN 2009-4086**

Introduction

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

** indicates a priority habitat under the Habitats Directive*

000479 Cloughmoyne SAC

8240 Limestone pavementsE

Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

NPWS Documents

Year :	2009
Title :	Ireland Red List No. 2: Non-marine molluscs
Author :	Byrne, A.; Moorkens, E.A.; Anderson, R.; Killeen, I.J.; Regan, E.C.
Series :	Ireland Red List series, NPWS
<hr/>	
Year :	2010
Title :	Ireland Red List No. 4: Butterflies
Author :	Regan, E.C.; Nelson, B.; Aldwell, B.; Bertrand, C.; Bond, K.; Harding, J.; Nash, D.; Nixon, D.; Wilson, C.J.
Series :	Ireland Red List series, NPWS
<hr/>	
Year :	2012
Title :	Ireland Red List No. 8: Bryophytes
Author :	Lockhart, N.; Hodgetts, N.; Holyoak, D.
Series :	Ireland Red List series, NPWS
<hr/>	
Year :	2013
Title :	National survey of limestone pavement and associated habitats in Ireland
Author :	Wilson, S.; Fernandez, F.
Series :	Irish Wildlife Manuals, No. 73
<hr/>	
Year :	2016
Title :	Ireland Red List No. 10: Vascular Plants
Author :	Wyse Jackson, M.; FitzPatrick, Ú.; Cole, E.; Jebb, M.; McFerran, D.; Sheehy Skeffington, M.; Wright, M.
Series :	Ireland Red List Series, NPWS

Other References

Year :	1995
Title :	Wood bitter-vetch <i>Vicia orobus</i> DC. on lake islands and limestone heath in Cos Galway (H16, H17) and Mayo (H26)
Author :	Roden, C.
Series :	Irish Naturalists' Journal, 25(4): 128-134

Spatial data sources

Year :	2013
Title :	National Survey of Limestone Pavement and Associated Habitats in Ireland distribution data
GIS Operations :	Dataset clipped to the SAC boundary. Expert opinion used as necessary to resolve any issues arising
Used For :	8240 (map 2)


Conservation Objectives for : Cloughmoyne SAC [000479]


8240 Limestone pavements


To restore the favourable conservation condition of Limestone pavements* in Cloughmoyne SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Limestone pavements* in Cloughmoyne SAC exhibits a good example of the shattered form of the habitat and occurs in intimate association with species-rich calcareous grassland, juniper (<i>Juniperus communis</i>) scrub and heath habitats. Therefore, these habitats cannot easily be mapped or considered separately. Wilson and Fernandez (2013) mapped the indicative area of limestone pavement, including mosaics with associated habitats, as 45.2ha in the SAC (see map 2). As part of the National Survey of Limestone Pavement and Associated Habitats (Wilson and Fernandez, 2013), the sub-site Cloughmoyne (site code NSLP26) was surveyed in detail. This survey should be consulted for further details
Habitat distribution	Occurrence	No decline. Map 2 shows indicative distribution, including mosaics with other habitats	See the notes for Habitat area above. Distribution based on data from Wilson and Fernandez (2013). This habitat can be split into exposed pavement and wooded pavement. In this SAC, the habitat is largely confined to hilly land in the east of the SAC where there are some typical clint and grike features in addition to mostly shattered pavement with loose limestone rubble (NPWS internal files)
Vegetation composition: positive indicator species	Number at a representative number of monitoring stops	At least seven positive indicator species present	Positive indicator species for exposed and wooded pavement are listed in Wilson and Fernandez (2013). Positive indicator species recorded by Wilson and Fernandez (2013) in exposed pavement in the SAC include herb-robert (<i>Geranium robertianum</i>), bloody crane's-bill (<i>G. sanguineum</i>), ivy (<i>Hedera helix</i>), wall lettuce (<i>Mycelis muralis</i>), wood sage (<i>Teucrium scorodonia</i>), wild madder (<i>Rubia peregrina</i>), blue moor-grass (<i>Sesleria caerulea</i>), burnet rose (<i>Rosa spinosissima</i>), juniper (<i>Juniperus communis</i>), maidenhair spleenwort (<i>Asplenium trichomanes</i>), wall-rue (<i>A. ruta-muraria</i>) and the mosses <i>Ctenidium molluscum</i> , <i>Fissidens dubius</i> , <i>Neckera crispa</i> and <i>Tortella tortuosa</i>
Vegetation composition: bryophyte layer	Percentage at a representative number of monitoring stops	Bryophyte cover at least 50% on wooded pavement	Attribute and target based on Wilson and Fernandez (2013)
Vegetation composition: negative indicator species	Percentage at a representative number of monitoring stops	Collective cover of negative indicator species on exposed pavement not more than 1%	Negative indicator species are listed in Wilson and Fernandez (2013). Negative indicator species for wooded pavement overlap with non-native species (below)
Vegetation composition: non-native species	Percentage at a representative number of monitoring stops	Cover of non-native species not more than 1% on exposed pavement; on wooded pavement not more than 10% with no regeneration	Attribute and target based on Wilson and Fernandez (2013). Cotoneaster (<i>Cotoneaster</i> spp.) was recorded in the Cloughmoyne sub-site (NSLP26) by Wilson and Fernandez (2013)
Vegetation composition: scrub	Percentage at a representative number of monitoring stops	Scrub cover no more than 25% of exposed pavement	Attribute and target based on Wilson and Fernandez (2013). Scrub encroachment was recorded as a negative impact in the habitat in this SAC by Wilson and Fernandez (2013)
Vegetation composition: bracken cover	Percentage at a representative number of monitoring stops	Bracken (<i>Pteridium aquilinum</i>) cover no more than 10% on exposed pavement	Attribute and target based on Wilson and Fernandez (2013). There is some encroachment of bracken onto areas of the limestone pavement habitat in parts of the SAC (Wilson and Fernandez, 2013; NPWS internal files)
Vegetation structure: woodland canopy	Percentage at a representative number of monitoring stops	Canopy cover on wooded pavement at least 30%	Attribute and target based on Wilson and Fernandez (2013)

Vegetation structure: dead wood	Occurrence in a representative number of monitoring stops	Sufficient quantity of dead wood on wooded pavement to provide habitat for saproxylic organisms	Dead wood is a valuable resource and an integral part of a healthy, functioning woodland ecosystem
Physical structure: disturbance	Occurrence in a representative number of monitoring stops	No evidence of grazing pressure on wooded pavement	Attribute and target based on Wilson and Fernandez (2013)
Indicators of local distinctiveness	Occurrence	Indicators of local distinctiveness are maintained	This includes species on the Flora (Protection) Order, 2015 (FPO) and/or Red Lists (Byrne et al., 2009; Regan et al., 2010; Lockhart et al., 2012; Wyse Jackson et al., 2016, etc.) and other rare or localised species, as well as archaeological and geological features, which often support distinctive species. Rare and threatened species recorded in the limestone pavement and associated calcareous grassland and heath habitats in the SAC include the FPO listed and Critically Endangered limestone fern (<i>Gymnocarpium robertianum</i>), the FPO listed and Vulnerable wood bitter-vetch (<i>Vicia orobus</i>), the Vulnerable green-winged orchid (<i>Anacamptis morio</i>) and the Near Threatened species spring gentian (<i>Gentiana verna</i>), autumn gentian (<i>G. amarella</i>), field gentian (<i>G. campestris</i>) and dense-flowered orchid (<i>Neotinea maculata</i>) (Roden, 1995; Wilson and Fernandez, 2013; Wyse Jackson et al., 2016; NPWS internal files)




Legend
 Cloughmoyne SAC 000479

 An Roinn Cultúir,
 Oidhreacht agus Gaeltachta
 Department of Culture,
 Heritage and the Gaeltacht

**MAP 1:
 CLOUGHMOYNE SAC
 CONSERVATION OBJECTIVES
 SAC DESIGNATION**
 Map to be read in conjunction with the NPWS Conservation Objectives Document.

**SITE CODE:
 SAC 000479; version 3.02. CO. MAYO**


0 90 180 270 360 Meters






The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
 Ordnance Survey Ireland Licence No OSI-NMA-014 © Ordnance Survey Ireland Government of Ireland

Níl sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbhreithnithe a déanamh ar theorainneacha na gceantar comharthaithe. Suirbhéarachta Ordonáis na hÉireann Ceadúnas Uimh OSI-NMA-014. © Suirbhéarachta Ordonáis na hÉireann Rialtas na hÉireann


Map Version 1
Date: Feb 2019



Legend

-  8240 Limestone pavements
-  Cloughmoyne SAC 000479
-  OSi Discovery Series County Boundary




An Roinn Cultúir,
Oidhreacht agus Gaeltachta
Department of Culture,
Heritage and the Gaeltacht

**MAP 2:
CLOUGHMOYNE SAC
CONSERVATION OBJECTIVES
LIMESTONE PAVEMENT**

Map to be read in conjunction with the NPWS Conservation Objectives Document.

**SITE CODE:
SAC 000479; version 3.02. CO. MAYO**



The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
Ordnance Survey Ireland Licence No OSI-NMA-014
© Ordnance Survey Ireland Government of Ireland
Níl sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbheithnithe a déanamh ar theorainneacha na gceantar comharthaithe. Suirbhéarachta Ordonáis na hÉireann Ceadúnas Uimh OSI-NMA-014 © Suirbhéarachta Ordonáis na hÉireann Riailtas na hÉireann



**Map Version 1
Date: Feb 2019**



Conservation objectives for Clyard Kettle-holes SAC [000480]

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Objective: To maintain or restore the favourable conservation condition of the Annex I habitat(s) and/or the Annex II species for which the SAC has been selected:

Code	Description
------	-------------

3180	Turloughs*
------	------------

7210	Calcareous fens with <i>Cladium mariscus</i> and species of the <i>Caricion davallianae</i> *
------	---

* denotes a priority habitat



Citation: NPWS (2020) Conservation objectives for Clyard Kettle-holes SAC [000480]. Generic Version 7.0. Department of Culture, Heritage and the Gaeltacht.



Conservation objectives for Cross Lough (Killadoon) SAC [000484]

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Objective: To maintain or restore the favourable conservation condition of the Annex I habitat(s) and/or the Annex II species for which the SAC has been selected:

Code	Description
------	-------------

1220	Perennial vegetation of stony banks
------	-------------------------------------

* denotes a priority habitat



Citation: NPWS (2020) Conservation objectives for Cross Lough (Killadoon) SAC [000484]. Generic Version 7.0. Department of Culture, Heritage and the Gaeltacht.

National Parks and Wildlife Service

Conservation Objectives Series

Corraun Plateau SAC 000485



An Roinn Ealaíon, Oidhreachta,
Gnóthaí Réigiúnacha, Tuaithe agus Gaeltachta

Department of Arts, Heritage,
Regional, Rural and Gaeltacht Affairs



**National Parks and Wildlife Service,
Department of Arts, Heritage, Regional, Rural and Gaeltacht Affairs,
7 Ely Place, Dublin 2, Ireland.
Web: www.npws.ie
E-mail: nature.conservation@ahg.gov.ie**

Citation:

**NPWS (2016) Conservation Objectives: Corraun Plateau SAC 000485. Version 1.
National Parks and Wildlife Service, Department of Arts, Heritage, Regional,
Rural and Gaeltacht Affairs.**

**Series Editor: Rebecca Jeffrey
ISSN 2009-4086**

Introduction

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

* indicates a priority habitat under the Habitats Directive

000485	Corraun Plateau SAC
4010	Northern Atlantic wet heaths with <i>Erica tetralix</i>
4030	European dry heaths
4060	Alpine and Boreal heaths
5130	<i>Juniperus communis</i> formations on heaths or calcareous grasslands
8110	Siliceous scree of the montane to snow levels (<i>Androsacetalia alpinae</i> and <i>Galeopsietalia ladani</i>)
8220	Siliceous rocky slopes with chasmophytic vegetation

Please note that this SAC adjoins Owenduff/Nepin Complex SAC (000534). See map 2. The conservation objectives for this site should be used in conjunction with those for adjacent sites as appropriate.

Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

NPWS Documents

Year :	2012
Title :	The Conservation Status of Juniper Formations in Ireland
Author :	Cooper, F.; Stone, R.E.; McEvoy, P.; Wilkins, T.; Reid, N.
Series :	Irish Wildlife Manual No. 63
<hr/>	
Year :	2014
Title :	Guidelines for a national survey and conservation assessment of upland vegetation and habitats in Ireland, Version 2.0
Author :	Perrin, P.M.; Barron, S.J.; Roche, J.R.; O'Hanrahan, B.
Series :	Irish Wildlife Manual No. 79
<hr/>	
Year :	2014
Title :	National survey of upland habitats (pilot survey phase, 2009-2010), site report no. 2: Corraun Plateau cSAC (000485), Co. Mayo (revision)
Author :	Roche, J.R.; Perrin, P.M.; Barron, S.J.; Daly, O.H.
Series :	Unpublished report to NPWS
<hr/>	
Year :	2014
Title :	National survey of upland habitats (phase 4, 2013-2014), summary report
Author :	Barron, S.J.; Perrin, P.M.
Series :	Unpublished report to NPWS
<hr/>	
Year :	2016
Title :	Corraun Plateau SAC (site code: 485) Conservation objectives supporting document- upland habitats V1
Author :	NPWS
Series :	Conservation objectives supporting document

Spatial data sources

Year : 2010
Title : National Survey of Upland Habitats
GIS Operations : Habitat dataset for site clipped to SAC boundary. Relevant QI selected and exported to new dataset. Expert opinion used as necessary to resolve any issues arising
Used For : 4010, 4030, 4060, 8110, 8220 (maps 3, 4, 5, 7 and 8)

Year : 2012
Title : The conservation status of juniper formations in Ireland
GIS Operations : Juniper formations polygon centroids clipped to SAC boundary
Used For : 5130 (map 6)

Conservation Objectives for : Corraun Plateau SAC [000485]

4010 Northern Atlantic wet heaths with *Erica tetralix*

To restore the favourable conservation condition of Northern Atlantic wet heaths with *Erica tetralix* in Corraun Plateau SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area increasing, subject to natural processes	Corraun Plateau SAC was surveyed as part of the National Survey of Upland Habitats (NSUH; see Roche et al., 2014 & Perrin et al., 2014). The total current area of wet heath in the SAC stated by Roche et al. (2014) is 2082.4ha. It is the most extensive Annex I habitat at the SAC covering 53.57% of the SAC. Roche et al. (2014) report obvious losses of habitat since 1995 of approximately 8.07ha. A summary of the mapping methodology and a brief discussion of restoration potential are presented in the uplands supporting document
Habitat distribution	Occurrence	No decline from current distribution, subject to natural processes. See map 3	Wet heath was recorded by Roche et al. (2014) throughout the SAC except on the higher ground. Extensive patches occur at on the eastern and southern slopes. A summary of the mapping methodology is presented in the uplands supporting document
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil nutrient status within natural range	See the uplands supporting document for further details
Community diversity	Abundance of variety of vegetation communities	Maintain variety of vegetation communities, subject to natural processes	Perrin et al. (2014) recorded five different wet heath communities within this SAC. Data on the abundance of these communities is reproduced in the uplands supporting document. Further information on these communities is presented in Perrin et al. (2014)
Vegetation composition: cross-leaved heath	Occurrence within 20m of a representative number of 2m x 2m monitoring stops	Presence of cross-leaved heath (<i>Erica tetralix</i>) near each monitoring stop	Based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation composition: positive indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of positive indicator species at least 50%	Based on Perrin et al. (2014). The list of positive indicator species for this habitat is presented in Perrin et al. (2014). Further details can be found in the uplands supporting document
Vegetation composition: lichens and bryophytes	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of <i>Cladonia</i> and <i>Sphagnum</i> species, <i>Racomitrium lanuginosum</i> and pleurocarpous mosses at least 10%	Based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation composition: ericoid species and crowberry	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of ericoid species and crowberry (<i>Empetrum nigrum</i>) at least 15%	Based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation composition: dwarf shrub species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of dwarf shrubs less than 75%	Based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation composition: negative indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of negative indicator species less than 1%	Based on Perrin et al. (2014). The list of negative indicator species is given in Perrin et al. (2014). See the uplands supporting document for further details
Vegetation composition: non-native species	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of non-native species less than 1%	Based on Perrin et al. (2014). See the uplands supporting document for further details. <i>Campylopus introflexus</i> was recorded within this habitat by Roche et al. (2014) and forming extensive carpets. A small population of <i>Rhododendron ponticum</i> was recorded from this habitat by Roche et al. (2014)

Vegetation composition: native trees and shrubs	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of scattered native trees and shrubs less than 20%	Based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation composition: bracken	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of bracken (<i>Pteridium aquilinum</i>) less than 10%	Based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation composition: soft rush	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of soft rush (<i>Juncus effusus</i>) less than 10%	Based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation structure: <i>Sphagnum</i> condition	Condition at a representative number of 2m x 2m monitoring stops	Less than 10% of the <i>Sphagnum</i> cover is crushed, broken and/or pulled up	Based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation structure: signs of browsing	Percentage of shoots browsed at a representative number of 2m x 2m monitoring stops	Less than 33% collectively of the last complete growing season's shoots of ericoids, crowberry (<i>Empetrum nigrum</i>) and bog-myrtle (<i>Myrica gale</i>) showing signs of browsing	Based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation structure: burning	Occurrence in local vicinity of a representative number of 2m x 2m monitoring stops	No signs of burning in sensitive areas, into the moss, liverwort or lichen layer or exposure of peat surface due to burning	Based on Perrin et al. (2014). The list of sensitive areas is presented in Perrin et al. (2014). See the uplands supporting document for further details
Physical structure: disturbed bare ground	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of disturbed bare ground less than 10%	Based on Perrin et al. (2014). See the uplands supporting document for further details
Physical structure: drainage	Percentage cover in local vicinity of a representative number of monitoring stops	Area showing signs of drainage from heavy trampling, tracking or ditches less than 10%	Based on Perrin et al. (2014). See the uplands supporting document for further details
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat and no decline in status of hepatic mats associated with this habitat	Roche et al. (2014) compiled and mapped existing rare and notable plant records for the SAC and added any new records collected during the NSUH survey. <i>Cladonia rangiferina</i> has been recorded from within the habitat. This species is listed on a preparatory list of rare and threatened lichens prepared by D. McFerran, National Museums Northern Ireland. This and any new records should be considered within this attribute. Hepatic mats of the <i>Calluna vulgaris-Herbertus aduncus</i> community were recorded within this habitat by Roche et al. (2014). No assessment of the conservation status of this community has been conducted but proposals for such an assessment are presented in Barron and Perrin (2014). See the uplands supporting document for further details

Conservation Objectives for : Corraun Plateau SAC [000485]

4030 European dry heaths

To restore the favourable conservation condition of European dry heaths in Corraun Plateau SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Corraun Plateau SAC was surveyed as part of the National Survey of Upland Habitats (NSUH; see Roche et al., 2014 & Perrin et al., 2014). The total current area of dry heath in the SAC stated by Roche et al. (2014) is 207.9ha. It covers 5.35% of the SAC. Roche et al. (2014) report no significant losses of area since 1995. A summary of the mapping methodology is presented in the uplands supporting document
Habitat distribution	Occurrence	No decline from current habitat distribution, subject to natural processes. See map 4	Dry heath was recorded by Roche et al. (2014) throughout the SAC, but was most abundant on the northern slopes of the SAC. A summary of the mapping methodology is presented in the uplands supporting document
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil nutrient status within natural range	See the uplands supporting document for further details
Community diversity	Abundance of variety of vegetation communities	Maintain variety of vegetation communities, subject to natural processes	Roche et al. (2014) recorded three different dry heath communities within this SAC. Data on the abundance of these communities is reproduced in the uplands supporting document. Further information on these communities is presented in Perrin et al. (2014)
Vegetation composition: lichens and bryophytes	Number of species at a representative number of 2m x 2m monitoring stops	Number of bryophyte or non-crustose lichen species present at each monitoring stop is at least three, excluding <i>Campylopus</i> and <i>Polytrichum</i> mosses	Based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation composition: positive indicator species	Number of species at a representative number of 2m x 2m monitoring stops	Number of positive indicator species present at each monitoring stop is at least two	Based on Perrin et al. (2014). The list of positive indicator species for this habitat, which is composed of dwarf shrubs, is presented in Perrin et al. (2014). See the uplands supporting document for further details
Vegetation composition: positive indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of positive indicator species at least 50% for siliceous dry heath and 50-75% for calcareous dry heath	Based on Perrin et al. (2014). The list of positive indicator species for this habitat, which is composed of dwarf shrubs, is presented in Perrin et al. (2014). See the uplands supporting document for further details
Vegetation composition: dwarf shrub composition	Percentage cover at a representative number of 2m x 2m monitoring stops	Proportion of dwarf shrub cover composed collectively of bog-myrtle (<i>Myrica gale</i>), creeping willow (<i>Salix repens</i>) and western gorse (<i>Ulex gallii</i>) is less than 50%	Based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation composition: negative indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of negative indicator species less than 1%	Based on Perrin et al. (2014). The list of negative indicator species is given in Perrin et al. (2014). See the uplands supporting document for further details
Vegetation composition: non-native species	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of non-native species less than 1%	Based on Perrin et al. (2014). See the uplands supporting document for further details. <i>Campylopus introflexus</i> was recorded within this habitat by Roche et al. (2014) but did not form extensive carpets
Vegetation structure: native trees and shrubs	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of scattered native trees and shrubs less than 20%	Based on Perrin et al. (2014). See the uplands supporting document for further details

Vegetation composition: bracken	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of bracken (<i>Pteridium aquilinum</i>) less than 10%	Based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation composition: soft rush	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of soft rush (<i>Juncus effusus</i>) less than 10%	Based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation structure: senescent ling	Percentage cover at a representative number of 2m x 2m monitoring stops	Senescent proportion of ling (<i>Calluna vulgaris</i>) cover less than 50%	Based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation structure: signs of browsing	Percentage of shoots browsed at a representative number of 2m x 2m monitoring stops	Less than 33% collectively of the last complete growing season's shoots of ericoids showing signs of browsing	Based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation structure: burning	Occurrence in local vicinity of a representative number of monitoring stops	No signs of burning in sensitive areas	Based on Perrin et al. (2014). The list of sensitive areas is presented in Perrin et al. (2014). See the uplands supporting document for further details
Vegetation structure: growth phases of ling	Percentage cover in local vicinity of a representative number of monitoring stops	Outside sensitive areas, all growth phases of ling (<i>Calluna vulgaris</i>) should occur throughout, with at least 10% of cover in the mature phase	Based on Perrin et al. (2014). See the uplands supporting document for further details
Physical structure: disturbed bare ground	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of disturbed bare ground less than 10%	Based on Perrin et al. (2014). See the uplands supporting document for further details
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat and no decline in status of hepatic mats associated with this habitat	Roche et al. (2014) compiled and mapped existing rare and notable plant records for the SAC and added any new records collected during the NSUH survey. No relevant species were recorded in this habitat, however, new records should be considered within this attribute. Hepatic mats of the <i>Calluna vulgaris-Herbertus aduncus</i> community were recorded within this habitat by Roche et al. (2014). No assessment of the conservation status of this community has been conducted but proposals for such an assessment are presented in Barron and Perrin (2014). See the uplands supporting document for further details

Conservation Objectives for : Corraun Plateau SAC [000485]

4060 Alpine and Boreal heaths

To restore the favourable conservation condition of Alpine and Boreal heaths in Corraun Plateau SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Corraun Plateau SAC was surveyed as part of the National Survey of Upland Habitats (NSUH; see Roche et al., 2014 & Perrin et al., 2014). The total current area of Alpine and Boreal heath in the SAC stated by Roche et al. (2014) is 278.3ha. It covers 7.16% of the SAC. Roche et al. (2014) report obvious losses of habitat since 1995 of approximately 0.45ha. A summary of the mapping methodology is presented in the uplands supporting document
Habitat distribution	Occurrence	No decline from current distribution, subject to natural processes. See map 5	Alpine and Boreal heath was recorded by Roche et al. (2014) on the high ground through the centre of the SAC, but was also recorded at lower elevations on the southern slopes. A summary of the mapping methodology is presented in the uplands supporting document
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil nutrient status within natural range	See the uplands supporting document for further details
Community diversity	Abundance of variety of vegetation communities	Maintain variety of vegetation communities, subject to natural processes	Roche et al. (2014) recorded three different Alpine and Boreal heath communities within this SAC. Data on the abundance of these communities is reproduced in the uplands supporting document. Further information on these communities is presented in Perrin et al. (2014)
Vegetation composition: lichens and bryophytes	Number of species at a representative number of 2m x 2m monitoring stops	Number of bryophyte or non-crustose lichen species present at each monitoring stop is at least three	Based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation composition: positive indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of positive indicator species at least 66%	Based on Perrin et al. (2014). The list of positive indicator species for this habitat is presented in Perrin et al. (2014). See the uplands supporting document for further details
Vegetation composition: dwarf-shrub species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of dwarf-shrub species at least 10%	Based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation composition: negative indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of negative indicator species less than 10%	Based on Perrin et al. (2014). The list of negative indicator species is given in Perrin et al. (2014). See the uplands supporting document for further details
Vegetation composition: non-native species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of non-native species less than 1%	Based on Perrin et al. (2014). See the uplands supporting document for further details. No non-native species were recorded within this habitat by Roche et al. (2014)
Vegetation structure: signs of grazing	Percentage of leaves browsed at a representative number of 2m x 2m monitoring stops	Less than 10% collectively of the live leaves of specific graminoids showing signs of grazing	Based on Perrin et al. (2014). See the uplands supporting document for further details including the list of specific graminoids
Vegetation structure: signs of browsing	Percentage of shoots browsed at a representative number of 2m x 2m monitoring stops	Less than 33% collectively of the last complete growing season's shoots of ericoids and crowberry (<i>Empetrum nigrum</i>) showing signs of browsing	Based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation structure: burning	Occurrence in local vicinity of a representative number of monitoring stops	No signs of burning within the habitat	Based on Perrin et al. (2014). See the uplands supporting document for further details

Physical structure: disturbed bare ground	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of disturbed bare ground less than 10%	Based on Perrin et al. (2014). See the uplands supporting document for further details
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat and no decline in status of hepatic mats associated with this habitat	Roche et al. (2014) compiled and mapped existing rare and notable plant records for the SAC and added any new records collected during the NSUH survey. <i>Cladonia rangiferina</i> has been recorded from within the habitat. This species is listed on a preparatory list of rare and threatened lichens prepared by D. McFerran, National Museums Northern Ireland. This and any new records should be considered within this attribute. Hepatic mats of the <i>Calluna vulgaris-Herbertus aduncus</i> community were recorded within this habitat by Roche et al. (2014). No assessment of the conservation status of this community has been conducted but proposals for such an assessment are presented in Barron and Perrin (2014). See the uplands supporting document for further details

Conservation Objectives for : Corraun Plateau SAC [000485]

5130 Juniperus communis formations on heaths or calcareous grasslands

To restore the favourable conservation condition of *Juniperus communis* formations on heaths or calcareous grasslands in Corraun Plateau SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	<i>Juniperus communis</i> formations on heaths or calcareous grasslands occurs in close association with heath habitats, particularly Alpine and Boreal heaths (4060). Cooper et al. (2012) recorded the habitat on the southern facing slopes of the central section of the SAC
Habitat distribution	Occurrence	No decline from current distribution, subject to natural processes. See map 6 for survey location	See notes for area above
Juniper population size	Number per formation	At least 50 plants per formation	To classify as a juniper formation, at least 50 plants should be present (Cooper et al., 2012). At least 500 plants were recorded by Cooper et al. (2012)
Vegetation composition: typical species	Number per formation	At least 50% of the listed positive indicator species for the relevant vegetation group present	Cooper et al. (2012) lists positive indicator species for five vegetation groups. The formation described at this SAC by Cooper et al. (2012) falls into vegetation group 4 (<i>Calluna vulgaris</i> / <i>Erica cinerea</i> group). See Cooper et al. (2012) for positive indicator species
Vegetation composition: negative indicator species	Occurrence per formation	Negative indicator species, particularly non-native invasive species, absent or under control	Negative indicator species listed by Cooper et al. (2012)
Vegetation structure: cone-bearing plants	Percentage per formation	At least 10% of juniper plants are bearing cones	Attribute and target based on Cooper et al. (2012)
Vegetation structure: seedling recruitment	Percentage per formation	At least 10% of juniper plants are seedlings	Attribute and target based on Cooper et al. (2012)
Vegetation structure: dead juniper	Percentage per formation	Mean percentage of each juniper plant dead less than 10%	Attribute and target based on Cooper et al. (2012)

Conservation Objectives for : Corraun Plateau SAC [000485]

8110 Siliceous scree of the montane to snow levels (*Androsacetalia alpinae* and *Galeopsietalia ladani*)

To restore the favourable conservation condition of Siliceous scree of the montane to snow levels (*Androsacetalia alpinae* and *Galeopsietalia ladani*) in Corraun Plateau SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Corraun Plateau SAC was surveyed as part of the National Survey of Upland Habitats (NSUH; see Roche et al., 2014 & Perrin et al., 2014). The total current area of siliceous scree in the SAC stated by Roche et al. (2014) is 30.3ha. This covers 0.78% of the SAC. Roche et al. (2014) report no significant losses of area since 1995. A summary of the mapping methodology is presented in the uplands supporting document
Habitat distribution	Occurrence	No decline from current distribution, subject to natural processes. See map 7	Siliceous scree was recorded by Roche et al. (2014) on the northern slopes of the SAC. A summary of the mapping methodology is presented in the uplands supporting document
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil nutrient status within natural range	See the uplands supporting document for further details
Vegetation composition: lichens and bryophytes	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of bryophytes and non-crustose lichen species at least 5%	Based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation composition: negative indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Proportion of vegetation composed of negative indicator species less than 1%	Based on Perrin et al. (2014). The list of negative indicator species is given in Perrin et al. (2014). See the uplands supporting document for further details
Vegetation composition: non-native species	Percentage cover at a representative number of 2m x 2m monitoring stops	Proportion of vegetation composed of non-native species less than 1%	Based on Perrin et al. (2014). See the uplands supporting document for further details. No non-native species were recorded within this habitat by Roche et al. (2014)
Vegetation composition: positive indicator species	Number of species in local vicinity of a representative number of monitoring stops	Number of positive indicator species present in vicinity of each monitoring stop in block scree is at least one	Based on Perrin et al. (2014). The list of positive indicator species for this habitat is presented in Perrin et al. (2014) and is the same as for 8220 Siliceous rocky slopes. Further details can be found in the uplands supporting document
Vegetation composition: grass species and dwarf shrubs	Percentage cover in local vicinity of a representative number of monitoring stops	Total cover of grass species and dwarf shrubs less than 20%	Based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation composition: bracken, native trees and scrub	Percentage cover in local vicinity of a representative number of monitoring stops	Total cover of bracken (<i>Pteridium aquilinum</i>), native trees and scrub less than 25%	Based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation structure: grazing and browsing	Percentage of leaves/shoots grazed/browsed at a representative number of 2m x 2m monitoring stops	Live leaves of forbs and shoots of dwarf shrubs showing signs of grazing or browsing collectively less than 50%	Based on Perrin et al. (2014). See the uplands supporting document for further details
Physical structure: disturbance	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Ground disturbed by human and animal paths, scree running, vehicles less than 10%	Based on Perrin et al. (2014). See the uplands supporting document for further details


Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat and no decline in status of hepatic mats associated with this habitat	Roche et al. (2014) compiled and mapped existing rare and notable plant records for the SAC and added any new records collected during the NSUH survey. No relevant species were recorded in this habitat, however, new records should be considered within this attribute. Hepatic mats of the <i>Calluna vulgaris-Herbertus aduncus</i> community were recorded within this habitat by Roche et al. (2014). No assessment of the conservation status of this community has been conducted but proposals for such an assessment are presented in Barron and Perrin (2014). See the uplands supporting document for further details
-------------------------------------	--------------------------------	--	---

Conservation Objectives for : Corraun Plateau SAC [000485]


8220 Siliceous rocky slopes with chasmophytic vegetation


To restore the favourable conservation condition of Siliceous rocky slopes with chasmophytic vegetation in Corraun Plateau SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Corraun Plateau SAC was surveyed as part of the National Survey of Upland Habitats (NSUH; see Roche et al., 2014 & Perrin et al., 2014). The total current area of siliceous rocky slopes in the SAC stated by Roche et al. (2014) is 15.0ha. This covers 0.39% of the SAC. A summary of the mapping methodology is presented in the uplands supporting document
Habitat distribution	Occurrence	No decline from current distribution, subject to natural processes. See map 8	Siliceous rocky slopes were recorded by Roche et al. (2014) with patches on the northern slopes and also in the south. A summary of the mapping methodology is presented in the uplands supporting document
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil nutrient status within natural range	See the uplands supporting document for further details
Vegetation composition: positive indicator species	Number of species at a representative number of monitoring stops	Number of positive indicator species present in vicinity of each monitoring stop is at least one	Based on Perrin et al. (2014). The list of positive indicator species for this habitat is presented in Perrin et al. (2014). Further details can be found in the uplands supporting document
Vegetation composition: non-native species	Percentage cover in local vicinity of a representative number of monitoring stops	Proportion of vegetation composed of non-native species less than 1%	Based on Perrin et al. (2014). See the uplands supporting document for further details. No non-native species were recorded within this habitat by Roche et al. (2014)
Vegetation composition: bracken, native trees and scrub	Percentage cover in local vicinity of a representative number of monitoring stops	Total cover of bracken (<i>Pteridium aquilinum</i>), native trees and scrub less than 25%	Based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation structure: grazing and browsing	Percentage of leaves/shoots grazed/browsed in local vicinity of a representative number of monitoring stops	Live leaves of forbs and shoots of dwarf shrubs showing signs of grazing or browsing collectively less than 50%	Based on Perrin et al. (2014). See the uplands supporting document for further details
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat and no decline in status of hepatic mats associated with this habitat	Roche et al. (2014) compiled and mapped existing rare and notable plant records for the SAC and added any new records collected during the NSUH survey. No relevant species were recorded in this habitat, however, new records should be considered within this attribute. See the uplands supporting document for further details




Legend

 Corraun Plateau SAC 000485



Legend

- Corraun Plateau SAC 000485
- Owenduff / Nephin Complex SAC 000534





Legend

Corraun Plateau SAC 000485

Cover of 4010 Northern Atlantic wet heaths with *Erica tetralix*


- 0%
- 0.1% - 20%
- 20.1% - 40%
- 40.1% - 60%
- 60.1% - 80%
- 80.1% - 100%






Legend

- Corraun Plateau SAC 000485
- OSi Discovery Series County Boundary
- Cover of 4060 Alpine and Boreal heaths**
- 0%
- 0.1% - 20%
- 20.1% - 40%
- 40.1% - 60%
- 60.1% - 80%
- 80.1% - 100%



Legend

- Corraun Plateau SAC 000485
- ★ 5130 *Juniperus communis* formations on heaths or calcareous grasslands Survey Sites




Legend

- Corraun Plateau SAC 000485
- OSi Discovery Series County Boundary

Cover of 8110 Siliceous scree of the montane to snow levels (*Androsacetalia alpinae* and *Galeopsietalia ladani*)

- 0%
- 0.1% - 20%
- 20.1% - 40%
- 40.1% - 60%
- 60.1% - 80%
- 80.1% - 100%



Legend

- Corraun Plateau SAC 000485
- OSi Discovery Series County Boundary

Cover of 8220 Siliceous rocky slopes with chasmophytic vegetation

- 0%
- 0.1% - 20%
- 20.1% - 40%
- 40.1% - 60%
- 60.1% - 80%
- 80.1% - 100%



Conservation objectives for Doocastle Turlough SAC [000492]

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Objective: To maintain or restore the favourable conservation condition of the Annex I habitat(s) and/or the Annex II species for which the SAC has been selected:

Code	Description
------	-------------

3180	Turloughs*
------	------------

* denotes a priority habitat



Citation: NPWS (2020) Conservation objectives for Doocastle Turlough SAC [000492]. Generic Version 7.0. Department of Culture, Heritage and the Gaeltacht.

National Parks and Wildlife Service

Conservation Objectives Series

Duvillaun Islands SAC 000495



An Roinn
Ealaíon, Oidhreachta agus Gaeltachta

Department of
Arts, Heritage and the Gaeltacht



**National Parks and Wildlife Service,
Department of Arts, Heritage and the Gaeltacht,**

7 Ely Place, Dublin 2, Ireland.

Web: www.npws.ie

E-mail: nature.conservation@ahg.gov.ie

Citation:

**NPWS (2013) Conservation Objectives: Duvillaun Islands SAC 000495. Version 1.
National Parks and Wildlife Service, Department of Arts, Heritage and the
Gaeltacht.**

Series Editor: Rebecca Jeffrey

ISSN 2009-4086

Introduction

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

** indicates a priority habitat under the Habitats Directive*

000495 Duvillaun Islands SAC

1364 Grey seal *Halichoerus grypus*

Please note that this SAC overlaps with Duvillaun Islands SPA (004111) and adjoins West Connacht Coast SAC (002998). See map 2. The conservation objectives for this site should be used in conjunction with those for overlapping and adjacent sites as appropriate.

Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

NPWS Documents

Year :	2003
Title :	Grey seal population status at islands in the Inishkea group, as determined from breeding ground surveys in 2002
Author :	O Cadhla, O.; Strong, D.
Series :	Unpublished report to NPWS
Year :	2004
Title :	Harbour seal population assessment in the Republic of Ireland: August 2003
Author :	Cronin, M.; Duck, C.; O Cadhla, O.; Nairn, R.; Strong, D.; O'Keefe, C.
Series :	Irish Wildlife Manual No. 11
Year :	2004
Title :	Summary of National Parks and Wildlife Service surveys for common (harbour) seals (<i>Phoca vitulina</i>) and grey seals (<i>Halichoerus grypus</i>), 1978 to 2003
Author :	Lyons, D.O.
Series :	Irish Wildlife Manual No. 13
Year :	2007
Title :	Grey seal moult population survey in the Republic of Ireland, 2007
Author :	O Cadhla, O.; Strong, D.
Series :	Unpublished report to NPWS
Year :	2008
Title :	An assessment of the breeding population of grey seals in the Republic of Ireland, 2005
Author :	O Cadhla, O.; Strong, D.; O'Keefe, C.; Coleman, M.; Cronin, M.; Duck, C.; Murray, T.; Dower, P.; Nairn, R.; Murphy, P.; Smiddy, P.; Saich, C.; Lyons, D.O.; Hiby, L.
Series :	Irish Wildlife Manual No. 34
Year :	2013
Title :	Duvillaun Islands SAC (site code: 495) Conservation objectives supporting document- marine species V1
Author :	NPWS
Series :	Conservation objectives supporting document

Other References

Year :	1998
Title :	Population biology of grey seals (<i>Halichoerus grypus</i> , Fabricius 1791) in western Ireland
Author :	Kiely, O.R.M.
Series :	Unpublished PhD thesis, National University of Ireland, University College Cork
Year :	1998
Title :	Grey seal (<i>Halichoerus grypus</i>) pup production at the Inishkea island group, Co. Mayo and the Blasket Islands, Co. Kerry
Author :	Kiely, O.; Myers, A.A.
Series :	Biology and Environment: Proc. Royal Ir. Acad. 98B (2): 113-122
Year :	2001
Title :	Grey seal interactions with fisheries in Irish coastal waters
Author :	BIM
Series :	Report to the European Commission DG XIV. Study 95/40

Year : 2007
Title : Aerial surveying of grey seal breeding colonies on the Blasket Islands, Co. Kerry, the Inishkeas group, Co. Mayo and the Donegal coast, Ireland
Author : Cronin, M.A.; Duck, C.D.; O Cadhla, O.
Series : J. Nat. Conserv. 15(2): 77-83

Spatial data sources


Year : 2005
Title : OSi Discovery series vector data
GIS Operations : High Water Mark (HWM) polyline feature class converted into polygon feature class; clipped to SAC boundary. Expert opinion used as necessary to resolve any issues arising
Used For : 1364 (map 3)


Year : 2013
Title : NPWS rare and threatened species database
GIS Operations : Dataset created from spatial references in database records. Expert opinion used as necessary to resolve any issues arising
Used For : 1364 (map 3)

1364 Grey seal *Halichoerus grypus*

To maintain the favourable conservation condition of Grey Seal in Duvillaun Islands SAC, which is defined by the following list of attributes and targets:


Attribute	Measure	Target	Notes
Access to suitable habitat	Number of artificial barriers	Species range within the site should not be restricted by artificial barriers to site use. See map 3	See marine supporting document for further details
Breeding behaviour	Breeding sites	Conserve the breeding sites in a natural condition. See map 3 for known sites	Attribute and target based on background knowledge of Irish breeding populations, comprehensive breeding surveys in 1995 (Kiely, 1998; Kiely and Myers, 1998), 1998 and 1999 (BIM, 2001), 2002 (Ó Cadhla and Strong, 2003) and 2005 (Ó Cadhla et al., 2008) and unpublished NPWS records including those reported by Lyons (2004). See marine supporting document for further details
Moulting behaviour	Moult haul-out sites	Conserve the moult haul-out sites in a natural condition. See map 3 for known sites	Attribute and target based on background knowledge of Irish breeding populations, review of data from Kiely (1998) and Lyons (2004), a national moult survey (Ó Cadhla and Strong, 2007) and unpublished NPWS records. See marine supporting document for further details
Resting behaviour	Resting haul-out sites	Conserve the resting haul-out sites in a natural condition. See map 3 for known sites	Attribute and target based on review data from Kiely (1998), BIM (2001), Lyons (2004), Cronin et al. (2004), Ó Cadhla et al. (2008) and unpublished NPWS records. See marine supporting document for further details
Disturbance	Level of impact	Human activities should occur at levels that do not adversely affect the grey seal population at the site	See marine supporting document for further details





Legend

- Duvillaun Islands SAC 000495
- West Connacht Coast SAC 002998
- Duvillaun Islands SPA 004111
- OSi Discovery Series County Boundaries



Legend

- Duvillaun Islands SAC 000495
- ★ 1364 Grey Seal - *Halichoerus grypus* breeding sites
- ★ 1364 Grey Seal - *Halichoerus grypus* moulting sites
- ★ 1364 Grey Seal - *Halichoerus grypus* resting sites
- 1364 Grey Seal - *Halichoerus grypus* habitat
- OSi Discovery Series County Boundaries



National Parks and Wildlife Service

Conservation Objectives Series

Flughany Bog SAC 000497



*An Roinn
Ealaíon, Oidhreachta agus Gaeltachta*

*Department of
Arts, Heritage and the Gaeltacht*



**National Parks and Wildlife Service,
Department of Arts, Heritage and the Gaeltacht,
7 Ely Place, Dublin 2, Ireland.
Web: www.npws.ie
E-mail: nature.conservation@ahg.gov.ie**

Citation:

**NPWS (2016) Conservation Objectives: Flughany Bog SAC 000497. Version 1.
National Parks and Wildlife Service, Department of Arts, Heritage and the
Gaeltacht.**

**Series Editor: Rebecca Jeffrey
ISSN 2009-4086**

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

** indicates a priority habitat under the Habitats Directive*

000497	Flughany Bog SAC
7110	Active raised bogsE
7120	Degraded raised bogs still capable of natural regeneration
7150	Depressions on peat substrates of the Rhynchosporion

Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

NPWS Documents

Year :	2014
Title :	Raised Bog Monitoring and Assessment Survey 2013
Author :	Fernandez, F.; Connolly K.; Crowley W.; Denyer J.; Duff K.; Smith G.
Series :	Irish Wildlife Manual No. 81
<hr/>	
Year :	2014
Title :	National raised bog SAC management plan
Author :	Department of Arts, Heritage and the Gaeltacht
Series :	Draft for consultation. 15 January 2014
<hr/>	
Year :	2014
Title :	Flughany Bog (SAC 000497), Co. Sligo, Site Report
Author :	Fernandez, F.; Connolly, K.; Crowley, W.; Denyer J.; Duff K.; Smith G.
Series :	Raised bog monitoring and assessment survey 2013
<hr/>	
Year :	2016
Title :	Flughany Bog SAC (site code: 497) Conservation objectives supporting document- raised bog habitats V1
Author :	NPWS
Series :	Conservation objectives supporting document

Other References

Year :	2011
Title :	Review and revision of empirical critical loads and dose-response relationships. Proceedings of an expert workshop, Noordwijkerhout, 23-25 June 2010
Author :	Bobbink, R.; Hettelingh, J.P.
Series :	RIVM report 680359002, Coordination Centre for Effects, National Institute for Public Health and the Environment (RIVM)
<hr/>	
Year :	2014
Title :	Nitrogen deposition and exceedance of critical loads for nutrient nitrogen in Irish grasslands
Author :	Henry, J.; Aherne, J.
Series :	Science of the Total Environment 470–471: 216–223

Spatial data sources

Year :	2014
Title :	Scientific Basis for Raised Bog Conservation in Ireland
GIS Operations :	RBSB13_SACs_ARB_DRB dataset, RBSB13_SACs_2012_HB dataset, RBSB13_SACs_DrainagePatterns_5k dataset and RBSB13_SAC_LIDAR_DTMs dataset clipped to SAC boundary. Expert opinion used as necessary to resolve any issues arising potential
Used For :	7110; digital elevation model; drainage patterns (maps 2 and 4)
<hr/>	
Year :	2013
Title :	Raised Bog Monitoring and Assessment Survey 2013
GIS Operations :	RBMA13_ecotope_map dataset clipped to SAC boundary. Appropriate ecotopes selected and exported to new dataset. Expert opinion used as necessary to resolve any issues arising
Used For :	7110 ecotopes (map 3)
<hr/>	

Conservation Objectives for : Flughany Bog SAC [000497]

7110 Active raised bogs

To restore the favourable conservation condition of Active raised bogs in Flughany Bog SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Restore area of active raised bog to 23.6ha, subject to natural processes	Active Raised Bog (ARB) habitat was mapped at 11.4ha by Fernandez et al. (2014). Area of Degraded Raised Bog (DRB) on the High Bog (HB) has been modelled as 13.1ha. See map 2. However, it is estimated that only 9.2ha is potentially restorable to ARB by drain blocking. The total potential ARB on the HB is therefore estimated to be 20.6ha. Eco-hydrological assessments of the cutover estimates that an additional 3.0ha of bog forming habitats could be restored. The long term target for ARB is therefore 23.6ha. See raised bog supporting document for further details on this and following attributes
Habitat distribution	Occurrence	Restore the distribution and variability of active raised bog across the SAC. See map 3 for distribution in 2012	ARB habitat at Flughany Bog is central and sub-central ecotopes and active flush. ARB currently occurs most abundantly on the south-eastern part of Flughany Bog. DRB occurs on both parts of the bog, which will require restoration measures. There is also potential for ARB restoration on cutover areas of the bog (see area target above)
High bog area	Hectares	No decline in extent of high bog necessary to support the development and maintenance of active raised bog. See map 2	The area of high bog within Flughany Bog SAC in 2012 (latest figure available) was 143.7ha (DAHG 2014)
Hydrological regime: water levels	Centimetres	Restore appropriate water levels throughout the site	For ARB, mean water level needs to be near or above the surface of the bog lawns for most of the year. Seasonal fluctuations should not exceed 20cm, and should only be 10cm below the surface, except for very short periods of time. Open water is often characteristic of soak systems
Hydrological regime: flow patterns	Flow direction; slope	Restore, where possible, appropriate high bog topography, flow directions and slopes. See map 4 for current situation	ARB depends on mean water levels being near or above the surface of bog lawns for most of the year. Long and gentle slopes are the most favourable to achieve these conditions. Changes to flow directions due to subsidence of bogs can radically change water regimes and cause drying out of high quality ARB areas and soak systems
Transitional areas between high bog and adjacent mineral soils (including cutover areas)	Hectares; distribution	Restore adequate transitional areas to support/protect active raised bog and the services it provides	No natural marginal habitats exist around the margins of the bog. Eco-hydrological assessments have evaluated the potential for ARB restoration on cutover areas (see note for habitat area attribute above)
Vegetation quality: central ecotope, active flush, soaks, bog woodland	Hectares	Restore 11.8ha of central ecotope/active flush/soaks/bog woodland as appropriate	At least 50% of ARB habitat should be high quality (i.e. central ecotope, active flush, soaks, bog woodland). Target area of active raised bog for the site has been set at 23.6ha (see area target above)
Vegetation quality: microtopographical features	Hectares	Restore adequate cover of high quality microtopographical features	High quality microtopography (hummocks, hollows and pools) is well developed in the southern part of Flughany Bog
Vegetation quality: bog moss (<i>Sphagnum</i>) species	Percentage cover	Restore adequate cover of bog moss (<i>Sphagnum</i>) species to ensure peat-forming capacity	<i>Sphagnum</i> cover varies naturally across Ireland with relatively high cover in the east to lower cover in the west. Hummock forming species such as <i>Sphagnum austini</i> are particularly good peat formers. <i>Sphagnum</i> cover and distribution also varies naturally across a site

Typical ARB species: flora	Occurrence	Restore, where appropriate, typical active raised bog flora	Typical flora species include widespread species, as well as those with more restricted distributions but typical of the habitat's subtypes or geographical range
Typical ARB species: fauna	Occurrence	Restore, where appropriate, typical active raised bog fauna	Typical fauna species include widespread species, as well as those with more restricted distributions but typical of the habitat's subtypes or geographical range
Elements of local distinctiveness	Occurrence	Maintain features of local distinctiveness, subject to natural processes	Flughany Bog is noted for the presence of a number of flush systems and associated swallow-holes
Negative physical indicators	Percentage cover	Negative physical features absent or insignificant	Negative physical indicators include: bare peat, algae dominated pools and hollows, marginal cracks, tear patterns, subsidence features such as dry mineral mounds /ridges emerging or expanding and evidence of burning
Vegetation composition: native negative indicator species	Percentage cover	Native negative indicator species at insignificant levels	Disturbance indicators include species indicative of conditions drying out such as abundant bog asphodel (<i>Narthecium ossifragum</i>), deergrass (<i>Trichophorum germanicum</i>) and harestalk cotton-grass (<i>Eriophorum vaginatum</i>) forming tussocks; abundant magellanic bog-moss (<i>Sphagnum magellanicum</i>) in pools previously dominated by <i>Sphagnum</i> species typical of very wet conditions (e.g. feathery bog-moss (<i>S. cuspidatum</i>)); and indicators of frequent burning events such as abundant <i>Cladonia floerkeana</i> and high cover of carnation sedge (<i>Carex panicea</i>) (particularly in true midlands raised bogs)
Vegetation composition: non-native invasive species	Percentage cover	Non-native invasive species at insignificant levels and not more than 1% cover	Most common non-native invasive species include lodgepole pine (<i>Pinus contorta</i>), rhododendron (<i>Rhododendron ponticum</i>), and pitcherplant (<i>Sarracenia purpurea</i>)
Air quality: nitrogen deposition	kg N/ha/year	Air quality surrounding bog close to natural reference conditions. The total N deposition should not exceed 5kg N/ha/yr	Change in air quality can result from fertiliser drift; adjacent quarry activities; or other atmospheric inputs. The critical load range for ombrotrophic bogs has been set as between 5 and 10kg N/ha/yr (Bobbink and Hettelingh, 2011). The latest N deposition figures for the area around Flughany Bog suggests that the current level is approximately 8.2kg N/ha/yr (Henry and Aherne, 2014)
Water quality	Hydrochemical measures	Water quality on the high bog and in transitional areas close to natural reference conditions	Water chemistry within raised bogs is influenced by atmospheric inputs (rainwater). However, within soak systems, water chemistry is influenced by other inputs such as focused flow or interaction with underlying substrates. Water chemistry in areas surrounding the high bog varies due to influences of different water types (bog water, regional groundwater, and run-off from surrounding mineral lands)

Conservation Objectives for : Flughany Bog SAC [000497]

7120 Degraded raised bogs still capable of natural regeneration

The long-term aim for Degraded raised bogs still capable of natural regeneration is that its peat-forming capability is re-established; therefore, the conservation objective for this habitat is inherently linked to that of Active raised bogs (7110) and a separate conservation objective has not been set in Flughany Bog SAC


Attribute	Measure	Target	Notes

Conservation Objectives for : Flughany Bog SAC [000497]

7150 Depressions on peat substrates of the Rhynchosporion


Depressions on peat substrates of the Rhynchosporion is an integral part of good quality Active raised bogs (7110) and thus a separate conservation objective has not been set for the habitat in Flughany Bog SAC

Attribute	Measure	Target	Notes
-----------	---------	--------	-------



Legend

 Flughany Bog SAC 000497

 *An Roinn Ealaíon, Oidhreacht agus Gaeltachta*
 Department of Arts, Heritage and the Gaeltacht

**MAP 1:
 FLUGHANY BOG SAC
 CONSERVATION OBJECTIVES
 SAC DESIGNATION**

Map to be read in conjunction with the NPWS Conservation Objectives Document.


**SITE CODE:
 SAC 000497; version 3.01. Co. Mayo / Sligo**

0 100 200 300 400 500 m

The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
 Ordnance Survey of Ireland Licence No EN 0059216. © Ordnance Survey of Ireland Government of Ireland.


Níl sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbheithniú a déanamh ar theorainneacha na gceantar comharthaíthe. Suirbhéarachtá Ordonáis na hÉireann Ceadúnas Uimh EN 0059216. © Suirbhéarachtá Ordonáis na hÉireann Rialtas na hÉireann.



**Map Version 1
 Date: Jan 2016**



Legend


- Flughany Bog SAC 000497
- High Bog Boundary
- Potential 7110 *Active Raised Bogs
- OSi Discovery Series County Boundary





Legend



- Flughany Bog SAC 000497
- High Bog Boundary
- OSi Discovery Series County Boundary
- Active Raised Bogs Ecotopes**
- Central ecotope
- Soaks / active flush
- Sub-central ecotope



Legend

- Flughany Bog SAC 000497
- OSi Discovery Series County Boundary
- High Bog Boundary
- Drainage Patterns
- Contours

Elevation



National Parks and Wildlife Service

Conservation Objectives Series

Glenamoy Bog Complex SAC 000500



An Roinn Ealaíon, Oidhreachta,
Gnóthaí Réigiúnacha, Tuaithe agus Gaeltachta

Department of Arts, Heritage,
Regional, Rural and Gaeltacht Affairs



**National Parks and Wildlife Service,
Department of Arts, Heritage, Regional, Rural and Gaeltacht Affairs,
7 Ely Place, Dublin 2, Ireland.
Web: www.npws.ie
E-mail: nature.conservation@ahg.gov.ie**

Citation:

**NPWS (2017) Conservation Objectives: Glenamoy Bog Complex SAC 000500.
Version 1. National Parks and Wildlife Service, Department of Arts, Heritage,
Regional, Rural and Gaeltacht Affairs.**

**Series Editor: Rebecca Jeffrey
ISSN 2009-4086**

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

* indicates a priority habitat under the Habitats Directive

000500	Glenamoy Bog Complex SAC
1106	Salmon <i>Salmo salar</i>
1230	Vegetated sea cliffs of the Atlantic and Baltic coasts
1393	Slender Green Feather-moss <i>Drepanocladus vernicosus</i>
1395	Petalwort <i>Petalophyllum ralfsii</i>
1528	Marsh Saxifrage <i>Saxifraga hirculus</i>
21A0	Machairs (* in Ireland)
3160	Natural dystrophic lakes and ponds
4010	Northern Atlantic wet heaths with <i>Calluna vulgaris</i>
5130	<i>Rhynchospora</i> formations on heaths or calcareous grasslands
7130	Blanket bogs (* if active bog)
7140	Transition mires and quaking bogs
7150	Depressions on peat substrates of the Rhynchosporion

Please note that this SAC overlaps with Blacksod Bay/Broad Haven SPA (004037) and Illanmaster SPA (004074) and is adjacent to Broadhaven Bay SAC (000472) and Slieve Fyagh Bog SAC (000542). See map 2. The conservation objectives for this site should be used in conjunction with those for the overlapping and adjacent sites as appropriate.

Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

NPWS Documents

Year :	1983
Title :	Report on the conservation value of Irish coastal sites: Machair in Ireland
Author :	Bassett, A.
Series :	Unpublished Report
Year :	1987
Title :	A survey to locate blanket bogs of scientific interest in County Mayo. Part I
Author :	Foss, P.; McGee, E.
Series :	A report commissioned by the Wildlife Service
Year :	1989
Title :	Survey to locate blanket bogs of scientific interest in Mayo. Part II
Author :	Douglas, C.; Garvey, L.; Kelly, L.; O'Sullivan, A.; Van Doorsleer, L.
Series :	A report commissioned by the Wildlife Service
Year :	1998
Title :	Biomar survey of Irish machair sites 1996
Author :	Crawford, I.; Bleasdale, A.; Conaghan, J.
Series :	Irish Wildlife Manual No. 3
Year :	2009
Title :	Coastal Monitoring Project 2004-2006
Author :	Ryle, T.; Murray, A.; Connolly, K.; Swann, M.
Series :	Unpublished report to NPWS
Year :	2011
Title :	National survey and assessment of the conservation status of Irish sea cliffs
Author :	Barron, S.J.; Delaney, A.; Perrin, P.M.; Martin, J.; O'Neill, F.
Series :	Irish Wildlife Manual No. 53
Year :	2012
Title :	The conservation status of juniper formations in Ireland
Author :	Cooper, F.; Stone, R.E.; McEvoy, P.; Wilkins, T.; Reid, N.
Series :	Irish Wildlife Manual No. 63
Year :	2012
Title :	Ireland Red List No. 8: Bryophytes
Author :	Lockhart, N.; Hodgetts, N.; Holyoak, D.
Series :	Ireland Red List series, NPWS
Year :	2013
Title :	Monitoring survey of Annex I sand dune habitats in Ireland
Author :	Delaney, A.; Devaney, F.M.; Martin, J.M.; Barron, S.J.
Series :	Irish Wildlife Manual No. 75
Year :	2013
Title :	The status of EU protected habitats and species in Ireland. Volume 2. Habitats assessments
Author :	NPWS
Series :	Conservation assessments

Year :	2013
Title :	The status of EU protected habitats and species in Ireland. Volume 3. Species assessments
Author :	NPWS
Series :	Conservation assessments
Year :	2014
Title :	Guidelines for a national survey and conservation assessment of upland vegetation and habitats in Ireland, Version 2.0
Author :	Perrin, P.M.; Barron, S.J.; Roche, J.R.; O'Hanrahan, B.
Series :	Irish Wildlife Manual No. 79
Year :	2015
Title :	Monitoring recommendations for Marsh Saxifrage (<i>Saxifraga hirculus</i> L.) in the Republic of Ireland
Author :	Muldoon, C.S.; Waldren, S.; Lynn, D.
Series :	Irish Wildlife Manual No. 88
Year :	2015
Title :	Habitats Directive Annex I lake habitats: a working interpretation for the purposes of site-specific conservation objectives and Article 17 reporting
Author :	O Connor, Á.
Series :	Unpublished document by NPWS
Year :	2015
Title :	Monitoring methods for <i>Petalophyllum ralfsii</i> (Wils.) Nees & Gottsche (Petalwort) in the Republic of Ireland
Author :	Campbell, C.; Hodgetts, N.; Lockhart, N.
Series :	Irish Wildlife Manual No. 90
Year :	2015
Title :	Monitoring methods for <i>Hamatocaulis vernicosus</i> (Mitt.) Hedenäs (Slender green feather-moss) in the Republic of Ireland
Author :	Campbell, C.; Hodgetts, N.; Lockhart, N.
Series :	Irish Wildlife Manual No. 91
Year :	2016
Title :	Ireland Red List No. 10: Vascular Plants
Author :	Wyse Jackson, M.; FitzPatrick, Ú.; Cole, E.; Jebb, M.; McFerran, D.; Sheehy Skeffington, M.; Wright, M.
Series :	Ireland Red Lists series, NPWS
Year :	2017
Title :	Glenamoy Bog Complex SAC (site code: 500) Conservation objectives supporting document-blanket bogs and associated habitats V1
Author :	NPWS
Series :	Conservation objectives supporting document
Year :	2017
Title :	Glenamoy Bog Complex SAC (site code: 500) Conservation objectives supporting document-coastal habitats V1
Author :	NPWS
Series :	Conservation objectives supporting document

Other References

Year :	1982
Title :	Eutrophication of waters. Monitoring assessment and control
Author :	OECD
Series :	OECD, Paris

Year :	1989
Title :	The genera <i>Scorpidium</i> and <i>Hamatocaulis</i> , gen. nov., in northern Europe
Author :	Hedenäs, L.
Series :	Lindbergia, 15: 8-36
Year :	1990
Title :	Ecology and conservation of Irish peatlands
Author :	Doyle, G.J. (Ed.)
Series :	Royal Irish Academy, Dublin
Year :	2000
Title :	Colour in Irish lakes
Author :	Free, G.; Allott, N.; Mills, P.; Kennelly, C.; Day, S.
Series :	Verhandlungen Internationale Vereinigung für theoretische und angewandte Limnologie, 27: 2620-2623
Year :	2005
Title :	National inventory of sea cliffs and coastal heaths
Author :	Browne, A.
Series :	Unpublished Report to NPWS
Year :	2006
Title :	The vegetation of Irish machair
Author :	Gaynor, K.
Series :	Biology and Environment: Proceedings of the Royal Irish Academy, vol 106B, No. 3: 311-321
Year :	2006
Title :	A reference-based typology and ecological assessment system for Irish lakes. Preliminary investigations. Final report. Project 2000-FS-1-M1 Ecological assessment of lakes pilot study to establish monitoring methodologies EU (WFD)
Author :	Free, G.; Little, R.; Tierney, D.; Donnelly, K.; Coroni, R.
Series :	EPA, Wexford
Year :	2008
Title :	Water Quality in Ireland 2004-2006
Author :	Clabby, K.J.; Bradley, C.; Craig, M.; Daly, D.; Lucey, J.; McGarrigle, M.; O'Boyle, S.; Tierney, D.; Bowman, J.
Series :	EPA, Wexford
Year :	2010
Title :	Water quality in Ireland 2007-2009
Author :	McGarrigle, M.; Lucey, J.; Ó Cinnéide, M.
Series :	EPA, Wexford
Year :	2011
Title :	Conservation biology of <i>Saxifraga hirculus</i> L. in Ireland
Author :	Muldoon, C.S.
Series :	Unpublished Ph.D. Thesis, Trinity College Dublin
Year :	2012
Title :	The impact of conifer plantation forestry on the ecology of peatland lakes
Author :	Drinan, T.J.
Series :	Unpublished Ph.D. thesis, University College Cork
Year :	2013
Title :	Conservation of selected legally protected and Red Listed bryophytes in Ireland
Author :	Campbell, C.
Series :	Unpublished Ph.D. Thesis, Trinity College Dublin

Year :	2013
Title :	Interpretation manual of European Union habitats- Eur 28
Author :	European Commission- DG Environment
Series :	European Commission
<hr/>	
Year :	2015
Title :	Water quality in Ireland 2010-2012
Author :	Bradley, C.; Byrne, C.; Craig, M.; Free, G.; Gallagher, T.; Kennedy, B.; Little, R.; Lucey, J.; Mannix, A.; McCreesh, P.; McDermott, G.; McGarrigle, M.; Ní Longphuirt, S.; O'Boyle, S.; Plant, C.; Tierney, D.; Trodd, W.; Webster, P.; Wilkes, R.; Wynne, C.
Series :	EPA, Wexford
<hr/>	
Year :	2016
Title :	The Status of Irish Salmon Stocks in 2015 with Precautionary Catch Advice for 2016
Author :	SSCS (Standing Scientific Committee on Salmon)
Series :	Independent Scientific Report to Inland Fisheries Ireland
<hr/>	

Spatial data sources

Year :	2011
Title :	National survey and assessment of the conservation status of Irish sea cliffs
GIS Operations :	Clipped to SAC boundary
Used For :	1230 (map 3)
<hr/>	
Year :	2013
Title :	Sand Dune Monitoring Project 2011. Version 1
GIS Operations :	QIs selected; clipped to SAC boundary. Expert opinion used as necessary to resolve any issues arising
Used For :	21A0 (map 3)
<hr/>	
Year :	2008
Title :	OSi 1:5000 IG vector dataset
GIS Operations :	WaterPolygons feature class clipped to the SAC boundary. Expert opinion used to identify Annex I habitat and to resolve any issues arising
Used For :	3160 (map 4)
<hr/>	
Year :	2017
Title :	NPWS rare and threatened species database
GIS Operations :	Dataset created from spatial references in database records. Expert opinion used as necessary to resolve any issues arising
Used For :	1393, 1395, 1528 (maps 5 and 6)
<hr/>	

1230 Vegetated sea cliffs of the Atlantic and Baltic coasts

To maintain the favourable conservation condition of Vegetated sea cliffs of the Atlantic and Baltic coasts in Glenamoy Bog Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat length	Kilometres	Area stable, subject to natural processes, including erosion. For the sub-site mapped (Glinsk), total length of cliff sections: 27.93km. See map 3	Based on data from the Irish Sea Cliff Survey (ISCS) (Barron et al., 2011). Cliffs are linear features and are therefore measured in kilometres. The sub-site Glinsk (ISCS site ID: 08005) was identified using a combination of aerial photos and the DCENR helicopter viewer. The length of cliff was measured (in sections) to give a total estimated area of 27.93km within the SAC. The length of cliff is likely to be underestimated. See the Glenamoy Bog Complex SAC conservation objectives supporting document for coastal habitats for further details
Habitat distribution	Occurrence	No decline, subject to natural processes. See map 3	Sea cliffs are known to occur along the coastline and extend along the north coast ranging in height from 20m to 253m, at Benwee Head. Hard cliffs have been noted in this SAC and it is estimated that all of the cliffs are of the hard type (Browne, 2005; Barron et al., 2011). See the coastal habitats supporting document for further details
Physical structure: functionality and hydrological regime	Occurrence of artificial barriers	No alteration to natural functioning of geomorphological and hydrological processes, including groundwater quality, due to artificial structures	Based on data from Barron et al. (2011). Maintaining natural geomorphological processes, including natural erosion, is important for the health of vegetated sea cliffs. Hydrological processes maintain flushes that can be associated with sea cliffs. Hydrological features such as gullies, streams and cascades were identified by the ISCS as occurring on the cliffs in Glenamoy Bog Complex SAC. See the coastal habitats supporting document for further details
Vegetation structure: zonation	Occurrence	Maintain range of sea cliff habitat zonation including transitional zones, subject to natural processes including erosion and succession	Based on data from Barron et al. (2011). Cliff top vegetation, which ranges from <i>Plantago</i> -dominated sward to coastal heath and maritime grassland, occurs adjacent to sea cliff vegetation in Glenamoy Bog Complex SAC. See the coastal habitats supporting document for further details
Vegetation structure: vegetation height	Centimetres	Maintain structural variation within sward	Based on data from Barron et al. (2011). See the coastal habitats supporting document for further details
Vegetation composition: typical species and sub-communities	Percentage cover at a representative number of monitoring stops	Maintain range of sub-communities with typical species listed in the Irish Sea Cliff Survey (Barron et al., 2011)	See the coastal habitats supporting document for further details
Vegetation composition: negative indicator species	Percentage	Negative indicator species (including non-native species) to represent less than 5% cover	Based on data from Barron et al. (2011). See the coastal habitats supporting document for further details
Vegetation composition: bracken and woody species	Percentage	Cover of bracken (<i>Pteridium aquilinum</i>) on grassland and/or heath less than 10%. Cover of woody species on grassland and/or heath less than 20%	Based on data from Barron et al. (2011). See the coastal habitats supporting document for further details

Conservation Objectives for : Glenamoy Bog Complex SAC [000500]

21A0 Machairs (* in Ireland)

To restore the favourable conservation condition of Machairs (* in Ireland) in Glenamoy Bog Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes including erosion and succession. For the sub-site mapped: Garter Hill - 234.0ha. See map 3	Based on data from the Sand Dunes Monitoring Project (SDM) (Delaney et al., 2013). Machair habitat was recorded and mapped at the sub-site Garter Hill (SDM site ID: 128) to give a total estimated area of 234.0ha within Glenamoy Bog Complex SAC. See the Glenamoy Bog Complex SAC conservation objectives supporting document for coastal habitats for further details
Habitat distribution	Occurrence	No decline or change in habitat distribution, subject to natural processes. See map 3 for known distribution	Based on data from Ryle et al. (2009) and Delaney et al. (2013). Machair occurs on the western coast of Glenamoy Bog Complex SAC. South of Benwee Head, the rocky coastline grades into an estuarine system which contains the machair system. The machair extends upslope to cover much of Garter Hill (158m) and is the highest recorded in Ireland. See the coastal habitats supporting document for further details
Physical structure: functionality and sediment supply	Presence/absence of physical barriers	Maintain the natural circulation of sediment and organic matter, without any physical obstructions	Based on data from Ryle et al. (2009) and Delaney et al. (2013). Physical barriers can lead to fossilisation or over-stabilisation of dunes, as well as beach starvation resulting in increased rates of erosion. There are some coastal protection (rock armour) works adjacent to the machair in the SAC, but these do not appear to have caused any change in erosion patterns. See the coastal habitats supporting document for further details
Physical structure: hydrological and flooding regime	Water table levels; groundwater fluctuations (metres)	Maintain natural hydrological regime	Based on data from Bassett (1983), Crawford et al. (1998), Gaynor (2006), Ryle et al. (2009) and Delaney et al. (2013). There are a number of streams and flushes that run through the machair edged by wet vegetation, which add to the diversity of species and help to retain the blowing sand. Several seeps and streams are present and tufa deposits have developed in some places. See the coastal habitats supporting document for further details
Vegetation structure: zonation	Occurrence	Maintain the range of coastal habitats including transitional zones, subject to natural processes including erosion and succession	Based on data from Ryle et al. (2009) and Delaney et al. (2013). A series of fixed dune ridges (up to 1.5m) and mobile dunes front the seaward side of the machair with low-lying ridges in the transition from fixed dune to machair. This particular machair is unusual in that it extends up from the sand dune system onto the slopes of Garter Hill, grading into heath and wet grassland at the top of the hill. Agricultural fields flank the east and western edges of the machair. See the coastal habitats supporting document for further details
Vegetation structure: bare ground	Percentage cover	Bare ground should not exceed 10% of machair habitat, subject to natural processes	Severe erosion was recorded in the 1996 machair survey (Crawford et al., 1998) and the site was regarded as a poor example of machair. It is likely that the quality of the habitat has declined even further as unsustainable grazing pressure and erosion persist on the commonage. The impacts of overgrazing have exacerbated the natural erosion with large areas of bare sand present in the transitional area between the fixed dune and machair habitats. The SDM concluded that overgrazing remains a problem, but is less severe than it was previously (Delaney et al., 2013). See the coastal habitats supporting document for further details

Vegetation structure: sward height	Centimetres	Maintain structural variation within sward	Based on data from Ryle et al. (2009) and Delaney et al. (2013). The sward is closely-cropped (2-5cm) due to the persistent overgrazing of this habitat over a long period of time and bare sand covers large areas. In the area of recovering machair, the average sward height was 2.06cm. See the coastal habitats supporting document for further details
Vegetation composition: typical species and sub-communities	Percentage cover at a representative number of monitoring stops	Maintain range of sub-communities with typical species listed in Delaney et al. (2013)	Based on data from Gaynor (2006), Ryle et al. (2009) and Delaney et al. (2013). The typical machair species diversity is poor in this SAC. See the coastal habitats supporting document for further details
Vegetation composition: negative indicator species	Percentage cover	Negative indicator species (including non-native species) to represent less than 5% cover	Based on data from Ryle et al. (2009) and Delaney et al. (2013). Negative indicators include non-native species, species indicative of changes in nutrient status and species not considered characteristic of the habitat. See the coastal habitats supporting document for further details
Vegetation composition: scrub/trees	Percentage cover	No more than 5% cover or under control	Based on data from Ryle et al. (2009) and Delaney et al. (2013). See the coastal habitats supporting document for further details
Vegetation composition: bryophytes	Percentage cover	Should always be at least an occasional component of the vegetation	Based on data from Ryle et al. (2009) and Delaney et al. (2013). The machair contains abundant mosses including <i>Bryum pseudotriquetrum</i> , <i>Calliergonella cuspidata</i> , <i>Homalothecium lutescens</i> , <i>Rhytidiadelphus squarrosus</i> and <i>Syntrichia ruralis</i> subsp. <i>ruraliformis</i> . The Near Threatened moss <i>Catocopium nigratum</i> (Lockhart et al., 2012), which is also listed on the Flora (Protection) Order, 2015 (FPO), is associated with flushed machair at Garter Hill. Petalwort (<i>Petalophyllum ralfsii</i>), a liverwort species listed on Annex II of the EU Habitats Directive and on the FPO, has been recorded on machair in this SAC (Campbell et al., 2015). See the conservation objective for petalwort (1395) and the coastal habitats supporting document for further details

Conservation Objectives for : Glenamoy Bog Complex SAC [000500]

3160 Natural dystrophic lakes and ponds

To maintain the favourable conservation condition of Natural dystrophic lakes and ponds in Glenamoy Bog Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	The blanket bog in Glenamoy Bog Complex SAC has very well-developed pool systems and some deep, open lakes. Lake habitat 3160 is likely to occur in all pools and lakes, including the largest, Lougherglass (38.9ha). Although there are more than 2,100 lake/pool polygons, not all pools are mapped in the 1:5,000 OSI data (see map 4). All lakes and pools are considered to be potential 3160. The habitat is of high conservation value in the SAC. For further information on the distribution, vegetation and morphology of the habitat in the SAC, see Foss and McGee (1987) and Douglas et al. (1989). Two measures of extent should be used: 1. the area of the lake itself and; 2. the extent of the vegetation communities/zones that typify the habitat. Further information relating to all attributes is provided in the lake habitats supporting document for the purposes of site-specific conservation objectives and Article 17 reporting (O Connor, 2015)
Habitat distribution	Occurrence	No decline, subject to natural processes	As noted above, the habitat is widespread and of high conservation value in the SAC (see map 4). All lake/pond polygons have been mapped as potential 3160
Typical species	Occurrence	Typical species present, in good condition, and demonstrating typical abundances and distribution	For lists of typical plant and invertebrate species, see the Article 17 habitat assessment for 3160 (NPWS, 2013) and O Connor (2015). Douglas et al. (1989) state larger pools at Glenamoy chiefly contain pipewort (<i>Eriocaulon aquaticum</i>), a Near Threatened species in Ireland (Wyse Jackson et al., 2016), with water lobelia (<i>Lobelia dortmanna</i>), lesser bladderwort (<i>Utricularia minor</i>), bogbean (<i>Menyanthes trifoliata</i>) and the bog mosses <i>Sphagnum auriculatum</i> and <i>S. cuspidatum</i> as frequent associates. Ungrazed island hummocks, with crowberry (<i>Empetrum nigrum</i>) and juniper (<i>Juniperus communis</i>), are frequent features of larger pools
Vegetation composition: characteristic zonation	Occurrence	All characteristic zones should be present, correctly distributed and in good condition	Further work is necessary to describe the characteristic zonation and other spatial patterns in lake habitat 3160 (see O Connor, 2015). Spatial patterns are likely to be relatively simple in 3160 lakes and ponds, with limited zonation
Vegetation distribution: maximum depth	Metres	Maintain maximum depth of vegetation, subject to natural processes	The maximum depth of vegetation is likely to be specific to the lake shoreline in question. Further work is necessary to develop indicative targets for lake habitat 3160. 3160 lakes and pools naturally have very clear water and, therefore, maximum depth can be large

Hydrological regime: water level fluctuations	Metres	Maintain appropriate natural hydrological regime necessary to support the habitat	Fluctuations in lake water level are typical in Ireland, but can be amplified by activities such as abstraction and drainage. Increased water level fluctuations can increase wave action, up-root vegetation, increase turbidity, alter the substratum and lead to release of nutrients from the sediment. The hydrological regime of the lakes and pools must be maintained so that the area, distribution and depth of the lake habitat and its constituent/characteristic vegetation zones and communities are not reduced. The hydrological regime of 3160 lakes and pools is integrally linked to that of the surrounding blanket bog, transition mire/quaking bog and other peatland habitats. Owing to their size and the sensitivity of peatland, 3160 lakes and pools can easily be damaged or destroyed by drainage
Lake substratum quality	Various	Maintain appropriate substratum type, extent and chemistry to support the vegetation	Research is required to further characterise the substratum types (particle size and origin) and substratum quality (notably pH, calcium, iron and nutrient concentrations) favoured by each of the five Annex I lake habitats in Ireland. It is likely that habitat 3160 is associated with nutrient-poor peat and silt substrates
Water quality: transparency	Metres	Maintain appropriate Secchi transparency. There should be no decline in Secchi depth/transparency	Transparency relates to light penetration and, hence, to the depth of colonisation of vegetation. It can be affected by phytoplankton blooms, water colour and turbidity. Specific targets have yet to be established for lake habitat 3160 (O Connor, 2015). Habitat 3160 is associated with very clear water. The OECD fixed boundary system set transparency targets for ultra-oligotrophic lakes of $\geq 12\text{m}$ annual mean Secchi disk depth, and $\geq 6\text{m}$ annual minimum Secchi disk depth
Water quality: nutrients	$\mu\text{g/l P}$; mg/l N	Maintain the concentration of nutrients in the water column at sufficiently low levels to support the habitat and its typical species	As a nutrient-poor habitat, oligotrophic and Water Framework Directive (WFD) 'high' status targets apply. Where a lake has nutrient concentrations that are lower than these targets, there should be no decline within class, i.e. no upward trend in nutrient concentrations. For 3160 lakes and pools, annual average total phosphorus (TP) concentration should be $\leq 5\mu\text{g/l TP}$, average annual total ammonia concentration should be $\leq 0.040\text{mg/l N}$ and annual 95th percentile for total ammonia should be $\leq 0.090\text{mg/l N}$. See also The European Communities Environmental Objectives (Surface Waters) Regulations 2009
Water quality: phytoplankton biomass	$\mu\text{g/l Chlorophyll } a$	Maintain appropriate water quality to support the habitat, including high chlorophyll <i>a</i> status	Oligotrophic and WFD 'high' status targets apply to lake habitat 3160. The average growing season (March-October) chlorophyll <i>a</i> concentration must be $< 5.8\mu\text{g/l}$ (The European Communities Environmental Objectives (Surface Waters) Regulations 2009). Where a lake has a chlorophyll <i>a</i> concentration that is lower than this target, there should be no decline within class, i.e. no upward trend in phytoplankton biomass. The OECD targets may be more appropriate for lake habitat 3160: annual average chlorophyll <i>a</i> concentration $< 1\mu\text{g/l}$ and annual peak chlorophyll <i>a</i> concentration $\leq 2.5\mu\text{g/l}$. See also The European Communities Environmental Objectives (Surface Waters) Regulations 2009
Water quality: phytoplankton composition	EPA phytoplankton composition metric	Maintain appropriate water quality to support the habitat, including high phytoplankton composition status	The Environmental Protection Agency (EPA) has developed a phytoplankton composition metric for nutrient enrichment of Irish lakes. As for other water quality indicators, lake habitat 3160 requires WFD high status

Water quality: attached algal biomass	Algal cover and EPA phytobenthos metric	Maintain trace/absent attached algal biomass (<5% cover) and high phytobenthos status	Nutrient enrichment can favour epiphytic and epipelic algae that can out-compete the submerged vegetation. The cover abundance of attached algae in 3160 lakes and ponds should, therefore, be trace/absent (<5% cover). EPA phytobenthos can be used as an indicator of changes in attached algal biomass. As for other water quality indicators, lake habitat 3160 requires high phytobenthos status
Water quality: macrophyte status	EPA macrophyte metric (The Free Index)	Maintain high macrophyte status	Nutrient enrichment can favour more competitive submerged macrophyte species that out-compete the typical and characteristic species for the lake habitat. The EPA monitors macrophyte status for WFD purposes using the 'Free Index'. The target for 3160 lakes and pools is high status or an Ecological Quality Ratio (EQR) for lake macrophytes of ≥ 0.90 , as defined in Schedule Five of the European Communities Environmental Objectives (Surface Waters) Regulations 2009
Acidification status	pH units; mg/l	Maintain appropriate water and sediment pH, alkalinity and cation concentrations to support the habitat, subject to natural processes	Acidification can impact on species abundance and composition in soft water lake habitats. Although European Commission (2013) describes habitat 3160 as having pH 3-6, Drinan (2012) found mean pHs of 5.16 and 5.62 in upland and lowland 3160 lakes, respectively. The target for lake habitat 3160 is pH >4.5 and <9.0 , in line with the surface water standards for soft waters (where water hardness is $\leq 100\text{mg/l}$ calcium carbonate). See Schedule Five of the European Communities Environmental Objectives (Surface Waters) Regulations 2009. The specific requirements of lake habitat 3160, in terms of water and sediment pH, alkalinity and cation concentration, have not been determined
Water colour	mg/l PtCo	Maintain appropriate water colour to support the habitat	Increased water colour and turbidity decrease light penetration and can reduce the area of available habitat for lake macrophytes, particularly at the lower euphotic depths. The primary source of increased water colour in Ireland is disturbance to peatland. No habitat-specific or national standards for water colour currently exist. Studies have shown median colour concentrations in Irish lakes of 38mg/l PtCo (Free et al., 2000) and 33mg/l PtCo (Free et al., 2006). It is likely that the water colour in all Irish lake habitats would naturally be $<50\text{mg/l}$ PtCo. Water colour can be very low ($<20\text{mg/l}$ PtCo or even $<10\text{mg/l}$ PtCo) in 3160 lakes and pools where the peatland in the lake's catchment is intact
Dissolved organic carbon (DOC)	mg/l	Maintain appropriate organic carbon levels to support the habitat	Dissolved (and particulate) organic carbon (OC) in the water column is linked to water colour and acidification (organic acids). Increasing DOC in water has been documented across the Northern Hemisphere, including afforested peatland catchments in Ireland. Damage and degradation of peatland, leading to decomposition of peat is likely to be the predominant source of OC in Ireland. OC in water promotes decomposition by fungi and bacteria that, in turn, releases dissolved nutrients. The increased biomass of decomposers can also impact directly on the characteristic lake communities through shading, competition, etc.
Turbidity	Nephelometric turbidity units/ mg/l SS/ other appropriate units	Maintain appropriate turbidity to support the habitat	Turbidity can significantly affect the quantity and quality of light reaching rooted and attached vegetation and can, therefore, impact on lake habitats. The settlement of higher loads of inorganic or organic material on lake vegetation communities may also have impacts on sensitive, delicate species. Turbidity can increase as a result of re-suspension of material within the lake, higher loads entering the lake, or eutrophication. Turbidity measurement and interpretation is challenging. As a result, it is likely to be difficult to set habitat-specific targets for turbidity in lakes

Fringing habitat: Hectares
area

Maintain the area and condition of fringing habitats necessary to support the natural structure and functioning of habitat 3160

3160 lakes and pools intergrade with blanket bog and heath communities in this SAC, often with transition mire/quaking bog at the interface. These habitats support the structure and functions of the lake habitat. Equally, the fringing habitats are also dependent on the lake/pool, particularly its water levels, and can support wetland communities and species of conservation concern

Conservation Objectives for : Glenamoy Bog Complex SAC [000500]

4010 Northern Atlantic wet heaths with *Erica tetralix*

To restore the favourable conservation condition of Northern Atlantic wet heaths with *Erica tetralix* in Glenamoy Bog Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Northern Atlantic wet heaths with <i>Erica tetralix</i> has not been mapped in detail for Glenamoy Bog Complex SAC but from current available data the total area of the qualifying habitat is estimated to be approximately 723ha, covering 6% of the SAC (NPWS internal files). Further details on this and the following attributes can be found in the Glenamoy Bog Complex SAC conservation objectives supporting document for blanket bogs and associated habitats
Habitat distribution	Occurrence	No decline, subject to natural processes	The habitat is documented to occur on hillsides within the SAC (NPWS internal files). Further information can be found within NPWS internal files and the blanket bogs and associated habitats supporting document
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil nutrient status within natural range	See the blanket bogs and associated habitats supporting document for further details
Community diversity	Abundance of variety of vegetation communities	Maintain variety of vegetation communities, subject to natural processes	The diversity of wet heath communities within this SAC is unknown. Further information on vegetation communities associated with this habitat is presented in Perrin et al. (2014)
Vegetation composition: cross-leaved heath	Occurrence within 20m of a representative number of monitoring stops	Cross-leaved heath (<i>Erica tetralix</i>) present within a 20m radius of each monitoring stop	Attribute and target based on Perrin et al. (2014)
Vegetation composition: positive indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of positive indicator species at least 50%	Attribute and target based on Perrin et al. (2014), where the list of positive indicator species for this habitat is also presented
Vegetation composition: lichens and bryophytes	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of <i>Cladonia</i> and <i>Sphagnum</i> species, <i>Racomitrium lanuginosum</i> and pleurocarpous mosses at least 10%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: ericoid species and crowberry	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of ericoid species and crowberry (<i>Empetrum nigrum</i>) at least 15%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: dwarf shrub species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of dwarf shrubs less than 75%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: negative indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of negative indicator species less than 1%	Attribute and target based on Perrin et al. (2014), where the list of negative indicator species for this habitat is also presented
Vegetation composition: non-native species	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of non-native species less than 1%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: native trees and shrubs	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of scattered native trees and shrubs less than 20%	Attribute and target based on Perrin et al. (2014)

Vegetation composition: bracken	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of bracken (<i>Pteridium aquilinum</i>) less than 10%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: soft rush	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of soft rush (<i>Juncus effusus</i>) less than 10%	Attribute and target based on Perrin et al. (2014)
Vegetation structure: <i>Sphagnum</i> condition	Condition at a representative number of 2m x 2m monitoring stops	Less than 10% of the <i>Sphagnum</i> cover is crushed, broken and/or pulled up	Attribute and target based on Perrin et al. (2014)
Vegetation structure: signs of browsing	Percentage of shoots browsed at a representative number of 2m x 2m monitoring stops	Less than 33% collectively of the last complete growing season's shoots of ericoids, crowberry (<i>Empetrum nigrum</i>) and bog-myrtle (<i>Myrica gale</i>) showing signs of browsing	Attribute and target based on Perrin et al. (2014)
Vegetation structure: burning	Occurrence in local vicinity of a representative number of monitoring stops	No signs of burning in sensitive areas, into the moss, liverwort or lichen layer or exposure of peat surface due to burning	Attribute and target based on Perrin et al. (2014), where the list of sensitive areas for this habitat is also presented
Physical structure: disturbed bare ground	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of disturbed bare ground less than 10%	Attribute and target based on Perrin et al. (2014)
Physical structure: drainage	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Area showing signs of drainage from heavy trampling, tracking or ditches less than 10%	Attribute and target based on Perrin et al. (2014)
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat	This includes species listed in the Flora (Protection) Order, 2015 and/or the red data lists (Lockhart et al., 2012; Wyse Jackson et al., 2016)

Conservation Objectives for : Glenamoy Bog Complex SAC [000500]

5130 Juniperus communis formations on heaths or calcareous grasslands

To maintain the favourable conservation condition of *Juniperus communis* formations on heaths or calcareous grasslands in Glenamoy Bog Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Occurrence	Area stable or increasing, subject to natural processes	<i>Juniperus communis</i> formations on heath or calcareous grasslands has not been surveyed in detail in Glenamoy Bog Complex SAC and thus the total area of the qualifying habitat is unknown. Douglas et al. (1989) noted juniper <i>Juniperus communis</i> as being fairly widespread throughout the blanket bog habitat (7130*), particularly at pool margins and on islands within pool lakes. It also occurs on the coastal fringe and grades into heath and grassland communities in places
Habitat distribution	Occurrence	No decline, subject to natural processes	See notes for area above
Juniper population size	Number per formation	At least 50 plants per formation	To classify as a juniper (<i>Juniperus communis</i>) formation, at least 50 plants should be present (Cooper et al., 2012)
Vegetation composition: typical species	Number per formation	At least 50% of the listed positive indicator species for the relevant vegetation group present	Attribute and target based on Cooper et al. (2012), where positive indicator species for five vegetation groups are listed
Vegetation composition: negative indicator species	Occurrence per formation	Negative indicator species, particularly non-native invasive species, absent or under control	Attribute and target based on Cooper et al. (2012), where the list of negative indicator species is presented
Vegetation structure: cone-bearing plants	Percentage per formation	At least 10% of juniper plants are bearing cones	Attribute and target based on Cooper et al. (2012)
Vegetation structure: seedling recruitment	Percentage per formation	At least 10% of juniper plants are seedlings	Attribute and target based on Cooper et al. (2012)
Vegetation structure: dead juniper	Percentage per formation	Mean percentage of each juniper plant dead less than 10%	Attribute and target based on Cooper et al. (2012)

Conservation Objectives for : Glenamoy Bog Complex SAC [000500]

7130 Blanket bogs (* if active bog)

To restore the favourable conservation condition of Blanket bogs (* if active bog) in Glenamoy Bog Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Blanket bog has not been mapped in detail for Glenamoy Bog Complex SAC but from current available data the total area of the qualifying habitat is estimated to be approximately 6,749ha, covering 52% of the SAC (NPWS internal files). Further details on this and the following attributes can be found in the Glenamoy Bog Complex SAC conservation objectives supporting document for blanket bogs and associated habitats
Habitat distribution	Occurrence	No decline, subject to natural processes	Blanket bog habitat occupies the gentle undulating plain that dominates the central areas of this SAC. It also extends uphill to cover the slopes of Maumakeogh and Benmore in the eastern sector of the SAC, and northward, out toward the sea cliffs of the north-west Mayo coastline (NPWS internal files). Important peatland sites within the SAC include Glenamoy Bog, Rathavisteen Bog, Maumkeogh Bog and Glencalry Bog. Further information can be found in Foss and McGee (1987), Douglas et al. (1989), NPWS internal files and the blanket bogs and associated habitats supporting document
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil nutrient status within natural range	See the blanket bogs and associated habitats supporting document for further details
Ecosystem function: peat formation	Active blanket bog as a proportion of the total area of Annex I blanket bog habitat	At least 99% of the total Annex I blanket bog area is active	See the blanket bogs and associated habitats supporting document for further details
Ecosystem function: hydrology	Flow direction, water levels, occurrence of drains and erosion gullies	Natural hydrology unaffected by drains and erosion	Further details and a brief discussion of restoration potential is presented in the blanket bogs and associated habitats supporting document
Community diversity	Abundance of variety of vegetation communities	Maintain variety of vegetation communities, subject to natural processes	A variety of blanket bog vegetation communities have been recorded in this SAC (Foss and McGee, 1987; Douglas et al., 1989; NPWS internal files), five of which correspond to communities recorded in the National Survey of Upland Habitats and listed in the provisional list of vegetation communities described in Perrin et al. (2014). Further information on vegetation communities associated with this habitat is presented in Perrin et al. (2014)
Vegetation composition: positive indicator species	Number of species at a representative number of 2m x 2m monitoring stops	Number of positive indicator species present at each monitoring stop is at least seven	Attribute and target based on Perrin et al. (2014), where the list of positive indicator species for this habitat is also presented
Vegetation composition: lichens and bryophytes	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of bryophytes or lichens, excluding <i>Sphagnum fallax</i> , at least 10%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: potential dominant species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of each of the potential dominant species less than 75%	Attribute and target based on Perrin et al. (2014). See the blanket bogs and associated habitats supporting document for the list of potential dominant species
Vegetation composition: negative indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of negative indicator species less than 1%	Attribute and target based on Perrin et al. (2014), where the list of negative indicator species for this habitat is also presented

Vegetation composition: non-native species	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of non-native species less than 1%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: native trees and shrubs	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of scattered native trees and shrubs less than 10%	Attribute and target based on Perrin et al. (2014)
Vegetation structure: <i>Sphagnum</i> condition	Condition at a representative number of 2m x 2m monitoring stops	Less than 10% of the <i>Sphagnum</i> cover is crushed, broken and/or pulled up	Attribute and target based on Perrin et al. (2014)
Vegetation structure: signs of browsing	Percentage of shoots browsed at a representative number of 2m x 2m monitoring stops	Last complete growing season's shoots of ericoids, crowberry (<i>Empetrum nigrum</i>) and bog-myrtle (<i>Myrica gale</i>) showing signs of browsing collectively less than 33%	Attribute and target based on Perrin et al. (2014)
Vegetation structure: burning	Occurrence in local vicinity of a representative number of monitoring stops	No signs of burning in sensitive areas, into the moss, liverwort or lichen layer or exposure of peat surface due to burning	Attribute and target based on Perrin et al. (2014), where the list of sensitive areas for this habitat is also presented
Physical structure: disturbed bare ground	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of disturbed bare ground less than 10%	Attribute and target based on Perrin et al. (2014)
Physical structure: drainage	Percentage area in local vicinity of a representative number of monitoring stops	Area showing signs of drainage from heavy trampling, tracking or ditches less than 10%	Attribute and target based on Perrin et al. (2014)
Physical structure: erosion	Percentage area in local vicinity of a representative number of monitoring stops	Less than 5% of the greater bog mosaic comprises erosion gullies and eroded areas	Attribute and target based on Perrin et al. (2014)
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat	This includes species listed in the Flora (Protection) Order, 2015 and/or the red data lists (Lockhart et al., 2012; Wyse Jackson et al., 2016)

Conservation Objectives for : Glenamoy Bog Complex SAC [000500]

7140 Transition mires and quaking bogs

To restore the favourable conservation condition of Transition mires and quaking bogs in Glenamoy Bog Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Transition mires and quaking bogs have not been mapped in detail for Glenamoy Bog Complex SAC and thus the total area of the qualifying habitat is unknown. Further details on this and the following attributes can be found in the Glenamoy Bog Complex SAC conservation objectives supporting document for blanket bogs and associated habitats
Habitat distribution	Occurrence	No decline, subject to natural processes	The habitat is documented to occur in areas where bog vegetation and base-rich flushes merge, and also at the interface between large pools/lakes and adjacent bog (NPWS internal files). Examples of this habitat can be found at Glenamoy Bog, Rathavisteen Bog and Glencalry Bog (Foss and McGee, 1987; Douglas et al., 1989; R. Hodd, pers. comm.). Further information can be found in Foss and McGee (1987), Douglas et al. (1989) and the blanket bogs and associated habitats supporting document
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil nutrient status within natural range	See the blanket bogs and associated habitats supporting document for further details
Community diversity	Abundance of variety of vegetation communities	Maintain variety of vegetation communities, subject to natural processes	Foss and McGee (1987), Douglas et al. (1989) and R. Hodd (pers. comm.) recorded a variety of transition mire vegetation communities in this SAC, two of which correspond to communities recorded in the National Survey of Upland Habitats and listed in the provisional list of vegetation communities described in Perrin et al. (2014). Further information on vegetation communities associated with this habitat is presented in Perrin et al. (2014)
Vegetation composition: number of positive indicator species	Number of species at a representative number of 2m x 2m monitoring stops	Number of positive indicator species at each monitoring stop is at least three for infilling pools and flushes and at least six for fens	Attribute and target based on Perrin et al. (2014), where the list of positive indicator species for this habitat is also presented
Vegetation composition: number of core positive indicator species	Number of species at a representative number of 2m x 2m monitoring stops	At least one core positive indicator species present	Attribute and target based on Perrin et al. (2014), where the list of positive indicator species for this habitat is also presented
Vegetation composition: cover of positive indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of positive indicator species is at least 25%	Attribute and target based on Perrin et al. (2014), where the list of positive indicator species for this habitat is also presented
Vegetation composition: negative indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of negative indicator species less than 1%	Attribute and target based on Perrin et al. (2014), where the list of negative indicator species for this habitat is also presented
Vegetation composition: non-native species	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of non-native species less than 1%	Attribute and target based on Perrin et al. (2014)
Vegetation structure: height	Percentage of leaves/shoots at a representative number of 2m x 2m monitoring stops	Proportion of live leaves and/or flowering shoots of vascular plants that are more than 15cm above the ground surface should be at least 50%	Attribute and target based on Perrin et al. (2014). This attribute is only applicable to fen and flush examples of the habitat, not to infilling pool examples

Physical structure: disturbed bare ground	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of disturbed bare ground less than 10%	Attribute and target based on Perrin et al. (2014)
Physical structure: drainage	Percentage area in local vicinity of a representative number of monitoring stops	Area showing signs of drainage from heavy trampling, tracking or ditches less than 10%	Attribute and target based on Perrin et al. (2014)
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat	This includes species listed in the Flora (Protection) Order, 2015 (FPO) and/or the red data lists (Lockhart et al., 2012; Wyse Jackson et al., 2016). The Annex II listed, FPO listed and Near Threatened slender green feather-moss (<i>Hamatocaulis (Drepanocladus) vernicosus</i>) and the Vulnerable <i>Tomentypnum nitens</i> (Lockhart et al., 2012) are known to occur within the SAC (Foss and McGee, 1987; Campbell et al., 2015; NPWS internal files), although the latter species cannot be specifically assigned to this habitat. See also the conservation objective for slender green feather-moss (1393)

Conservation Objectives for : Glenamoy Bog Complex SAC [000500]

7150 Depressions on peat substrates of the Rhynchosporion

To restore the favourable conservation condition of Depressions on peat substrates of the Rhynchosporion in Glenamoy Bog Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Depressions on peat substrates of the Rhynchosporion has not been mapped in detail for Glenamoy Bog Complex SAC and thus the total area of the qualifying habitat is unknown. Further details on this and the following attributes can be found in the Glenamoy Bog Complex SAC conservation objectives supporting document for blanket bogs and associated habitats
Habitat distribution	Occurrence	No decline, subject to natural processes	This habitat is typically confined to relatively small areas but is best represented around pool margins and in wet hollows in the SAC (NPWS internal files). Examples of this habitat can be found at Glenamoy Bog (Douglas et al., 1989). Further information can be found within Douglas et al. (1989), NPWS internal files and the blanket bogs and associated habitats supporting document
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil nutrient status within natural range	See the blanket bogs and associated habitats supporting document for further details
Vegetation composition: positive indicator species	Number of species at a representative number of 2m x 2m monitoring stops	Number of positive indicator species at each monitoring stop is at least five	Attribute and target based on Perrin et al. (2014), where the list of positive indicator species for this habitat is also presented
Vegetation composition: <i>Rhynchospora</i> spp.	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of white beaked sedge (<i>Rhynchospora alba</i>) and brown beaked sedge (<i>R. fusca</i>) at least 10%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: potential dominant species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of each of the potential dominant species individually less than 35%	Attribute and target based on Perrin et al. (2014). See the blanket bogs and associated habitats supporting document for the list of potential dominant species
Vegetation composition: negative indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of negative indicator species less than 1%	Attribute and target based on Perrin et al. (2014), where the list of negative indicator species for this habitat is also presented
Vegetation composition: non-native species	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of non-native species less than 1%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: native trees and shrubs	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of scattered native trees and shrubs less than 10%	Attribute and target based on Perrin et al. (2014)
Vegetation structure: <i>Sphagnum</i> condition	Condition at a representative number of 2m x 2m monitoring stops	Less than 10% of the <i>Sphagnum</i> cover is crushed, broken and/or pulled up	Attribute and target based on Perrin et al. (2014)
Vegetation structure: signs of browsing	Percentage of shoots browsed at a representative number of 2m x 2m monitoring stops	Last complete growing season's shoots of ericoids, crowberry (<i>Empetrum nigrum</i>) and bog-myrtle (<i>Myrica gale</i>) showing signs of browsing collectively less than 33%	Attribute and target based on Perrin et al. (2014)

Vegetation structure: burning	Occurrence in local vicinity of a representative number of monitoring stops	No signs of burning in sensitive areas, into the moss, liverwort or lichen layer or exposure of peat surface due to burning	Attribute and target based on Perrin et al. (2014), where the list of sensitive areas for this habitat is also presented
Physical structure: disturbed bare ground	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of disturbed bare ground less than 10%	Attribute and target based on Perrin et al. (2014)
Physical structure: drainage	Percentage area in local vicinity of a representative number of monitoring stops	Area showing signs of drainage from heavy trampling, tracking or ditches less than 10%	Attribute and target based on Perrin et al. (2014)
Physical structure: erosion	Percentage area in local vicinity of a representative number of monitoring stops	Less than 5% of the greater bog mosaic comprises erosion gullies and eroded areas	Attribute and target based on Perrin et al. (2014)
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat	This includes species listed in the Flora (Protection) Order, 2015 and/or the red data lists (Lockhart et al., 2012; Wyse Jackson et al., 2016). The Near Threatened brown beak-sedge (<i>Rhynchospora fusca</i>) (Wyse Jackson et al., 2016) was recorded in this habitat by Doyle (1990)

Conservation Objectives for : Glenamoy Bog Complex SAC [000500]

1106 Salmon *Salmo salar*

To restore the favourable conservation condition of Atlantic Salmon in Glenamoy Bog Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Distribution: extent of anadromy	Percentage of river accessible	100% of river channels down to second order accessible from estuary	Artificial barriers block salmons' upstream migration, thereby limiting species to lower stretches and restricting access to spawning areas
Adult spawning fish	Number	Conservation limit (CL) for each system consistently exceeded	A conservation limit (CL) is defined by the North Atlantic Salmon Conservation Organisation (NASCO) as "the spawning stock level that produces long-term average maximum sustainable yield as derived from the adult to adult stock and recruitment relationship". The target is based on the Standing Scientific Committee on Salmon (SSCS) annual model output of CL attainment levels. See SSCS (2016). Attainment of CL estimates are derived from direct counts of adults (rod catch, fish counter) or indirectly by fry abundance counts. The Glenamoy River is currently below CL, meeting 87% of CL
Salmon fry abundance	Number of fry/5 minutes electrofishing	Maintain or exceed 0+ fry mean catchment-wide abundance threshold value. Currently set at 17 salmon fry/5 minutes sampling	The target is the threshold value for rivers currently exceeding their conservation limit (CL)
Out-migrating smolt abundance	Number	No significant decline	Smolt abundance can be negatively affected by a number of impacts such as estuarine pollution, predation and sea lice (<i>Lepeophtheirus salmonis</i>)
Number and distribution of redds	Number and occurrence	No decline in number and distribution of spawning redds due to anthropogenic causes	Salmon spawn in clean gravels
Water quality	EPA Q value	At least Q4 at all sites sampled by EPA	Q values based on triennial water quality surveys carried out by the Environmental Protection Agency (EPA)

Conservation Objectives for : Glenamoy Bog Complex SAC [000500]

1393

Slender Green Feather-moss *Drepanocladus vernicosus*

To maintain the favourable conservation condition of Slender Green Feather-moss (Shining Sickle-moss) in Glenamoy Bog Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Distribution	Number and geographical spread of populations	No decline, subject to natural processes. See map 5 for known location at Rathavisteen	(Please note that <i>Drepanocladus vernicosus</i> was reclassified as <i>Hamatocaulis vernicosus</i> by Hedenäs (1989)). The known population of slender green feather-moss (<i>Hamatocaulis vernicosus</i>) in Glenamoy Bog Complex SAC is at Rathavisteen, where it occurs at the bases of tussocky vegetation on the upper eastern margin of a marsh/fen which is a floating scragh. Data from NPWS survey by Neil Lockhart in 1999 (NPWS internal files). See also Campbell et al. (2015)
Population size	Number of individuals	No decline, subject to natural processes	The density was estimated by Lockhart in 1999 to be c.100 shoots per square metre, giving a shoot count of c.1000 shoots (NPWS internal files). See Campbell et al. (2015) for further details
Area of suitable habitat	Hectares	No decline, subject to natural processes	The full extent of suitable habitat in Glenamoy Bog Complex SAC is unknown. A small patch (c.10m x 1m, i.e. 0.001ha), was recorded in Rathavisteen by Lockhart in 1999 (NPWS internal files). See Campbell et al. (2015) for further details
Hydrological conditions: water table level	Metres	Maintain suitable hydrological conditions	Slender green feather-moss (<i>Hamatocaulis vernicosus</i>) is mostly confined to mesotrophic fens, a transitional habitat between acid bog and base-rich fen. This appears to occur in at least two forms in Ireland: upland transitional flushes, where the plants can occur in lawns that rise and fall with fluctuating water table levels, such as at Rathavisteen; and wet lowland sedge meadows, where plants can be inundated in winter, but may be subject to some desiccation in the summer. Based on Campbell (2013) and Campbell et al. (2015)
Vegetation composition: tree cover	Percentage cover in a representative number of 2m x 2m monitoring plots	Mean percentage tree cover should be less than 15%	Slender green feather-moss (<i>Hamatocaulis vernicosus</i>) grows in moss-dominated, open communities, generally with a low cover of trees and shrubs. See Campbell et al. (2015) for further details
Vegetation composition: shrub cover	Percentage cover in a representative number of 2m x 2m monitoring plots	Mean percentage shrub cover should be less than 20%	Slender green feather-moss (<i>Hamatocaulis vernicosus</i>) grows in moss-dominated, open communities, generally with a low cover of trees and shrubs. See Campbell et al. (2015) for further details
Vegetation composition: grass cover	Percentage cover in a representative number of 2m x 2m monitoring plots	Mean percentage grass species cover should be less than 25%	Slender green feather-moss (<i>Hamatocaulis vernicosus</i>) grows in moss-dominated, open communities, generally with a low cover of grasses, maintained by a low grazing intensity by cattle at Rathavisteen. See Campbell et al. (2015) for further details
Vegetation composition: bryophyte cover	Percentage cover in a representative number of 2m x 2m monitoring plots	Mean percentage bryophyte cover should be more than 50%	The part of the marsh/fen at Rathavisteen where slender green feather-moss (<i>Hamatocaulis vernicosus</i>) occurs is a floating scragh, dominated mostly by <i>Sphagnum</i> spp. with lenses of tussocky sedges and grasses when surveyed by Lockhart in 1999 (NPWS internal files). See also Campbell et al. (2015)
Vegetation composition: cover of <i>Calliergonella cuspidata</i>	Percentage cover in a representative number of 2m x 2m monitoring plots	Mean percentage cover of <i>Calliergonella cuspidata</i> should be less than 15%	<i>Calliergonella cuspidata</i> , a moss species often associated with high nutrient conditions, is usually present, but with low cover and never dominant. See Campbell et al. (2015) for further details
Vegetation structure: vegetation height	Centimetres in a representative number 2m x 2m monitoring plots	Mean vegetation height should not exceed 40cm	See Campbell et al. (2015) for further details

Conservation Objectives for : Glenamoy Bog Complex SAC [000500]

1395 Petalwort *Petalophyllum ralfsii*

To maintain the favourable conservation condition of Petalwort in Glenamoy Bog Complex SAC, which is defined by the following list of attributes and targets:


Attribute	Measure	Target	Notes
Distribution	Number and geographical spread of populations	No decline, subject to natural processes. See map 5 for recorded locations	The known population of petalwort (<i>Petalophyllum ralfsii</i>) in Glenamoy Bog Complex SAC is at Garter Hill, where it occurs on banks of water tracks and also on the sides of low sandhills in wetter flushed parts of the machair. Data from NPWS surveys (NPWS internal files) and Campbell (2013). See also Campbell et al. (2015) for further details
Area of suitable habitat	Hectares	No decline, subject to natural processes	The area of occupancy, estimated from polygons drawn around GPS co-ordinates taken from NPWS surveys and Campbell (2013), was 211,604m ² . However, only about 70% of this area is actually suitable habitat for petalwort (<i>Petalophyllum ralfsii</i>), which is 148,123m ² , i.e. c.14.8ha. See Campbell et al. (2015) for further details
Hydrological conditions: soil moisture	Occurrence of damp soil conditions	Maintain hydrological conditions so that the substrate is kept moist and damp throughout the year, but is not subject to prolonged inundation by flooding in winter	Petalwort (<i>Petalophyllum ralfsii</i>) grows on damp sandy substrate. Based on Campbell (2013) and Campbell et al. (2015)
Hydrological conditions: water table level	Centimetres in a representative number of 1m x 1m monitoring plots	Mean groundwater level should not be more than 80cm from ground surface	See Campbell et al. (2015) for further details
Physical structure: bare soil	Percentage cover in a representative number of 1m x 1m monitoring plots	Mean percentage cover of bare soil should be more than 5%	At Garter Hill, petalwort (<i>Petalophyllum ralfsii</i>) grows in compacted, sandy ground. See Campbell et al. (2015) for further details
Vegetation structure: vegetation height	Centimetres in a representative number of 1m x 1m monitoring plots	Mean vegetation height should be less than 6cm	At Garter Hill, petalwort (<i>Petalophyllum ralfsii</i>) habitat is maintained by rabbit (<i>Oryctolagus cuniculus</i>) and sheep grazing and some trampling. Campbell (2013) recorded a mean height of vegetation of 3cm at Garter Hill. See Campbell et al. (2015) for further details
Vegetation composition: shrub cover	Percentage cover in a representative number of 1m x 1m monitoring plots	Mean percentage shrub cover should be less than 25%	See Campbell et al. (2015) for further details
Vegetation composition: grass cover	Percentage cover in a representative number of 1m x 1m monitoring plots	Mean percentage grass species cover should be less than 60%	See Campbell et al. (2015) for further details


Conservation Objectives for : Glenamoy Bog Complex SAC [000500]

1528 Marsh Saxifrage *Saxifraga hirculus*

To maintain the favourable conservation condition of Marsh Saxifrage in Glenamoy Bog Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Distribution	Number and geographical spread of populations	No loss in geographical spread and number of populations, subject to natural processes. See map 6 for 1km grid square locations	The two known populations of marsh saxifrage (<i>Saxifraga hirculus</i>) in Glenamoy Bog Complex SAC occur at Barroosky and Aghoo. See Muldoon (2011) and Muldoon et al. (2015) for further details
Population size: number of rosettes	Number	Maintain the size of the known populations, subject to natural processes. The target numbers of rosettes are: at least 52,000 at Barroosky and at least 960 at Aghoo	The number of rosettes recorded by Muldoon (2011) were: 65,000 at Barroosky and 1,200 at Aghoo. The target figures are a 20% reduction of the recorded numbers to allow for a margin of error and variability over monitoring seasons. See Muldoon et al. (2015) for further details
Population size: area of occupancy	Hectares	Maintain the area of occupancy of the known populations, subject to natural processes. The target areas are: at least 0.2025ha at Barroosky and at least 0.017ha at Aghoo	The areas of occupancy estimated by Muldoon (2011) were: 2,250m ² (0.2250ha) at Barroosky and at 189m ² (0.019ha) at Aghoo. The target area figures are a 10% reduction of the recorded areas to allow for a margin of error. See Muldoon et al. (2015) for further details
Hydrological conditions: water level	Occurrence of high or fluctuating water levels	Maintain the appropriate natural hydrological regime necessary to support the habitat for the species	In Ireland, marsh saxifrage (<i>Saxifraga hirculus</i>) is now restricted to mineral flushes in blanket bog where rising groundwater forms small streams and seepage areas suitable for the species. Based on Muldoon (2011) and Muldoon et al. (2015)
Vegetation composition: positive indicator species	Occurrence in a representative number of 1m x 1m monitoring stops	Knotted pearlwort (<i>Sagina nodosa</i>) should be present in at least two of five 1m x 1m monitoring stops	The presence of the positive indicator species knotted pearlwort (<i>Sagina nodosa</i>) should be maintained (Muldoon, 2011; Muldoon et al., 2015)
Vegetation composition: negative indicator species	Mean percentage cover in five 1m x 1m monitoring stops	Mean percentage cover of purple moor-grass (<i>Molinia caerulea</i>) should not exceed 5%; mean percentage cover of Yorkshire fog (<i>Holcus lanatus</i>) should not exceed 15%	Low cover of the negative indicator species purple moor-grass (<i>Molinia caerulea</i>) and Yorkshire fog (<i>Holcus lanatus</i>) should be maintained. Part of the surrounding bog at the Barroosky site has been reclaimed for low grade agricultural use. No improvements (e.g. fertilizer inputs) should be carried out on the site. Monitoring of purple moor-grass levels in particular is recommended at Barroosky. See Muldoon (2011) and Muldoon et al. (2015) for further details
Vegetation structure: sward height	Centimetres	Maintain a mean vegetation height of less than 15cm	See Muldoon (2011) and Muldoon et al. (2015) for further details
Vegetation structure: grazing level	Evidence of grazing	Maintain grazing at light to moderate levels to ensure an open vegetation structure and to allow flowering to occur	The surrounding bog at the flush at Barroosky is overgrazed with high levels of poaching. Low grazing was identified as an issue at Aghoo. See Muldoon (2011) and Muldoon et al. (2015) for further details













Legend


- Glenamoy Bog Complex SAC 000500
- Broadhaven Bay SAC 000472
- Slieve Fyagh Bog SAC 000542
- Blacksod Bay/Broad Haven SPA 004037
- Illanmaster SPA 004074

Irish Sea Cliff Survey Site ID: 08005

SDM Site Code: 128


Legend

-  Glenamoy Bog Complex SAC 000500
-  Sand Dune Monitoring Project Survey Area
- Qualifying Interests**
-  1230 Vegetated sea cliffs of the Atlantic and Baltic coasts
-  21AO Machairs
- Non-qualifying interests**
-  2120 Shifting dunes along the shoreline with *Ammophila arenaria* ("white dunes")
-  2130 Fixed dunes with herbaceous vegetation ("grey dunes")
-  2190 Humid dune slacks
-  OSi Discovery Series County Boundary




Legend

- Glenamoy Bog Complex SAC 000500
- Potential 3160 Potential natural dystrophic lakes and ponds
- OSi Discovery Series County Boundary






Legend

- Glenamoy Bog Complex SAC 000500
- OSi Discovery Series County Boundary
- ★ 1393 Slender green feather moss *Drepanocladus (Hamatocaulis) vernicosus* (Mitt.) Warnst.
- ★ 1395 Petalwort *Petalophyllum ralfsii* (Wills.) Nees and Gott.



Legend

-  1528 Marsh saxifrage *Saxifraga hirculus*
-  Glenamoy Bog Complex SAC 000500
-  OSi Discovery Series County Boundary




An Roinn Ealaíon, Oidhreachta,
 Gnóthai Réigiúnacha, Tuaithe agus Gaeltachta
 Department of Arts, Heritage,
 Regional, Rural and Gaeltacht Affairs

**MAP 6:
 GLENAMOY BOG COMPLEX SAC
 CONSERVATION OBJECTIVES
 MARSH SAXIFRAGE**

Map to be read in conjunction with the NPWS Conservation Objectives Document.

**SITE CODE:
 SAC 000500; version 3.01.
 CO. MAYO**



The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
 Ordnance Survey of Ireland Licence No EN 0059216. © Ordnance Survey of Ireland Government of Ireland
 Níl sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbheithnithe a déanamh ar theorainneacha na gceantar comharthaite. Suirbhéarachta Ordonáis na hÉireann Ceadúnas



**Map Version 1
 Date: Oct 2016**



Conservation objectives for Greaghans Turlough SAC [000503]

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Objective: To maintain or restore the favourable conservation condition of the Annex I habitat(s) and/or the Annex II species for which the SAC has been selected:

Code	Description
------	-------------

3180	Turloughs*
------	------------

* denotes a priority habitat



Citation: NPWS (2020) Conservation objectives for Greaghans Turlough SAC [000503]. Generic Version 7.0. Department of Culture, Heritage and the Gaeltacht.



Conservation objectives for Kilglassan/Caheravoostia Turlough Complex SAC [000504]

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Objective: To maintain or restore the favourable conservation condition of the Annex I habitat(s) and/or the Annex II species for which the SAC has been selected:

Code	Description
------	-------------

3180	Turloughs*
------	------------

* denotes a priority habitat



Citation: NPWS (2020) Conservation objectives for Kilglassan/Caheravoostia Turlough Complex SAC [000504]. Generic Version 7.0. Department of Culture, Heritage and the Gaeltacht.

National Parks and Wildlife Service

Conservation Objectives Series

Inishkea Islands SAC 000507



*An Roinn
Ealaíon, Oidhreachta agus Gaeltachta*

*Department of
Arts, Heritage and the Gaeltacht*



**National Parks and Wildlife Service,
Department of Arts, Heritage and the Gaeltacht,**

7 Ely Place, Dublin 2, Ireland.

Web: www.npws.ie

E-mail: nature.conservation@ahg.gov.ie

Citation:

**NPWS (201) Conservation Objectives: Inishkea Islands SAC 000507. Version 1.
National Parks and Wildlife Service, Department of Arts, Heritage and the
Gaeltacht.**

Series Editor: Rebecca Jeffrey

ISSN 2009-4086

Introduction

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

** indicates a priority habitat under the Habitats Directive*

000507	Inishkea Islands SAC
1364	Grey Seal <i>Halichoerus grypus</i>
1395	Petalwort <i>Petalophyllum ralfsii</i>
21A0	Machairs (* in Ireland)

Please note that this SAC overlaps with Inishkea Islands SPA (004004) and adjoins West Connacht Coast SAC (002998). See map 2. The conservation objectives for this site should be used in conjunction with those for overlapping and adjacent sites as appropriate.

Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

NPWS Documents

Year :	2003
Title :	Grey seal population status at islands in the Inishkea group, as determined from breeding ground surveys in 2002
Author :	Ó Cadhla, O.; Strong, D.
Series :	Unpublished report to NPWS
Year :	2004
Title :	Harbour seal population assessment in the Republic of Ireland: August 2003
Author :	Cronin, M.; Duck, C.; O Cadhla, O.; Nairn, R.; Strong, D.; O'Keeffe, C.
Series :	Irish Wildlife Manual No. 11
Year :	2004
Title :	Summary of National Parks and Wildlife Service surveys for common (harbour) seals (<i>Phoca vitulina</i>) and grey seals (<i>Halichoerus grypus</i>), 1978 to 2003
Author :	Lyons, D.O.
Series :	Irish Wildlife Manual No. 13
Year :	2004
Title :	Aerial surveying of grey seal breeding colonies on the Blasket Islands, Co. Kerry, the Inishkea Group, Co. Mayo and the Donegal coast during the 2003 breeding season
Author :	Cronin, M.; Ó Cadhla, O.
Series :	Unpublished report to NPWS
Year :	2007
Title :	Grey seal moult population survey in the Republic of Ireland, 2007
Author :	Ó Cadhla, O.; Strong, D.
Series :	Unpublished report to NPWS
Year :	2008
Title :	An assessment of the breeding population of grey seals in the Republic of Ireland, 2005
Author :	O Cadhla, O.; Strong, D.; O'Keeffe, C.; Coleman, M.; Cronin, M.; Duck, C.; Murray, T.; Dower, P.; Nairn, R.; Murphy, P.; Smiddy, P.; Saich, C.; Lyons, D.O.; Hiby, L.
Series :	Irish Wildlife Manual No. 34
Year :	2009
Title :	Coastal Monitoring Project 2004-2006
Author :	Ryle, T.; Murray, A.; Connolly, K.; Swann, M.
Series :	Unpublished report to NPWS
Year :	2013
Title :	Monitoring of the breeding population of grey seals in Ireland, 2009 - 2012
Author :	Ó Cadhla, O.; Keena, T.; Strong, D.; Duck, C.; Hiby, L.
Series :	Irish Wildlife Manual No. 74
Year :	2013
Title :	An aerial survey of harbour seals in Ireland. Part 1: Lough Foyle to Galway Bay. August 2011
Author :	Duck, C; Morris, C.
Series :	Unpublished report to NPWS
Year :	2015
Title :	Inishkea Islands SAC (site code: 507) Conservation objectives supporting document- coastal habitats V1
Author :	NPWS
Series :	Conservation objectives supporting document

Year : 2015
Title : Inishkea Islands SAC (site code: 507) Conservation objectives supporting document- marine species V1
Author : NPWS
Series : Conservation objectives supporting document

Other References

Year : 1983
Title : The grey seal (*Halichoerus grypus*) in Ireland
Author : Summers, C.F.
Series : Unpublished Report to the Minister for Fisheries, Forestry and Wildlife

Year : 1998
Title : Population biology of grey seals (*Halichoerus grypus*, Fabricius 1791) in western Ireland
Author : Kiely, O.R.M.
Series : Unpublished PhD thesis, National University of Ireland, University College Cork

Year : 1998
Title : Grey seal (*Halichoerus grypus*) pup production at the Inishkea island group, Co. Mayo and the Blasket Islands, Co. Kerry
Author : Kiely, O.; Myers, A.A.
Series : Biology and Environment: Proc. Royal Ir. Acad. 98B (2): 113-122

Year : 2001
Title : Grey seal interactions with fisheries in Irish coastal waters
Author : BIM
Series : Report to the European Commission DG XIV. Study 95/40

Year : 2006
Title : The vegetation of Irish machair
Author : Gaynor, K.
Series : Biology and Environment: Proceedings of the Royal Irish Academy, vol 106B, No. 3: 311-321

Year : 2007
Title : Aerial surveying of grey seal breeding colonies on the Blasket Islands, Co. Kerry, the Inishkeas group, Co. Mayo and the Donegal coast, Ireland
Author : Cronin, M.A.; Duck, C.D.; Ó Cadhla, O.
Series : J. Nat. Conserv. 15(2): 77-83

Year : 2008
Title : The phytosociology and conservation value of Irish sand dunes
Author : Gaynor, K.
Series : Unpublished PhD thesis, National University of Ireland, Dublin

Year : 2013
Title : Conservation of selected legally protected and Red Listed bryophytes in Ireland
Author : Campbell, C.
Series : Unpublished Ph.D. Thesis, Trinity College Dublin

Spatial data sources

Year : 2009
Title : Coastal Monitoring Project 2004-2006. Version 1
GIS Operations : QIs selected; clipped to SAC boundary; overlapping regions with Saltmarsh CO data investigated and resolved with expert opinion used
Used For : 21A0 (map 3)

Year : 2015
Title : NPWS rare and threatened species database
GIS Operations : Dataset created from spatial references in database records. Expert opinion used as necessary to resolve any issues arising
Used For : 1364, 1395 (maps 4 and 5)

Conservation Objectives for : Inishkea Islands SAC [000507]

21A0 Machairs (* in Ireland)

To restore the favourable conservation condition of Machairs in Inishkea Islands SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes including erosion and succession. For sub-site mapped: Inishkea Islands - 115.65ha. See map 3	Based on data from the Coastal Monitoring Project (CMP) (Ryle et al., 2009). One sub-site was mapped, giving a total estimated area of 115.65ha. See coastal habitats supporting document for further details
Habitat distribution	Occurrence	No decline, subject to natural processes. See map 3	Based on data from Ryle et al. (2009). Iniskea North is low-lying and dominated by machair vegetation. At Inishkea South there is a small area of machair in the north-eastern part of the island. See coastal habitats supporting document for further details
Physical structure: functionality and sediment supply	Presence/ absence of physical barriers	Maintain the natural circulation of sediment and organic matter, without any physical obstructions	Based on data from Ryle et al. (2009). The CMP noted that the machair at Inishkea North was severely eroded, which had been exacerbated by overgrazing. See coastal habitats supporting document for further details
Vegetation structure: zonation	Occurrence	Maintain the range of coastal habitats including transitional zones, subject to natural processes including erosion and succession	Based on data from Ryle et al. (2009). In places, the machair and heath habitats grade into saltmarsh, particularly along their fringes. Other habitats present within the SAC include open marine water, lowland dry grassland, wet grassland, boulder beaches, shingle and bedrock shores. See coastal habitats supporting document for further details
Vegetation structure: bare ground	Percentage cover	Bare ground should not exceed 10% of Machair habitat, subject to natural processes	Based on data from Ryle et al. (2009). <i>Petalophyllum ralfsii</i> was recorded on the north island along a well worn sheep path- see the conservation objective for 1395. Bare ground cover greater than 10% was recorded by the CMP in monitoring stops on Inishkea South. See coastal habitats supporting document for further details
Vegetation structure: sward height	Centimeters	Maintain structural variation within sward	Based on data from Gaynor (2008) and Ryle et al. (2009). The machair on the Iniskea Islands was overgrazed. Inishkea North and South are grazed by rabbits (<i>Oryctolagus cuniculus</i>), barnacle geese (<i>Branta leucopsis</i>) and sheep. See coastal habitats supporting document for further details
Vegetation composition: typical species and sub-communities	Percentage cover at a representative number of monitoring stops	Maintain range of sub-communities with typical species listed in Ryle et al. (2009)	Based on data from Gaynor (2006) and Ryle et al. (2009). See coastal habitats supporting document for further details
Vegetation composition: negative indicator species	Percentage cover	Negative indicator species (including non-natives) to represent less than 5% cover	Based on data from Ryle et al. (2009). Negative indicators include non-native species, species indicative of changes in nutrient status and species not considered characteristic of the habitat. On Inishkea North, the negative indicators Yorkshire fog (<i>Holcus lanatus</i>) and ragwort (<i>Senecio jacobaea</i>) were noted by the CMP. On Inishkea South, thistles (<i>Cirsium</i> spp.) were present. See coastal habitats supporting document for further details
Vegetation composition: scrub/trees	Percentage cover	No more than 5% cover or under control	Based on data from Ryle et al. (2009). See coastal habitats supporting document for further details
Vegetation composition: bryophytes	Percentage cover	Should always be at least an occasional component of the vegetation	Based on data from Ryle et al. (2009). See coastal habitats supporting document for further details

Conservation Objectives for : Inishkea Islands SAC [000507]**1364 Grey Seal *Halichoerus grypus*****To maintain the favourable conservation condition of Grey Seal in Inishkea Islands SAC, which is defined by the following list of attributes and targets:**


Attribute	Measure	Target	Notes
Access to suitable habitat	Number of artificial barriers	Species range within the SAC should not be restricted by artificial barriers to site use. See map 4	See marine supporting document for further details
Breeding behaviour	Breeding sites	Conserve the breeding sites in a natural condition. See map 4 for known sites	Attribute and target based on background knowledge of Irish breeding populations, repeated breeding surveys in 1995, 1996 (Kiely, 1998; Kiely and Myers, 1998), 1998 and 1999 (BIM, 2001), 2002 (Ó Cadhla and Strong, 2003), 2003 (Cronin and Ó Cadhla, 2004; Cronin et al., 2007), 2005 (Ó Cadhla et al., 2008) and 2011 (Ó Cadhla et al., 2013), and unpublished NPWS records including those reported by Summers (1983) and Lyons (2004). See marine supporting document for further details
Moult behaviour	Moult haul-out sites	Conserve the moult haul-out sites in a natural condition. See map 4 for known sites	Attribute and target based on background knowledge of Irish populations, on review of data from Kiely (1998) and Lyons (2004), a national moult survey (Ó Cadhla and Strong, 2007) and unpublished NPWS records. See marine supporting document for further details
Resting behaviour	Resting haul-out sites	Conserve the resting haul-out sites in a natural condition. See map 4 for known sites	Attribute and target based on review of data from Kiely (1998), BIM (2001), Lyons (2004), Cronin et al. (2004), Ó Cadhla et al. (2008), Duck and Morris (2013) and unpublished NPWS. See marine supporting document for further details
Disturbance	Level of impact	Human activities should occur at levels that do not adversely affect the grey seal population at the site	See marine supporting document for further details

Conservation Objectives for : Inishkea Islands SAC [000507]

1395 Petalwort *Petalophyllum ralfsii*


To maintain the favourable conservation condition of Petalwort in Inishkea Islands SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Distribution of populations	Number and geographical spread of populations	No decline. See map 5 for recorded location	The population at North Inishkea was recorded at intervals along a well-worn sheep track traversing the lower edge of machair plain, c.100m from the sea by N. Lockhart of NPWS in July 1998. The islands have not been comprehensively searched for the species since that time
Population size	Number of individuals	No decline. The population has been estimated at a minimum of 7 thalli	Lockhart recorded a minimum of 7 thalli in July 1998
Area of suitable habitat	Hectares	No decline. Area of suitable habitat at North Inishkea currently unknown, but thought to be very small, c.0.00003ha	The extent of suitable habitat around the recorded location at North Inishkea has not been accurately measured using GPS, but is known to be very small (c.0.25m ²)
Hydrological conditions: soil moisture	Occurrence of damp soil conditions	Maintain hydrological conditions so that substrate is kept moist and damp throughout the year, but not subject to prolonged inundation by flooding in winter	<i>Petalophyllum ralfsii</i> grows in damp sand. Based on Campbell (2013)
Vegetation: open structure	Height and percentage cover of vegetation	Maintain open, low vegetation, with a high percentage cover of bryophytes (small acocarps and liverwort turf) and bare ground	<i>Petalophyllum ralfsii</i> grows in compacted, sandy ground, maintained by grazing and trampling (by sheep). Only recorded along sheep track but plenty of other potentially suitable habitat available (Lockhart, 1998, unpublished data)



Legend

 Inishkea Islands SAC 000507




SAC 000507


SPA 004004

SAC 002998




Legend


- Inishkea Islands SAC 000507
- West Connacht Coast SAC 002998
- Inishkea Islands SPA 004004






Legend


-  Inishkea Islands SAC 000507
-  21A0 *Machairs
-  OSi Discovery Series County Boundary



Legend


- Inishkea Islands SAC 000507
- ★ 1364 Grey Seal - *Halichoerus grypus* breeding sites
- ★ 1364 Grey Seal - *Halichoerus grypus* moulting sites
- ★ 1364 Grey Seal - *Halichoerus grypus* resting sites
- 1364 Grey Seal - *Halichoerus grypus* habitats
- OSi Discovery Series County Boundary





Legend

- Inishkea Islands SAC 000507
- 1395 Petalwort - *Petalophyllum ralfsii*
- OSi Discovery Series County Boundary



National Parks and Wildlife Service

Conservation Objectives Series

Lackan Saltmarsh and Kilcummin Head SAC 000516



An Roinn Ealaíon, Oidhreachta,
Gnóthaí Réigiúnacha, Tuaithe agus Gaeltachta

Department of Arts, Heritage,
Regional, Rural and Gaeltacht Affairs



**National Parks and Wildlife Service,
Department of Arts, Heritage, Regional, Rural and Gaeltacht Affairs,
7 Ely Place, Dublin 2, Ireland.**

**Web: www.npws.ie
E-mail: nature.conservation@ahg.gov.ie**

Citation:

**NPWS (2016) Conservation Objectives: Lackan Saltmarsh and Kilcummin Head
SAC 000516. Version 1. National Parks and Wildlife Service, Department of Arts,
Heritage, Regional, Rural and Gaeltacht Affairs.**

**Series Editor: Rebecca Jeffrey
ISSN 2009-4086**

Introduction

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

* indicates a priority habitat under the Habitats Directive

000516	Lackan Saltmarsh and Kilcummin Head SAC
1310	Ulex and other annuals colonising mud and sand
1330	Atlantic salt meadows (<i>Glauco-Puccinellietalia maritimae</i>)
1410	Mediterranean salt meadows (<i>Juncetalia maritimi</i>)
2120	Shifting dunes along the shoreline with Cladonia (white dunes)
2130	Fixed coastal dunes with herbaceous vegetation (grey dunes)E

Please note that this SAC overlaps with Killala Bay/Moy Estuary SPA (004036). See map 2. The conservation objectives for this site should be used in conjunction with those for the overlapping site as appropriate.

Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

NPWS Documents

Year :	2007
Title :	Saltmarsh Monitoring Project 2006
Author :	McCorry, M.
Series :	Unpublished report to NPWS
<hr/>	
Year :	2009
Title :	Coastal Monitoring Project 2004-2006
Author :	Ryle, T.; Murray, A.; Connolly, K.; Swann, M.
Series :	Unpublished report to NPWS
<hr/>	
Year :	2009
Title :	Saltmarsh monitoring project 2007-2008
Author :	McCorry, M.; Ryle, T.
Series :	Unpublished report to NPWS
<hr/>	
Year :	2013
Title :	Monitoring survey of Annex I sand dune habitats in Ireland
Author :	Delaney, A.; Devaney, F.M.; Martin, J.M.; Barron, S.J.
Series :	Irish Wildlife Manual No. 75
<hr/>	
Year :	2016
Title :	Lacken Saltmarsh and Kilcummin Head SAC (site code: 516) Conservation objectives supporting document- coastal habitats V1
Author :	NPWS
Series :	Conservation objectives supporting document

Other References

Year :	2008
Title :	The phytosociology and conservation value of Irish sand dunes
Author :	Gaynor, K.
Series :	Unpublished PhD thesis, National University of Ireland, Dublin

Spatial data sources

Year :	Revision 2010
Title :	Saltmarsh Monitoring Project 2007-2008. Version 1
GIS Operations :	QIs selected; clipped to SAC boundary; overlapping regions with Coastal CO data investigated and resolved with expert opinion used
Used For :	1310, 1330, 1410 (map 3)
<hr/>	
Year :	2009
Title :	Coastal Monitoring Project 2004-2006. Version 1
GIS Operations :	QIs selected; clipped to SAC boundary; overlapping regions with Saltmarsh CO data investigated and resolved with expert opinion used
Used For :	2120, 2130 (map 4)
<hr/>	

Conservation Objectives for : Lackan Saltmarsh and Kilcummin Head SAC [000516]

1310 **Salicornia and other annuals colonising mud and sand**

To restore the favourable conservation condition of *Salicornia* and other annuals colonising mud and sand in Lackan Saltmarsh and Kilcummin Head SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes, including erosion and succession. For the sub-site mapped: Lackan - 0.001ha	Based on data from the Saltmarsh Monitoring Project (SMP) (McCorry, 2007; McCorry and Ryle, 2009). <i>Salicornia</i> and other annuals colonising mud and sand was surveyed at the sub-site Lackan (site ID: SMP0022) to give a total estimated area of 0.001ha in Lackan Saltmarsh and Kilcummin Head SAC. This extent is too small to be mapped. NB further unsurveyed areas may be present within the SAC. See the Lackan Saltmarsh and Kilcummin Head SAC conservation objectives supporting document for coastal habitats for further details
Habitat distribution	Occurrence	No decline or change in habitat distribution, subject to natural processes	Based on data from McCorry (2007) and McCorry and Ryle (2009). <i>Salicornia</i> is an annual species, so its distribution can vary significantly from year to year. See the coastal habitats supporting document for further details
Physical structure: sediment supply	Presence/absence of physical barriers	Maintain, or where necessary restore, natural circulation of sediments and organic matter, without any physical obstructions	Based on data from McCorry (2007) and McCorry and Ryle (2009). Sediment supply is particularly important for this pioneer saltmarsh community, as its distribution depends on accretion rates. Within the estuary and along the margins of the Cloonalaghan River, sediments originating from the river have built up to form an extensive saltmarsh (Ryle et al., 2009). See the coastal habitats supporting document for further details
Physical structure: creeks and pans	Occurrence	Maintain creek and pan structure, subject to natural processes, including erosion and succession	Based on data from McCorry (2007) and McCorry and Ryle (2009). Creeks deliver sediment throughout the saltmarsh system. At Lackan, the creek network is well-developed and many of the creeks contain very soft mud and are unusually deep. See the coastal habitats supporting document for further details
Physical structure: flooding regime	Hectares flooded; frequency	Maintain natural tidal regime	Based on data from McCorry (2007) and McCorry and Ryle (2009). This pioneer saltmarsh community requires regular tidal inundation. See the coastal habitats supporting document for further details
Vegetation structure: zonation	Occurrence	Maintain the range of coastal habitats including transitional zones, subject to natural processes including erosion and succession	Based on data from McCorry (2007) and McCorry and Ryle (2009). See the coastal habitats supporting document for further details
Vegetation structure: sward height	Centimetres	Maintain structural variation within sward	Based on data from McCorry (2007) and McCorry and Ryle (2009). See the coastal habitats supporting document for further details
Vegetation structure: vegetation cover	Percentage cover at a representative number of monitoring stops	Maintain more than 90% of the area outside of creeks vegetated	Based on data from McCorry (2007) and McCorry and Ryle (2009). See the coastal habitats supporting document for further details
Vegetation composition: typical species and sub-communities	Percentage cover	Maintain the presence of species-poor communities with typical species listed in McCorry and Ryle (2009)	Based on data from McCorry (2007) and McCorry and Ryle (2009). There is frequent glasswort (<i>Salicornia</i> sp.) and occasional annual sea-blite (<i>Suaeda maritima</i>) associated with some areas. See the coastal habitats supporting document for further details
Vegetation composition: negative indicator species - <i>Spartina anglica</i>	Hectares	There is no record of common cordgrass (<i>Spartina anglica</i>) in the SAC and its establishment should be prevented	Based on data from McCorry (2007) and McCorry and Ryle (2009). No common cordgrass (<i>Spartina anglica</i>) was recorded in this habitat in the SAC. See the coastal habitats supporting document for further details

1330 Atlantic salt meadows (*Glauco-Puccinellietalia maritima*)

To maintain the favourable conservation condition of Atlantic salt meadows (*Glauco-Puccinellietalia maritima*) in Lackan Saltmarsh and Kilcummin Head SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes, including erosion and succession. For the sub-site (Lackan) and potential areas mapped: 32.70ha. See map 3	Based on data from the Saltmarsh Monitoring Project (SMP) (McCorry, 2007; McCorry and Ryle, 2009). The sub-site Lackan (site ID: SMP0022) that supports Atlantic Salt Meadows (ASM) was mapped (32.43ha) and additional areas of potential ASM habitat (0.27ha) were identified from an examination of aerial photographs, giving a total estimated area of 32.70ha within Lackan Saltmarsh and Kilcummin Head SAC. NB further unsurveyed areas may be present within the SAC. See the Lackan Saltmarsh and Kilcummin Head SAC conservation objectives supporting document for coastal habitats for further details
Habitat distribution	Occurrence	No decline or change in habitat distribution, subject to natural processes. See map 3 for known distribution	Based on data from McCorry (2007) and McCorry and Ryle (2009). The saltmarsh is mostly contained in one large main unit. A band of saltmarsh extends along the north-western and north-eastern shorelines of Lackan Bay, which eventually narrows out and transitions to sand dune and sandy beach habitats. NB further unsurveyed areas may be present within the SAC. See the coastal habitats supporting document for further details
Physical structure: sediment supply	Presence/absence of physical barriers	Maintain natural circulation of sediments and organic matter, without any physical obstructions	Based on data from McCorry (2007) and McCorry and Ryle (2009). Erosion and accretion mainly affects the ASM at this SAC. See the coastal habitats supporting document for further details
Physical structure: creeks and pans	Occurrence	Maintain creek and pan structure, subject to natural processes, including erosion and succession	Based on data from McCorry (2007) and McCorry and Ryle (2009). The original creek network has been affected by drainage and some of the channels in the mid-eastern part of the saltmarsh have been artificially deepened and straightened in the past. See the coastal habitats supporting document for further details
Physical structure: flooding regime	Hectares flooded; frequency	Maintain natural tidal regime	Based on data from McCorry (2007) and McCorry and Ryle (2009). There have been drainage and land reclamation works in the past with regularly-spaced drains across the north-western section of the saltmarsh linking with drains from adjacent wet grassland on slopes to the Cloonalaghan River. See the coastal habitats supporting document for further details
Vegetation structure: zonation	Occurrence	Maintain the range of coastal habitats including transitional zones, subject to natural processes including erosion and succession	Based on data from McCorry (2007) and McCorry and Ryle (2009). Natural transitions occur between saltmarsh types as well as to other coastal habitats such as sand dunes. See the coastal habitats supporting document for further details
Vegetation structure: sward height	Centimetres	Maintain structural variation within sward	Based on data from McCorry (2007) and McCorry and Ryle (2009). Sheep grazing has created a typical low sward (1-2cm high). See the coastal habitats supporting document for further details
Vegetation structure: vegetation cover	Percentage cover at a representative number of monitoring stops	Maintain more than 90% of the area outside of creeks vegetated	Based on data from McCorry (2007) and McCorry and Ryle (2009). There are vehicle tracks and wheel ruts on the ASM at the north-western and north-eastern corners of the saltmarsh where minor roads allow access to the sandflats and Lackan Bay. See the coastal habitats supporting document for further details

Vegetation composition: typical species and sub-communities	Percentage cover at a representative number of monitoring stops	Maintain range of sub-communities with typical species listed in McCorry and Ryle (2009)	Based on data from McCorry (2007) and McCorry and Ryle (2009). ASM vegetation is dominated by a thrift (<i>Armeria maritima</i>) and sea plantain (<i>Plantago maritima</i>) sward. See the coastal habitats supporting document for further details
Vegetation composition: negative indicator species - <i>Spartina anglica</i>	Hectares	There is no record of common cordgrass (<i>Spartina anglica</i>) in the SAC and its establishment should be prevented	Based on data from McCorry (2007) and McCorry and Ryle (2009). No common cordgrass (<i>Spartina anglica</i>) was recorded in this habitat in the SAC. See the coastal habitats supporting document for further details

Conservation Objectives for : Lackan Saltmarsh and Kilcummin Head SAC [000516]

1410 Mediterranean salt meadows (*Juncetalia maritimi*)

To restore the favourable conservation condition of Mediterranean salt meadows (*Juncetalia maritimi*) in Lackan Saltmarsh and Kilcummin Head SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes, including erosion and succession. For the sub-site (Lackan): 65.03ha. See map 3	Based on data from the Saltmarsh Monitoring Project (SMP) (McCorry, 2007; McCorry and Ryle, 2009). The sub-site Lackan (site ID: SMP0022) that supports Mediterranean Salt Meadows (MSM) was mapped to give a total estimated area of 65.03ha within Lackan Saltmarsh and Kilcummin Head SAC. NB further unsurveyed areas may be present within the SAC. See the Lackan Saltmarsh and Kilcummin Head SAC conservation objectives supporting document for coastal habitats for further details
Habitat distribution	Occurrence	No decline or change in habitat distribution, subject to natural processes. See map 3 for known distribution	Based on data from McCorry (2007) and McCorry and Ryle (2009). MSM habitat dominates the western side of Cloonalaghan River and the southern part of the saltmarsh. NB further unsurveyed areas may be present within the SAC. See the coastal habitats supporting document for further details
Physical structure: sediment supply	Presence/absence of physical barriers	Maintain natural circulation of sediments and organic matter, without any physical obstructions	Based on data from McCorry (2007) and McCorry and Ryle (2009). Some minor erosion and accretion occurs within the MSM further up the Cloonalaghan River channel from the ASM habitat. See the coastal habitats supporting document for further details
Physical structure: creeks and pans	Occurrence	Maintain creek and pan structure, subject to natural processes, including erosion and succession	Based on data from McCorry (2007) and McCorry and Ryle (2009). The creek and pan topography in the MSM is very well-developed with frequent pans and a dense network of creeks. See the coastal habitats supporting document for further details
Physical structure: flooding regime	Hectares flooded; frequency	Maintain natural tidal regime	Based on data from McCorry (2007) and McCorry and Ryle (2009). Mediterranean salt meadow is found high up in the saltmarsh but requires occasional tidal inundation. See the coastal habitats supporting document for further details
Vegetation structure: zonation	Occurrence	Maintain the range of coastal habitats including transitional zones, subject to natural processes including erosion and succession	Based on data from McCorry (2007) and McCorry and Ryle (2009). Natural transitions occur between saltmarsh types as well as to other coastal habitats such as sand dunes. See the coastal habitats supporting document for further details
Vegetation structure: sward height	Centimetres	Maintain structural variation in the sward	Based on data from McCorry (2007) and McCorry and Ryle (2009). The grazing level is low in the MSM as the dense patches of sea rush (<i>Juncus maritimus</i>) present protect the other vegetation. See the coastal habitats supporting document for further details
Vegetation structure: vegetation cover	Percentage cover at a representative number of monitoring stops	Maintain more than 90% of the area outside of creeks vegetated	Based on data from McCorry (2007) and McCorry and Ryle (2009). The MSM habitat has suffered some damage due to heavy cattle poaching. See the coastal habitats supporting document for further details
Vegetation composition: typical species and sub-communities	Percentage cover at a representative number of monitoring stops	Maintain range of sub-communities with typical species listed in McCorry and Ryle (2009)	Based on data from McCorry (2007) and McCorry and Ryle (2009). Sea rush (<i>Juncus maritimus</i>) occurs on slightly elevated sites and its sharp stems protect succulent plants such as common scurvygrass (<i>Cochlearia officinalis</i>) and sea aster (<i>Aster tripolium</i>) from grazing. Sea club-rush (<i>Bolboschoenus maritimus</i>) and common reed (<i>Phragmites australis</i>) are present in the ditches. This limited species diversity is typical of MSM habitat. See the coastal habitats supporting document for further details

Vegetation composition: negative indicator species - <i>Spartina anglica</i>	Hectares	There is no record of common cordgrass (<i>Spartina anglica</i>) in the SAC and its establishment should be prevented	Based on data from McCorry (2007) and McCorry and Ryle (2009). No common cordgrass (<i>Spartina anglica</i>) was recorded in this habitat in the SAC. See the coastal habitats supporting document for further details
--	----------	---	--

Conservation Objectives for : Lackan Saltmarsh and Kilcummin Head SAC [000516]

2120 Shifting dunes along the shoreline with *Ammophila arenaria* (white dunes)

To restore the favourable conservation condition of Shifting dunes along the shoreline with *Ammophila arenaria* (white dunes) in Lackan Saltmarsh and Kilcummin Head SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes, including erosion and succession. For the sub-site mapped: Lackan (including Rathlackan) - 2.82ha. See map 4	Based on data from the Coastal Monitoring Project (CMP) (Ryle et al., 2009). Shifting dunes along the shoreline with <i>Ammophila arenaria</i> was mapped at the sub-site Lackan (including Rathlackan; CMP site ID: 129) to give a total estimated area of 2.82ha within Lackan Saltmarsh and Kilcummin Head SAC. This habitat is very difficult to measure in view of its dynamic nature. See the Lackan Saltmarsh and Kilcummin Head SAC conservation objectives supporting document for coastal habitats for further details
Habitat distribution	Occurrence	No decline or change in habitat distribution, subject to natural processes. See map 4 for known distribution	Based on data from Ryle et al. (2009). See the coastal habitats supporting document for further details
Physical structure: functionality and sediment supply	Presence/absence of physical barriers	Maintain the natural circulation of sediment and organic matter, without any physical obstructions	Based on data from Ryle et al. (2009). Dunes are naturally dynamic systems that require continuous supply and circulation of sand. Marram grass (<i>Ammophila arenaria</i>) reproduces vegetatively and requires constant accretion of fresh sand to maintain active growth encouraging further accretion. The sandhills at the Rathlackan sub-site, on the north-west side of Lackan Saltmarsh and Kilcummin Head SAC, are badly eroded, which has resulted in the availability of sediment that may be re-worked to form temporary foredune habitat. There appears to have been some attempts at dune protection through the planting of marram grass and lyme-grass (<i>Leymus arenarius</i>) on heaped banks of sand and cobbles. See the coastal habitats supporting document for further details
Vegetation structure: zonation	Occurrence	Maintain the range of coastal habitats including transitional zones, subject to natural processes including erosion and succession	Based on data from Gaynor (2008) and Ryle et al. (2009). Mobile dunes at Rathlackan extend around the seaward edge of the spit. Behind the dunes, there are sheltered intertidal sandflats which in turn are backed by extensive saltmarsh. See the coastal habitats supporting document for further details
Vegetation composition: plant health of dune grasses	Percentage cover	More than 95% of marram grass (<i>Ammophila arenaria</i>) and/or lyme-grass (<i>Leymus arenarius</i>) should be healthy (i.e. green plant parts above ground and flowering heads present)	Based on data from Ryle et al. (2009). Although mobile dunes occur along the full northern edge of the spit in the SAC, the characteristic vegetation of marram (<i>Ammophila arenaria</i>) is frequently quite sparse and/or has an unhealthy appearance, reflecting the general lack of sediment mobility along the seaward edge of the dunes. Only at the western tip of the spit, where accreting or locally recycled sediment accumulates, is there a substantial band of healthy marram. See the coastal habitats supporting document for further details
Vegetation composition: typical species and sub-communities	Percentage cover at a representative number of monitoring stops	Maintain the presence of species-poor communities dominated by marram grass (<i>Ammophila arenaria</i>) and/or lyme-grass (<i>Leymus arenarius</i>)	Based on data from Ryle et al. (2009). The mobile dune habitat at Rathlackan is characterised by the presence of marram grass (<i>Ammophila arenaria</i>). Lyme-grass (<i>Leymus arenarius</i>) is also present in places. See the coastal habitats supporting document for further details
Vegetation composition: negative indicator species	Percentage cover	Negative indicator species (including non-native species) to represent less than 5% cover	Based on data from Ryle et al. (2009). Negative indicators include non-native species, species indicative of changes in nutrient status and species not considered characteristic of the habitat. Sea buckthorn (<i>Hippophae rhamnoides</i>) should be absent or effectively controlled. See the coastal habitats supporting document for further details


Conservation Objectives for : Lackan Saltmarsh and Kilcummin Head SAC [000516]

2130 Fixed coastal dunes with herbaceous vegetation (grey dunes)


To restore the favourable conservation condition of Fixed coastal dunes with herbaceous vegetation (grey dunes)* in Lackan Saltmarsh and Kilcummin Head SAC, which is defined by the following list of attributes and targets:


Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes, including erosion and succession. For sub-site mapped: Lackan (including Rathlackan) - 95.18ha. See map 4	Based on data from the Coastal Monitoring Project (CMP) (Ryle et al., 2009). Fixed coastal dunes with herbaceous vegetation was mapped at the sub-site Lackan (including Rathlackan; CMP site ID: 129) to give a total estimated area of 95.18ha within Lackan Saltmarsh and Kilcummin Head SAC. See the Lackan Saltmarsh and Kilcummin Head SAC conservation objectives supporting document for coastal habitats for further details
Habitat distribution	Occurrence	No decline or change in habitat distribution, subject to natural processes. See map 4 for known distribution	Based on data from Ryle et al. (2009). See the coastal habitats supporting document for further details
Physical structure: functionality and sediment supply	Presence/absence of physical barriers	Maintain the natural circulation of sediment and organic matter, without any physical obstructions	Based on data from Ryle et al. (2009). Physical barriers can lead to fossilisation or over-stabilisation of dunes, as well as beach starvation resulting in increased rates of erosion. The north-facing (seaward) side of the Lackan dunes has a highly eroded dune face which, coupled with the lack of any substantially accreting habitat and no significant foredune development, suggests the system is being depleted of sediment. See the coastal habitats supporting document for further details
Vegetation structure: zonation	Occurrence	Maintain the range of coastal habitats including transitional zones, subject to natural processes including erosion and succession	Based on data from Ryle et al. (2009). The outer zone of Lackan Saltmarsh and Kilcummin Head SAC is dominated by a sand dune system and a sandy beach. The sand dunes are dominated by fixed dunes. Behind the dunes, there are sheltered intertidal sandflats which in turn are backed by extensive saltmarsh. See the coastal habitats supporting document for further details
Vegetation structure: bare ground	Percentage cover	Bare ground should not exceed 10% of fixed dune habitat, subject to natural processes	Based on data from Gaynor (2008) and Ryle et al. (2009). See the coastal habitats supporting document for further details
Vegetation structure: sward height	Centimetres	Maintain structural variation within sward	Based on data from Gaynor (2008) and Ryle et al. (2009). Different levels of grazing have resulted in varying sward heights in the fixed dune habitat at this SAC. See the coastal habitats supporting document for further details
Vegetation composition: typical species and sub-communities	Percentage cover at a representative number of monitoring stops	Maintain the range of sub-communities with typical species listed in Delaney et al. (2013)	Based on data from Gaynor (2008) and Ryle et al. (2009). The more commonly noted species in the fixed dunes included sand sedge (<i>Carex arenaria</i>), glaucous sedge (<i>C. flacca</i>), red fescue (<i>Festuca rubra</i>), lady's bedstraw (<i>Galium verum</i>), cat's ear (<i>Hypochaeris radicata</i>), common bird's-foot trefoil (<i>Lotus corniculatus</i>), field wood-rush (<i>Luzula campestris</i>), mouse-ear-hawkweed (<i>Pilosella officinarum</i>), ribwort plantain (<i>Plantago lanceolata</i>), yellow-rattle (<i>Rhinanthus minor</i>), wild thyme (<i>Thymus polytrichus</i>) and Germander speedwell (<i>Veronica chamaedrys</i>). See the coastal habitats supporting document for further details

Vegetation composition: negative indicator species	Percentage cover	Negative indicator species (including non-native species) to represent less than 5% cover	Based on data from Gaynor (2008) and Ryle et al. (2009). Negative indicators include non-native species, species indicative of changes in nutrient status and species not considered characteristic of the habitat. Sea buckthorn (<i>Hippophae rhamnoides</i>) should be absent or effectively controlled. At Lackan Saltmarsh and Kilcummin Head SAC, the localised proliferation of species such as creeping thistle (<i>Cirsium arvense</i>), spear thistle (<i>C. vulgare</i>) and common ragwort (<i>Senecio jacobaea</i>) in the fixed dunes may be indicative of recent overgrazing and intensive management. See the coastal habitats supporting document for further details
Vegetation composition: scrub/trees	Percentage cover	No more than 5% cover or under control	Based on data from Ryle et al. (2009). At Lackan Saltmarsh and Kilcummin Head SAC, there were occasional stunted hawthorn (<i>Crataegus monogyna</i>) shrubs in the fixed dune grassland, although the total shrub and tree cover was insignificant. See the coastal habitats supporting document for further details



Legend

 Lackan Saltmarsh and Kilcummin Head SAC 000516

 An Poinn Ealaíon, Oidhreacht, Gnóthaí Réigiúnacha, Tuaithe agus Gaeltachta
 Department of Arts, Heritage, Regional, Rural and Gaeltacht Affairs

**MAP 1:
 LACKAN SALTMARSH AND KILCUMMIN HEAD SAC
 CONSERVATION OBJECTIVES
 SAC DESIGNATION**

Map to be read in conjunction with the NPWS Conservation Objectives Document.

**SITE CODE:
 SAC 000516; version 3.
 CO. MAYO**




0 0.25 0.5 0.75 1 km

The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
 Ordnance Survey of Ireland Licence No EN 0059216. © Ordnance Survey of Ireland Government of Ireland.

Níl sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbhreithniú a déanamh ar theorainneacha na gceantar comharthaithe. Suirbhéarachtá Ordonáis na hÉireann Ceadúnas Uimh EN 0059216. © Suirbhéarachtá Ordonáis na hÉireann Rialtas na hÉireann.


**Map Version 1
 Date: Oct 2016**

© Ordnance Survey Ireland



Legend

- Lackan Saltmarsh and Kilcummin Head SAC 000516
- Killala Bay/Moy Estuary SPA 004036
- OSi Discovery Series County Boundary

An Poinn Ealaíon, Oidhreacht, Gnóthai Réigiúnacha, Tuaithe agus Gaeltachta
 Department of Arts, Heritage, Regional, Rural and Gaeltacht Affairs

MAP 2:
LACKAN SALTMARSH AND KILCUMMIN HEAD SAC
CONSERVATION OBJECTIVES
OVERLAPPING DESIGNATIONS


Map to be read in conjunction with the NPWS Conservation Objectives Document.

SITE CODE:
SAC 000516; version 3. SPA 004036; version 3.
CO. MAYO

The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
 Ordnance Survey of Ireland Licence No EN 0059216. © Ordnance Survey of Ireland Government of Ireland.


Níl sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbheithnihe a déanamh ar theorainneacha na gceantar comharthaithe. Suirbhéarachta Ordonáis na hÉireann Ceadúnas Uimh EN 0059216. © Suirbhéarachta Ordonáis na hÉireann Rialtas na hÉireann.

Map Version 1
Date: Oct 2016



Legend

- Lackan Saltmarsh and Kilcummin Head SAC 000516
- Saltmarsh Monitoring Project Study Area
- 1330 / 1410 Atlantic salt meadows (*Glauco-Puccinellietalia maritimae*) / Mediterranean salt meadows (*Juncetalia maritimi*)
- 1330 Atlantic salt meadows (*Glauco-Puccinellietalia maritimae*)
- 1410 Mediterranean salt meadows (*Juncetalia maritimi*)
- Potential 1330 Atlantic salt meadows (*Glauco-Puccinellietalia maritimae*)
- OSi Discovery Series County Boundary



Legend

- Lackan Saltmarsh and Kilcummin Head SAC 000516
- Coastal Monitoring Project Survey Area

Qualifying Interests

- 2120, Shifting dunes along the shoreline with *Ammophila arenaria* ("white dunes")
- 2130, Fixed dunes with herbaceous vegetation ("grey dunes")

Non Qualifying Interests

- 2110 Embryonic shifting dunes
- OSi Discovery Series County Boundary

National Parks and Wildlife Service

Conservation Objectives Series

Lough Gall Bog SAC 000522



An Roinn Ealaíon, Oidhreachta,
Gnóthaí Réigiúnacha, Tuaithe agus Gaeltachta

Department of Arts, Heritage,
Regional, Rural and Gaeltacht Affairs



**National Parks and Wildlife Service,
Department of Arts, Heritage, Regional, Rural and Gaeltacht Affairs,
7 Ely Place, Dublin 2, Ireland.
Web: www.npws.ie
E-mail: nature.conservation@ahg.gov.ie**

Citation:

**NPWS (2017) Conservation Objectives: Lough Gall Bog SAC 000522. Version 1.
National Parks and Wildlife Service, Department of Arts, Heritage, Regional,
Rural and Gaeltacht Affairs.**

**Series Editor: Rebecca Jeffrey
ISSN 2009-4086**

Introduction

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

** indicates a priority habitat under the Habitats Directive*

000522	Lough Gall Bog SAC
7130	Blanket bogs (* if active bog)
7150	Depressions on peat substrates of the Rhynchosporion

Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

NPWS Documents

Year :	1987
Title :	A survey to locate blanket bogs of scientific interest in County Mayo. Part I
Author :	Foss, P.; McGee, E.
Series :	A report commissioned by the Wildlife Service
<hr/>	
Year :	2012
Title :	Ireland Red List No. 8: Bryophytes
Author :	Lockhart, N.; Hodgetts, N.; Holyoak, D.
Series :	Ireland Red List series, NPWS
<hr/>	
Year :	2014
Title :	Guidelines for a national survey and conservation assessment of upland vegetation and habitats in Ireland, Version 2.0
Author :	Perrin, P.M.; Barron, S.J.; Roche, J.R.; O'Hanrahan, B.
Series :	Irish Wildlife Manual No. 79
<hr/>	
Year :	2016
Title :	Ireland Red List No. 10: Vascular Plants
Author :	Wyse Jackson, M.; FitzPatrick, Ú.; Cole, E.; Jebb, M.; McFerran, D.; Sheehy Skeffington, M.; Wright, M.
Series :	Ireland Red Lists series, NPWS
<hr/>	
Year :	2017
Title :	Lough Gall Bog SAC (site code: 522) Conservation objectives supporting document- blanket bog and associated habitats V1
Author :	NPWS
Series :	Conservation objectives supporting document

Other References

Conservation Objectives for : Lough Gall Bog SAC [000522]

7130 Blanket bogs (* if active bog)

To maintain the favourable conservation condition of Blanket bogs (* if active bog) in Lough Gall Bog SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Blanket bog has not been mapped in detail for Lough Gall Bog SAC but from current available data the total area of the qualifying habitat is estimated to be approximately 297ha, covering 82% of the SAC (NPWS internal files). Further details on this and the following attributes can be found in the Lough Gall Bog SAC conservation objectives supporting document for blanket bogs and associated habitats
Habitat distribution	Occurrence	No decline, subject to natural processes	The habitat is documented to occur throughout the SAC and is the dominant habitat present. Further information can be found within Foss and McGee (1987), NPWS internal files and the blanket bogs and associated habitats supporting document
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil nutrient status within natural range	See the blanket bogs and associated habitats supporting document for further details
Ecosystem function: peat formation	Active blanket bog as a proportion of the total area of Annex I blanket bog habitat	At least 99% of the total Annex I blanket bog area is active	See the blanket bogs and associated habitats supporting document for further details
Ecosystem function: hydrology	Flow direction, water levels, occurrence of drains and erosion gullies	Natural hydrology unaffected by drains and erosion	Further details and a brief discussion of restoration potential is presented in the blanket bogs and associated habitats supporting document
Community diversity	Abundance of variety of vegetation communities	Maintain variety of vegetation communities, subject to natural processes	A variety of blanket bog vegetation communities have been recorded in this SAC (Foss and McGee, 1987; NPWS internal files), one of which corresponds to a community recorded in the National Survey of Upland Habitats and listed in the provisional list of vegetation communities described in Perrin et al. (2014). Further information on vegetation communities associated with this habitat is presented in Perrin et al. (2014)
Vegetation composition: positive indicator species	Number of species at a representative number of 2m x 2m monitoring stops	Number of positive indicator species present at each monitoring stop is at least seven	Attribute and target based on Perrin et al. (2014), where the list of positive indicator species for this habitat is also presented
Vegetation composition: lichens and bryophytes	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of bryophytes or lichens, excluding <i>Sphagnum fallax</i> , at least 10%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: potential dominant species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of each of the potential dominant species less than 75%	Attribute and target based on Perrin et al. (2014). See the blanket bogs and associated habitats supporting document for the list of potential dominant species
Vegetation composition: negative indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of negative indicator species less than 1%	Attribute and target based on Perrin et al. (2014), where the list of negative indicator species for this habitat is also presented
Vegetation composition: non-native species	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of non-native species less than 1%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: native trees and shrubs	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of scattered native trees and shrubs less than 10%	Attribute and target based on Perrin et al. (2014)

Vegetation structure: <i>Sphagnum</i> condition	Condition at a representative number of 2m x 2m monitoring stops	Less than 10% of the <i>Sphagnum</i> cover is crushed, broken and/or pulled up	Attribute and target based on Perrin et al. (2014)
Vegetation structure: signs of browsing	Percentage of shoots browsed at a representative number of 2m x 2m monitoring stops	Last complete growing season's shoots of ericoids, crowberry (<i>Empetrum nigrum</i>) and bog-myrtle (<i>Myrica gale</i>) showing signs of browsing collectively less than 33%	Attribute and target based on Perrin et al. (2014)
Vegetation structure: burning	Occurrence in local vicinity of a representative number of monitoring stops	No signs of burning in sensitive areas, into the moss, liverwort or lichen layer or exposure of peat surface due to burning	Attribute and target based on Perrin et al. (2014), where the list of sensitive areas for this habitat is also presented
Physical structure: disturbed bare ground	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of disturbed bare ground less than 10%	Attribute and target based on Perrin et al. (2014)
Physical structure: drainage	Percentage area in local vicinity of a representative number of monitoring stops	Area showing signs of drainage from heavy trampling, tracking or ditches less than 10%	Attribute and target based on Perrin et al. (2014)
Physical structure: erosion	Percentage area in local vicinity of a representative number of monitoring stops	Less than 5% of the greater bog mosaic comprises erosion gullies and eroded areas	Attribute and target based on Perrin et al. (2014)
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat	This includes species listed in the Flora (Protection) Order, 2015 (FPO) and/or the red data lists (Lockhart et al., 2012; Wyse Jackson et al., 2016). The FPO listed and Vulnerable marsh clubmoss (<i>Lycopodiella inundata</i>) (Wyse Jackson et al., 2016) has been recorded within the SAC (NPWS internal files), but this species cannot be specifically assigned to blanket bog




Conservation Objectives for : Lough Gall Bog SAC [000522]

7150 Depressions on peat substrates of the Rhynchosporion


To restore the favourable conservation condition of Depressions on peat substrates of the Rhynchosporion in Lough Gall Bog SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Depressions on peat substrates of the Rhynchosporion has not been mapped in detail for Lough Gall Bog SAC and thus the total area of the qualifying habitat is unknown. Further details on this and the following attributes can be found in the Lough Gall Bog SAC conservation objectives supporting document for blanket bogs and associated habitats
Habitat distribution	Occurrence	No decline, subject to natural processes	The habitat is documented to occur scattered throughout blanket bog areas and is well-represented in locations with deep quaking peat (NPWS internal files). Foss and McGee (1987) note the presence of this habitat in the north-east of the SAC. Further information can be found in Foss and McGee (1987), NPWS internal files and the blanket bogs and associated habitats supporting document
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil nutrient status within natural range	See the blanket bogs and associated habitats supporting document for further details
Vegetation composition: positive indicator species	Number of species at a representative number of 2m x 2m monitoring stops	Number of positive indicator species at each monitoring stop is at least five	Attribute and target based on Perrin et al. (2014), where the list of positive indicator species for this habitat is also presented
Vegetation composition: <i>Rhynchospora</i> spp.	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of white beaked sedge (<i>Rhynchospora alba</i>) and brown beaked sedge (<i>R. fusca</i>) at least 10%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: potential dominant species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of each of the potential dominant species individually less than 35%	Attribute and target based on Perrin et al. (2014). See the blanket bogs and associated habitats supporting document for the list of potential dominant species
Vegetation composition: negative indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of negative indicator species less than 1%	Attribute and target based on Perrin et al. (2014), where the list of negative indicator species for this habitat is also presented
Vegetation composition: non-native species	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of non-native species less than 1%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: native trees and shrubs	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of scattered native trees and shrubs less than 10%	Attribute and target based on Perrin et al. (2014)
Vegetation structure: <i>Sphagnum</i> condition	Condition at a representative number of 2m x 2m monitoring stops	Less than 10% of the <i>Sphagnum</i> cover is crushed, broken and/or pulled up	Attribute and target based on Perrin et al. (2014)
Vegetation structure: signs of browsing	Percentage of shoots browsed at a representative number of 2m x 2m monitoring stops	Last complete growing season's shoots of ericoids, crowberry (<i>Empetrum nigrum</i>) and bog-myrtle (<i>Myrica gale</i>) showing signs of browsing collectively less than 33%	Attribute and target based on Perrin et al. (2014)

Vegetation structure: burning	Occurrence in local vicinity of a representative number of monitoring stops	No signs of burning in sensitive areas, into the moss, liverwort or lichen layer or exposure of peat surface due to burning	Attribute and target based on Perrin et al. (2014), where the list of sensitive areas for this habitat is also presented
Physical structure: disturbed bare ground	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of disturbed bare ground less than 10%	Attribute and target based on Perrin et al. (2014)
Physical structure: drainage	Percentage area in local vicinity of a representative number of monitoring stops	Area showing signs of drainage from heavy trampling, tracking or ditches less than 10%	Attribute and target based on Perrin et al. (2014)
Physical structure: erosion	Percentage area in local vicinity of a representative number of monitoring stops	Less than 5% of the greater bog mosaic comprises erosion gullies and eroded areas	Attribute and target based on Perrin et al. (2014)
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat	This includes species listed in the Flora (Protection) Order, 2015 (FPO) and/or the red data lists (Lockhart et al., 2012; Wyse Jackson et al., 2016). The FPO and Vulnerable marsh clubmoss (<i>Lycopodiella inundata</i>) (Wyse Jackson et al., 2016) is present within the SAC (NPWS internal files), but this species cannot be specifically assigned to this habitat



Legend

 Lough Gall Bog SAC 000522

An Roinn Ealaíon, Oidhreacht, Gnóthai Réigiúnacha, Tuaithe agus Gaeltachta
 Department of Arts, Heritage, Regional, Rural and Gaeltacht Affairs

**MAP 1:
 LOUGH GALL BOG SAC
 CONSERVATION OBJECTIVES
 SAC DESIGNATION**

Map to be read in conjunction with the NPWS Conservation Objectives Document.

**SITE CODE:
 SAC 000522; version 1.01
 Co. Mayo**

0 0.2 0.4 0.6 0.8 km

The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
 Ordnance Survey of Ireland Licence No EN 0059216. © Ordnance Survey of Ireland Government of Ireland.

Níl sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbheithnihe a déanamh ar theorainneacha na gceantar comharthaithe. Suirbhéarachta Ordonáis na hÉireann Ceadúnas Uimh EN 0059216. © Suirbhéarachta Ordonáis na hÉireann Rialtas na hÉireann.



**Map Version 1
 Date: Jan 2017**

© Ordnance Survey Ireland



Conservation objectives for Shrule Turlough SAC [000525]

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Objective: To maintain or restore the favourable conservation condition of the Annex I habitat(s) and/or the Annex II species for which the SAC has been selected:

Code	Description
------	-------------

3180	Turloughs*
------	------------

* denotes a priority habitat



Citation: NPWS (2020) Conservation objectives for Shrule Turlough SAC [000525]. Generic Version
7.0. Department of Culture, Heritage and the Gaeltacht.

National Parks and Wildlife Service

Conservation Objectives Series

Moore Hall (Lough Carra) SAC 000527



An Roinn Cultúir,
Oidhreacht agus Gaeltachta
Department of Culture,
Heritage and the Gaeltacht

**National Parks and Wildlife Service,
Department of Culture, Heritage and the Gaeltacht,
90 King Street North, Dublin 7, D07 N7CV, Ireland.**

**Web: www.npws.ie
E-mail: nature.conservation@chg.gov.ie**

Citation:

**NPWS (2018) Conservation Objectives: Moore Hall (Lough Carra) SAC 000527.
Version 1. National Parks and Wildlife Service, Department of Culture, Heritage
and the Gaeltacht.**

**Series Editor: Rebecca Jeffrey
ISSN 2009-4086**

Introduction

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

** indicates a priority habitat under the Habitats Directive*

000527 Moore Hall (Lough Carra) SAC

1303 Lesser Horseshoe Bat *Rhinolophus hipposideros*

Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

NPWS Documents

Year :	2006
Title :	Bat mitigation guidelines for Ireland
Author :	Kelleher, C.; Marnell, F.
Series :	Irish Wildlife Manual No. 25
<hr/>	
Year :	2018
Title :	Conservation objectives supporting document – lesser horseshoe bat (<i>Rhinolophus hipposideros</i>)
Author :	NPWS
Series :	Conservation objectives supporting document
<hr/>	

Other References

Year :	2007
Title :	Protecting and managing underground sites for bats
Author :	Mitchell-Jones, A.J.; Bihari, Z.; Masing, M.; Rodrigues, L.
Series :	EUROBATS Publication Series No. 2
<hr/>	
Year :	2008
Title :	The lesser horseshoe bat conservation handbook
Author :	Schofield, H.W.
Series :	The Vincent Wildlife Trust
<hr/>	

Spatial data sources


Year :	2018
Title :	NPWS lesser horseshoe bat database
GIS Operations :	Roosts identified, clipped to SAC boundary. Expert opinion used as necessary to resolve any issues arising
Used For :	1303 (map 2)
<hr/>	
Year :	2007
Title :	Forest Inventory and Planning System (FIPS)
GIS Operations :	Dataset clipped to 2.5km buffer centred on roost locations
Used For :	1303 (map 2)
<hr/>	

Conservation Objectives for : Moore Hall (Lough Carra) SAC [000527]

1303 Lesser Horseshoe Bat *Rhinolophus hipposideros*

To maintain the favourable conservation condition of Lesser Horseshoe Bat in Moore Hall (Lough Carra) SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Population per roost	Number	Minimum number of 129 bats for the summer roost (roost id. 684 in NPWS database); minimum number of 455 bats for the winter roost (roost id. 684). See map 2	A figure of 100 bats for summer roosts and 50 bats for winter roosts was set as a minimum qualifying standard (MQS) when SACs were being selected for lesser horseshoe bat (<i>Rhinolophus hipposideros</i>). NPWS conduct annual counts at each qualifying roost. Qualified means from the most recent available five years of data (collected between 2012-2017) have been calculated whereby the year with the highest maximum count and the year with the lowest maximum count were removed and the mean of the remaining years was calculated. This mean is set as the target figure for the roost. The roost in Moore Hall (Lough Carra) SAC (roost id. 684 in NPWS database) is a qualifying summer and winter roost and, as such, separate targets have been set for each season using the summer and winter count data. See the conservation objectives supporting document for lesser horseshoe bat (NPWS, 2018) for further information on all attributes and targets
Winter roosts	Condition	No decline	Moore Hall (Lough Carra) SAC has been selected for lesser horseshoe bat because of the presence of one internationally important winter roost (roost id. 684 in NPWS database). Damage or disturbance to the roost or to the habitat immediately surrounding it will lead to a decline in its condition (Mitchell-Jones et al., 2007)
Summer roosts	Condition	No decline	Moore Hall (Lough Carra) SAC has been selected for lesser horseshoe bat because of the presence of one internationally important summer roost (roost id. 684 in NPWS database). Damage or disturbance to the roost or to the habitat immediately surrounding it will lead to a decline in its condition (Kelleher and Marnell, 2006)
Extent of potential foraging habitat	Hectares	No significant decline within 2.5km of qualifying roosts	Lesser horseshoe bats normally forage in woodlands/scrub within 2.5km of their roosts (Schofield, 2008). See map 2 which shows a 2.5km zone around the above roosts and identifies potential foraging grounds
Linear features	Kilometres	No significant loss within 2.5km of qualifying roosts. See map 2	This species follows commuting routes from its roost to its foraging grounds. Lesser horseshoe bats will not cross open ground. Consequently, linear features such as hedgerows, treelines and stone walls provide vital connectivity for this species within 2.5km around each roost (Schofield, 2008)
Light pollution	Lux	No significant increase in artificial light intensity adjacent to named roosts or along commuting routes within 2.5km of those roosts. See map 2	Lesser horseshoe bats are very sensitive to light pollution and will avoid brightly lit areas. Inappropriate lighting around roosts may cause abandonment; lighting along commuting routes may cause preferred foraging areas to be abandoned, thus increasing energetic costs for bats (Schofield, 2008)




Legend
 Moore Hall (Lough Carra) SAC 000527

 An Roinn Cultúir,
 Oidhreacht agus Gaeltachta
 Department of Culture,
 Heritage and the Gaeltacht

MAP 1:
MOORE HALL (LOUGH CARRA) SAC
CONSERVATION OBJECTIVES
SAC DEIS GNÁTHA
 Map to be read in conjunction with the NPWS Conservation Objectives Document.

SITE CODE
SAC 000527; version 3.01 CO MAYO

0 20 40 60 80 Meters



The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
 Ordnance Survey of Ireland Licence No EN 0059216. © Ordnance Survey of Ireland and Government of Ireland

Ní snáiteoraí nneacha ar nal éarscáil eanna ach nod garshú omhach g near áta. Féad ar at hbréithithe a déana mh ar theorá nneacha na gceantar comhartáithe. Súrbhéaracht a Odonás na hÉireann Ceadúnas U mh EN 0059216. © Súrbhéaracht a Odonás na hÉireann Rátas na hÉireann




Map Version 1
Date: June 2018

SAC000527








SAC000527

Bat Site Code: 684



Legend

-  Moore Hall (Lough Carra) SAC000527
-  OS Discovery Series County Boundary
- 1303 Lesser Horseshoe Bat *Rhinolophus hipposideros***
-  Roost Location
-  Bat Site Code 684 Foraging Range
-  Potential Foraging Grounds



An Roinn Cultúir,
Oidhreacht agus Gaeltachta
Department of Culture,
Heritage and the Gaeltacht

**MAP 2:
MOORE HALL (LOUGH CARRA) SAC
CONSERVATION OBJECTIVES ADJACENT
LESSER HORSESHOE BAT**

Map to be read in conjunction with the NPWS Conservation Objectives Document.

**SITE CODE
SAC 000527; version 3.01.
CQ MAYO**

0 0.45 0.9 1.35 1.8 2.25 Kilometers



The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
Ordnance Survey of Ireland Licence No EN0059216. © Ordnance Survey of Ireland and Government of Ireland
Ní sna teoráinneacha ar na léarscáil eanna ach nod garshú omhach ginearálta. Féad ar a bhfeithithe a déana mh ar theoráinneacha na gceart ar comhartáithe. Súr bhéaracht a Ordónás na hÉireann Ceadúnas U mh EN0059216. © Súr bhéaracht a Ordónás na hÉireann Rádas na hÉireann



**Map Version 1
Date: June 2018**



Conservation objectives for Oldhead Wood SAC [000532]

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Objective: To maintain or restore the favourable conservation condition of the Annex I habitat(s) and/or the Annex II species for which the SAC has been selected:

Code	Description
------	-------------

4030	European dry heaths
------	---------------------

91A0	Old sessile oak woods with <i>Ilex</i> and <i>Blechnum</i> in the British Isles
------	---

* denotes a priority habitat



Citation: NPWS (2020) Conservation objectives for Oldhead Wood SAC [000532]. Generic Version 7.0.
Department of Culture, Heritage and the Gaeltacht.

National Parks and Wildlife Service

Conservation Objectives Series

Owenduff/Nephin Complex SAC 000534



An Roinn Ealaíon, Oidhreachta,
Gnóthaí Réigiúnacha, Tuaithe agus Gaeltachta

Department of Arts, Heritage,
Regional, Rural and Gaeltacht Affairs



**National Parks and Wildlife Service,
Department of Arts, Heritage, Regional, Rural and Gaeltacht Affairs,
7 Ely Place, Dublin 2, Ireland.
Web: www.npws.ie
E-mail: nature.conservation@ahg.gov.ie**

Citation:

**NPWS (2017) Conservation Objectives: Owenduff/Nephin Complex SAC 000534.
Version 1. National Parks and Wildlife Service, Department of Arts, Heritage,
Regional, Rural and Gaeltacht Affairs.**

**Series Editor: Rebecca Jeffrey
ISSN 2009-4086**

Introduction

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

* indicates a priority habitat under the Habitats Directive

000534	Owenduff/Nephin Complex SAC
1106	Salmon <i>Salmo salar</i>
1355	Otter <i>Lutra lutra</i>
1393	Slender Green Feather-moss <i>Drepanocladus vernicosus</i>
1528	Marsh Saxifrage <i>Saxifraga hirculus</i>
3110	Oligotrophic waters containing very few minerals of sandy plains (Littorelletalia uniflorae)
3130	Oligotrophic to mesotrophic standing waters with vegetation of the Littorelletea uniflorae and/or Isoeto-Nanojuncetea
3160	Natural dystrophic lakes and ponds
3260	Water courses of plain to montane levels with the Ranunculion fluitantis and Callitriche-Batrachion vegetation
4010	Northern Atlantic wet heaths with Oxalidaceae Carex
4060	Alpine and Boreal heaths
5130	R formations on heaths or calcareous grasslands
7130	Blanket bogs (* if active bog)
7140	Transition mires and quaking bogs

Please note that this SAC overlaps with Owenduff/Nephin Complex SPA (004098) and is adjacent to Carrowmore Lake Complex SAC (000476), Corraun Plateau SAC (000485), Clew Bay Complex SAC (001482) and Blacksod Bay/Broad Haven SPA (004037). See map 2. The conservation objectives for this site should be used in conjunction with those for the overlapping and adjacent sites as appropriate.

Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

NPWS Documents

Year :	1987
Title :	The vegetation of Irish rivers
Author :	Heuff, H.
Series :	Unpublished report to NPWS
Year :	1987
Title :	A survey to locate blanket bogs of scientific interest in County Mayo. Part I
Author :	Foss, P.; McGee, E.
Series :	A report commissioned by the Wildlife Service
Year :	1989
Title :	Survey to locate blanket bogs of scientific interest in Mayo. Part II
Author :	Douglas, C.; Garvey, L.; Kelly, L.; O'Sullivan, A.; Van Doorsleer, L.
Series :	A report commissioned by the Wildlife Service
Year :	1999
Title :	A survey of the rare and protected flora of County Mayo
Author :	McKee, A-M.
Series :	Unpublished report to Duchas
Year :	2006
Title :	Otter survey of Ireland 2004/2005
Author :	Bailey, M.; Rochford, J.
Series :	Irish Wildlife Manual No. 23
Year :	2006
Title :	Conservation Plan for 2006-2011. Owenduff/Nepin Complex cSAC and SPA Site Codes 000534 and 004098 Co. Mayo
Author :	NPWS
Series :	Conservation Plan
Year :	2007
Title :	Supporting documentation for the Habitats Directive Conservation Status Assessment - backing documents. Article 17 forms and supporting maps
Author :	NPWS
Series :	Unpublished report to NPWS
Year :	2012
Title :	The conservation status of juniper formations in Ireland
Author :	Cooper, F.; Stone, R.E.; McEvoy, P.; Wilkins, T.; Reid, N.
Series :	Irish Wildlife Manual No. 63
Year :	2012
Title :	Ireland Red List No. 8: Bryophytes
Author :	Lockhart, N.; Hodgetts, N.; Holyoak, D.
Series :	Ireland Red List series, NPWS
Year :	2013
Title :	National otter survey of Ireland 2010/12
Author :	Reid, N.; Hayden, B.; Lundy, M.G.; Pietravalle, S.; McDonald, R.A.; Montgomery, W.I.
Series :	Irish Wildlife Manual No. 76

Year :	2013
Title :	A survey of red grouse (<i>Lagopus lagopus scoticus</i>) in the Owenduff/Nephin Complex Special Protection Area, County Mayo
Author :	Murray, T.; Clotworthy, C.; Bleasdale, A.
Series :	Irish Wildlife Manual No. 77
Year :	2013
Title :	The status of EU protected habitats and species in Ireland. Volume 2. Habitats assessments
Author :	NPWS
Series :	Conservation assessments
Year :	2014
Title :	Guidelines for a national survey and conservation assessment of upland vegetation and habitats in Ireland, Version 2.0
Author :	Perrin, P.M.; Barron, S.J.; Roche, J.R.; O'Hanrahan, B.
Series :	Irish Wildlife Manual No. 79
Year :	2015
Title :	Monitoring recommendations for Marsh Saxifrage (<i>Saxifraga hirculus</i> L.) in the Republic of Ireland
Author :	Muldoon, C.S.; Waldren, S.; Lynn, D.
Series :	Irish Wildlife Manual No. 88
Year :	2015
Title :	Habitats Directive Annex I lake habitats: a working interpretation for the purposes of site-specific conservation objectives and Article 17 reporting
Author :	O Connor, Á.
Series :	Unpublished document by NPWS
Year :	2015
Title :	Monitoring methods for <i>Hamatocaulis vernicosus</i> (Mitt.) Hedenäs (Slender green feather-moss) in the Republic of Ireland
Author :	Campbell, C.; Hodgetts, N.; Lockhart, N.
Series :	Irish Wildlife Manual No. 91
Year :	2016
Title :	Ireland Red List No. 10: Vascular Plants
Author :	Wyse Jackson, M.; FitzPatrick, Ú.; Cole, E.; Jebb, M.; McFerran, D.; Sheehy Skeffington, M.; Wright, M.
Series :	Ireland Red Lists series, NPWS
Year :	2017
Title :	Owenduff/Nephin Complex SAC (site code: 534) Conservation objectives supporting document- blanket bogs and associated habitats V1
Author :	NPWS
Series :	Conservation objectives supporting document

Other References

Year :	1982
Title :	Otter survey of Ireland
Author :	Chapman, P.J.; Chapman, L.L.
Series :	Unpublished report to Vincent Wildlife Trust
Year :	1982
Title :	Eutrophication of waters. Monitoring assessment and control
Author :	OECD
Series :	OECD, Paris

Year :	1989
Title :	The genera <i>Scorpidium</i> and <i>Hamatocaulis</i> , gen. nov., in northern Europe
Author :	Hedenäs, L.
Series :	Lindbergia, 15: 8-36
Year :	1989
Title :	Three new localities for <i>Saxifraga hirculus</i> L. in Ireland
Author :	Lockhart, N.
Series :	Irish Naturalists' Journal, 23(2): 65-69
Year :	1991
Title :	The spatial organization of otters (<i>Lutra lutra</i>) in Shetland
Author :	Kruuk, H.; Moorhouse, A.
Series :	Journal of Zoology, 224: 41-57
Year :	1998
Title :	Studies in Irish Limnology
Author :	Giller, P.S. (ed.)
Series :	Marine Institute, Dublin
Year :	1998
Title :	Studies of Irish Rivers and Lakes
Author :	Moriarty, C. (ed.)
Series :	Marine Institute, Dublin
Year :	2000
Title :	Colour in Irish lakes
Author :	Free, G.; Allott, N.; Mills, P.; Kennelly, C.; Day, S.
Series :	Verhandlungen Internationale Vereinigung für theoretische und angewandte Limnologie, 27: 2620-2623
Year :	2002
Title :	Deterioration of Atlantic soft water macrophyte communities by acidification, eutrophication and alkalinisation
Author :	Arts, G.H.P.
Series :	Aquatic Botany, 73: 373-393
Year :	2003
Title :	Ecology of watercourses characterised by Ranunculion fluitantis and Callitriche-Batrachion vegetation
Author :	Hatton-Ellis, T.W.; Grieve, N.
Series :	Conserving Natura 2000 Rivers Ecology Series No. 11. English Nature, Peterborough
Year :	2005
Title :	Lichens. An illustrated guide to the British and Irish species
Author :	Dobson, F.S.
Series :	The Richmond Publishing Co. Ltd., Slough
Year :	2006
Title :	Otters - ecology, behaviour and conservation
Author :	Kruuk, H.
Series :	Oxford University Press

Year : 2006

Title : A reference-based typology and ecological assessment system for Irish lakes. Preliminary investigations. Final report. Project 2000-FS-1-M1 Ecological assessment of lakes pilot study to establish monitoring methodologies EU (WFD)

Author : Free, G.; Little, R.; Tierney, D.; Donnelly, K.; Coroni, R.

Series : EPA, Wexford

Year : 2008

Title : Water Quality in Ireland 2004-2006

Author : Clabby, K.J.; Bradley, C.; Craig, M.; Daly, D.; Lucey, J.; McGarrigle, M.; O'Boyle, S.; Tierney, D.; Bowman, J.

Series : EPA, Wexford

Year : 2009

Title : The identification, characterization and conservation value of isoetid lakes in Ireland

Author : Free, G.; Bowman, J.; McGarrigle, M.; Little, R.; Coroni, R.; Donnelly, K.; Tierney, D.; Trodd, W.

Series : Aquatic Conservation: Marine and Freshwater Ecosystems, 19(3): 264–273

Year : 2010

Title : Otter tracking study of Roaringwater Bay

Author : De Jongh, A.; O'Neill, L.

Series : Unpublished draft report to NPWS

Year : 2010

Title : Water quality in Ireland 2007-2009

Author : McGarrigle, M.; Lucey, J.; Ó Cinnéide, M.

Series : EPA, Wexford

Year : 2011

Title : Conservation biology of *Saxifraga hirculus* L. in Ireland

Author : Muldoon, C.S.

Series : Unpublished Ph.D. Thesis, Trinity College Dublin

Year : 2012

Title : Rare and threatened bryophytes of Ireland

Author : Lockhart, N.; Hodgetts, N.; Holyoak, D.

Series : National Museums Northern Ireland

Year : 2012

Title : The impact of conifer plantation forestry on the ecology of peatland lakes

Author : Drinan, T.J.

Series : Unpublished Ph.D. thesis, University College Cork

Year : 2012

Title : Lichens of Ireland. An illustrated introduction to over 250 species

Author : Whelan, P.

Series : The Collins Press, Wilton, Cork

Year : 2013

Title : Conservation of selected legally protected and Red Listed bryophytes in Ireland

Author : Campbell, C.

Series : Unpublished Ph.D. Thesis, Trinity College Dublin

Year : 2013

Title : Management strategies for the protection of high status water bodies

Author : Ní Chatháin, B.; Moorkens, E.; Irvine, K.

Series : Strive Report Series No. 99. EPA, Wexford

Year :	2013
Title :	Interpretation manual of European Union habitats- Eur 28
Author :	European Commission- DG Environment
Series :	European Commission
Year :	2014
Title :	The impact of rural land management changes on soil hydraulic properties and runoff processes: results from experimental plots in upland UK
Author :	Marshall, M.R.; Ballard, C.E.; Frogbrook, Z.L.; Solloway, I.; McIntyre, N.; Reynolds, B.; Wheater, H.S.
Series :	Hydrological Processes, 28: 2617–2629
Year :	2014
Title :	Identifying the role of environmental drivers in organic carbon export from a forested peat catchment
Author :	Ryder, E.; de Eyto, E.; Dillane, M.; Poole, R.; Jennings, E.
Series :	Science of the Total Environment, 490: 28–36.
Year :	2015
Title :	Water quality in Ireland 2010-2012
Author :	Bradley, C.; Byrne, C.; Craig, M.; Free, G.; Gallagher, T.; Kennedy, B.; Little, R.; Lucey, J.; Mannix, A.; McCreesh, P.; McDermott, G.; McGarrigle, M.; Ní Longphuirt, S.; O'Boyle, S.; Plant, C.; Tierney, D.; Trodd, W.; Webster, P.; Wilkes, R.; Wynne, C.
Series :	EPA, Wexford
Year :	2015
Title :	The fecundity of wild Irish Atlantic salmon <i>Salmo salar</i> L. and its application for stock assessment purposes
Author :	de Eyto, E.; White, J.; Boylan, P.; Clarke, B.; Cotter, D.; Doherty, D.; Gargan, P.; Kennedy, R.; McGinnity, P.; O'Maoiléidigh, N.; O'Higgins, K.
Series :	Fisheries Research, 164: 159–169.
Year :	2016
Title :	A narrative for conserving freshwater and wetland habitats in England
Author :	Mainstone, C.; Hall, R.; Diack, I.
Series :	Natural England Research Reports Number 064
Year :	2016
Title :	The Status of Irish Salmon Stocks in 2015 with Precautionary Catch Advice for 2016
Author :	SSCS (Standing Scientific Committee on Salmon)
Series :	Independent Scientific Report to Inland Fisheries Ireland

Spatial data sources

Year :	2008
Title :	OSi 1:5000 IG vector dataset
GIS Operations :	WaterPolygons feature class clipped to the SAC boundary. Expert opinion used to identify Annex I habitats and to resolve any issues arising
Used For :	3110, 3160 (map 3)
<hr/>	
Year :	2017
Title :	NPWS rare and threatened species database
GIS Operations :	Dataset created from spatial references in database records. Expert opinion used as necessary to resolve any issues arising
Used For :	1393, 1528 (maps 4 and 5)
<hr/>	
Year :	2005
Title :	OSi Discovery series vector data
GIS Operations :	Creation of 10m buffer on the terrestrial side of river banks data; creation of 20m buffer applied to canal centreline data. Creation of 20m buffer applied to river and stream centreline data; These datasets combined with the derived OSi 1:5000 vector lake buffer data. Overlapping regions investigated and resolved; resulting dataset clipped to SAC boundary. Expert opinion used as necessary to resolve any issues arising
Used For :	1355 (no map)
<hr/>	
Year :	2010
Title :	OSi 1:5000 IG vector dataset
GIS Operations :	Creation of 80m buffer on the aquatic side of lake data; creation of 10m buffer on the terrestrial side of lake data. These datasets combined with the derived OSi Discovery Series river and canal datasets. Overlapping regions investigated and resolved; resulting dataset clipped to SAC boundary. Expert opinion used as necessary to resolve any issues arising
Used For :	1355 (no map)
<hr/>	

Conservation Objectives for : Owenduff/Nephin Complex SAC [000534]

3110 Oligotrophic waters containing very few minerals of sandy plains (Littorelletalia uniflorae)

To maintain the favourable conservation condition of Oligotrophic waters containing very few minerals of sandy plains (Littorelletalia uniflorae) in Owenduff/Nephin Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Lake habitat 3110 is considered likely to occur in Lough Feeagh and other larger lakes (e.g. Bunaveela, Anaffrin) in Owenduff/Nephin Complex SAC (see map 3). The SAC was formerly selected for lake habitat 3130, based on an older interpretation of that habitat where it was associated with uplands (see O Connor, 2015). In line with Article 17 reporting (NPWS, 2013), all lakes larger than 1ha were mapped as potential 3110. Lake habitat 3160 is likely to co-occur with this habitat in many lakes in the SAC, particularly at higher altitude (above 200m), owing to the base-poor geology (quartzite and schist) and blanket peats. Two measures of extent should be used: 1. the area of the lake itself and; 2. the extent of the vegetation communities/zones that typify the habitat. Further information relating to all attributes is provided in the lake habitats supporting document for the purposes of site-specific conservation objectives and Article 17 reporting (O Connor, 2015)
Habitat distribution	Occurrence	No decline, subject to natural processes	As noted above, all lakes larger than 1ha have been mapped as potential 3110 (see map 3)
Typical species	Occurrence	Typical species present, in good condition, and demonstrating typical abundances and distribution	For lists of typical plant species, see the Article 17 habitat assessment for 3110 (NPWS, 2013) and the lake habitats supporting document (O Connor, 2015). Douglas et al. (1989) provide some records for lake macrophytes from the SAC. Lough Feeagh is a Water Framework Directive (WFD) monitoring lake and regular macrophyte surveys are conducted by the Environmental Protection Agency (EPA). EPA data indicate Lough Feeagh has limited submerged vegetation: quillwort (<i>Isoetes lacustris</i>), shoreweed (<i>Littorella uniflora</i>) and bulbous rush (<i>Juncus bulbosus</i>). There has been extensive study of the Burrishoole catchment, including Lough Feeagh, by the Marine Institute (formerly the Salmon Research Agency). This had concentrated particularly on fish and water quality, but also acidification, climate change, organic carbon, etc. (e.g. Cross et al., 1998 in Giller, 1998; Whelan et al., 1998 in Moriarty, 1998; Ryder et al., 2014; de Eyto et al., 2015)
Vegetation composition: characteristic zonation	Occurrence	All characteristic zones should be present, correctly distributed and in good condition	Further work is necessary to describe the characteristic zonation and other spatial patterns in lake habitat 3110 (see O Connor, 2015)
Vegetation distribution: maximum depth	Metres	Maintain maximum depth of vegetation, subject to natural processes	The maximum depth of vegetation is likely to be specific to the lake shoreline in question. Further work is necessary to develop indicative targets for lake habitat 3110. Maximum depth should be large in lakes in the SAC within undisturbed peatland and uplands; however, pressures such as overgrazing, forestry and peat-cutting may have reduced vegetation depth in some lakes. Data on macrophyte depth in Lough Feeagh will be available from EPA monitoring

Hydrological regime: water level fluctuations	Metres	Maintain/restore appropriate natural hydrological regime necessary to support the habitat	Fluctuations in lake water level are typical in Ireland, but can be amplified by activities such as abstraction, drainage and overgrazing. Increased water level fluctuations can increase wave action, up-root vegetation, increase turbidity, alter the substratum and lead to release of nutrients from the sediment. The hydrological regime of the lakes must be maintained so that the area, distribution and depth of the lake habitat and its constituent/characteristic vegetation zones and communities are not reduced. It is likely that the hydrological regimes of many of the lakes are still altered owing to historic overgrazing (faster run-off, higher flood peaks, lower base flows, etc.; see Marshall et al., 2014)
Lake substratum quality	Various	Maintain appropriate substratum type, extent and chemistry to support the vegetation	Research is required to further characterise the substratum types (particle size and origin) and substratum quality (notably pH, calcium, iron and nutrient concentrations) favoured by each of the five Annex I lake habitats in Ireland. It is likely that lake habitat 3110 is associated with a range of nutrient-poor substrates, from stones, cobble and gravel, through sands, silt, clay and peat. Substratum particle size is likely to vary with depth and along the shoreline within a single lake
Water quality: transparency	Metres	Maintain appropriate Secchi transparency. There should be no decline in Secchi depth/transparency	Transparency relates to light penetration and, hence, to the depth of colonisation of vegetation. It can be affected by phytoplankton blooms, water colour and turbidity. Specific targets have yet to be established for lake habitat 3110 (O Connor, 2015). Habitat 3110 is associated with very clear water. The OECD fixed boundary system set transparency targets for oligotrophic lakes of $\geq 6\text{m}$ annual mean Secchi disk depth, and $\geq 3\text{m}$ annual minimum Secchi disk depth. Free et al. (2009) found high isoetid abundance in lakes with Secchi depths of more than 3m
Water quality: nutrients	$\mu\text{g/l P}$; mg/l N	Maintain the concentration of nutrients in the water column at sufficiently low levels to support the habitat and its typical species	As a nutrient poor-habitat, oligotrophic and WFD 'high' status targets apply. Where a lake has nutrient concentrations that are lower than these targets, there should be no decline within class, i.e. no upward trend in nutrient concentrations. For lake habitat 3110, annual average total phosphorus (TP) concentration should be $\leq 10\mu\text{g/l TP}$, average annual total ammonia concentration should be $\leq 0.040\text{mg/l N}$ and annual 95th percentile for total ammonia should be $\leq 0.090\text{mg/l N}$. See also The European Communities Environmental Objectives (Surface Waters) Regulations 2009. Lough Feeagh passed the nutrient conditions target in 2007-09 and 2010-12 (McGarrigle et al., 2010; Bradley et al., 2015)
Water quality: phytoplankton biomass	$\mu\text{g/l Chlorophyll } a$	Maintain appropriate water quality to support the habitat, including high chlorophyll <i>a</i> status	Oligotrophic and WFD 'high' status targets apply to lake habitat 3110. Where a lake has a chlorophyll <i>a</i> concentration that is lower than this target, there should be no decline within class, i.e. no upward trend in phytoplankton biomass. The average growing season (March-October) chlorophyll <i>a</i> concentration must be $< 5.8\mu\text{g/l}$. The annual average chlorophyll <i>a</i> concentration should be $< 2.5\mu\text{g/l}$ and the annual peak chlorophyll <i>a</i> concentration should be $\leq 8.0\mu\text{g/l}$. See also The European Communities Environmental Objectives (Surface Waters) Regulations 2009. Lough Feeagh passed the target and had high chlorophyll <i>a</i> status in 2007-09 and 2010-12 (McGarrigle et al., 2010; Bradley et al., 2015)
Water quality: phytoplankton composition	EPA phytoplankton composition metric	Maintain appropriate water quality to support the habitat, including high phytoplankton composition status	The EPA has developed a phytoplankton composition metric for nutrient enrichment of Irish lakes. As for other water quality indicators, lake habitat 3110 requires WFD high status

Water quality: attached algal biomass	Algal cover and EPA phytobenthos metric	Maintain trace/absent attached algal biomass (<5% cover) and high phytobenthos status	Nutrient enrichment can favour epiphytic and epipelic algae that can out-compete the submerged vegetation. The cover abundance of attached algae in lake habitat 3110 should, therefore, be trace/absent (<5% cover). EPA phytobenthos can be used as an indicator of changes in attached algal biomass. As for other water quality indicators, habitat 3110 requires high phytobenthos status
Water quality: macrophyte status	EPA macrophyte metric (The Free Index)	Restore high macrophyte status	Nutrient enrichment can favour more competitive submerged macrophyte species that out-compete the typical and characteristic species for the lake habitat. The EPA monitors macrophyte status for WFD purposes using the 'Free Index'. The target for lake habitat 3110 is high status or an Ecological Quality Ratio (EQR) for lake macrophytes of ≥ 0.90 , as defined in Schedule Five of the European Communities Environmental Objectives (Surface Waters) Regulations 2009. Lough Feeagh failed the target in 2010-12, having good macrophyte status (Bradley et al., 2015)
Acidification status	pH units; mg/l	Maintain appropriate water and sediment pH, alkalinity and cation concentrations to support the habitat, subject to natural processes	Acidification can impact on species abundance and composition in soft water lake habitats. In Europe, acidification of isoetid lakes can lead to loss of isoetids and dominance by submerged <i>Sphagnum</i> mosses and <i>Juncus bulbosus</i> (Arts, 2002). The specific requirements of lake habitat 3110, in terms of water and sediment pH, alkalinity and cation concentration, have not been determined. For lake habitat 3110, and adopting a precautionary approach based on Arts (2002), minimum pH should not be <5.5 pH units. Maximum pH should be <9.0 pH units, in line with the surface water standards established for soft waters (where water hardness is ≤ 100 mg/l calcium carbonate). See Schedule Five of the European Communities Environmental Objectives (Surface Waters) Regulations 2009. See McGarrigle et al. (2010) and Bradley et al. (2015) for WFD acidification status in the 2007-09 and 2010-12 periods
Water colour	mg/l PtCo	Maintain/restore appropriate water colour to support the habitat	Increased water colour and turbidity decrease light penetration and can reduce the area of available habitat for lake macrophytes, particularly at the lower euphotic depths. The primary source of increased water colour in Ireland is disturbance to peatland. No habitat-specific or national standards for water colour currently exist. Studies have shown median colour concentrations in Irish lakes of 38mg/l PtCo (Free et al., 2000) and 33mg/l PtCo (Free et al., 2006). It is likely that the water colour in all Irish lake habitats would naturally be <50mg/l PtCo. Water colour can be very low (<20mg/l PtCo or even <10mg/l PtCo) in lakes with habitat 3110, where the peatland in the lake's catchment is intact. Free et al. (2006) reported colour of 80mg/l PtCo in Lough Feeagh. Overgrazing and other peatland degradation is likely to have increased colour in some lakes in Owenduff/Nephin Complex SAC

Dissolved organic carbon (DOC)	mg/l	Maintain/restore appropriate organic carbon levels to support the habitat	Dissolved (and particulate) organic carbon (OC) in the water column is linked to water colour and acidification (organic acids). Increasing DOC in water has been documented across the Northern Hemisphere, including afforested peatland catchments in Ireland. Damage and degradation of peatland, leading to decomposition of peat is likely to be the predominant source of OC in Ireland. OC in water promotes decomposition by fungi and bacteria that, in turn, releases dissolved nutrients. The increased biomass of decomposers can also impact directly on the characteristic lake communities through shading, competition, etc. Ryder et al. (2014) investigated OC losses from forestry in the Burrishoole. Overgrazing and other peatland degradation is also likely to have increased DOC in some lakes in Owenduff/Nephin Complex SAC
Turbidity	Nephelometric turbidity units/ mg/l SS/ other appropriate units	Maintain/restore appropriate turbidity to support the habitat	Turbidity can significantly affect the quantity and quality of light reaching rooted and attached vegetation and can, therefore, impact on lake habitats. The settlement of higher loads of inorganic or organic material on lake vegetation communities may also have impacts on sensitive, delicate species. Turbidity can increase as a result of re-suspension of material within the lake, higher loads entering the lake, or eutrophication. Turbidity measurement and interpretation is challenging. As a result, it is likely to be difficult to set habitat-specific targets for turbidity in lakes. Increased loads of fine organic and inorganic particles from overgrazing may have increased turbidity in lakes in Owenduff/Nephin Complex SAC
Fringing habitat: area and condition	Hectares	Maintain the area and condition of fringing habitats necessary to support the natural structure and functioning of habitat 3110	Most lake shorelines have fringing habitats of reedswamp, other swamp, fen, marsh or wet woodland that intergrade with and support the structure and functions of the lake habitat. In this SAC, lake shorelines are likely to have acid grassland, swamp, heath, blanket bog and rock communities. Fringing habitats are dependent on the lake, particularly its water levels, and support wetland communities and species of conservation concern. Many of the fringing wetland habitats support higher invertebrate and plant species richness than the lake habitats themselves

Conservation Objectives for : Owenduff/Nephin Complex SAC [000534]

3160 Natural dystrophic lakes and ponds

To maintain the favourable conservation condition of Natural dystrophic lakes and ponds in Owenduff/Nephin Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Owenduff/Nephin Complex SAC has both lowland blanket bog pool systems and upland lakes with habitat 3160. The habitat is likely to be found in many lakes in the SAC, where it may co-occur with lake habitat 3110, and all lakes, with the exception of Lough Feeagh, have been mapped as potential 3160 (see map 3). Many of the bog pools are not mapped in the 1:5,000 OSi data (map 3). Lake habitat 3160 is of high conservation value in the SAC. For further information on the distribution, vegetation and morphology of the habitat in the SAC, see Foss and McGee (1987) and Douglas et al. (1989). Two measures of extent should be used: 1. the area of the lake itself and; 2. the extent of the vegetation communities/zones that typify the habitat. Further information relating to all attributes is provided in the lake habitats supporting document for the purposes of site-specific conservation objectives and Article 17 reporting (O Connor, 2015)
Habitat distribution	Occurrence	No decline, subject to natural processes	As noted above, all lakes and ponds in the SAC, with the exception of Lough Feeagh, have been mapped as potential 3160 (see map 3). Atlantic blanket bog pools, including interconnecting pool systems, were recorded at Uggool, Sheeanmore and Altnabrocky, Owenglass West and East, Bellagaravaun, and other areas of the SAC (Foss and McGee, 1987; Douglas et al., 1989)
Typical species	Occurrence	Typical species present, in good condition, and demonstrating typical abundances and distribution	For lists of typical plant and invertebrate species, see the Article 17 habitat assessment for 3160 (NPWS, 2013) and O Connor (2015). Douglas et al. (1989) recorded many-stalked spike-rush (<i>Eleocharis multicaulis</i>), bogbean (<i>Menyanthes trifoliata</i>) and the bog mosses <i>Sphagnum auriculatum</i> and <i>S. cuspidatum</i> in pools, and some water lobelia (<i>Lobelia dortmanna</i>) and pipewort (<i>Eriocaulon aquaticum</i>), the latter of which is Near Threatened in Ireland (Wyse Jackson et al., 2016). Great sundew (<i>Drosera anglica</i>) and round-leaved sundew (<i>D. rotundifolia</i>) were abundant in shallower interconnecting pools. Bulbous rush (<i>Juncus bulbosus</i>) was recorded in lakes with rock basins
Vegetation composition: characteristic zonation	Occurrence	All characteristic zones should be present, correctly distributed and in good condition	Further work is necessary to describe the characteristic zonation and other spatial patterns in lake habitat 3160 (see O Connor, 2015). Spatial patterns are likely to be relatively simple in 3160 lakes and ponds, with limited zonation
Vegetation distribution: maximum depth	Metres	Maintain maximum depth of vegetation, subject to natural processes	The maximum depth of vegetation is likely to be specific to the lake shoreline in question. Further work is necessary to develop indicative targets for lake habitat 3160. 3160 lakes and pools naturally have very clear water and, therefore, maximum depth can be large within undisturbed peatland and uplands. Pressures such as overgrazing, forestry and peat-cutting may have reduced vegetation depth in some lakes in Owenduff/Nephin Complex SAC

Hydrological regime: water level fluctuations	Metres	Maintain/restore appropriate natural hydrological regime necessary to support the habitat	Natural water level fluctuations can be amplified by activities such as abstraction, drainage and overgrazing. Increased fluctuations can increase wave action, up-root vegetation, increase turbidity, alter the substratum and lead to release of nutrients from the sediment. The hydrological regime of the lakes and pools must be maintained so that the area, distribution and depth of the lake habitat and its constituent/characteristic vegetation zones and communities are not reduced. The hydrological regime of 3160 lakes and pools is integrally linked to that of the surrounding blanket bog, transition mire/quaking bog and other peatland habitats. Owing to their size and the sensitivity of peatland, 3160 lakes and pools can easily be damaged or destroyed by drainage. It is likely that the hydrological regimes of lakes and pools may still be altered owing to historic overgrazing (faster run-off, higher flood peaks, lower base flows, etc.; see Marshall et al., 2014)
Lake substratum quality	Various	Maintain appropriate substratum type, extent and chemistry to support the vegetation	Research is required to further characterise the substratum types (particle size and origin) and substratum quality (notably pH, calcium, iron and nutrient concentrations) favoured by each of the five Annex I lake habitats in Ireland. It is likely that lake habitat 3160 is associated with nutrient-poor peat and silt substrates
Water quality: transparency	Metres	Maintain appropriate Secchi transparency. There should be no decline in Secchi depth/transparency	Transparency relates to light penetration and, hence, to the depth of colonisation of vegetation. It can be affected by phytoplankton blooms, water colour and turbidity. Specific targets have yet to be established for lake habitat 3160. Lake habitat 3160 is associated with very clear water. The OECD fixed boundary system set transparency targets for ultra-oligotrophic lakes of $\geq 12\text{m}$ annual mean Secchi disk depth, and $\geq 6\text{m}$ annual minimum Secchi disk depth
Water quality: nutrients	$\mu\text{g/l P}$; mg/l N	Maintain the concentration of nutrients in the water column at sufficiently low levels to support the habitat and its typical species	As a nutrient-poor habitat, oligotrophic and Water Framework Directive (WFD) 'high' status targets apply. Where a lake has nutrient concentrations that are lower than these targets, there should be no decline within class, i.e. no upward trend in nutrient concentrations. For 3160 lakes and pools, annual average total phosphorus (TP) concentration should be $\leq 5\mu\text{g/l TP}$, average annual total ammonia concentration should be $\leq 0.040\text{mg/l N}$ and annual 95th percentile for total ammonia should be $\leq 0.090\text{mg/l N}$. See also The European Communities Environmental Objectives (Surface Waters) Regulations 2009
Water quality: phytoplankton biomass	$\mu\text{g/l Chlorophyll } a$	Maintain appropriate water quality to support the habitat, including high chlorophyll <i>a</i> status	Oligotrophic and WFD 'high' status targets apply to lake habitat 3160. The average growing season (March-October) chlorophyll <i>a</i> concentration must be $< 5.8\mu\text{g/l}$ (The European Communities Environmental Objectives (Surface Waters) Regulations 2009). Where a lake has a chlorophyll <i>a</i> concentration that is lower than this target, there should be no decline within class, i.e. no upward trend in phytoplankton biomass. The OECD targets may be more appropriate for habitat 3160: annual average chlorophyll <i>a</i> concentration $< 1\mu\text{g/l}$ and annual peak chlorophyll <i>a</i> concentration $\leq 2.5\mu\text{g/l}$. See also The European Communities Environmental Objectives (Surface Waters) Regulations 2009
Water quality: phytoplankton composition	EPA phytoplankton composition metric	Maintain appropriate water quality to support the habitat, including high phytoplankton composition status	The Environmental Protection Agency (EPA) has developed a phytoplankton composition metric for nutrient enrichment of Irish lakes. As for other water quality indicators, lake habitat 3160 requires WFD high status

Water quality: attached algal biomass	Algal cover and EPA phytobenthos metric	Maintain trace/absent attached algal biomass (<5% cover) and high phytobenthos status	Nutrient enrichment can favour epiphytic and epipelic algae that can out-compete the submerged vegetation. The cover abundance of attached algae in 3160 lakes and ponds should, therefore, be trace/absent (<5% cover). EPA phytobenthos can be used as an indicator of changes in attached algal biomass. As for other water quality indicators, lake habitat 3160 requires high phytobenthos status
Water quality: macrophyte status	EPA macrophyte metric (The Free Index)	Maintain high macrophyte status	Nutrient enrichment can favour more competitive submerged macrophyte species that out-compete the typical and characteristic species for the lake habitat. The EPA monitors macrophyte status for WFD purposes using the 'Free Index'. The target for 3160 lakes and pools is high status or an Ecological Quality Ratio (EQR) for lake macrophytes of ≥ 0.90 , as defined in Schedule Five of the European Communities Environmental Objectives (Surface Waters) Regulations 2009
Acidification status	pH units; mg/l	Maintain appropriate water and sediment pH, alkalinity and cation concentrations to support the habitat, subject to natural processes	Acidification can impact on species abundance and composition in soft water lake habitats. Although European Commission (2013) describes lake habitat 3160 as having pH 3-6, Drinan (2012) found mean pH values of 5.16 and 5.62 in upland and lowland 3160 lakes, respectively. The target for lake habitat 3160 is pH >4.5 and <9.0, in line with the surface water standards for soft waters (where water hardness is $\leq 100\text{mg/l}$ calcium carbonate). See Schedule Five of the European Communities Environmental Objectives (Surface Waters) Regulations 2009. The specific requirements of habitat 3160, in terms of water and sediment pH, alkalinity and cation concentration, have not been determined
Water colour	mg/l PtCo	Maintain/restore appropriate water colour to support the habitat	Increased water colour and turbidity decrease light penetration and can reduce the area of available habitat for lake macrophytes, particularly at the lower euphotic depths. The primary source of increased water colour in Ireland is disturbance to peatland. No habitat-specific or national standards for water colour currently exist. Studies have shown median colour concentrations in Irish lakes of 38mg/l PtCo (Free et al., 2000) and 33mg/l PtCo (Free et al., 2006). It is likely that the water colour in all Irish lake habitats would naturally be <50mg/l PtCo. Water colour can be very low (<20mg/l PtCo or even <10mg/l PtCo) in 3160 lakes and pools where the peatland in the lake's catchment is intact. Overgrazing and other peatland degradation is likely to have increased colour in some lakes and pools in Owenduff/Nephin Complex SAC
Dissolved organic carbon (DOC)	mg/l	Maintain/restore appropriate organic carbon levels to support the habitat	Dissolved (and particulate) organic carbon (OC) in the water column is linked to water colour and acidification (organic acids). Increasing DOC in water has been documented across the Northern Hemisphere, including afforested peatland catchments in Ireland. Damage and degradation of peatland, leading to decomposition of peat is likely to be the predominant source of OC in Ireland. OC in water promotes decomposition by fungi and bacteria that, in turn, releases dissolved nutrients. The increased biomass of decomposers can also impact directly on the characteristic lake communities through shading, competition, etc. Overgrazing and other peatland degradation is likely to have increased DOC in some lakes and pools in the SAC

Turbidity	Nephelometric turbidity units/ mg/l SS/ other appropriate units	Maintain/restore appropriate turbidity to support the habitat	Turbidity can significantly affect the quantity and quality of light reaching rooted and attached vegetation and can, therefore, impact on lake habitats. The settlement of higher loads of inorganic or organic material on lake vegetation communities may also have impacts on sensitive, delicate species. Turbidity can increase as a result of re-suspension of material within the lake, higher loads entering the lake, or eutrophication. Turbidity measurement and interpretation is challenging. As a result, it is likely to be difficult to set habitat-specific targets for turbidity in lakes. Increased loads of fine organic and inorganic particles from overgrazing may have increased turbidity in lakes in Owenduff/Nephe Complex SAC
Fringing habitat: area and condition	Hectares	Maintain the area and condition of fringing habitats necessary to support the natural structure and functioning of habitat 3160	Lakes with 3160, particularly in uplands, are likely to be fringed by acid grassland, heath and rock communities. 3160 pools intergrade with blanket bog communities in Owenduff/Nephe Complex SAC. Spring-fed flushes are also a feature of the SAC. Quaking bog is also associated with pool systems in the SAC. Fringing habitats support the structure and functions of the lake/pool habitat. They are also dependent on the lake/pool, particularly its water levels, and can support wetland communities and species of conservation concern

Conservation Objectives for : Owenduff/Nephin Complex SAC [000534]

3260 Water courses of plain to montane levels with the *Ranunculus fluitantis* and *Callitriche-Batrachion* vegetation

To maintain the favourable conservation condition of Water courses of plain to montane levels with the *Ranunculus fluitantis* and *Callitriche-Batrachion* vegetation in Owenduff/Nephin Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Kilometres	Area stable or increasing, subject to natural processes	The description of habitat 3260 covers from upland rivers with bryophytes and macroalgae to lowland depositing rivers with pondweeds and starworts. Owenduff/Nephin Complex SAC was selected for highly oligotrophic, base-poor rivers, with limited aquatic vegetation. The main rivers in the SAC are the Owenduff and its tributaries to the south, and parts of the Owenmore and tributaries to the north-east. The Owenduff system was rated as of unique conservation importance and had communities dominated by mosses, liverworts and algae (Heuff, 1987). It is likely that most streams and rivers in the SAC have been negatively impacted by overgrazing in the Nephins and Nephin Begs (see NPWS, 2006; Murray et al., 2013)
Habitat distribution	Occurrence	No decline, subject to natural processes	Further study is needed of Irish sub-types and their conservation value to interpret the broad description of 3260 (European Commission, 2013). As noted above, the SAC was selected for a species-poor sub-type dominated by bryophytes and algae and with limited vascular plants. The uncommon river lichens <i>Ephebe lanata</i> and <i>Porpidia hydrophila</i> are known from the Altaconey and Srahmore rivers in the SAC (see Dobson (2005) and Whelan (2012) for notes on ecology). <i>Bryum riparium</i> , an endangered bryophyte of damp rock near streams and waterfalls occurs in the SAC (Lockhart et al., 2012). Ivy-leaved bellflower (<i>Wahlenbergia hederacea</i>) occurs at a number of sites along the Owenduff (including near Lagduff and Srahduff) and on the Tarsaghaunmore tributary (see McKee, 1999). This is an important outlying population of a Near Threatened species (Wyse Jackson et al., 2016) otherwise concentrated in the south-west and east
Hydrological regime: river flow	Metres per second	Maintain appropriate hydrological regimes	High conservation value sub-types are associated with natural hydrology. A natural flow regime is required for both plant communities and channel geomorphology to be in favourable condition, exhibiting typical dynamics for the river type (Hatton-Ellis and Grieve, 2003). For many sub-types, high flows are required to maintain the substratum necessary for the characteristic species. Flow variation can be particularly important, with high and flood flows being critical to the hydromorphology. Peatlands also have slow-flowing or ponded streams and rivers, with biotic communities likely to resemble those in associated lakes. Many of the rivers and streams in the SAC are naturally very flashy, although some more ponded and slow-flowing stretches occur in areas of relatively flat bog. It is likely that the hydrological regimes of many of the rivers are still altered owing to historic overgrazing (faster run-off, higher flood peaks, lower base flows, etc.; see Marshall et al., 2014)

Hydrological regime: groundwater discharge	Metres per second	Maintain appropriate hydrological regime	The groundwater contribution to rivers in the SAC is likely to be small, owing to the geology and dominance of blanket peat soils. Even small groundwater contributions, however, can significantly alter the hydrochemistry, particularly where there is basic bedrock and/or subsoils
Substratum composition: particle size range	Millimetres	Maintain appropriate substratum particle size range, quantity and quality, subject to natural processes	Many of the high conservation value sub-types are dominated by coarse substrata, and it is likely that bedrock, boulders, cobbles and coarse gravels were naturally abundant in many rivers in this SAC. The size and distribution of particles is largely determined by the river flow. The chemical composition (particularly minerals and nutrients) of the substratum is also important. The quality of finer sediment particles is a notable driver for rooted plant communities. The geomorphology, including channel shape and substratum, of many streams and rivers in the SAC are likely to be significantly altered as a result of overgrazing
Water quality	Various	Maintain appropriate water quality to support the natural structure and functioning of the habitat	The specific targets may vary among sub-types. The rivers within Owenduff/Nephin Complex SAC are naturally very nutrient-poor and, therefore, typically require Water Framework Directive high status, in terms of nutrient and oxygenation standards, and EQRs (Ecological Quality Ratios) for macroinvertebrates and phytobenthos. The occurrence of high status river sites downstream of areas of previously severe overgrazing damage is unexpected and suggests the metrics may not be sensitive to such impacts. See also The European Communities Environmental Objectives (Surface Waters) Regulations 2009, Environmental Protection Agency (EPA) river water quality reports (e.g. Bradley et al., 2015) and Ni Chatháin et al. (2013)
Typical species	Occurrence	Typical species of the relevant habitat sub-types should be present and in good condition	The sub-types of this habitat are poorly understood and their typical species have not yet been fully defined. The typical species may include higher plants, bryophytes, macroalgae and microalgae, and invertebrates. As noted above, rare lichens, bryophytes and the vascular plant species ivy-leaved bellflower (<i>Wahlenbergia hederacea</i>) are associated with rivers, streams and riparian areas in Owenduff/Nephin Complex SAC
Floodplain connectivity: area	Hectares	The area of active floodplain at, and upstream of, the habitat, necessary to support all sub-types of the habitat, should be maintained	River connectivity with the floodplain is important for the functioning of this habitat. Channels with a naturally functioning floodplain are better able to maintain habitat and water quality (Hatton-Ellis and Grieve, 2003). Floodplain connectivity is particularly important in terms of sediment sorting and nutrient deposition. High conservation value rivers are intimately connected to floodplain habitats and function as important wildlife corridors, connecting otherwise isolated or fragmented habitats in the wider countryside (Hatton-Ellis and Grieve, 2003; Mainstone et al., 2016). The hydro-morphological impacts associated with overgrazing may have impacted on floodplain connectivity in Owenduff/Nephin Complex SAC

Fringing habitats: Hectares
area and condition

Maintain the area and
condition of fringing
habitats necessary to
support the habitat and its
sub-types

Riparian habitats (including those along lake shores), particularly natural/semi-natural woodlands and wetlands, are an integral part of the structure and functioning of river systems, even where they do not form part of a natural floodplain. Fringing habitats can contribute to the aquatic food web (e.g. allochthonous matter such as leaf fall), provide habitat (refuge and resources) for certain life-stages of fish, birds and aquatic invertebrates, assist in the settlement of fine suspended material, protect banks from erosion and contribute to nutrient cycling. Shade may also be important in suppressing algal growth in enriched rivers and moderating temperatures. Equally, fringing habitats are dependent on rivers/lakes, particularly their water levels, and support wetland communities and species of conservation concern. See Mainstone et al. (2016). Rivers and streams in Owenduff/Nepin Complex SAC are often fringed by acid wet grassland, and also blanket bog, heath and flush/poor fen

4010 Northern Atlantic wet heaths with *Erica tetralix*

To restore the favourable conservation condition of Northern Atlantic wet heaths with *Erica tetralix* in Owenduff/Nephin Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Northern Atlantic wet heaths with <i>Erica tetralix</i> has not been mapped in detail for Owenduff/Nephin Complex SAC, but from current available data the total area of the qualifying habitat is estimated to be approximately 4,524ha, covering 17% of the SAC (NPWS internal files). Further details on this and the following attributes can be found in the Owenduff/Nephin Complex SAC conservation objectives supporting document for blanket bogs and associated habitats
Habitat distribution	Occurrence	No decline, subject to natural processes	The habitat is documented to occur in mosaic with blanket bog within the SAC and is present on the lower slopes of mountains (NPWS, 2006). Further information can be found within NPWS (2006) and the blanket bogs and associated habitats supporting document
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil nutrient status within natural range	See the blanket bogs and associated habitats supporting document for further details
Community diversity	Abundance of variety of vegetation communities	Maintain variety of vegetation communities, subject to natural processes	A variety of wet heath vegetation communities have been noted within the SAC (NPWS, 2006), two of which correspond to communities recorded in the National Survey of Upland Habitats and listed in the provisional list of vegetation communities described in Perrin et al. (2014). Further information on vegetation communities associated with this habitat is presented in Perrin et al. (2014)
Vegetation composition: cross-leaved heath	Occurrence within 20m of a representative number of monitoring stops	Cross-leaved heath (<i>Erica tetralix</i>) present within a 20m radius of each monitoring stop	Attribute and target based on Perrin et al. (2014)
Vegetation composition: positive indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of positive indicator species at least 50%	Attribute and target based on Perrin et al. (2014), where the list of positive indicator species for this habitat is also presented
Vegetation composition: lichens and bryophytes	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of <i>Cladonia</i> and <i>Sphagnum</i> species, <i>Racomitrium lanuginosum</i> and pleurocarpous mosses at least 10%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: ericoid species and crowberry	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of ericoid species and crowberry (<i>Empetrum nigrum</i>) at least 15%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: dwarf shrub species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of dwarf shrubs less than 75%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: negative indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of negative indicator species less than 1%	Attribute and target based on Perrin et al. (2014), where the list of negative indicator species for this habitat is also presented
Vegetation composition: non-native species	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of non-native species less than 1%	Attribute and target based on Perrin et al. (2014). Rhododendron (<i>Rhododendron ponticum</i>) was recorded from wet heaths in the SAC (NPWS internal files)

Vegetation composition: native trees and shrubs	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of scattered native trees and shrubs less than 20%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: bracken	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of bracken (<i>Pteridium aquilinum</i>) less than 10%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: soft rush	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of soft rush (<i>Juncus effusus</i>) less than 10%	Attribute and target based on Perrin et al. (2014)
Vegetation structure: <i>Sphagnum</i> condition	Condition at a representative number of 2m x 2m monitoring stops	Less than 10% of the <i>Sphagnum</i> cover is crushed, broken and/or pulled up	Attribute and target based on Perrin et al. (2014)
Vegetation structure: signs of browsing	Percentage of shoots browsed at a representative number of 2m x 2m monitoring stops	Less than 33% collectively of the last complete growing season's shoots of ericoids, crowberry (<i>Empetrum nigrum</i>) and bog-myrtle (<i>Myrica gale</i>) showing signs of browsing	Attribute and target based on Perrin et al. (2014)
Vegetation structure: burning	Occurrence in local vicinity of a representative number of monitoring stops	No signs of burning in sensitive areas, into the moss, liverwort or lichen layer or exposure of peat surface due to burning	Attribute and target based on Perrin et al. (2014), where the list of sensitive areas for this habitat is also presented
Physical structure: disturbed bare ground	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of disturbed bare ground less than 10%	Attribute and target based on Perrin et al. (2014)
Physical structure: drainage	Percentage area in local vicinity of a representative number of monitoring stops	Area showing signs of drainage from heavy trampling, tracking or ditches less than 10%	Attribute and target based on Perrin et al. (2014)
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat	This includes species listed in the Flora (Protection) Order, 2015 (FPO) and/or the red data lists (Lockhart et al., 2012; Wyse Jackson et al., 2016). The FPO listed and Vulnerable marsh clubmoss (<i>Lycopodiella inundata</i>) (Wyse Jackson et al., 2016) is present within the SAC (NPWS, 2006), but cannot be assigned specifically to wet heath

Conservation Objectives for : Owenduff/Nephin Complex SAC [000534]

4060 Alpine and Boreal heaths

To restore the favourable conservation condition of Alpine and Boreal heaths in Owenduff/Nephin Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Alpine and Boreal heaths have not been mapped in detail for Owenduff/Nephin Complex SAC, but from current available data the total area of the qualifying habitat is estimated to be approximately 1,150ha, covering 4% of the SAC (NPWS internal files). Further details on this and the following attributes can be found in the Owenduff/Nephin Complex SAC conservation objectives supporting document for blanket bogs and associated habitats
Habitat distribution	Occurrence	No decline, subject to natural processes	The habitat occurs on summits and ridges above 400-500m where it forms a mosaic with bare rock (NPWS internal files). Further information can be found within NPWS internal files and the blanket bogs and associated habitats supporting document
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil nutrient status within natural range	See the blanket bogs and associated habitats supporting document for further details
Community diversity	Abundance of variety of vegetation communities	Maintain variety of vegetation communities, subject to natural processes	The diversity of Alpine and Boreal heath communities within this SAC is unknown. Further information on vegetation communities associated with this habitat is presented in Perrin et al. (2014)
Vegetation composition: lichens and bryophytes	Number of species at a representative number of 2m x 2m monitoring stops	Number of bryophyte or non-crustose lichen species present at each monitoring stop is at least three	Attribute and target based on Perrin et al. (2014)
Vegetation composition: positive indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of positive indicator species at least 66%	Attribute and target based on Perrin et al. (2014), where the list of positive indicator species for this habitat is also presented
Vegetation composition: dwarf shrub species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of dwarf shrub species at least 10%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: negative indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of negative indicator species less than 10%	Attribute and target based on Perrin et al. (2014), where the list of negative indicator species for this habitat is also presented
Vegetation composition: non-native species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of non-native species less than 1%	Attribute and target based on Perrin et al. (2014)
Vegetation structure: signs of grazing	Percentage of leaves grazed at a representative number of 2m x 2m monitoring stops	Less than 10% collectively of the live leaves of specific graminoids showing signs of grazing	Attribute and target based on Perrin et al. (2014). See the blanket bogs and associated habitats supporting document for the list of specific graminoids
Vegetation structure: signs of browsing	Percentage of shoots browsed at a representative number of 2m x 2m monitoring stops	Less than 33% collectively of the last complete growing season's shoots of ericoids and crowberry (<i>Empetrum nigrum</i>) showing signs of browsing	Attribute and target based on Perrin et al. (2014)
Vegetation structure: burning	Occurrence in local vicinity of a representative number of monitoring stops	No signs of burning within the habitat	Attribute and target based on Perrin et al. (2014)

Physical structure: disturbed bare ground	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of disturbed bare ground less than 10%	Attribute and target based on Perrin et al. (2014)
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat	This includes species listed in the Flora (Protection) Order, 2015 and/or the red data lists (Lockhart et al., 2012; Wyse Jackson et al., 2016). The Near Threatened Alpine clubmoss (<i>Diphasiastrum alpinum</i>) (Wyse Jackson et al., 2016) was recorded from the Nephin Beg Range by Praeger (NPWS, 2006). This species is known to be associated with this habitat type

Conservation Objectives for : Owenduff/Nephin Complex SAC [000534]

5130 Juniperus communis formations on heaths or calcareous grasslands

To maintain the favourable conservation condition of *Juniperus communis* formations on heaths or calcareous grasslands in Owenduff/Nephin Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	<i>Juniperus communis</i> formations on heaths or calcareous grasslands habitat has not been mapped in detail for Owenduff/Nephin Complex SAC and thus the total area of the qualifying habitat is unknown. It has been noted that the habitat is rare within the SAC (Foss and McGee, 1987; Douglas et al., 1990) and is largely confined to ungrazed islands within larger dystrophic and oligotrophic lakes, and may also occur near well-drained areas of bog surrounding rock outcrops in the SAC, and often occurs in a mosaic with wet heath (NPWS, 2006; NPWS internal files)
Habitat distribution	Occurrence	No decline, subject to natural processes	See notes on habitat area above
Juniper population size	Number per formation	At least 50 plants per formation	To classify as a juniper (<i>Juniperus communis</i>) formation, at least 50 plants should be present (Cooper et al., 2012)
Vegetation composition: typical species	Number per formation	At least 50% of the listed positive indicator species for the relevant vegetation group present	Attribute and target based on Cooper et al. (2012), where positive indicator species for five vegetation groups are listed
Vegetation composition: negative indicator species	Occurrence per formation	Negative indicator species, particularly non-native invasive species, absent or under control	Attribute and target based on Cooper et al. (2012), where the list of negative indicator species is presented. Rhododendron (<i>Rhododendron ponticum</i>) has been reported from some of the lake islands that support juniper (<i>Juniperus communis</i>) scrub in the SAC (NPWS internal files)
Vegetation structure: cone-bearing plants	Percentage per formation	At least 10% of juniper plants are bearing cones	Attribute and target based on Cooper et al. (2012)
Vegetation structure: seedling recruitment	Percentage per formation	At least 10% of juniper plants are seedlings	Attribute and target based on Cooper et al. (2012)
Vegetation structure: dead juniper	Percentage per formation	Mean percentage of each juniper plant dead less than 10%	Attribute and target based on Cooper et al. (2012)

Conservation Objectives for : Owenduff/Nephin Complex SAC [000534]

7130 Blanket bogs (* if active bog)

To restore the favourable conservation condition of Blanket bogs in Owenduff/Nephin Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Blanket bog has not been mapped in detail for Owenduff/Nephin Complex SAC, but from current available data the total area of the qualifying habitat is estimated to be approximately 18,393ha, covering 68% of the SAC (NPWS internal files). Further details on this and the following attributes can be found in the Owenduff/Nephin Complex SAC conservation objectives supporting document for blanket bogs and associated habitats
Habitat distribution	Occurrence	No decline, subject to natural processes	The habitat covers most of the western and northern parts of the SAC, as well as much of the upland areas in the east and south. Large areas of intact blanket bog are also present in the centre of the SAC. Further information can be found within Foss and McGee (1987), Douglas et al. (1989), NPWS internal files and the blanket bogs and associated habitats supporting document
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil nutrient status within natural range	See the blanket bogs and associated habitats supporting document for further details
Ecosystem function: peat formation	Active blanket bog as a proportion of the total area of Annex I blanket bog habitat	At least 99% of the total Annex I blanket bog area is active	See the blanket bogs and associated habitats supporting document for further details
Ecosystem function: hydrology	Flow direction, water levels, occurrence of drains and erosion gullies	Natural hydrology unaffected by drains and erosion	Further details and a brief discussion of restoration potential is presented in the blanket bogs and associated habitats supporting document
Community diversity	Abundance of variety of vegetation communities	Maintain variety of vegetation communities, subject to natural processes	A variety of blanket bog vegetation communities have been recorded in this SAC (Foss and McGee, 1987; Douglas et al., 1989; NPWS internal files), five of which correspond to communities recorded in the National Survey of Upland Habitats and listed in the provisional list of vegetation communities described in Perrin et al. (2014). Further information on vegetation communities associated with this habitat is presented in Perrin et al. (2014)
Vegetation composition: positive indicator species	Number of species at a representative number of 2m x 2m monitoring stops	Number of positive indicator species present at each monitoring stop is at least seven	Attribute and target based on Perrin et al. (2014), where the list of positive indicator species for this habitat is also presented
Vegetation composition: lichens and bryophytes	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of bryophytes or lichens, excluding <i>Sphagnum fallax</i> , at least 10%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: potential dominant species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of each of the potential dominant species less than 75%	Attribute and target based on Perrin et al. (2014). See the blanket bogs and associated habitats supporting document for the list of potential dominant species
Vegetation composition: negative indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of negative indicator species less than 1%	Attribute and target based on Perrin et al. (2014), where the list of negative indicator species for this habitat is also presented
Vegetation composition: non-native species	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of non-native species less than 1%	Attribute and target based on Perrin et al. (2014). Rhododendron (<i>Rhododendron ponticum</i>) and the non-native moss <i>Campylopus introflexus</i> are present within blanket bog in the SAC (NPWS, 2006; NPWS internal files)

Vegetation composition: native trees and shrubs	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of scattered native trees and shrubs less than 10%	Attribute and target based on Perrin et al. (2014)
Vegetation structure: <i>Sphagnum</i> condition	Condition at a representative number of 2m x 2m monitoring stops	Less than 10% of the <i>Sphagnum</i> cover is crushed, broken and/or pulled up	Attribute and target based on Perrin et al. (2014)
Vegetation structure: signs of browsing	Percentage of shoots browsed at a representative number of 2m x 2m monitoring stops	Last complete growing season's shoots of ericoids, crowberry (<i>Empetrum nigrum</i>) and bog-myrtle (<i>Myrica gale</i>) showing signs of browsing collectively less than 33%	Attribute and target based on Perrin et al. (2014)
Vegetation structure: burning	Occurrence in local vicinity of a representative number of monitoring stops	No signs of burning in sensitive areas, into the moss, liverwort or lichen layer or exposure of peat surface due to burning	Attribute and target based on Perrin et al. (2014), where the list of sensitive areas for this habitat is also presented
Physical structure: disturbed bare ground	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of disturbed bare ground less than 10%	Attribute and target based on Perrin et al. (2014)
Physical structure: drainage	Percentage area in local vicinity of a representative number of monitoring stops	Area showing signs of drainage from heavy trampling, tracking or ditches less than 10%	Attribute and target based on Perrin et al. (2014)
Physical structure: erosion	Percentage area in local vicinity of a representative number of monitoring stops	Less than 5% of the greater bog mosaic comprises erosion gullies and eroded areas	Attribute and target based on Perrin et al. (2014)
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat	Several rare and threatened species are present in blanket bog flushes in the SAC including the Annex II and Annex IV listed, Flora (Protection) Order, 2015 (FPO) and Near Threatened (NT) marsh saxifrage (<i>Saxifraga hirculus</i>), the FPO and Vulnerable (VU) marsh clubmoss (<i>Lycopodiella inundata</i>), the FPO and NT bog orchid (<i>Hammarbya paludosa</i>), the Annex II listed, FPO and NT slender green feather-moss (<i>Hamatocaulis (Drepanocladus) vernicosus</i>) and the VU moss <i>Tomentypnum nitens</i> (NPWS, 2006; Lockhart et al., 2012; Campbell et al., 2015; Muldoon et al., 2015; Wyse Jackson et al., 2016; NPWS internal files). Although some of these species cannot be assigned specifically to blanket bog habitat (i.e. they are flush/fen species) they do occur in association with the habitat. The NT brown beak-sedge (<i>Rhynchospora fusca</i>) (Wyse Jackson et al., 2016) is also present within the SAC (NPWS, 2006), but cannot be assigned specifically to blanket bog

Conservation Objectives for : Owenduff/Nephin Complex SAC [000534]

7140 Transition mires and quaking bogs

To restore the favourable conservation condition of Transition mires and quaking bogs in Owenduff/Nephin Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Transition mires and quaking bogs have not been mapped in detail for Owenduff/Nephin Complex SAC and thus the total area of the qualifying habitat is unknown. Further details on this and the following attributes can be found in the Owenduff/Nephin Complex SAC conservation objectives supporting document for blanket bogs and associated habitats
Habitat distribution	Occurrence	No decline, subject to natural processes	The habitat occurs in locations where bog vegetation merges with base-rich flushes, and at the interface between water bodies and adjacent bog. Examples can be found at Owenglass West, Uggool, Sheeanmore and Lagduff. Further information can be found within Foss and McGee (1987), Douglas et al. (1989), NPWS (2006) and the blanket bogs and associated habitats supporting document
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil nutrient status within natural range	See the blanket bogs and associated habitats supporting document for further details
Community diversity	Abundance of variety of vegetation communities	Maintain variety of vegetation communities, subject to natural processes	Douglas et al. (1989) recorded one transition mire vegetation community that corresponds to a community recorded in the National Survey of Upland Habitats and listed in the provisional list of vegetation communities described in Perrin et al. (2014). Further information on vegetation communities associated with this habitat is presented in Perrin et al. (2014)
Vegetation composition: number of positive indicator species	Number of species at a representative number of 2m x 2m monitoring stops	Number of positive indicator species at each monitoring stop is at least three for infilling pools and flushes and at least six for fens	Attribute and target based on Perrin et al. (2014), where the list of positive indicator species for this habitat is also presented
Vegetation composition: number of core positive indicator species	Number of species at a representative number of 2m x 2m monitoring stops	At least one core positive indicator species present	Attribute and target based on Perrin et al. (2014), where the list of positive indicator species for this habitat is also presented
Vegetation composition: cover of positive indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of positive indicator species is at least 25%	Attribute and target based on Perrin et al. (2014), where the list of positive indicator species for this habitat is also presented
Vegetation composition: negative indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of negative indicator species less than 1%	Attribute and target based on Perrin et al. (2014), where the list of negative indicator species for this habitat is also presented
Vegetation composition: non-native species	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of non-native species less than 1%	Attribute and target based on Perrin et al. (2014)
Vegetation structure: height	Percentage of leaves/shoots at a representative number of 2m x 2m monitoring stops	Proportion of live leaves and/or flowering shoots of vascular plants that are more than 15cm above the ground surface should be at least 50%	Attribute and target based on Perrin et al. (2014). This attribute is only applicable to fen and flush examples of the habitat, not to infilling pool examples

Physical structure: disturbed bare ground	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of disturbed bare ground less than 10%	Attribute and target based on Perrin et al. (2014)
Physical structure: drainage	Percentage area in local vicinity of a representative number of monitoring stops	Area showing signs of drainage from heavy trampling, tracking or ditches less than 10%	Attribute and target based on Perrin et al. (2014)
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat	This includes species listed in the Flora (Protection) Order, 2015 (FPO) and/or the red data lists (Lockhart et al., 2012; Wyse Jackson et al., 2016). The FPO listed and Vulnerable marsh clubmoss (<i>Lycopodiella inundata</i>), the FPO listed and Near Threatened bog orchid (<i>Hammarbya paludosa</i>), the Near Threatened brown beak-sedge (<i>Rhynchospora fusca</i>) (Wyse Jackson et al., 2016) and the Vulnerable moss <i>Tomentypnum nitens</i> (Lockhart et al., 2012) are present within the SAC (NPWS, 2006; NPWS internal files), but cannot be assigned specifically to transition mires. The Annex II and FPO listed and Near Threatened slender green feather-moss (<i>Hamatocaulis (Drepanocladus) vernicosus</i>) (Lockhart et al., 2012) occurs in the habitat in the SAC (Campbell et al., 2015). See also the conservation objective for slender green feather-moss (1393)

Conservation Objectives for : Owenduff/Nephin Complex SAC [000534]

1106 Salmon *Salmo salar*

To restore the favourable conservation condition of Atlantic Salmon in Owenduff/Nephin Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Distribution: extent of anadromy	Percentage of river accessible	100% of river channels down to second order accessible from estuary	Artificial barriers block salmons' upstream migration, thereby limiting species to lower stretches and restricting access to spawning areas
Adult spawning fish	Number	Conservation limit (CL) for each system consistently exceeded	A conservation limit (CL) is defined by the North Atlantic Salmon Conservation Organisation (NASCO) as "the spawning stock level that produces long-term average maximum sustainable yield as derived from the adult to adult stock and recruitment relationship". The target is based on the Standing Scientific Committee on Salmon (SSCS) annual model output of CL attainment levels. See SSCS (2016). Attainment of CL estimates are derived from direct counts of adults (rod catch, fish counter) or indirectly by fry abundance counts. The Owenmore River is currently below CL
Salmon fry abundance	Number of fry/5 minutes electrofishing	Maintain or exceed 0+ fry mean catchment-wide abundance threshold value. Currently set at 17 salmon fry/5 minutes sampling	The target is the threshold value for rivers currently exceeding their conservation limit (CL)
Out-migrating smolt abundance	Number	No significant decline	Smolt abundance can be negatively affected by a number of impacts such as estuarine pollution, predation and sea lice (<i>Lepeophtheirus salmonis</i>)
Number and distribution of redds	Number and occurrence	No decline in number and distribution of spawning redds due to anthropogenic causes	Salmon spawn in clean gravels
Water quality	EPA Q value	At least Q4 at all sites sampled by EPA	Q values based on triennial water quality surveys carried out by the Environmental Protection Agency (EPA)

Conservation Objectives for : Owenduff/Nephin Complex SAC [000534]

1355 Otter *Lutra lutra*

To maintain the favourable conservation condition of Otter in Owenduff/Nephin Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Distribution	Percentage positive survey sites	No significant decline	Measure based on standard otter survey technique. Favourable Conservation Status (FCS) target, based on 1980/81 survey findings, is 88% in SACs. Current range is estimated at 93.6% (Reid et al., 2013)
Extent of terrestrial habitat	Hectares	No significant decline. Area mapped and calculated as 840.63ha along river banks/lake shoreline/ around pools	No field survey. Areas mapped to include 10m terrestrial buffer along shorelines and river banks identified as critical for otters (NPWS, 2007)
Extent of freshwater (river) habitat	Kilometres	No significant decline. Length mapped and calculated as 382.65km	No field survey. River length calculated on the basis that otters will utilise freshwater habitats from estuary to headwaters (Chapman and Chapman, 1982)
Extent of freshwater (lake) habitat	Hectares	No significant decline. Area mapped and calculated as 540.66ha	No field survey. Area mapped based on evidence that otters tend to forage within 80m of the shoreline (NPWS, 2007)
Couching sites and holts	Number	No significant decline	Otters need lying up areas throughout their territory where they are secure from disturbance (Kruuk and Moorhouse, 1991; Kruuk, 2006)
Fish biomass available	Kilograms	No significant decline	Broad diet that varies locally and seasonally, but dominated by fish, in particular salmonids, eels and sticklebacks in freshwater (Bailey and Rochford, 2006; Reid et al., 2013)
Barriers to connectivity	Number	No significant increase	Otters will regularly commute across stretches of open water up to 500m e.g. between the mainland and an island; between two islands; across an estuary (De Jongh and O'Neill, 2010). It is important that such commuting routes are not obstructed

1393 Slender Green Feather-moss *Drepanocladus vernicosus*

To maintain the favourable conservation condition of Slender Green Feather-moss (Shining Sickie-moss) in Owenduff/Nephin Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Distribution	Number and geographical spread of populations	No decline, subject to natural processes. See map 4 for known location at Ugool	(Please note that <i>Drepanocladus vernicosus</i> was reclassified as <i>Hamatocaulis vernicosus</i> by Hedenäs (1989)). The known population of slender green feather-moss (<i>Hamatocaulis vernicosus</i>) in Owenduff/Nephin Complex SAC occurs in a flush within the blanket bog at Ugool, in the vicinity of marsh saxifrage (<i>Saxifraga hirculus</i>). Data from NPWS survey by N. Lockhart in 1999 (NPWS internal files). See also Campbell et al. (2015)
Population size	Number of individuals	No decline, subject to natural processes	Lockhart estimated the population to be c.320 shoots (NPWS internal files). This is likely to be an underestimate. See Campbell et al. (2015) for further details
Area of suitable habitat	Hectares	No decline, subject to natural processes	The extent of occupancy for the species at Ugool was estimated by Lockhart to be one square metre; however, only about 4% of this area was suitable i.e. 0.04m ² (c.0.00004ha) (NPWS internal files). This is likely to be an underestimate. See Campbell et al. (2015) for further details
Hydrological conditions: water table level	Metres	Maintain suitable hydrological conditions	Slender green feather-moss (<i>Hamatocaulis vernicosus</i>) is mostly confined to mesotrophic fens, a transitional habitat between acid bog and base-rich fen. This appears to occur in at least two forms in Ireland: upland transitional flushes, where the plants can occur in lawns that rise and fall with fluctuating water table levels, such as at Ugool; and wet lowland sedge meadows, where plants can be inundated in winter, but may be subject to some desiccation in the summer. Based on Campbell (2013) and Campbell et al. (2015)
Vegetation composition: tree cover	Percentage cover in a representative number of 2m x 2m monitoring plots	Mean percentage tree cover should be less than 15%	Slender green feather-moss (<i>Hamatocaulis vernicosus</i>) grows in moss-dominated, open communities, generally with a low cover of trees and shrubs. See Campbell et al. (2015) for further details
Vegetation composition: shrub cover	Percentage cover in a representative number of 2m x 2m monitoring plots	Mean percentage shrub cover should be less than 20%	Slender green feather-moss (<i>Hamatocaulis vernicosus</i>) grows in moss-dominated, open communities, generally with a low cover of trees and shrubs. See Campbell et al. (2015) for further details
Vegetation composition: grass cover	Percentage cover in a representative number of 2m x 2m monitoring plots	Mean percentage grass species cover should be less than 25%	Slender green feather-moss (<i>Hamatocaulis vernicosus</i>) grows in moss-dominated, open communities, generally with a low cover of grasses, maintained by a low grazing intensity by sheep at Ugool. See Campbell et al. (2015) for further details
Vegetation composition: bryophyte cover	Percentage cover in a representative number of 2m x 2m monitoring plots	Mean percentage bryophyte cover should be more than 50%	In 1999, Lockhart recorded slender green feather-moss (<i>Hamatocaulis vernicosus</i>) at the edge of spring-dominated vegetation, which occurs at the edge of a swelling lawn of mosses, with <i>Aneura pinguis</i> , <i>Cratoneuron filicinum</i> , <i>Palustriella commutata</i> , <i>Philonotis fontana</i> , <i>Scorpidium revolvens</i> and <i>Warnstorfia exannulata</i> (NPWS internal files). See Campbell et al. (2015) for further details
Vegetation composition: cover of <i>Calliergonella cuspidata</i>	Percentage cover in a representative number of 2m x 2m monitoring plots	Mean percentage cover of <i>Calliergonella cuspidata</i> should be less than 15%	<i>Calliergonella cuspidata</i> , a moss species often associated with high nutrient conditions, is usually present, but with low cover and never dominant. See also Campbell et al. (2015)


Vegetation structure: vegetation height	Centimetres in a representative number 2m x 2m monitoring plots	Mean vegetation height should not exceed 40cm	See Campbell et al. (2015) for further details
---	---	---	--

Conservation Objectives for : Owenduff/Nephin Complex SAC [000534]

1528 Marsh Saxifrage *Saxifraga hirculus*


To maintain the favourable conservation condition of Marsh Saxifrage in Owenduff/Nephin Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Distribution	Number and geographical spread of populations	No loss in geographical spread and number of populations, subject to natural processes. See map 5 for 1km grid square locations	Marsh saxifrage (<i>Saxifraga hirculus</i>) is known to occur in the Owenduff/Nephin Complex SAC in five flushes at Sheean A, Sheean B, Sheean C, Sheean D and Uggool. See Lockhart (1989), Muldoon (2011) and Muldoon et al. (2015) for further details
Population size: number of rosettes	Number	Maintain the size of each known population, subject to natural processes. The target numbers of rosettes are: at least 151,200 rosettes at Sheean A, at least 36,000 rosettes at Sheean B, at least 104,000 rosettes at Sheean C, at least 19,200 rosettes at Sheean D and at least 24,000 rosettes at Uggool	The number of rosettes recorded by Muldoon (2011) were: 189,000 at Sheean A, 45,000 at Sheean B, 130,000 at Sheean C, 24,000 at Sheean D and 30,000 at Uggool. The target figures are a 20% reduction of the recorded number to allow for a margin of error and variability over monitoring seasons. See Muldoon et al. (2015) for further details
Population size: area of occupancy	Hectares	Maintain the area of occupancy of each known population, subject to natural processes. The target areas of occupancy are: at least 0.162ha at Sheean A, at least 0.042ha at Sheean B, at least 0.078ha at Sheean C, at least 0.051ha at Sheean D and at least 0.029ha at Uggool	The areas of occupancy for the species estimated by Muldoon (2011) were: 1,800m ² (0.1800ha) at Sheean A, 470m ² (0.047ha) at Sheean B, 870m ² (0.087ha) at Sheean C, 570m ² (0.057ha) at Sheean D and 315m ² (0.032ha) at Uggool. The target area figures are a 10% reduction of the recorded areas to allow for a margin of error. See Muldoon et al. (2015) for further details
Hydrological conditions: water level	Occurrence of high or fluctuating water levels	Maintain the appropriate natural hydrological regime necessary to support the habitat for the species	In Ireland, marsh saxifrage (<i>Saxifraga hirculus</i>) is now restricted to mineral flushes in blanket bog where rising groundwater forms small streams and seepage areas suitable for the species. Based on Muldoon (2011) and Muldoon et al. (2015)
Vegetation composition: positive indicator species	Occurrence in a number of 1m x 1m monitoring stops	Knotted pearlwort (<i>Sagina nodosa</i>) should be present in at least two of five 1m x 1m monitoring stops	The presence of the positive indicator species knotted pearlwort (<i>Sagina nodosa</i>) should be maintained (Muldoon, 2011; Muldoon et al., 2015)
Vegetation composition: negative indicator species	Mean percentage cover in five 1m x 1m monitoring stops	Mean percentage cover of purple moor-grass (<i>Molinia caerulea</i>) should not exceed 5%; mean percentage cover of Yorkshire fog (<i>Holcus lanatus</i>) should not exceed 15%	Low cover of the negative indicator species purple moor-grass (<i>Molinia caerulea</i>) and Yorkshire fog (<i>Holcus lanatus</i>) should be maintained. Cover of Yorkshire fog was greater than 15% at Uggool (Muldoon, 2011). See Muldoon et al. (2015) for further details
Vegetation structure: sward structure	Centimetres in five 1m x 1m monitoring stops	Maintain a mean vegetation height of less than 15cm	See Muldoon (2011) and Muldoon et al. (2015) for further details
Vegetation structure: grazing level	Evidence of grazing	Maintain grazing at light to moderate levels to ensure an open vegetation structure and to allow flowering to occur	See Muldoon (2011) and Muldoon et al. (2015) for further details



Legend

 Owenduff/Nephin Complex SAC 000534




An Roinn Ealaíon, Oidhreacht, Gnóthaí Réigiúnacha, Tuaithe agus Gaeltachta
 Department of Arts, Heritage, Regional, Rural and Gaeltacht Affairs

**MAP 1:
 OWENDUFF/NEPHIN COMPLEX SAC
 CONSERVATION OBJECTIVES
 SAC DESIGNATION**


Map to be read in conjunction with the NPWS Conservation Objectives Document.

**SITE CODE:
 SAC 000534; version 1.09. CO. MAYO**




The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision. Ordnance Survey of Ireland Licence No EN 0059216. © Ordnance Survey of Ireland Government of Ireland.

Nil sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbheithníthe a déanamh ar theorainneacha na gceantar comharthaithe. Suirbhéarachta Ordonáis na hÉireann Ceadúnas Uimh EN 0059216. © Suirbhéarachta Ordonáis na hÉireann Rialtas na hÉireann.



**Map Version 1
 Date: March 2017**


© Ordnance Survey Ireland



Legend

- Owenduff/Nephin Complex SAC 000534
- Carrowmore Lake Complex SAC 000476
- Corraun Plateau SAC 000485
- Clew Bay Complex SAC 001482
- Owenduff/Nephin Complex SPA 004098
- Blacksod Bay/Broad Haven SPA 004037

© Ordnance Survey Ireland






Legend

- Owenduff/Nephin Complex SAC 000534
- OSi Discovery Series County Boundary




Indicative Lake Habitats



- Potential 3110 Potential oligotrophic waters containing very few minerals of sandy plains: *Littorelletalia uniflorae*
- Potential 3110 / Potential 3160 Potential oligotrophic waters containing very few minerals of sandy plains: *Littorelletalia uniflorae* / Potential natural dystrophic lakes and ponds
- Potential 3160 Potential natural dystrophic lakes and ponds








Legend

-  1393 Slender Green Feather-moss *Hamatocaulis vernicosus*
-  Owenduff/Nephin Complex SAC 000534
-  OSi Discovery Series County Boundary



Legend

-  1538 Marsh Saxifrage *Saxifraga hirculus*
-  Owenduff/Nephin Complex SAC 000534
-  OSi Discovery Series County Boundary



Conservation objectives for Skealaghan Turlough SAC [000541]

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Objective: To maintain or restore the favourable conservation condition of the Annex I habitat(s) and/or the Annex II species for which the SAC has been selected:

Code	Description
------	-------------

3180	Turloughs*
------	------------

* denotes a priority habitat



Citation: NPWS (2020) Conservation objectives for Skealaghan Turlough SAC [000541]. Generic Version 7.0. Department of Culture, Heritage and the Gaeltacht.

National Parks and Wildlife Service

Conservation Objectives Series

Slieve Fyagh Bog SAC 000542



An Roinn Ealaíon, Oidhreachta,
Gnóthaí Réigiúnacha, Tuaithe agus Gaeltachta

Department of Arts, Heritage,
Regional, Rural and Gaeltacht Affairs



**National Parks and Wildlife Service,
Department of Arts, Heritage, Regional, Rural and Gaeltacht Affairs,
7 Ely Place, Dublin 2, Ireland.
Web: www.npws.ie
E-mail: nature.conservation@ahg.gov.ie**

Citation:

**NPWS (2016) Conservation Objectives: Slieve Fyagh Bog SAC 000542. Version 1.
National Parks and Wildlife Service, Department of Arts, Heritage, Regional,
Rural and Gaeltacht Affairs.**

**Series Editor: Rebecca Jeffrey
ISSN 2009-4086**

Introduction

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

** indicates a priority habitat under the Habitats Directive*

000542 Slieve Fyagh Bog SAC

7130 Blanket bogs (* if active bog)

Please note that this SAC adjoins Carrowmore Lake Complex SAC (000476) and Glenamoy Bog Complex SAC (000500). See map 2. The conservation objectives for this site should be used in conjunction with those for the adjacent sites as appropriate.

Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

NPWS Documents

Year :	1987
Title :	A survey to locate blanket bogs of scientific interest in County Mayo. Part I
Author :	Foss, P.; McGee, E.
Series :	A report commissioned by the Wildlife Service
Year :	1989
Title :	Survey to locate blanket bogs of scientific interest in Mayo. Part II
Author :	Doughlas, C.; Garvey, L.; Kelly, L.; O'Sullivan, A; Van Doorsleer, L.
Series :	A report commissioned by the Wildlife Service
Year :	2012
Title :	Ireland Red List no. 8: Bryophytes
Author :	Lockhart, N.; Hodgetts, N.; Holyoak, D.
Series :	Ireland Red List series, NPWS
Year :	2013
Title :	The status of EU protected habitats and species in Ireland. Volume 2. Habitats assessments
Author :	NPWS
Series :	Conservation assessments
Year :	2014
Title :	Guidelines for a national survey and conservation assessment of upland vegetation and habitats in Ireland, Version 2.0
Author :	Perrin, P.M.; Barron, S.J.; Roche, J.R.; O'Hanrahan, B.
Series :	Irish Wildlife Manual No. 79
Year :	2016
Title :	Slieve Fyagh Bog SAC (site code: 542) Conservation objectives supporting document- raised bog habitats V1
Author :	NPWS
Series :	Conservation objectives supporting document

Other References

Year :	1988
Title :	The Irish red data book 1. Vascular plants
Author :	Curtis, T.G.F; McGough, H.N.
Series :	Wildlife Service, Dublin


Conservation Objectives for : Slieve Fyagh Bog SAC [000542]

7130 Blanket bogs (* if active bog)


To restore the favourable conservation condition of Blanket bogs in Slieve Fyagh Bog SAC, which is defined by the following list of attributes and targets:


Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Blanket bog has not been mapped in detail for Slieve Fyagh Bog SAC, but from current available data the total area of the qualifying habitat is approximately 1,700 ha. Further information can be found in Foss and McGee (1987) and Douglas et al. (1989). Further details on this and the following attributes can be found in the Slieve Fyagh Bog SAC conservation objectives supporting document for upland habitats
Habitat distribution	Occurrence	No decline, subject to natural processes	Extensive areas of blanket bogs were recorded by Foss and McGee (1987) and Douglas et al. (1989) throughout this SAC. Further information can be found within these sources and the uplands supporting document
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil nutrient status within natural range	See the uplands supporting document for further details
Ecosystem function: peat formation	Active blanket bog as a proportion of the total area of Annex I blanket bog habitat	At least 99% of the total Annex I blanket bog area is active	See the uplands supporting document for further details
Ecosystem function: hydrology	Flow direction, water levels, occurrence of drains and erosion gullies	Natural hydrology unaffected by drains and erosion	Further details and a brief discussion of restoration potential is presented in the upland habitats supporting document
Community diversity	Abundance of variety of vegetation communities	Maintain variety of vegetation communities, subject to natural processes	Douglas et al. (1989) recorded different active blanket bogs communities within this SAC. Further information on vegetation communities associated with this habitat is presented in Perrin et al. (2014)
Vegetation composition: positive indicator species	Number of species at a representative number of 2m x 2m monitoring stops	Number of positive indicator species at each monitoring stop is at least seven	Based on Perrin et al. (2014), where the list of positive indicator species for this habitat is also presented. See the uplands supporting document for further details
Vegetation composition: lichens and bryophytes	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of bryophytes or lichens, excluding <i>Sphagnum fallax</i> , at least 10%	Based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation composition: potential dominant species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of each of the potential dominant species less than 75%	Based on Perrin et al. (2014). See the uplands supporting document for further details, including the list of potentially dominant species
Vegetation composition: negative indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of negative indicator species less than 1%	Based on Perrin et al. (2014), where the list of negative indicator species for this habitat is also presented. See the uplands supporting document for further details
Vegetation composition: non-native species	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of non-native species less than 1%	Based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation composition: native trees and scrub	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of scattered native trees and shrubs less than 10%	Based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation structure: <i>Sphagnum</i> condition	Condition at a representative number of 2m x 2m monitoring stops	Less than 10% of the <i>Sphagnum</i> cover is crushed, broken and/or pulled up	Based on Perrin et al. (2014). See the uplands supporting document for further details

Vegetation structure: signs of browsing	Percentage of shoots browsed at a representative number of 2m x 2m monitoring stops	Last complete growing season's shoots of ericoids, crowberry (<i>Empetrum nigrum</i>) and bog-myrtle (<i>Myrica gale</i>) showing signs of browsing collectively less than 33%	Based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation structure: burning	Occurrence in local vicinity of a representative number of monitoring stops	No signs of burning in sensitive areas, into the moss, liverwort or lichen layer or exposure of peat surface due to burning	Based on Perrin et al. (2014), where the list of sensitive areas is also presented. See the uplands supporting document for further details
Physical structure: disturbed bare ground	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of disturbed bare ground less than 10%	Based on Perrin et al. (2014). See the uplands supporting document for further details
Physical structure: drainage	Percentage area in local vicinity of a representative number of monitoring stops	Area showing signs of drainage from heavy trampling, tracking or ditches less than 10%	Based on Perrin et al. (2014). See the uplands supporting document for further details
Physical structure: erosion	Occurrence in local vicinity of a representative number of monitoring stops	Less than 5% of the greater bog mosaic comprises erosion gullies and eroded areas	Based on Perrin et al. (2014). See the uplands supporting document for further details
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat	This includes species listed in the Flora (Protection) Order, 2015 and/or the red data lists, Curtis and McGough (1988) and Lockhart et al. (2012). See the uplands supporting document for further details



Legend

 Slieve Fyagh Bog SAC 000542



Legend

- Slieve Fyagh Bog SAC 000542
- Carrowmore Lake Complex SAC 000476
- Glenamoy Bog Complex SAC 000500

An Roinn Ealaíon, Oidhreachta, Gnóthaí Réigiúnacha, Tuairthe agus Gaeltachta
 Department of Arts, Heritage, Regional, Rural and Gaeltacht Affairs

MAP 2:
SLIEVE FYAGH BOG SAC
CONSERVATION OBJECTIVES
ADJOINING / ADJACENT AND
OVERLAPPING DESIGNATIONS

Map to be read in conjunction with the NPWS Conservation Objectives Document.

SITE CODE:
SAC 000542; version 3.01. SAC 000500; version 3.01.
SAC 000476; version 3. CO. MAYO.

0 1 2 3 4 5 km

The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
 Ordnance Survey of Ireland Licence No EN 0059216. © Ordnance Survey of Ireland Government of Ireland

Níl sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbhreithniú a déanamh ar theorainneacha na gceantar comharthaithe. Suirbhéarachta Ordonáis na hÉireann Ceadúnas Uimh EN 0059216. © Suirbhéarachta Ordonáis na hÉireann Rialtas na hÉireann

Map Version 1
Date: July 2016

National Parks and Wildlife Service

Conservation Objectives Series

Bellanagare Bog SAC 000592



An Roinn
Ealaíon, Oidhreachta agus Gaeltachta

Department of
Arts, Heritage and the Gaeltacht



**National Parks and Wildlife Service,
Department of Arts, Heritage and the Gaeltacht,**

7 Ely Place, Dublin 2, Ireland.

Web: www.npws.ie

E-mail: nature.conservation@ahg.gov.ie

Citation:

**NPWS (201) Conservation Objectives: Bellanagare Bog SAC 000592. Version 1.
National Parks and Wildlife Service, Department of Arts, Heritage and the
Gaeltacht.**

Series Editor: Rebecca Jeffrey

ISSN 2009-4086

Introduction

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

** indicates a priority habitat under the Habitats Directive*

000592 Bellanagare Bog SAC

7110 Active raised bogsE

7120 Degraded raised bogs still capable of natural regeneration

7150 Depressions on peat substrates of the Rhynchosporion

Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

NPWS Documents

Year :	2014
Title :	Raised Bog Monitoring and Assessment Survey 2013
Author :	Fernandez, F.; Connolly K.; Crowley W.; Denyer J.; Duff K.; Smith G.
Series :	Irish Wildlife Manual No. 81
<hr/>	
Year :	2014
Title :	National raised bog SAC management plan
Author :	Department of Arts, Heritage and the Gaeltacht
Series :	Draft for consultation. 15 January 2014
<hr/>	
Year :	2014
Title :	Bellanagare Bog (SAC 000592), Co. Roscommon, Site Report
Author :	Fernandez, F.; Connolly, K.; Crowley, W.; Denyer J.; Duff K.; Smith G.
Series :	Raised bog monitoring and assessment survey 2013
<hr/>	
Year :	2015
Title :	Bellanagare Bog SAC (site code: 592) Conservation objectives supporting document- raised bog habitats V1
Author :	NPWS
Series :	Conservation objectives supporting document

Other References

Year :	2011
Title :	Review and revision of empirical critical loads and dose-response relationships. Proceedings of an expert workshop, Noordwijkerhout, 23-25 June 2010
Author :	Bobbink, R.; Hettelingh, J.P.
Series :	RIVM report 680359002, Coordination Centre for Effects, National Institute for Public Health and the Environment (RIVM)
<hr/>	
Year :	2014
Title :	Nitrogen deposition and exceedance of critical loads for nutrient nitrogen in Irish grasslands
Author :	Henry, J.; Aherne, J.
Series :	Science of the Total Environment 470–471: 216–223

Spatial data sources

Year :	2014
Title :	Scientific Basis for Raised Bog Conservation in Ireland
GIS Operations :	RBSB13_SACs_ARB_DRB dataset, RBSB13_SACs_2012_HB dataset, RBSB13_SACs_DrainagePatterns_5k dataset and RBSB13_SAC_LIDAR_DTMs dataset clipped to SAC boundary. Expert opinion used as necessary to resolve any issues arising
Used For :	potential 7110; digital elevation model; drainage patterns (maps 2 and 4)
<hr/>	
Year :	2013
Title :	Raised Bog Monitoring and Assessment Survey 2013
GIS Operations :	RBMA13_ecotope_map dataset clipped to SAC boundary. Appropriate ecotopes selected and exported to new dataset. Expert opinion used as necessary to resolve any issues arising
Used For :	7110 ecotopes (map 3)
<hr/>	

Conservation Objectives for : Bellanagare Bog SAC [000592]

7110 Active raised bogs

To restore the favourable conservation condition of Active raised bogs in Bellanagare Bog SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Restore area of active raised bog to 139.1ha, subject to natural processes	Active Raised Bog (ARB) habitat was mapped at 49.6ha by Fernandez et al. (2014). Area of Degraded Raised Bog (DRB) on the High Bog (HB) has been modelled as 105.4ha. See map 2. However, it is estimated that only 73.8ha is potentially restorable to ARB by drain blocking. The total potential ARB on the HB is therefore estimated to be 123.4ha. Eco-hydrological assessments of the cutover estimates that an additional 15.7ha of bog forming habitats could be restored. The long term target for ARB is therefore 139.1ha. See raised bog supporting document for further details on this and following attributes
Habitat distribution	Occurrence	Restore the distribution and variability of active raised bog across the SAC. See map 3 for distribution in 2013	The ARB habitat at Bellanagare includes central and sub-central ecotope, as well as active flush and a soak system. There is also potential for ARB restoration on cutover areas of the bog (see area target above)
High bog area	Hectares	No decline in extent of high bog necessary to support the development and maintenance of active raised bog. See map 2	The area of high bog within Bellanagare Bog SAC in 2012 (latest figure available) was 879.1ha (DAHG 2014)
Hydrological regime: water levels	Centimetres	Restore appropriate water levels throughout the site	For ARB, mean water level needs to be near or above the surface of the bog lawns for most of the year. Seasonal fluctuations should not exceed 20cm, and should only be 10cm below the surface, except for very short periods of time. Open water is often characteristic of soak systems
Hydrological regime: flow patterns	Flow direction; slope	Restore, where possible, appropriate high bog topography, flow directions and slopes. See map 4 for current situation	ARB depends on mean water levels being near or above the surface of bog lawns for most of the year. Long and gentle slopes are the most favourable to achieve these conditions. Changes to flow directions due to subsidence of bogs can radically change water regimes and cause drying out of high quality ARB areas and soak systems
Transitional areas between high bog and adjacent mineral soils (including cutover areas)	Hectares; distribution	Restore adequate transitional areas to support/protect active raised bog and the services it provides	ARB is threatened due to effects of past drainage and peat cutting around margins of Bellanagare Bog. No natural marginal habitats exist along these margins. Eco-hydrological assessments have evaluated the potential for ARB restoration on cutover areas (see note for habitat area attribute above)
Vegetation quality: central ecotope, active flush, soaks, bog woodland	Hectares	Restore 69.6ha of central ecotope/active flush/soaks/bog woodland as appropriate	At least 50% of ARB habitat should be high quality (i.e. central ecotope, active flush, soaks, bog woodland). Target area of active raised bog for the site has been set at 139.1ha (see area target above)
Vegetation quality: microtopographical features	Hectares	Restore adequate cover of high quality microtopographical features	Hummock and hollow microtopography is well developed on Bellanagare Bog
Vegetation quality: bog moss (<i>Sphagnum</i>) species	Percentage cover	Restore adequate cover of bog moss (<i>Sphagnum</i>) species to ensure peat-forming capacity	<i>Sphagnum</i> cover varies naturally across Ireland with relatively high cover in the east to lower cover in the west. Hummock forming species such as <i>Sphagnum austinii</i> are particularly good peat formers. <i>Sphagnum</i> cover and distribution also varies naturally across a site

Typical ARB species: flora	Occurrence	Restore, where appropriate, typical active raised bog flora	Typical flora species include widespread species, as well as those with more restricted distributions but typical of the habitat's subtypes or geographical range
Typical ARB species: fauna	Occurrence	Restore, where appropriate, typical active raised bog fauna	Typical fauna species include widespread species, as well as those with more restricted distributions but typical of the habitat's subtypes or geographical range
Elements of local distinctiveness	Occurrence	Maintain features of local distinctiveness, subject to natural processes	High quality ARB is present in the form of active flush, particularly a flush in the mid-eastern lobe, which is essentially an infilling lake or soak where the <i>Sphagnum</i> cover is 76-90%
Negative physical indicators	Percentage cover	Negative physical features absent or insignificant	Negative physical indicators include: bare peat, algae dominated pools and hollows, marginal cracks, tear patterns, subsidence features such as dry mineral mounds /ridges emerging or expanding and evidence of burning
Vegetation composition: native negative indicator species	Percentage cover	Native negative indicator species at insignificant levels	Disturbance indicators include species indicative of conditions drying out such as abundant bog asphodel (<i>Narthecium ossifragum</i>), deergrass (<i>Trichophorum germanicum</i>) and haretail cotton-grass (<i>Eriophorum vaginatum</i>) forming tussocks; abundant magellanic bog-moss (<i>Sphagnum magellanicum</i>) in pools previously dominated by <i>Sphagnum</i> species typical of very wet conditions (e.g. feathery bog-moss (<i>S. cuspidatum</i>)); and indicators of frequent burning events such as abundant <i>Cladonia floerkeana</i> and high cover of carnation sedge (<i>Carex panicea</i>) (particularly in true midlands raised bogs)
Vegetation composition: non-native invasive species	Percentage cover	Non-native invasive species at insignificant levels and not more than 1% cover	Most common non-native invasive species include lodgepole pine (<i>Pinus contorta</i>), rhododendron (<i>Rhododendron ponticum</i>), and pitcherplant (<i>Sarracenia purpurea</i>)
Air quality: nitrogen deposition	kg N/ha/year	Air quality surrounding bog close to natural reference conditions. The total N deposition should not exceed 5kg N/ha/yr	Change in air quality can result from fertiliser drift; adjacent quarry activities; or other atmospheric inputs. The critical load range for ombrotrophic bogs has been set as between 5 and 10kg N/ha/yr (Bobbink and Hettelingh, 2011). The latest N deposition figures for the area around Bellanagare Bog suggests that the current level is approximately 10.1kg N/ha/yr (Henry and Aherne, 2014)
Water quality	Hydrochemical measures	Water quality on the high bog and in transitional areas close to natural reference conditions	Water chemistry within raised bogs is influenced by atmospheric inputs (rainwater). However, within soak systems, water chemistry is influenced by other inputs such as focused flow or interaction with underlying substrates. Water chemistry in areas surrounding the high bog varies due to influences of different water types (bog water, regional groundwater, and run-off from surrounding mineral lands)

Conservation Objectives for : Bellanagare Bog SAC [000592]

7120 Degraded raised bogs still capable of natural regeneration

The long-term aim for Degraded raised bogs still capable of natural regeneration is that its peat-forming capability is re-established; therefore, the conservation objective for this habitat is inherently linked to that of Active raised bogs (7110) and a separate conservation objective has not been set in Bellanagare Bog SAC


Attribute	Measure	Target	Notes
-----------	---------	--------	-------

Conservation Objectives for : Bellanagare Bog SAC [000592]


7150 Depressions on peat substrates of the Rhynchosporion


Depressions on peat substrates of the Rhynchosporion is an integral part of good quality Active raised bogs (7110) and thus a separate conservation objective has not been set for the habitat in Bellanagare Bog SAC


Attribute	Measure	Target	Notes
------------------	----------------	---------------	--------------






Legend

 Bellanagare Bog SAC 000592






Legend

-  Bellanagare Bog SAC 000592
-  High Bog Boundary
-  Potential 7110 *Active Raised Bog






Legend

- Bellanagare Bog SAC 000592
- High Bog Boundary
- Active Raised Bog Ecotopes**
- Central ecotope
- Soaks / active flush
- Sub-central ecotope





Legend

- Bellanagare Bog SAC 000592
- High Bog Boundary
- Drainage Patterns
- Contours

Elevation

128.41 m

91.17 m

National Parks and Wildlife Service

Conservation Objectives Series

Callow Bog SAC 000595



*An Roinn
Ealaíon, Oidhreachta agus Gaeltachta*

*Department of
Arts, Heritage and the Gaeltacht*



**National Parks and Wildlife Service,
Department of Arts, Heritage and the Gaeltacht,
7 Ely Place, Dublin 2, Ireland.
Web: www.npws.ie
E-mail: nature.conservation@ahg.gov.ie**

Citation:

**NPWS (201) Conservation Objectives: Callow Bog SAC 000595. Version 1.
National Parks and Wildlife Service, Department of Arts, Heritage and the
Gaeltacht.**

**Series Editor: Rebecca Jeffrey
ISSN 2009-4086**

Introduction

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

** indicates a priority habitat under the Habitats Directive*

000595 Callow Bog SAC

7110 Active raised bogsE

7120 Degraded raised bogs still capable of natural regeneration

7150 Depressions on peat substrates of the Rhynchosporion

Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

NPWS Documents

Year :	2014
Title :	Raised Bog Monitoring and Assessment Survey 2013
Author :	Fernandez, F.; Connolly K.; Crowley W.; Denyer J.; Duff K.; Smith G.
Series :	Irish Wildlife Manual No. 81
<hr/>	
Year :	2014
Title :	National raised bog SAC management plan
Author :	Department of Arts, Heritage and the Gaeltacht
Series :	Draft for consultation. 15 January 2014
<hr/>	
Year :	2014
Title :	Callow Bog (SAC 000595), Co. Roscommon, Site Report
Author :	Fernandez, F.; Connolly, K.; Crowley, W.; Denyer J.; Duff K.; Smith G.
Series :	Raised bog monitoring and assessment survey 2013
<hr/>	
Year :	2016
Title :	Callow Bog SAC (site code: 595) Conservation objectives supporting document- raised bog habitats V1
Author :	NPWS
Series :	Conservation objectives supporting document

Other References

Year :	2011
Title :	Review and revision of empirical critical loads and dose-response relationships. Proceedings of an expert workshop, Noordwijkerhout, 23-25 June 2010
Author :	Bobbink, R.; Hettelingh, J.P.
Series :	RIVM report 680359002, Coordination Centre for Effects, National Institute for Public Health and the Environment (RIVM)
<hr/>	
Year :	2014
Title :	Nitrogen deposition and exceedance of critical loads for nutrient nitrogen in Irish grasslands
Author :	Henry, J.; Aherne, J.
Series :	Science of the Total Environment 470–471: 216–223

Spatial data sources

Year :	2014
Title :	Scientific Basis for Raised Bog Conservation in Ireland
GIS Operations :	RBSB13_SACs_ARB_DRB dataset, RBSB13_SACs_2012_HB dataset, RBSB13_SACs_DrainagePatterns_5k dataset and RBSB13_SAC_LIDAR_DTMs dataset clipped to SAC boundary. Expert opinion used as necessary to resolve any issues arising
Used For :	potential 7110; digital elevation model; drainage patterns (maps 2 and 4)
<hr/>	
Year :	2013
Title :	Raised Bog Monitoring and Assessment Survey 2013
GIS Operations :	RBMA13_ecotope_map dataset clipped to SAC boundary. Appropriate ecotopes selected and exported to new dataset. Expert opinion used as necessary to resolve any issues arising
Used For :	7110 ecotopes (map 3)
<hr/>	

Conservation Objectives for : Callow Bog SAC [000595]

7110 Active raised bogs

To restore the favourable conservation condition of Active raised bogs in Callow Bog SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Restore area of active raised bog to 53.4ha, subject to natural processes	Active Raised Bog (ARB) habitat was mapped at 11.4ha by Fernandez et al. (2014). Area of Degraded Raised Bog (DRB) on the High Bog (HB) has been modelled as 48.4ha. See map 2. However, it is estimated that only 33.9ha is potentially restorable to ARB by drain blocking. The total potential ARB on the HB is therefore estimated to be 45.3ha. Eco-hydrological assessments of the cutover estimates that an additional 8.1ha of bog forming habitats could be restored. The long term target for ARB is therefore 53.4ha. See raised bog supporting document for further details on this and following attributes
Habitat distribution	Occurrence	Restore the distribution and variability of active raised bog across the SAC. See map 3 for distribution in 2012	ARB occurs in four of the six sections of Callow Bog. DRB occurs on all sections of the bog, which will require restoration measures. There is also potential for ARB restoration on cutover areas of the bog (see area target above)
High bog area	Hectares	No decline in extent of high bog necessary to support the development and maintenance of active raised bog. See map 2	The area of high bog within Callow Bog SAC in 2012 (latest figure available) was 352ha (DAHG 2014)
Hydrological regime: water levels	Centimetres	Restore appropriate water levels throughout the site	For ARB, mean water level needs to be near or above the surface of the bog lawns for most of the year. Seasonal fluctuations should not exceed 20cm, and should only be 10cm below the surface, except for very short periods of time. Open water is often characteristic of soak systems
Hydrological regime: flow patterns	Flow direction; slope	Restore, where possible, appropriate high bog topography, flow directions and slopes. See map 4 for current situation	ARB depends on mean water levels being near or above the surface of bog lawns for most of the year. Long and gentle slopes are the most favourable to achieve these conditions. Changes to flow directions due to subsidence of bogs can radically change water regimes and cause drying out of high quality ARB areas and soak systems
Transitional areas between high bog and adjacent mineral soils (including cutover areas)	Hectares; distribution	Restore adequate transitional areas to support/protect active raised bog and the services it provides	ARB is threatened due to effects of past drainage and peat-cutting around the margins of Callow Bog. Eco-hydrological assessments have evaluated the potential for ARB restoration on cutover areas (see note for habitat area attribute above)
Vegetation quality: central ecotope, active flush, soaks, bog woodland	Hectares	Restore 26.7ha of central ecotope/active flush/soaks/bog woodland as appropriate	At least 50% of ARB habitat should be high quality (i.e. central ecotope, active flush, soaks, bog woodland). Target area of active raised bog for the site has been set at 53.4ha (see area target above)
Vegetation quality: microtopographical features	Hectares	Restore adequate cover of high quality microtopographical features	High quality microtopography (hummocks, hollows and pools) is well developed in the ARB areas of Callow Bog
Vegetation quality: bog moss (<i>Sphagnum</i>) species	Percentage cover	Restore adequate cover of bog moss (<i>Sphagnum</i>) species to ensure peat-forming capacity	<i>Sphagnum</i> cover varies naturally across Ireland with relatively high cover in the east to lower cover in the west. Hummock forming species such as <i>Sphagnum austini</i> are particularly good peat formers. <i>Sphagnum</i> cover and distribution also varies naturally across a site
Typical ARB species: flora	Occurrence	Restore, where appropriate, typical active raised bog flora	Typical flora species include widespread species, as well as those with more restricted distributions but typical of the habitat's subtypes or geographical range

Typical ARB species: fauna	Occurrence	Restore, where appropriate, typical active raised bog fauna	Typical fauna species include widespread species, as well as those with more restricted distributions but typical of the habitat's subtypes or geographical range
Elements of local distinctiveness	Occurrence	Maintain features of local distinctiveness, subject to natural processes	These include features of geological, topographical, archaeological and hydrological interest as well as noteworthy species of flora and fauna
Negative physical indicators	Percentage cover	Negative physical features absent or insignificant	Negative physical indicators include: bare peat, algae dominated pools and hollows, marginal cracks, tear patterns, subsidence features such as dry mineral mounds /ridges emerging or expanding and evidence of burning
Vegetation composition: native negative indicator species	Percentage cover	Native negative indicator species at insignificant levels	Disturbance indicators include species indicative of conditions drying out such as abundant bog asphodel (<i>Narthecium ossifragum</i>), deergrass (<i>Trichophorum germanicum</i>) and harestail cotton-grass (<i>Eriophorum vaginatum</i>) forming tussocks; abundant magellanic bog-moss (<i>Sphagnum magellanicum</i>) in pools previously dominated by <i>Sphagnum</i> species typical of very wet conditions (e.g. feathery bog-moss (<i>S. cuspidatum</i>)); and indicators of frequent burning events such as abundant <i>Cladonia floerkeana</i> and high cover of carnation sedge (<i>Carex panicea</i>) (particularly in true midlands raised bogs)
Vegetation composition: non-native invasive species	Percentage cover	Non-native invasive species at insignificant levels and not more than 1% cover	Most common non-native invasive species include lodgepole pine (<i>Pinus contorta</i>), rhododendron (<i>Rhododendron ponticum</i>), and pitcherplant (<i>Sarracenia purpurea</i>)
Air quality: nitrogen deposition	kg N/ha/year	Air quality surrounding bog close to natural reference conditions. The total N deposition should not exceed 5kg N/ha/yr	Change in air quality can result from fertiliser drift; adjacent quarry activities; or other atmospheric inputs. The critical load range for ombrotrophic bogs has been set as between 5 and 10kg N/ha/yr (Bobbink and Hettelingh, 2011). The latest N deposition figures for the area around Callow Bog suggests that the current level is approximately 9.1kg N/ha/yr (Henry and Aherne, 2014)
Water quality	Hydrochemical measures	Water quality on the high bog and in transitional areas close to natural reference conditions	Water chemistry within raised bogs is influenced by atmospheric inputs (rainwater). However, within soak systems, water chemistry is influenced by other inputs such as focused flow or interaction with underlying substrates. Water chemistry in areas surrounding the high bog varies due to influences of different water types (bog water, regional groundwater and run-off from surrounding mineral lands)

Conservation Objectives for : Callow Bog SAC [000595]

7120 Degraded raised bogs still capable of natural regeneration

The long-term aim for Degraded raised bogs still capable of natural regeneration is that its peat-forming capability is re-established; therefore, the conservation objective for this habitat is inherently linked to that of Active raised bogs (7110) and a separate conservation objective has not been set in Callow Bog SAC


Attribute	Measure	Target	Notes

Conservation Objectives for : Callow Bog SAC [000595]


7150 Depressions on peat substrates of the Rhynchosporion


Depressions on peat substrates of the Rhynchosporion is an integral part of good quality Active raised bogs (7110) and thus a separate conservation objective has not been set for the habitat in Callow Bog SAC

Attribute	Measure	Target	Notes
------------------	----------------	---------------	--------------



Legend

 Callow Bog SAC 000595

 *An Roinn Ealaíon, Oidhreachta agus Gaeltachta*
 Department of Arts, Heritage and the Gaeltacht

**MAP 1:
 CALLOW BOG SAC
 CONSERVATION OBJECTIVES
 SAC DESIGNATION**


Map to be read in conjunction with the NPWS Conservation Objectives Document.

**SITE CODE:
 SAC 000595; version 3. Co. Roscommon**


0 250 500 750 1,000 m

The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
 Ordnance Survey of Ireland Licence No EN 0059216. © Ordnance Survey of Ireland Government of Ireland.

Níl sna teorainneacha ar na léarscáilleana ach nod garshuíomhach ginearálta. Féadfar athbheithniú a déanamh ar theorainneacha na gceantar comharthaite. Suirbhéarachta Ordonáis na hÉireann Ceadúnas Uimh EN 0059216. © Suirbhéarachta Ordonáis na hÉireann Rialtas na hÉireann.




**Map Version 1
 Date: Jan 2016**



Legend


- Callow Bog SAC 000595
- High Bog Boundary
- Potential *7110 Active Raised Bogs
- OSi Discovery Series County Boundary





Legend



- Callow Bog SAC 000595
- High Bog Boundary
- OSi Discovery Series County Boundary
- Active Raised Bogs Ecotopes**
- Central ecotope
- Sub-central ecotope



Legend

- Callow Bog SAC 000595
- OSi Discovery Series County Boundary
- High Bog Boundary
- Drainage Patterns
- Contours

Elevation



National Parks and Wildlife Service

Conservation Objectives Series

Carrowbehy/Caher Bog SAC 000597



*An Roinn
Ealaíon, Oidhreachta agus Gaeltachta*

*Department of
Arts, Heritage and the Gaeltacht*



**National Parks and Wildlife Service,
Department of Arts, Heritage and the Gaeltacht,**

7 Ely Place, Dublin 2, Ireland.

Web: www.npws.ie

E-mail: nature.conservation@ahg.gov.ie

Citation:

**NPWS (201) Conservation Objectives: Carrowbehy/Caher Bog SAC 000597.
Version 1. National Parks and Wildlife Service, Department of Arts, Heritage
and the Gaeltacht.**

Series Editor: Rebecca Jeffrey

ISSN 2009-4086

Introduction

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

** indicates a priority habitat under the Habitats Directive*

000597	Carrowbehy/Caher Bog SAC
7110	Active raised bogsE
7120	Degraded raised bogs still capable of natural regeneration
7150	Depressions on peat substrates of the Rhynchosporion

Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

NPWS Documents

Year :	2014
Title :	Raised Bog Monitoring and Assessment Survey 2013
Author :	Fernandez, F.; Connolly K.; Crowley W.; Denyer J.; Duff K.; Smith G.
Series :	Irish Wildlife Manual No. 81
<hr/>	
Year :	2014
Title :	National raised bog SAC management plan
Author :	Department of Arts, Heritage and the Gaeltacht
Series :	Draft for consultation. 15 January 2014
<hr/>	
Year :	2014
Title :	Carrowbehy Bog (SAC 000597), Co. Roscommon, Site Report
Author :	Fernandez, F.; Connolly, K.; Crowley, W.; Denyer J.; Duff K.; Smith G.
Series :	Raised bog monitoring and assessment survey 2013
<hr/>	
Year :	2015
Title :	Carrowbehy/Caher Bog SAC (site code: 597) Conservation objectives supporting document- raised bog habitats V1
Author :	NPWS
Series :	Conservation objectives supporting document

Other References

Year :	2011
Title :	Review and revision of empirical critical loads and dose-response relationships. Proceedings of an expert workshop, Noordwijkerhout, 23-25 June 2010
Author :	Bobbink, R.; Hettelingh, J.P.
Series :	RIVM report 680359002, Coordination Centre for Effects, National Institute for Public Health and the Environment (RIVM)
<hr/>	
Year :	2014
Title :	Nitrogen deposition and exceedance of critical loads for nutrient nitrogen in Irish grasslands
Author :	Henry, J.; Aherne, J.
Series :	Science of the Total Environment 470–471: 216–223

Spatial data sources

Year :	2014
Title :	Scientific Basis for Raised Bog Conservation in Ireland
GIS Operations :	RBSB13_SACs_ARB_DRB dataset, RBSB13_SACs_2012_HB dataset, RBSB13_SACs_DrainagePatterns_5k dataset and RBSB13_SAC_LIDAR_DTMs dataset clipped to SAC boundary. Expert opinion used as necessary to resolve any issues arising potential
Used For :	7110; digital elevation model; drainage patterns (maps 2 and 4)
<hr/>	
Year :	2013
Title :	Raised Bog Monitoring and Assessment Survey 2013
GIS Operations :	RBMA13_ecotope_map dataset clipped to SAC boundary. Appropriate ecotopes selected and exported to new dataset. Expert opinion used as necessary to resolve any issues arising
Used For :	7110 ecotopes (map 3)
<hr/>	

7110 Active raised bogs

To restore the favourable conservation condition of Active raised bogs in Carrowbehy/Caher Bog SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Restore area of active raised bog to 92.3ha, subject to natural processes	Active Raised Bog (ARB) habitat was mapped at 69.9ha by Fernandez et al. (2014). Area of Degraded Raised Bog (DRB) on the High Bog (HB) has been modelled as 35.6ha. See map 2. However, it is estimated that only 17.8ha is potentially restorable to ARB by drain blocking. The total potential ARB on the HB is therefore estimated to be 87.7ha. Eco-hydrological assessments of the cutover estimates that an additional 4.6ha of bog forming habitats could be restored. The long term target for ARB is therefore 92.3ha. See raised bog supporting document for further details on this and following attributes
Habitat distribution	Occurrence	Restore the distribution and variability of active raised bog across the SAC. See map 3 for distribution in 2012	ARB currently occurs on all four sections of Carrowbehy/Caher bog. DRB also occurs on these four parts, which will require restoration measures. There is also potential for ARB restoration on cutover areas of the bog (see area target above)
High bog area	Hectares	No decline in extent of high bog necessary to support the development and maintenance of active raised bog. See map 2	The area of high bog within Carrowbehy/Caher Bog SAC in 2012 (latest figure available) was 204.6ha (DAHG 2014)
Hydrological regime: water levels	Centimetres	Restore appropriate water levels throughout the site	For ARB, mean water level needs to be near or above the surface of the bog lawns for most of the year. Seasonal fluctuations should not exceed 20cm, and should only be 10cm below the surface, except for very short periods of time
Hydrological regime: flow patterns	Flow direction; slope	Restore, where possible, appropriate high bog topography, flow directions and slopes. See map 4 for current situation	ARB depends on mean water levels being near or above the surface of bog lawns for most of the year. Long and gentle slopes are the most favourable to achieve these conditions. Changes to flow directions due to subsidence of bogs can radically change water regimes and cause drying out of high quality ARB areas and soak systems
Transitional areas between high bog and adjacent mineral soils (including cutover areas)	Hectares; distribution	Restore adequate transitional areas to support/protect active raised bog and the services it provides	Studies suggest that the ARB is threatened due to water loss arising from past drainage and turf cutting, in particular along the southern margin of Carrowbehy/Caher Bog. The most intact margins of the bog include a narrow transitional zone between the bog and an esker to the north, while an example of a poor fen is found in a slight depression beside a major drainage channel at the north-eastern end of the SAC. Eco-hydrological assessments have evaluated the potential for ARB restoration on cutover areas (see note for habitat area attribute above)
Vegetation quality: central ecotope, active flush, soaks, bog woodland	Hectares	Maintain at least 57.7ha of central ecotope/active flush/soaks/bog woodland as appropriate	At least 50% of ARB habitat should be central ecotope/active flush/soaks/bog woodland. Target area of active raised bog for the site has been set at 92.3ha (see area target above). As the current area (57.7ha) of central ecotope/active flush/soaks exceeds the 50% target, this area is set as the target
Vegetation quality: microtopographical features	Hectares	Restore adequate cover of high quality microtopographical features	Hummock and hollow microtopography is well developed on Carrowbehy/Caher bog where interconnecting linear pools occur

Vegetation quality: bog moss (<i>Sphagnum</i>) species	Percentage cover	Restore adequate cover of bog moss (<i>Sphagnum</i>) species to ensure peat-forming capacity	<i>Sphagnum</i> cover varies naturally across Ireland with relatively high cover in the east to lower cover in the west. Hummock forming species such as <i>Sphagnum austini</i> are particularly good peat formers. <i>Sphagnum</i> cover and distribution also varies naturally across a site
Typical ARB species: flora	Occurrence	Restore, where appropriate, typical active raised bog flora	Typical flora species include widespread species, as well as those with more restricted distributions but typical of the habitat's subtypes or geographical range
Typical ARB species: fauna	Occurrence	Restore, where appropriate, typical active raised bog fauna	Typical fauna species include widespread species, as well as those with more restricted distributions but typical of the habitat's subtypes or geographical range
Elements of local distinctiveness	Occurrence	Maintain features of local distinctiveness, subject to natural processes	Infilling lakes, swallow holes, a number of flush/soak systems are the main features of local distinctiveness on Carrowbehy/Caher Bog
Negative physical indicators	Percentage cover	Negative physical features absent or insignificant	Negative physical indicators include: bare peat, algae dominated pools and hollows, marginal cracks, tear patterns, subsidence features such as dry mineral mounds /ridges emerging or expanding, and burning evidence
Vegetation composition: native negative indicator species	Percentage cover	Native negative indicator species at insignificant levels	Disturbance indicators include species indicative of conditions drying out such as abundant bog asphodel (<i>Narthecium ossifragum</i>), deergrass (<i>Trichophorum germanicum</i>) and harestalk cotton-grass (<i>Eriophorum vaginatum</i>) forming tussocks; abundant magellanic bog-moss (<i>Sphagnum magellanicum</i>) in pools previously dominated by <i>Sphagnum</i> species typical of very wet conditions (e.g. feathery bog-moss (<i>S. cuspidatum</i>)); and indicators of frequent burning events such as abundant <i>Cladonia floerkeana</i> and high cover of carnation sedge (<i>Carex panicea</i>) (particularly in true midlands raised bogs)
Vegetation composition: non-native invasive species	Percentage cover	Non-native invasive species at insignificant levels and not more than 1% cover	Most common non-native invasive species include lodgepole pine (<i>Pinus contorta</i>), rhododendron (<i>Rhododendron ponticum</i>), and pitcherplant (<i>Sarracenia purpurea</i>)
Air quality: nitrogen deposition	kg N/ha/year	Air quality surrounding bog close to natural reference conditions. The total N deposition should not exceed 5kg N/ha/yr	Change in air quality can result from fertiliser drift; adjacent quarry activities; or other atmospheric inputs. The critical load range for ombrotrophic bogs has been set as between 5kg N/ha/yr (Bobbink & Hettelingh 2011). The latest N deposition figures for the area around Carrowbehy/Caher bog suggests that the current level is approximately 9.8kg N/ha/yr (Henry and Aherne, 2014)
Water quality	Hydrochemical measures	Water quality on the high bog and in transitional areas close to natural reference conditions	Water chemistry within raised bogs is influenced by atmospheric inputs (rainwater). However, within soak systems, water chemistry is influenced by other inputs such as focused flow or interaction with underlying substrates. Water chemistry in areas surrounding the high bog varies due to influences of different water types (bog water, regional groundwater, and run-off from surrounding mineral lands)

Conservation Objectives for : Carrowbehy/Caher Bog SAC [000597]

7120 Degraded raised bogs still capable of natural regeneration

The long-term aim for Degraded raised bogs still capable of natural regeneration is that its peat-forming capability is re-established; therefore, the conservation objective for this habitat is inherently linked to that of Active raised bogs (7110) and a separate conservation objective has not been set in Carrowbehy/Caher Bog SAC


Attribute	Measure	Target	Notes
------------------	----------------	---------------	--------------

Conservation Objectives for : Carrowbehy/Caher Bog SAC [000597]


7150 Depressions on peat substrates of the Rhynchosporion


Depressions on peat substrates of the Rhynchosporion is an integral part of good quality Active raised bogs (7110) and thus a separate conservation objective has not been set for the habitat in Carrowbehy/Caher Bog SAC

Attribute	Measure	Target	Notes
-----------	---------	--------	-------



Legend

 Carrowbey/Caher Bog SAC 000597

 *An Roinn Ealaíon, Oidhreachta agus Gaeltachta*
 Department of Arts, Heritage and the Gaeltacht

MAP 1:
CARROWBEHY/CAHER BOG SAC
CONSERVATION OBJECTIVES
SAC DESIGNATION

Map to be read in conjunction with the NPWS Conservation Objectives Document.

SITE CODE:
SAC 000597; version 3.01. Co. Roscommon


0 200 400 600 800 1,000 m

The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
 Ordnance Survey of Ireland Licence No EN 0059214. © Ordnance Survey of Ireland Government of Ireland.

Níl sna teorainneacha ar na léarscáilleana ach nod garshuíomhach ginearálta. Féadfar athbheithnithe a déanamh ar theorainneacha na gceantar comharthaite. Suirbhéaracha Ordonáis na hÉireann Ceadúnas Uimh EN 0059214. © Suirbhéaracha Ordonáis na hÉireann Rialtas na hÉireann.



Map Version 1
Date: Oct 2015



Legend

- Carrowbehy/Caher Bog SAC 000597
- High Bog Boundary
- Potential 7110 *Active Raised Bogs
- OSi Discovery Series County Boundary

MAP 2:
CARROWBEHY/CAHER BOG SAC
CONSERVATION OBJECTIVES
EXTENT OF POTENTIAL
ACTIVE RAISED BOGS


Map to be read in conjunction with the NPWS Conservation Objectives Document.

SITE CODE:
SAC 000597; version 3.01. Co. Roscommon




The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
 Ordnance Survey of Ireland Licence No EN 0059214. © Ordnance Survey of Ireland Government of Ireland.

Níl sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbheithnithe a déanamh ar theorainneacha na gceantar comharthaite. Suirbhéarachta Ordonáis na hÉireann Ceadúnas Uimh EN 0059214. © Suirbhéarachta Ordonáis na hÉireann Rialtas na hÉireann.




Map Version 1
 Date: Oct 2015




Legend

-  Carrowbehy/Caher Bog SAC 000597
-  OSi Discovery Series County Boundary
-  High Bog Boundary

Active Raised Bog Ecotopes

-  Central ecotope
-  Soaks / active flush
-  Sub-central ecotope



Legend

- Carrowbehy/Caher Bog SAC 000597
- OSi Discovery Series County Boundary
- High Bog Boundary
- Drainage Patterns
- Contours

Elevation

98.22 m

78.74 m

National Parks and Wildlife Service

Conservation Objectives Series

Cloonchambers Bog SAC 000600



An Roinn
Ealaíon, Oidhreachta agus Gaeltachta

Department of
Arts, Heritage and the Gaeltacht



**National Parks and Wildlife Service,
Department of Arts, Heritage and the Gaeltacht,**

7 Ely Place, Dublin 2, Ireland.

Web: www.npws.ie

E-mail: nature.conservation@ahg.gov.ie

Citation:

**NPWS (201) Conservation Objectives: Cloonchambers Bog SAC 000600.
Version 1. National Parks and Wildlife Service, Department of Arts, Heritage
and the Gaeltacht.**

Series Editor: Rebecca Jeffrey

ISSN 2009-4086

Introduction

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

** indicates a priority habitat under the Habitats Directive*

000600	Cloonchambers Bog SAC
7110	Active raised bogsE
7120	Degraded raised bogs still capable of natural regeneration
7150	Depressions on peat substrates of the Rhynchosporion

Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

NPWS Documents

Year :	2014
Title :	Raised Bog Monitoring and Assessment Survey 2013
Author :	Fernandez, F.; Connolly K.; Crowley W.; Denyer J.; Duff K.; Smith G.
Series :	Irish Wildlife Manual No. 81
<hr/>	
Year :	2014
Title :	National raised bog SAC management plan
Author :	Department of Arts, Heritage and the Gaeltacht
Series :	Draft for consultation. 15 January 2014
<hr/>	
Year :	2014
Title :	Cloonchambers Bog (SAC 000600), Co. Roscommon, Site Report
Author :	Fernandez, F.; Connolly, K.; Crowley, W.; Denyer J.; Duff K.; Smith G.
Series :	Raised bog monitoring and assessment survey 2013
<hr/>	
Year :	2016
Title :	Cloonchambers Bog SAC (site code: 600) Conservation objectives supporting document- raised bog habitats V1
Author :	NPWS
Series :	Conservation objectives supporting document

Other References

Year :	2011
Title :	Review and revision of empirical critical loads and dose-response relationships. Proceedings of an expert workshop, Noordwijkerhout, 23-25 June 2010
Author :	Bobbink, R.; Hettelingh, J.P.
Series :	RIVM report 680359002, Coordination Centre for Effects, National Institute for Public Health and the Environment (RIVM)
<hr/>	
Year :	2014
Title :	Nitrogen deposition and exceedance of critical loads for nutrient nitrogen in Irish grasslands
Author :	Henry, J.; Aherne, J.
Series :	Science of the Total Environment 470–471: 216–223

Spatial data sources

Year :	2014
Title :	Scientific Basis for Raised Bog Conservation in Ireland
GIS Operations :	RBSB13_SACs_ARB_DRB dataset, RBSB13_SACs_2012_HB dataset, RBSB13_SACs_DrainagePatterns_5k dataset and RBSB13_SAC_LIDAR_DTMs dataset clipped to SAC boundary. Expert opinion used as necessary to resolve any issues arising
Used For :	potential 7110; digital elevation model; drainage patterns (maps 2 and 4)
<hr/>	
Year :	2013
Title :	Raised Bog Monitoring and Assessment Survey 2013
GIS Operations :	RBMA13_ecotope_map dataset clipped to SAC boundary. Appropriate ecotopes selected and exported to new dataset. Expert opinion used as necessary to resolve any issues arising
Used For :	7110 ecotopes (map 3)
<hr/>	

Conservation Objectives for : Cloonchambers Bog SAC [000600]

7110 Active raised bogs

To restore the favourable conservation condition of Active raised bogs in Cloonchambers Bog SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Restore area of active raised bog to 32.1ha, subject to natural processes	Active Raised Bog (ARB) habitat was mapped at 7.7ha in 2012 by Fernandez et al. (2014). Area of Degraded Raised Bog (DRB) on the High Bog (HB) has been modelled as 42.2ha. See map 2. However, it is estimated that only 21.1ha is potentially restorable to ARB by drain blocking. The total potential ARB on the HB is therefore estimated to be 28.8ha. Eco-hydrological assessments of the cutover estimates that an additional 3.3ha of bog forming habitats could be restored. The long term target for ARB is therefore 32.1ha. See raised bog supporting document for further details on this and following attributes
Habitat distribution	Occurrence	Restore the distribution and variability of active raised bog across the SAC. See map 3 for distribution in 2012	ARB occurs at a number of locations on Cloonchambers Bog. DRB occurs on all lobes of the bog, which will require restoration measures. There is also potential for ARB restoration on cutover areas of the bog (see area target above)
High bog area	Hectares	No decline in extent of high bog necessary to support the development and maintenance of active raised bog. See map 2	The area of high bog within Cloonchambers Bog SAC in 2012 (latest figure available) was 195.8ha (DAHG 2014)
Hydrological regime: water levels	Centimetres	Restore appropriate water levels throughout the site	For ARB, mean water level needs to be near or above the surface of the bog lawns for most of the year. Seasonal fluctuations should not exceed 20cm, and should only be 10cm below the surface, except for very short periods of time. Open water is often characteristic of soak systems
Hydrological regime: flow patterns	Flow direction; slope	Restore, where possible, appropriate high bog topography, flow directions and slopes. See map 4 for current situation	ARB depends on mean water levels being near or above the surface of bog lawns for most of the year. Long and gentle slopes are the most favourable to achieve these conditions. Changes to flow directions due to subsidence of bogs can radically change water regimes and cause drying out of high quality ARB areas and soak systems
Transitional areas between high bog and adjacent mineral soils (including cutover areas)	Hectares; distribution	Restore adequate transitional areas to support/protect active raised bog and the services it provides	ARB is threatened due to effects of past drainage and peat-cutting around margins of Cloonchambers Bog. No natural marginal habitats exist along these margins. Eco-hydrological assessments have evaluated the potential for ARB restoration on cutover areas (see note for habitat area attribute above)
Vegetation quality: central ecotope, active flush, soaks, bog woodland	Hectares	Restore 16.1ha of central ecotope/active flush/soaks/bog woodland as appropriate	At least 50% of ARB habitat should be high quality (i.e. central ecotope, active flush, soaks, bog woodland). Target area of active raised bog for the SAC has been set at 32.1ha (see area target above)
Vegetation quality: microtopographical features	Hectares	Restore adequate cover of high quality microtopographical features	High quality microtopography (hummocks and hollows) is moderately well developed on some areas of Cloonchambers Bog
Vegetation quality: bog moss (<i>Sphagnum</i>) species	Percentage cover	Restore adequate cover of bog moss (<i>Sphagnum</i>) species to ensure peat-forming capacity	<i>Sphagnum</i> cover varies naturally across Ireland with relatively high cover in the east to lower cover in the west. Hummock forming species such as <i>Sphagnum austinii</i> are particularly good peat formers. <i>Sphagnum</i> cover and distribution also varies naturally across a site

Typical ARB species: flora	Occurrence	Restore, where appropriate, typical active raised bog flora	Typical flora species include widespread species, as well as those with more restricted distributions but typical of the habitat's subtypes or geographical range
Typical ARB species: fauna	Occurrence	Restore, where appropriate, typical active raised bog fauna	Typical fauna species include widespread species, as well as those with more restricted distributions but typical of the habitat's subtypes or geographical range
Elements of local distinctiveness	Occurrence	Maintain features of local distinctiveness, subject to natural processes	Cloonchambers Bog is noted for the presence of an extensive flush system with fen vegetation, a rare feature of Irish raised bogs
Negative physical indicators	Percentage cover	Negative physical features absent or insignificant	Negative physical indicators include: bare peat, algae dominated pools and hollows, marginal cracks, tear patterns, subsidence features such as dry mineral mounds /ridges emerging or expanding and evidence of burning
Vegetation composition: native negative indicator species	Percentage cover	Native negative indicator species at insignificant levels	Disturbance indicators include species indicative of conditions drying out such as abundant bog asphodel (<i>Narthecium ossifragum</i>), deergrass (<i>Trichophorum germanicum</i>) and harestail cotton-grass (<i>Eriophorum vaginatum</i>) forming tussocks; abundant magellanic bog-moss (<i>Sphagnum magellanicum</i>) in pools previously dominated by <i>Sphagnum</i> species typical of very wet conditions (e.g. feathery bog-moss (<i>S. cuspidatum</i>)); and indicators of frequent burning events such as abundant <i>Cladonia floerkeana</i> and high cover of carnation sedge (<i>Carex panicea</i>) (particularly in true midlands raised bogs)
Vegetation composition: non-native invasive species	Percentage cover	Non-native invasive species at insignificant levels and not more than 1% cover	Most common non-native invasive species include lodgepole pine (<i>Pinus contorta</i>), rhododendron (<i>Rhododendron ponticum</i>), and pitcherplant (<i>Sarracenia purpurea</i>)
Air quality: nitrogen deposition	kg N/ha/year	Air quality surrounding bog close to natural reference conditions. The total N deposition should not exceed 5kg N/ha/yr	Change in air quality can result from fertiliser drift; adjacent quarry activities; or other atmospheric inputs. The critical load range for ombrotrophic bogs has been set as between 5 and 10kg N/ha/yr (Bobbink and Hettelingh, 2011). The latest N deposition figures for the area around Cloonchambers Bog suggests that the current level is approximately 10.2kg N/ha/yr (Henry and Aherne, 2014)
Water quality	Hydrochemical measures	Water quality on the high bog and in transitional areas close to natural reference conditions	Water chemistry within raised bogs is influenced by atmospheric inputs (rainwater). However, within soak systems, water chemistry is influenced by other inputs such as focused flow or interaction with underlying substrates. Water chemistry in areas surrounding the high bog varies due to influences of different water types (bog water, regional groundwater and run-off from surrounding mineral lands)

Conservation Objectives for : Cloonchambers Bog SAC [000600]

7120 Degraded raised bogs still capable of natural regeneration

The long-term aim for Degraded raised bogs still capable of natural regeneration is that its peat-forming capability is re-established; therefore, the conservation objective for this habitat is inherently linked to that of Active raised bogs (7110) and a separate conservation objective has not been set in Cloonchambers Bog SAC


Attribute	Measure	Target	Notes
------------------	----------------	---------------	--------------


Conservation Objectives for : Cloonchambers Bog SAC [000600]

7150 Depressions on peat substrates of the Rhynchosporion

Depressions on peat substrates of the Rhynchosporion is an integral part of good quality Active raised bogs (7110) and thus a separate conservation objective has not been set for the habitat in Cloonchambers Bog SAC

Attribute	Measure	Target	Notes
-----------	---------	--------	-------



Legend
 Cloonchambers Bog SAC 000600

An Roinn
 Ealaíon, Oidhreachta agus Gaeltachta
 Department of
 Arts, Heritage and the Gaeltacht

MAP 1:
CLOONCHAMBERS BOG SAC
CONSERVATION OBJECTIVES
SAC DESIGNATION
 Map to be read in conjunction with the NPWS Conservation Objectives Document.


SITE CODE:
SAC 000600; version 3. Co. Roscommon

0 100 200 300 400 500 m

The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
 Ordnance Survey of Ireland Licence No EN 0059216. © Ordnance Survey of Ireland Government of Ireland.


Níl sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbheithníthe a déanamh ar theorainneacha na gceantar comharthaíthe. Suirbhéarachta Ordonáis na hÉireann Ceadúnas Uimh EN 0059216. © Suirbhéarachta Ordonáis na hÉireann Rialtas na hÉireann.


Map Version 1
Date: Jan 2016



Legend


- Cloonchambers Bog SAC 000600
- High Bog Boundary
- Potential 7110 *Active Raised Bogs






Legend

- Cloonchambers Bog SAC 000600
- High Bog Boundary
- OSi Discovery Series County Boundary
- Active Raised Bogs Ecotopes**
- Soaks / active flush
- Sub-central ecotope







Legend

- Cloonchambers Bog SAC 000600
- High Bog Boundary
- Drainage Patterns
- Contours

Elevation



National Parks and Wildlife Service

Conservation Objectives Series

Derrinea Bog SAC 000604



*An Roinn
Ealaíon, Oidhreachta agus Gaeltachta*

*Department of
Arts, Heritage and the Gaeltacht*



**National Parks and Wildlife Service,
Department of Arts, Heritage and the Gaeltacht,**

7 Ely Place, Dublin 2, Ireland.

Web: www.npws.ie

E-mail: nature.conservation@ahg.gov.ie

Citation:

**NPWS (201) Conservation Objectives: Derrinea Bog SAC 000604. Version 1.
National Parks and Wildlife Service, Department of Arts, Heritage and the
Gaeltacht.**

Series Editor: Rebecca Jeffrey

ISSN 2009-4086

Introduction

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

** indicates a priority habitat under the Habitats Directive*

000604	Derrinea Bog SAC
7110	Active raised bogsE
7120	Degraded raised bogs still capable of natural regeneration
7150	Depressions on peat substrates of the Rhynchosporion

Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

NPWS Documents

Year :	2014
Title :	Raised Bog Monitoring and Assessment Survey 2013
Author :	Fernandez, F.; Connolly K.; Crowley W.; Denyer J.; Duff K.; Smith G.
Series :	Irish Wildlife Manual No. 81
<hr/>	
Year :	2014
Title :	National raised bog SAC management plan
Author :	Department of Arts, Heritage and the Gaeltacht
Series :	Draft for consultation. 15 January 2014
<hr/>	
Year :	2014
Title :	Derrinea Bog (SAC 000604), Co. Roscommon, Site Report
Author :	Fernandez, F.; Connolly, K.; Crowley, W.; Denyer J.; Duff K.; Smith G.
Series :	Raised bog monitoring and assessment survey 2013
<hr/>	
Year :	2015
Title :	Derrinea Bog SAC (site code: 604) Conservation objectives supporting document- raised bog habitats V1
Author :	NPWS
Series :	Conservation objectives supporting document

Other References

Year :	2011
Title :	Review and revision of empirical critical loads and dose-response relationships. Proceedings of an expert workshop, Noordwijkerhout, 23-25 June 2010
Author :	Bobbink, R.; Hettelingh, J.P.
Series :	RIVM report 680359002, Coordination Centre for Effects, National Institute for Public Health and the Environment (RIVM)
<hr/>	
Year :	2014
Title :	Nitrogen deposition and exceedance of critical loads for nutrient nitrogen in Irish grasslands
Author :	Henry, J.; Aherne, J.
Series :	Science of the Total Environment 470–471: 216–223

Spatial data sources

Year :	2014
Title :	Scientific Basis for Raised Bog Conservation in Ireland
GIS Operations :	RBSB13_SACs_ARB_DRB dataset, RBSB13_SACs_2012_HB dataset, RBSB13_SACs_DrainagePatterns_5k dataset and RBSB13_SAC_LIDAR_DTMs dataset clipped to SAC boundary. Expert opinion used as necessary to resolve any issues arising
Used For :	potential 7110; digital elevation model; drainage patterns (maps 2 and 4)
<hr/>	
Year :	2013
Title :	Raised Bog Monitoring and Assessment Survey 2013
GIS Operations :	RBMA13_ecotope_map dataset clipped to SAC boundary. Appropriate ecotopes selected and exported to new dataset. Expert opinion used as necessary to resolve any issues arising 7110 ecotopes (map 4)
Used For :	7110 ecotopes (map 3)
<hr/>	

Conservation Objectives for : Derrinea Bog SAC [000604]

7110 Active raised bogs

To restore the favourable conservation condition of Active raised bogs in Derrinea Bog SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Restore area of active raised bog to 24.7ha, subject to natural processes	Active Raised Bog (ARB) habitat was mapped at 17.1ha by Fernandez et al. (2014). Area of Degraded Raised Bog (DRB) on the High Bog (HB) has been modelled as 9.8ha. However, it has been estimated that only 6.8ha is potentially restorable to ARB by drain blocking. The total potential ARB on the HB is therefore estimated to be 23.9ha. See map 2. Eco-hydrological assessments of the cutover estimates that an additional 0.8ha of bog forming habitats could be restored. The long term target for ARB is therefore 24.7ha. See raised bog supporting document for further details on this and following attributes
Habitat distribution	Occurrence	Restore the distribution and variability of active raised bog across the SAC. See map 3 for distribution in 2012	ARB currently occurs in the central area of Derrinea Bog. DRB surrounds the ARB areas, which will require restoration measures. There is also potential for ARB restoration on cutover areas of the bog (see area target above)
High bog area	Hectares	No decline in extent of high bog necessary to support the development and maintenance of active raised bog. See map 2	The area of high bog within Derrinea Bog SAC in 2012 (latest figure available) was 53.8ha (DAHG 2014)
Hydrological regime: water levels	Centimetres	Restore appropriate water levels throughout the site	For ARB, mean water level needs to be near or above the surface of the bog lawns for most of the year. Seasonal fluctuations should not exceed 20cm, and should only be 10cm below the surface, except for very short periods of time. Open water is often characteristic of soak systems
Hydrological regime: flow patterns	Flow direction; slope	Restore, where possible, appropriate high bog topography, flow directions and slopes. See map 4 for current situation	ARB depends on mean water levels being near or above the surface of bog lawns for most of the year. Long and gentle slopes are the most favourable to achieve these conditions. Changes to flow directions due to subsidence of bogs can radically change water regimes and cause drying out of high quality ARB areas and soak systems
Transitional areas between high bog and adjacent mineral soils (including cutover areas)	Hectares; distribution	Restore adequate transitional areas to support/protect active raised bog and the services it provides	Only remnant semi-natural exist along the eastern and northern margin of Derrinea Bog. Eco-hydrological assessments have evaluated the potential for ARB restoration on cutover areas (see note for habitat area attribute above)
Vegetation quality: central ecotope, active flush, soaks, bog woodland	Hectares	Restore 12.4ha of central ecotope/active flush/soaks/bog woodland	At least 50% of ARB habitat should be central ecotope/active flush/soaks/bog woodland. Target area of active raised bog for the site has been set at 24.7ha (see area target above)
Vegetation quality: microtopographical features	Hectares	Restore adequate cover of high quality microtopographical features	High quality microtopography (hummocks, hollows and pools) is well developed in the central part of Derrinea Bog
Vegetation quality: bog moss (<i>Sphagnum</i>) species	Percentage cover	Restore adequate cover of bog moss (<i>Sphagnum</i>) species to ensure peat-forming capacity	<i>Sphagnum</i> cover varies naturally across Ireland with relatively high cover in the east to lower cover in the west. Hummock forming species such as <i>Sphagnum austinii</i> are particularly good peat formers. <i>Sphagnum</i> cover and distribution also varies naturally across a site
Typical ARB species: flora	Occurrence	Restore, where appropriate, typical active raised bog flora	Typical flora species include widespread species, as well as those with more restricted distributions but typical of the habitat's subtypes or geographical range

Typical ARB species: fauna	Occurrence	Restore, where appropriate, typical active raised bog fauna	Typical fauna species include widespread species, as well as those with more restricted distributions but typical of the habitat's subtypes or geographical range
Elements of local distinctiveness	Occurrence	Maintain features of local distinctiveness, subject to natural processes	Derrinea Bog supports three pools, two of which are larger and these are infilling with rafts of Rhynchosporion vegetation. Surrounding the pools, quaking flats have good bog moss cover
Negative physical indicators	Percentage cover	Negative physical features absent or insignificant	Negative physical indicators include: bare peat, algae dominated pools and hollows, marginal cracks, tear patterns, subsidence features such as dry mineral mounds /ridges emerging or expanding, and evidence of burning
Vegetation composition: native negative indicator species	Percentage cover	Native negative indicator species at insignificant levels	Disturbance indicators include species indicative of conditions drying out such as abundant bog asphodel (<i>Nartheicum ossifragum</i>), deergrass (<i>Trichophorum germanicum</i>) and haretail cotton-grass (<i>Eriophorum vaginatum</i>) forming tussocks; abundant magellanic bog-moss (<i>Sphagnum magellanicum</i>) in pools previously dominated by <i>Sphagnum</i> species typical of very wet conditions (e.g. feathery bog-moss (<i>S. cuspidatum</i>)); and indicators of frequent burning events such as abundant <i>Cladonia floerkeana</i> and high cover of carnation sedge (<i>Carex panicea</i>) (particularly in true midlands raised bogs)
Vegetation composition: non-native invasive species	Percentage cover	Non-native invasive species at insignificant levels and not more than 1% cover	Most common non-native invasive species include lodgepole pine (<i>Pinus contorta</i>), rhododendron (<i>Rhododendron ponticum</i>), and pitcherplant (<i>Sarracenia purpurea</i>). No non-native invasive species were recorded on Derrinea Bog following a survey in 2012 (Fernandez et al. 2014)
Air quality: nitrogen deposition	kg N/ha/year	Air quality surrounding bog close to natural reference conditions. The total N deposition should not exceed 5kg N/ha/yr	Change in air quality can result from fertiliser drift; adjacent quarry activities; or other atmospheric inputs. The critical load range for ombrotrophic bogs has been set as between 5 and 10kg N/ha/yr (Bobbink and Hettelingh 2011). The latest N deposition figures for the area around Derrinea Bog suggests that the current level is approximately 9.4kg N/ha/yr (Henry and Aherne 2014)
Water quality	Hydrochemical measures	Water quality on the high bog and in transitional areas close to natural reference conditions	Water chemistry within raised bogs is influenced by atmospheric inputs (rainwater). However, within soak systems, water chemistry is influenced by other inputs such as focused flow or interaction with underlying substrates. Water chemistry in areas surrounding the high bog varies due to influences of different water types (bog water, regional groundwater, and run-off from surrounding mineral lands)

Conservation Objectives for : Derrinea Bog SAC [000604]

7120 Degraded raised bogs still capable of natural regeneration

The long-term aim for Degraded raised bogs still capable of natural regeneration is that its peat-forming capability is re-established; therefore, the conservation objective for this habitat is inherently linked to that of Active raised bogs (7110) and a separate conservation objective has not been set in Derrinea Bog SAC


Attribute	Measure	Target	Notes
-----------	---------	--------	-------

Conservation Objectives for : Derrinea Bog SAC [000604]


7150 Depressions on peat substrates of the Rhynchosporion

Depressions on peat substrates of the Rhynchosporion is an integral part of good quality Active raised bogs (7110) and thus a separate conservation objective has not been set for the habitat in Derrinea Bog SAC

Attribute	Measure	Target	Notes
-----------	---------	--------	-------



Legend

 Derrinea Bog SAC 000604

*An Roinn
Ealaíon, Oidhreachta agus Gaeltachta
Department of
Arts, Heritage and the Gaeltacht*

**MAP 1:
DERRINEA BOG SAC
CONSERVATION OBJECTIVES
SAC DESIGNATION**

Map to be read in conjunction with the NPWS Conservation Objectives Document.

**SITE CODE:
SAC 000604; version 3.01. Co. Roscommon**

0 100 200 300 400 500 m


The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
Ordnance Survey of Ireland Licence No EN 0059214. © Ordnance Survey of Ireland Government of Ireland.

Níl sna teorainneacha ar na léarscáileanna ach nod garshuimhach ginearálta. Féadfar athbheithníthe a déanamh ar theorainneacha na gceantar comharthaíthe. Suirbhéarachta Ordonáis na hÉireann Ceadúnas Uimh EN 0059214. © Suirbhéarachta Ordonáis na hÉireann Rialtas na hÉireann.

N




**Map Version 1
Date: Oct 2015**










Legend

- Derrinea Bog SAC 000604
- High Bog Boundary
- Potential 7110 *Active Raised Bogs
- OSi Discovery Series County Boundary



Legend

-  Derrinea Bog SAC 000604
-  OSi Discovery Series County Boundary
-  High Bog Boundary
- Active Raised Bog Ecotopes**
-  Central ecotope
-  Soaks / active flush
-  Sub-central ecotope







Legend

- Derrinea Bog SAC 000604
- OSi Discovery Series County Boundary
- High Bog Boundary
- Drainage Patterns
- Countours

Elevation



National Parks and Wildlife Service

Conservation Objectives Series

Errit Lough SAC 000607



An Roinn
Cultúir, Oidhreachta agus Gaeltachta
Department of
Culture, Heritage and the Gaeltacht



**National Parks and Wildlife Service,
Department of Culture, Heritage and the Gaeltacht,
7 Ely Place, Dublin 2, Ireland.
Web: www.npws.ie
E-mail: nature.conservation@ahg.gov.ie**

Citation:

**NPWS (2017) Conservation Objectives: Errit Lough SAC 000607. Version 1.
National Parks and Wildlife Service, Department of Culture, Heritage and the
Gaeltacht.**

**Series Editor: Rebecca Jeffrey
ISSN 2009-4086**

Introduction

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

** indicates a priority habitat under the Habitats Directive*

000607 Errit Lough SAC

3140 Hard oligo-mesotrophic waters with benthic vegetation of Chara spp.

Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

NPWS Documents

Year :	1984
Title :	The vegetation of Irish lakes
Author :	Heuff, H.
Series :	Unpublished report to NPWS
Year :	2013
Title :	A survey of the benthic macrophytes of three hard-water lakes: Lough Bunny, Lough Carra and Lough Owel
Author :	Roden, C.; Murphy, P.
Series :	Irish Wildlife Manual No. 70
Year :	2013
Title :	The status of EU protected habitats and species in Ireland. Volume 2. Habitats assessments
Author :	NPWS
Series :	Conservation assessments
Year :	2015
Title :	Habitats Directive Annex I lake habitats: a working interpretation for the purposes of site-specific conservation objectives and Article 17 reporting
Author :	O Connor, Á.
Series :	Unpublished document by NPWS

Other References

Year :	1982
Title :	Eutrophication of waters. Monitoring assessment and control
Author :	OECD
Series :	OECD, Paris
Year :	2000
Title :	Colour in Irish lakes
Author :	Free, G.; Allott, N.; Mills, P.; Kennelly, C.; Day, S.
Series :	Verhandlungen Internationale Vereinigung für theoretische und angewandte Limnologie, 27: 2620-2623
Year :	2006
Title :	A reference-based typology and ecological assessment system for Irish lakes. Preliminary investigations. Final report. Project 2000-FS-1-M1 Ecological assessment of lakes pilot study to establish monitoring methodologies EU (WFD)
Author :	Free, G.; Little, R.; Tierney, D.; Donnelly, K.; Coroni, R.
Series :	EPA, Wexford
Year :	in prep.
Title :	Monitoring of hard-water lakes in Ireland using charophytes and other macrophytes
Author :	Roden, C.; Murphy, P.
Series :	Unpublished report to NPWS

Spatial data sources

Year : 2008

Title : OSi 1:5000 IG vector dataset

GIS Operations : WaterPolygons feature class clipped to the SAC boundary. Expert opinion used to identify Annex I habitat and to resolve any issues arising

Used For : 3140 (map 2)

Conservation Objectives for : Errit Lough SAC [000607]

3140 Hard oligo-mesotrophic waters with benthic vegetation of *Chara* spp.


To maintain the favourable conservation condition of Hard oligo-mesotrophic waters with benthic vegetation of *Chara* spp. in Errit Lough SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Errit Lough is a marl lake containing lake habitat 3140. Its vegetation was surveyed in August 1978 (Heuff, 1984) and again in 2012 (Roden and Murphy, in prep.). Two measures of extent should be used: 1. the area of the lake itself and; 2. the extent of the vegetation communities/zones that typify the habitat. Further information relating to all attributes is provided in the lake habitats supporting document for the purposes of site-specific conservation objectives and Article 17 reporting (O Connor, 2015)
Habitat distribution	Occurrence	No decline, subject to natural processes	As noted above, lake habitat 3140 occurs in Errit Lough in the SAC. See map 2
Vegetation composition: typical species	Occurrence	Typical species present, in good condition, and demonstrating typical abundances and distribution	For lists of 3140 typical species (cyanobacteria, algae, higher plants and water beetles), see the Article 17 habitat assessment for lake habitat 3140 (NPWS, 2013) and the lake habitats supporting document (O Connor, 2015). Heuff (1984) includes general descriptions of the vegetation of Errit Lough and data for three relevés. <i>Chara</i> spp. dominate and vascular species include <i>Littorella uniflora</i> , <i>Nuphar lutea</i> and <i>Potamogeton perfoliatus</i> (Heuff, 1984). In 2012, Roden and Murphy (in prep.) recorded the following species in Errit Lough: <i>Chara aculeolata</i> , <i>C. contraria</i> , <i>C. curta</i> , <i>C. rudis</i> , <i>C. virgata</i> , <i>Elodea canadensis</i> , <i>Equisetum fluviatile</i> , <i>Littorella uniflora</i> , <i>Nuphar lutea</i> , <i>Potamogeton gramineus</i> , <i>P. praelongus</i> , <i>P. zizii</i> , <i>Schoenoplectus lacustris</i> and <i>Utricularia</i> sp.
Vegetation composition: characteristic zonation	Occurrence	All characteristic zones should be present, correctly distributed and in good condition	The characteristic zonation of 3140 is described in Roden and Murphy (2013). Heuff (1984) recorded the zonation at Errit Lough: sparse reedbeds on the sheltered south-western shore, a <i>Chara aculeolata</i> zone to 1m water depth, with <i>C. rudis</i> at the southern end of the lake, a <i>C. curta</i> -dominated zone with some <i>Littorella uniflora</i> from 1-2m, a <i>C. globularis</i> zone from 2-3m followed by a sparse band of <i>Nuphar lutea</i> , and <i>Potamogeton perfoliatus</i> at 4m. Roden and Murphy (in prep.) stated charophyte zones were not well-developed in Errit in 2012, with only <i>C. aculeolata</i> and <i>C. rudis</i> forming substantial beds and no charophytes below 3m. Krustenstein was well-developed on rocks in shallow water. <i>Littorella uniflora</i> and <i>Scorpidium</i> sp. were recorded at some shallow-water stations, rather than the typical charophytes (Roden and Murphy, in prep.). The condition of Errit may have deteriorated 1978 to 2012

Vegetation distribution: maximum depth	Metres	Maintain/restore maximum depth of vegetation, subject to natural processes	Maximum vegetation depth is expected to be deep in clear, hard water lakes, and extremely clear marl lakes can have charophyte vegetation to more than 9m (e.g. Lough Rea has charophytes to 10-11m, Coolorta >9m) (Roden and Murphy, in prep.). The indicative target of >6m for lake habitat 3140 may need to be modified based on the habitat sub-type/form and/or the specific lake in question (Roden and Murphy, 2013, in prep.). In this SAC, the maximum depth of vegetation at Errit Lough was 4m in 1978 (Heuff, 1984) and 3-3.5m in 2012 (Roden and Murphy, in prep.). The water is highly coloured in Errit Lough, and this appears to limit vegetation development in deeper water. The areas of drained peatland (for turf-cutting and conifer forest) in the catchment of Errit Lough are likely to artificially increase the lake's water colour
Hydrological regime: water level fluctuations	Metres	Maintain appropriate hydrological regime necessary to support the habitat	The hydrological regime of lakes with habitat 3140 is driven by groundwater flows. Groundwater can discharge directly to the lake, via springs or seepages, or to in-flowing rivers. Fluctuations in lake water level are typical in Ireland, but can be amplified by activities such as abstraction and drainage. Increased water level fluctuations can increase wave action and turbidity, up-root vegetation, alter the substratum and lead to nutrient release from sediment. The hydrological regime, particularly the groundwater contribution, must be maintained so that the area, distribution and depth of the lake habitat and its constituent/characteristic vegetation zones and communities are not reduced
Lake substratum quality	Various	Maintain appropriate substratum type, extent and chemistry to support the vegetation	The hard water lake habitat (3140) is associated with a range of base-rich substratum types, from marl and limestone bedrock, through rocks, cobbles, gravel, muds and even peat. Further research into substratum quality (notably calcium, iron and nutrient concentrations) in the hard water lake habitat would be beneficial. The exposed eastern shore of Errit Lough is stony/rocky, while much of the lake has sandy marl (Heuff, 1984). Roden and Murphy (in prep.) also recorded peat
Water quality: transparency	Metres	Maintain/restore appropriate Secchi transparency. There should be no decline in Secchi depth/transparency	Transparency relates to light penetration and, hence, to the depth of colonisation of vegetation. It can be affected by phytoplankton blooms, water colour and turbidity. A target of >6m has been set for hard water lakes (3140) (Roden and Murphy, in prep.). The OECD fixed boundary system set transparency targets for oligotrophic lakes of $\geq 6\text{m}$ annual mean Secchi disk depth and $\geq 3\text{m}$ annual minimum Secchi disk depth. Hard water lakes typically have high transparency, particularly in the very clear and typical marl forms (Roden and Murphy, in prep.). Secchi depth at Errit Lough was 2.7m in 2001 (Free et al., 2006) and 3m in 2012 (Roden and Murphy, in prep.)
Water quality: nutrients	$\mu\text{g/l P}$; mg/l N	Maintain the concentration of nutrients in the water column at sufficiently low levels to support the habitat and its typical species	Lake habitat 3140 is typically associated with high water quality, as demonstrated by naturally low dissolved nutrients. The target for Errit Lough is Water Framework Directive (WFD) High Status or oligotrophic (OECD, 1982). Annual average total phosphorus (TP) concentration should be $\leq 10\mu\text{g/l}$ TP, average annual total ammonia concentration should be $\leq 0.04\text{mg/l N}$ and annual 95th percentile for total ammonia should be $\leq 0.09\text{mg/l N}$. Where nutrient concentrations are lower than the targets, there should be no upward trend in concentrations. See also The European Communities Environmental Objectives (Surface Waters) Regulations 2009. Free et al. (2006) measured $< 10\mu\text{g/l}$ TP in Errit Lough in April 2001


Water quality: phytoplankton biomass	µg/l Chlorophyll <i>a</i>	Maintain appropriate water quality to support the habitat, including high chlorophyll <i>a</i> status	Lake habitat 3140 is associated with high water quality, as demonstrated by naturally low algal growth. As for nutrients, the default target is WFD High Status or oligotrophic (OECD, 1982). Average growing season (March-October) chlorophyll <i>a</i> concentration must be <5.8µg/l. Annual average chlorophyll <i>a</i> concentration should be <2.5µg/l and the annual peak should be <8.0µg/l. Where chlorophyll <i>a</i> concentrations are lower than the targets, there should be no upward trend in phytoplankton biomass. See also The European Communities Environmental Objectives (Surface Waters) Regulations 2009. Free et al. (2006) measured 6.3µg/l chlorophyll <i>a</i> in Errit Lough in April 2001
Water quality: phytoplankton composition	EPA phytoplankton composition metric	Maintain appropriate water quality to support the habitat, including high phytoplankton composition status	The Environmental Protection Agency (EPA) has developed a phytoplankton composition metric for nutrient enrichment of Irish lakes. As for other water quality indicators, the default target for lake habitat 3140 is WFD high status
Water quality: attached algal biomass	Algal cover and EPA phytobenthos metric	Maintain trace/absent attached algal biomass (<5% cover) and high phytobenthos status	Nutrient enrichment can favour epiphytic and epipelic algae that can out-compete the submerged vegetation. The cover abundance of attached algae in hard water lakes (3140) should, therefore, be trace/absent (<5% cover). EPA phytobenthos status can be used as an indicator of changes in attached algal biomass. As for other water quality indicators, the default target for lake habitat 3140 is high phytobenthos status
Water quality: macrophyte status	EPA macrophyte metric (The Free Index)	Maintain high macrophyte status	Nutrient enrichment can favour more competitive submerged macrophyte species that out-compete the typical and characteristic species for hard water lakes (3140). The EPA monitors macrophyte status for WFD purposes using the 'Free Index'. The target for lake habitat 3140 is high status or an Ecological Quality Ratio (EQR) for lake macrophytes of ≥0.90, as defined in Schedule Five of the European Communities Environmental Objectives (Surface Waters) Regulations 2009
Acidification status	pH units; mg/l	Maintain appropriate water and sediment pH, alkalinity and cation concentrations to support the habitat, subject to natural processes	The specific requirements of lake habitat 3140, in terms of water and sediment pH, alkalinity and cation concentration, have not been fully determined. Acidification is not considered a threat to lake habitat 3140; however, eutrophication can lead to at least temporary increases in pH to toxic levels (>9/9.5 pH units). Maximum pH should be <9.0 pH units, in line with the surface water standards. See Schedule Five of the European Communities Environmental Objectives (Surface Waters) Regulations 2009
Water colour	mg/l PtCo	Restore appropriate water colour to support the habitat	Increased colour decreases light penetration and reduces the area of macrophyte habitat, particularly at the lower euphotic depths. Higher colour also appears to favour angiosperms over charophytes in hard water lakes (Roden and Murphy, in prep.). The primary source of increased colour in Ireland is peatland disturbance. No habitat-specific or national standards for water colour exist. Studies have shown median colour concentrations in Irish lakes of 38mg/l PtCo (Free et al., 2000) and 33mg/l PtCo (Free et al., 2006). Lake habitat 3140 is typically associated with very clear waters and expected colour would be <10mg/l PtCo or, more likely, <5mg/l PtCo. Free et al. (2006) recorded colour of 46mg/l PtCo in Errit Lough and Roden and Murphy (in prep.) noted high water colour in 2012, a very shallow euphotic zone, low Secchi depth and stated peat-staining from disturbed bogland may have reduced water transparency

Dissolved organic carbon (DOC)	mg/l	Maintain/restore appropriate organic carbon levels to support the habitat	Dissolved (and particulate) organic carbon (OC) in the water column is linked to water colour and acidification (organic acids). Increasing DOC in water has been documented across the Northern Hemisphere, including afforested peatland catchments in Ireland. Damage and degradation of peatland, leading to decomposition of peat is likely to be the predominant source of OC in Ireland. OC in water promotes decomposition by fungi and bacteria that, in turn, releases dissolved nutrients. The increased biomass of decomposers can also impact directly on the characteristic lake communities through shading, competition, etc. As noted above, increased water colour, low transparency and shallow vegetation zones at Errit Lough appear to be linked to peatland disturbance
Turbidity	Nephelometric turbidity units/ mg/l SS/ other appropriate unit	Maintain appropriate turbidity to support the habitat	Turbidity can significantly affect the quantity and quality of light reaching rooted and attached vegetation and can, therefore, impact on lake habitats. The settlement of higher loads of inorganic or organic material on lake vegetation communities may also have impacts on sensitive, delicate species. Turbidity can increase as a result of re-suspension of material within the lake, higher loads entering the lake, or eutrophication. Turbidity measurement and interpretation is challenging. As a result, it is likely to be difficult to set habitat-specific targets for turbidity in lakes
Fringing habitat: area and condition	Hectares	Maintain the area and condition of fringing habitats necessary to support the natural structure and functioning of habitat 3140	Most lake shorelines have fringing habitats of reedswamp, other swamp, fen, marsh or wet woodland that intergrade with and support the structure and functions of the lake habitat. Equally, fringing habitats are dependent on the lake, particularly its water levels, and support wetland communities and species of conservation concern. Many of the fringing wetland habitats support higher invertebrate and plant species richness than the lake habitats themselves. Fringing fen habitats can be particularly important around hard water lakes, notably the Annex I habitats alkaline fen, <i>Cladium</i> fen and petrifying springs (habitat codes 7230, 7210 and 7220 respectively). Sparse reedbeds (<i>Schoenoplectus lacustris</i> , <i>Phragmites australis</i>) and marginal swamp (<i>Carex rostrata</i> , <i>Eleocharis palustris</i>) occur on the south-west shore of Errit Lough (Heuff, 1984). Deciduous woodland, heath and wet grassland also fringe Errit Lough



Legend

 Errit Lough SAC 000607



An Roinn
Cultúir, Oidhreacht agus Gaeltachta
Department of
Culture, Heritage and the Gaeltacht

**MAP 1:
ERRIT LOUGH SAC
CONSERVATION OBJECTIVES
SAC DESIGNATION**

Map to be read in conjunction with the NPWS Conservation Objectives Document.

**SITE CODE:
SAC 000607 ; version 3. CO. ROSCOMMON**

0 100 200 300 400 Meters


The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
Ordnance Survey of Ireland Licence No EN 0059216. © Ordnance Survey of Ireland Government of Ireland

Níl sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbhreithnithe a déanamh ar theorainneacha na gceantar comharthaite. Suirbhéarachtá Ordonáis na hÉireann Ceadúnas Uimh EN 0059216. © Suirbhéarachtá Ordonáis na hÉireann Rialtas na hÉireann

N




**Map Version 1
Date: Nov 2017**



Legend

- 3140 Hard oligo-mesotrophic waters with benthic vegetation of Chara spp.
- Errit Lough SAC 000607
- OSi Discovery Series County Boundary




An Roinn
Cultúir, Oidhreacht agus Gaeltachta
Department of
Culture, Heritage and the Gaeltacht

**MAP 2:
ERRIT LOUGH SAC
CONSERVATION OBJECTIVES
INDICATIVE LAKE HABITATS**

Map to be read in conjunction with the NPWS Conservation Objectives Document.


**SITE CODE:
SAC 000607; version 3.
CO. ROSCOMMON**

0 125 250 375 500 Meters



The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
Ordnance Survey of Ireland Licence No EN 0059216. © Ordnance Survey of Ireland Government of Ireland

Níl sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbheithnithe a déanamh ar theorainneacha na gceantar comharthaíthe. Suirbhéarachta Ordonáis na hÉireann Ceadúnas Uimh EN 0059216. © Suirbhéarachta Ordonáis na hÉireann Rialtas na hÉireann



**Map Version 1
Date: Nov 2017**

National Parks and Wildlife Service

Conservation Objectives Series

Cloonshanville Bog SAC 000614



*An Roinn
Ealaíon, Oidhreachta agus Gaeltachta*

*Department of
Arts, Heritage and the Gaeltacht*



**National Parks and Wildlife Service,
Department of Arts, Heritage and the Gaeltacht,**

7 Ely Place, Dublin 2, Ireland.

Web: www.npws.ie

E-mail: nature.conservation@ahg.gov.ie

Citation:

**NPWS (2016) Conservation Objectives: Cloonshanville Bog SAC 000614. Version
1. National Parks and Wildlife Service, Department of Arts, Heritage and the
Gaeltacht.**

Series Editor: Rebecca Jeffrey

ISSN 2009-4086

Introduction

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

** indicates a priority habitat under the Habitats Directive*

000614	Cloonshanville Bog SAC
7110	Active raised bogs*
7120	Degraded raised bogs still capable of natural regeneration
7150	Depressions on peat substrates of the Rhynchosporion
91D0	Bog woodland*

Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

NPWS Documents

Year :	2014
Title :	Raised Bog Monitoring and Assessment Survey 2013
Author :	Fernandez, F.; Connolly K.; Crowley W.; Denyer J.; Duff K.; Smith G.
Series :	Irish Wildlife Manual No. 81
<hr/>	
Year :	2014
Title :	National raised bog SAC management plan
Author :	Department of Arts, Heritage and the Gaeltacht
Series :	Draft for consultation. 15 January 2014
<hr/>	
Year :	2014
Title :	Cloonshanville Bog (SAC 000614), Co. Roscommon, Site Report
Author :	Fernandez, F.; Connolly, K.; Crowley, W.; Denyer J.; Duff K.; Smith G.
Series :	Raised bog monitoring and assessment survey 2013
<hr/>	
Year :	2016
Title :	Cloonshanville Bog SAC (000614) Conservation objectives supporting document- raised bog habitats V1
Author :	NPWS
Series :	Conservation objectives supporting document

Other References

Year :	2011
Title :	Review and revision of empirical critical loads and dose-response relationships. Proceedings of an expert workshop, Noordwijkerhout, 23-25 June 2010
Author :	Bobbink, R.; Hettelingh, J.P.
Series :	RIVM report 680359002, Coordination Centre for Effects, National Institute for Public Health and the Environment (RIVM)
<hr/>	
Year :	2014
Title :	Nitrogen deposition and exceedance of critical loads for nutrient nitrogen in Irish grasslands
Author :	Henry, J.; Aherne, J.
Series :	Science of the Total Environment 470–471: 216–223

Spatial data sources

Year :	2014
Title :	Scientific Basis for Raised Bog Conservation in Ireland
GIS Operations :	RBSB13_SACs_ARB_DRB dataset, RBSB13_SACs_2012_HB dataset, RBSB13_SACs_DrainagePatterns_5k dataset and RBSB13_SAC_LIDAR_DTMs dataset clipped to SAC boundary. Expert opinion used as necessary to resolve any issues arising
Used For :	potential 7110; digital elevation model; drainage patterns (maps 2 and 4)
<hr/>	
Year :	2013
Title :	Raised Bog Monitoring and Assessment Survey 2013
GIS Operations :	RBMA13_ecotope_map dataset clipped to SAC boundary. Appropriate ecotopes selected and exported to new dataset. Expert opinion used as necessary to resolve any issues arising
Used For :	7110 ecotopes (map 3)
<hr/>	

Conservation Objectives for : Cloonshanville Bog SAC [000614]

7110 Active raised bogs

To restore the favourable conservation condition of Active raised bogs in Cloonshanville Bog SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Restore area of active raised bog to 44.0ha, subject to natural processes	Active Raised Bog (ARB) habitat was mapped at 20.1ha in 2012 by Fernandez et al. (2014). Area of Degraded Raised Bog (DRB) on the High Bog (HB) has been modelled as 45.2ha. See map 2. However, it is estimated that only 22.6ha is potentially restorable to ARB by drain blocking. The total potential ARB on the HB is therefore estimated to be 42.7ha. Eco-hydrological assessments of the cutover estimates that an additional 1.3ha of bog forming habitats could be restored. The long term target for ARB is therefore 44.0ha. See raised bog supporting document for further details on this and following attributes
Habitat distribution	Occurrence	Restore the distribution and variability of active raised bog across the SAC. See map 3 for distribution in 2012	ARB, including bog woodland, covers 20.1 ha (13.7%) of the high bog area. A small area of central ecotope is found in the north-west corner of the bog. Sub-central ecotope of somewhat lower quality is more widespread across the bog. The most notable actively peat-forming feature of the site is an extensive active flush that bisects the high bog, this area features two areas of bog woodland
High bog area	Hectares	No decline in extent of high bog necessary to support the development and maintenance of active raised bog. See map 2	The area of high bog within Cloonshanville Bog SAC in 2012 (latest figure available) was 146.4ha (DAHG 2014)
Hydrological regime: water levels	Centimetres	Restore appropriate water levels throughout the site	For ARB, mean water level needs to be near or above the surface of the bog lawns for most of the year. Seasonal fluctuations should not exceed 20cm, and should only be 10cm below the surface, except for very short periods of time. Open water is often characteristic of soak systems
Hydrological regime: flow patterns	Flow direction; slope	Restore, where possible, appropriate high bog topography, flow directions and slopes. See map 4 for current situation	ARB depends on mean water levels being near or above the surface of bog lawns for most of the year. Long and gentle slopes are the most favourable to achieve these conditions. Changes to flow directions due to subsidence of bogs can radically change water regimes and cause drying out of high quality ARB areas and soak systems
Transitional areas between high bog and adjacent mineral soils (including cutover areas)	Hectares; distribution	Restore adequate transitional areas to support/protect active raised bog and the services it provides	ARB is threatened due to effects of past drainage and peat-cutting around margins of Cloonshanville Bog. An interesting area of fen occurs in a former lake basin to the east of the high bog
Vegetation quality: central ecotope, active flush, soaks, bog woodland	Hectares	Restore 22.0ha of central ecotope/active flush/soaks/bog woodland as appropriate	At least 50% of ARB habitat should be high quality (i.e. central ecotope, active flush, soaks, bog woodland). Target area of active raised bog for the site has been set at 44.0ha (see area target above)
Vegetation quality: microtopographical features	Hectares	Restore adequate cover of high quality microtopographical features	High quality microtopography (hummocks, hollows and pools) is well developed on Cloonshanville Bog
Vegetation quality: bog moss (<i>Sphagnum</i>) species	Percentage cover	Restore adequate cover of bog moss (<i>Sphagnum</i>) species to ensure peat-forming capacity	<i>Sphagnum</i> cover varies naturally across Ireland with relatively high cover in the east to lower cover in the west. Hummock forming species such as <i>Sphagnum austinii</i> are particularly good peat formers. <i>Sphagnum</i> cover and distribution also varies naturally across a site

Typical ARB species: flora	Occurrence	Restore, where appropriate, typical active raised bog flora	Typical flora species include widespread species, as well as those with more restricted distributions but typical of the habitat's subtypes or geographical range
Typical ARB species: fauna	Occurrence	Restore, where appropriate, typical active raised bog fauna	Typical fauna species include widespread species, as well as those with more restricted distributions but typical of the habitat's subtypes or geographical range
Elements of local distinctiveness	Occurrence	Maintain features of local distinctiveness, subject to natural processes	The large flush and associated bog woodland are features of local distinctiveness on Cloonshanville Bog
Negative physical indicators	Percentage cover	Negative physical features absent or insignificant	Negative physical indicators include: bare peat, algae dominated pools and hollows, marginal cracks, tear patterns, subsidence features such as dry mineral mounds /ridges emerging or expanding and evidence of burning
Vegetation composition: native negative indicator species	Percentage cover	Native negative indicator species at insignificant levels	Disturbance indicators include species indicative of conditions drying out such as abundant bog asphodel (<i>Narthecium ossifragum</i>), deergrass (<i>Trichophorum germanicum</i>) and harestail cotton-grass (<i>Eriophorum vaginatum</i>) forming tussocks; abundant magellanic bog-moss (<i>Sphagnum magellanicum</i>) in pools previously dominated by <i>Sphagnum</i> species typical of very wet conditions (e.g. feathery bog-moss (<i>S. cuspidatum</i>)); and indicators of frequent burning events such as abundant <i>Cladonia floerkeana</i> and high cover of carnation sedge (<i>Carex panicea</i>) (particularly in true midlands raised bogs)
Vegetation composition: non-native invasive species	Percentage cover	Non-native invasive species at insignificant levels and not more than 1% cover	Most common non-native invasive species include lodgepole pine (<i>Pinus contorta</i>), rhododendron (<i>Rhododendron ponticum</i>), and pitcherplant (<i>Sarracenia purpurea</i>)
Air quality: nitrogen deposition	kg N/ha/year	Air quality surrounding bog close to natural reference conditions. The total N deposition should not exceed 5kg N/ha/yr	Change in air quality can result from fertiliser drift; adjacent quarry activities; or other atmospheric inputs. The critical load range for ombrotrophic bogs has been set as between 5 and 10kg N/ha/yr (Bobbink and Hettelingh, 2011). The latest N deposition figures for the area around Cloonshanville Bog suggests that the current level is approximately 9.6kg N/ha/yr (Henry and Aherne, 2014)
Water quality	Hydrochemical measures	Water quality on the high bog and in transitional areas close to natural reference conditions	Water chemistry within raised bogs is influenced by atmospheric inputs (rainwater). However, within soak systems, water chemistry is influenced by other inputs such as focused flow or interaction with underlying substrates. Water chemistry in areas surrounding the high bog varies due to influences of different water types (bog water, regional groundwater and run-off from surrounding mineral lands)

Conservation Objectives for : Cloonshanville Bog SAC [000614]

7120 Degraded raised bogs still capable of natural regeneration

The long-term aim for Degraded raised bogs still capable of natural regeneration is that its peat-forming capability is re-established; therefore, the conservation objective for this habitat is inherently linked to that of Active raised bogs (7110) and a separate conservation objective has not been set in Cloonshanville Bog SAC

Attribute	Measure	Target	Notes
------------------	----------------	---------------	--------------

Conservation Objectives for : Cloonshanville Bog SAC [000614]

7150 Depressions on peat substrates of the Rhynchosporion

Depressions on peat substrates of the Rhynchosporion is an integral part of good quality Active raised bogs (7110) and thus a separate conservation objective has not been set for the habitat in Cloonshanville Bog SAC



Attribute	Measure	Target	Notes
-----------	---------	--------	-------


Conservation Objectives for : Cloonshanville Bog SAC [000614]

91D0 Bog woodland

To maintain the favourable conservation condition of Bog woodland in Cloonshanville Bog SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes. At least 2.2ha. See map 4	Bog woodland occurs on Cloonshanville Bog and is regarded as a component of the Active Raised Bog (ARB) habitat (7110) of that bog. Thus, the conservation objective and supporting document for ARB (7110) are also relevant to this habitat and common attributes have not been repeated here. The latest survey for bog woodland in Cloonshanville Bog SAC is reported in Fernandez et al. (2014)
Habitat distribution	Occurrence	No decline, subject to natural processes. See map 4	Bog woodland occurs in extensive active flush that occurs at the centre of the high bog at Cloonshanville
Vegetation composition: positive indicator species	Number in a representative number of monitoring stops	Birch (<i>Betula pubescens</i>), bog moss (<i>Sphagnum</i> species) and at least five other species present	Bog woodland is typically species-poor but with a characteristic and distinctive flora. Positive indicator species are listed in bog woodland monitoring survey (Cross and Lynn, 2013)
Vegetation composition: negative indicator species	Percentage cover at a representative number of monitoring stops	Both native and non-native invasive species absent or under control. Total cover should be less than 10%	Negative indicator species include bracken (<i>Pteridium aquilinum</i>) and bramble (<i>Rubus fruticosus</i>), which can become invasive if the site begins drying out
Woodland structure: cover and height of birch	Percentage cover and metres at a representative number of monitoring stops	A minimum 30% cover of birch (<i>Betula pubescens</i>) with a median canopy height of 4m	Attribute and target based on Cross and Lynn (2013)
Woodland structure: dwarf shrub cover	Percentage cover at a representative number of monitoring stops	Dwarf shrub cover not more than 50%	Attribute and target based on Cross and Lynn (2013)
Woodland structure: ling cover	Percentage cover at a representative number of monitoring stops	Ling (<i>Calluna vulgaris</i>) cover not more than 40%	Attribute and target based on Cross and Lynn (2013)
Woodland structure: bryophyte cover	Percentage cover at a representative number of monitoring stops	Bryophyte cover at least 50%, with bog moss (<i>Sphagnum</i> spp.) cover at least 25%	Attribute and target based on Cross and Lynn (2013)
Woodland structure: tree size classes	Occurrence	Each size class present	Size classes are defined in Cross and Lynn (2013). The presence of all size classes suggests that a woodland has good structural variety with trees of varying ages
Woodland structure: senescent and dead wood	Occurrence	Senescent or dead wood present	Mature and veteran trees and dead wood are important for bryophytes, lichens, saproxylic organisms and some bird species. Their retention within a woodland is important to ensure continuity of habitats/niches and propagule sources over time. However, as birch (<i>Betula pubescens</i>) trees seldom exceed 30cm in diameter in this habitat and dead wood rots quickly and is engulfed by bog mosses (<i>Sphagnum</i> spp.), volume of dead wood may not be as high in bog woodland as in other woodland types



Legend
 Cloonshanville Bog SAC 000614

An Roinn Ealaíon, Oidhreacht agus Gaeltachta
 Department of Arts, Heritage and the Gaeltacht

MAP 1:
CLOONSHANVILLE BOG SAC
CONSERVATION OBJECTIVES
SAC DESIGNATION
 Map to be read in conjunction with the NPWS Conservation Objectives Document.


SITE CODE:
SAC 000614; version 3.01. Co. Roscommon

0 100 200 300 400 500 m

The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
 Ordnance Survey of Ireland Licence No EN 0059216. © Ordnance Survey of Ireland Government of Ireland.


Níl sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbheithníthe a déanamh ar theorainneacha na gceantar comharthaíthe. Suirbhéarachtá Ordonáis na hÉireann Ceadúnas Uimh EN 0059216. © Suirbhéarachtá Ordonáis na hÉireann Rialtas na hÉireann.


Map Version 1
Date: Jan 2016



Legend


- Cloonshanville Bog SAC 000614
- High Bog Boundary
- Potential 7110 *Active Raised Bogs






Legend

- Cloonshanville Bog SAC 000614
- High Bog Boundary
- Active Raised Bogs Ecotopes**
- Bog Woodland
- Central ecotope
- Soaks / active flush
- Sub-central ecotope







Legend

- Cloonshanville Bog SAC 000614
- High Bog Boundary
- Drainage Patterns
- Contours

Elevation



National Parks and Wildlife Service

Conservation Objectives Series

Lough Hoe Bog SAC 000633



An Roinn
Cultúir, Oidhreachta agus Gaeltachta
Department of
Culture, Heritage and the Gaeltacht



**National Parks and Wildlife Service,
Department of Culture, Heritage and the Gaeltacht,
7 Ely Place, Dublin 2, Ireland.
Web: www.npws.ie
E-mail: nature.conservation@ahg.gov.ie**

Citation:

**NPWS (2017) Conservation Objectives: Lough Hoe Bog SAC 000633. Version 1.
National Parks and Wildlife Service, Department of Culture, Heritage and the
Gaeltacht.**

**Series Editor: Rebecca Jeffrey
ISSN 2009-4086**

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

** indicates a priority habitat under the Habitats Directive*

000633	Lough Hoe Bog SAC
1013	Geyer's Whorl Snail <i>Vertigo geyeri</i>
1092	White-clawed Crayfish <i>Austropotamobius pallipes</i>
3110	Oligotrophic waters containing very few minerals of sandy plains (<i>Littorelletalia uniflorae</i>)
7130	Blanket bogs (* if active bog)

Please note that this SAC is adjacent to River Moy SAC (002298). See map 2. The conservation objectives for this site should be used in conjunction with those for the adjacent site as appropriate.

Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

NPWS Documents

Year :	1989
Title :	A survey to locate blanket bogs of scientific interest in County Kerry and County Sligo
Author :	Douglas, C.; Garvey, L.; Kelly, L.; O'Sullivan, A.
Series :	Unpublished report to NPWS
Year :	1998
Title :	Conservation management of the white-clawed crayfish, <i>Austropotamobius pallipes</i>
Author :	Reynolds, J.D.
Series :	Irish Wildlife Manual No. 1
Year :	2005
Title :	Conservation Plan for 2005-2010. Lough Hoe Bog cSAC Site Code 000633 Cos. Sligo and Mayo
Author :	NPWS
Series :	Conservation Plan
Year :	2010
Title :	A technical manual for monitoring white-clawed crayfish (<i>Austropotamobius pallipes</i>) in Irish lakes
Author :	Reynolds, J., O'Connor, W., O'Keeffe, C.; Lynn, D.
Series :	Irish Wildlife Manual No.45
Year :	2011
Title :	Monitoring and condition assessment of populations of <i>Vertigo geyeri</i> , <i>Vertigo angustior</i> and <i>Vertigo moulinsiana</i> in Ireland
Author :	Moorkens, E.; Killeen, I.
Series :	Irish Wildlife Manual No. 55
Year :	2012
Title :	Ireland Red List No. 8: Bryophytes
Author :	Lockhart, N.; Hodgetts, N.; Holyoak, D.
Series :	Ireland Red List series, NPWS
Year :	2013
Title :	The status of EU protected habitats and species in Ireland. Volume 2. Habitats assessments
Author :	NPWS
Series :	Conservation assessments
Year :	2014
Title :	Guidelines for a national survey and conservation assessment of upland vegetation and habitats in Ireland, Version 2.0
Author :	Perrin, P.M.; Barron, S.J.; Roche, J.R.; O'Hanrahan, B.
Series :	Irish Wildlife Manual No. 79
Year :	2015
Title :	Habitats Directive Annex I lake habitats: a working interpretation for the purposes of site-specific conservation objectives and Article 17 reporting
Author :	O Connor, Á.
Series :	Unpublished document by NPWS
Year :	2016
Title :	Ireland Red List No. 10: Vascular Plants
Author :	Wyse Jackson, M.; FitzPatrick, Ú.; Cole, E.; Jebb, M.; McFerran, D.; Sheehy Skeffington, M.; Wright, M.
Series :	Ireland Red Lists series, NPWS

Year : 2017
Title : Lough Hoe Bog SAC (site code: 633) Conservation objectives supporting document- blanket bogs and associated habitats V1
Author : NPWS
Series : Conservation objectives supporting document

Other References

Year : 1982
Title : Eutrophication of waters. Monitoring assessment and control
Author : OECD
Series : OECD, Paris

Year : 1996
Title : Notes on some non-marine Mollusca from Co. Sligo and Co. Leitrim, including a new site for *Vertigo geyeri* Lindholm
Author : Cawley, M.
Series : Irish Naturalists' Journal, 25: 183-185

Year : 2000
Title : Colour in Irish lakes
Author : Free, G.; Allott, N.; Mills, P.; Kennelly, C.; Day, S.
Series : Verhandlungen Internationale Vereinigung für theoretische und angewandte Limnologie, 27: 2620-2623

Year : 2002
Title : A survey of the white-clawed crayfish (*Austropotamobius pallipes*) Lereboullet and of water quality in two catchments of eastern Ireland
Author : Demers, A.; Reynolds, J.D.
Series : Bulletin Francais de la Peche et de la Pisciculture, 367: 729-740

Year : 2002
Title : Deterioration of Atlantic soft water macrophyte communities by acidification, eutrophication and alkalinisation
Author : Arts, G.H.P.
Series : Aquatic Botany, 73: 373-393

Year : 2006
Title : A reference-based typology and ecological assessment system for Irish lakes. Preliminary investigations. Final report. Project 2000-FS-1-M1 Ecological assessment of lakes pilot study to establish monitoring methodologies EU (WFD)
Author : Free, G.; Little, R.; Tierney, D.; Donnelly, K.; Coroni, R.
Series : EPA, Wexford

Year : 2008
Title : Water Quality in Ireland 2004-2006
Author : Clabby, K.J.; Bradley, C.; Craig, M.; Daly, D.; Lucey, J.; McGarrigle, M.; O'Boyle, S.; Tierney, D.; Bowman, J.
Series : EPA, Wexford

Year : 2009
Title : The identification, characterization and conservation value of isoetid lakes in Ireland
Author : Free, G.; Bowman, J.; McGarrigle, M.; Little, R.; Coroni, R.; Donnelly, K.; Tierney, D.; Trodd, W.
Series : Aquatic Conservation: Marine and Freshwater Ecosystems, 19(3): 264-273

Year : 2010
Title : Water quality in Ireland 2007-2009
Author : McGarrigle, M.; Lucey, J.; Ó Cinnéide, M.
Series : EPA, Wexford

Year : 2015
Title : Water quality in Ireland 2010-2012
Author : Bradley, C.; Byrne, C.; Craig, M.; Free, G.; Gallagher, T.; Kennedy, B.; Little, R.; Lucey, J.; Mannix, A.; McCreesh, P.; McDermott, G.; McGarrigle, M.; Ní Longphuirt, S.; O'Boyle, S.; Plant, C.; Tierney, D.; Trodd, W.; Webster, P.; Wilkes, R.; Wynne, C.
Series : EPA, Wexford

Spatial data sources

Year : 2008
Title : OSi 1:5000 IG vector dataset
GIS Operations : WaterPolygons feature class clipped to the SAC boundary. Expert opinion used to identify Annex I habitat and to resolve any issues arising
Used For : 3110 (map 3)

Year : 2017
Title : NPWS rare and threatened species database
GIS Operations : Dataset created from spatial references in database records. Expert opinion used as necessary to resolve any issues arising
Used For : 1013, 1092 (maps 4 and 5)

Conservation Objectives for : Lough Hoe Bog SAC [000633]

3110 Oligotrophic waters containing very few minerals of sandy plains (Littorelletalia uniflorae)

To maintain the favourable conservation condition of Oligotrophic waters containing very few minerals of sandy plains (Littorelletalia uniflorae) in Lough Hoe Bog SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Lake habitat 3110 is likely to occur in larger upland lakes in the SAC, such as Loughs Hoe, Alone, Fosseá and Nalackagh. The exact distribution of 3110 is unknown, as no specific information on lake vegetation is available. Habitat 3110 may co-occur with lake habitat 3160 in upland lakes, which is also likely to occur in smaller lakes and ponds. There are also calcareous influences- Lough Talt has marginal calcareous springs and may be dominated by lake habitat 3140 (nb 3140 and 3160 are not qualifying interests). In line with Article 17 reporting (NPWS, 2013), all lakes larger than 1ha have been mapped as 'potential 3110' (see map 3). Two measures of extent should be used: 1. the area of the lake itself and; 2. the extent of the vegetation communities/zones that typify the habitat. Further information relating to all attributes is provided in the lake habitats supporting document for the purposes of site-specific conservation objectives and Article 17 reporting (O Connor, 2015)
Habitat distribution	Occurrence	No decline, subject to natural processes	As noted above, the exact distribution of lake habitat 3110 in Lough Hoe Bog SAC is not known. On map 3, all lakes larger than 1ha (based on 1:5,000 data), other than Lough Talt, have been mapped as potential 3110. All of these are above 200m altitude, with seven lakes over 300m
Typical species	Occurrence	Typical species present, in good condition, and demonstrating typical abundances and distribution	For lists of typical plant species, see the Article 17 habitat assessment for lake habitat 3110 (NPWS, 2013) and the lake habitats supporting document (O Connor, 2015). Douglas et al. (1989) recorded bottle sedge (<i>Carex rostrata</i>), water horsetail (<i>Equisetum fluviatile</i>), bulbous rush (<i>Juncus bulbosus</i>), water lobelia (<i>Lobelia dortmanna</i>), bogbean (<i>Menyanthes trifoliata</i>), common reed (<i>Phragmites australis</i>), bog pondweed (<i>Potamogeton polygonifolius</i>) and common club-rush (<i>Schoenoplectus lacustris</i>), amongst others, in the 3110 lakes in Lough Hoe Bog SAC
Vegetation composition: characteristic zonation	Occurrence	All characteristic zones should be present, correctly distributed and in good condition	Further work is necessary to describe the characteristic zonation and other spatial patterns in lake habitat 3110 (see O Connor, 2015)
Vegetation distribution: maximum depth	Metres	Maintain maximum depth of vegetation, subject to natural processes	The maximum depth of vegetation is likely to be specific to the lake shoreline in question. Further work is necessary to develop indicative targets for lake habitat 3110. Water clarity is expected to be high in upland 3110 lakes, resulting in a large maximum depth of vegetation
Hydrological regime: water level fluctuations	Metres	Maintain appropriate natural hydrological regime necessary to support the habitat	Fluctuations in lake water level are typical in Ireland, but can be amplified by activities such as abstraction and drainage. Increased water level fluctuations can increase wave action, up-root vegetation, increase turbidity, alter the substratum and lead to release of nutrients from the sediment. The hydrological regime of the lakes must be maintained so that the area, distribution and depth of the lake habitat and its constituent/characteristic vegetation zones and communities are not reduced

Lake substratum quality	Various	Maintain appropriate substratum type, extent and chemistry to support the vegetation	Research is required to further characterise the substratum types (particle size and origin) and substratum quality (notably pH, calcium, iron and nutrient concentrations) favoured by each of the five Annex I lake habitats in Ireland. It is likely that lake habitat 3110 is associated with a range of nutrient-poor substrates, from stones, cobble and gravel, through sands, silt, clay and peat. Substratum particle size is likely to vary with depth and along the shoreline within a single lake
Water quality: transparency	Metres	Maintain appropriate Secchi transparency. There should be no decline in Secchi depth/transparency	Transparency relates to light penetration and, hence, to the depth of colonisation of vegetation. It can be affected by phytoplankton blooms, water colour and turbidity. Specific targets have yet to be established for lake habitat 3110 (O Connor, 2015). Habitat 3110 is associated with very clear water, particularly upland examples. The OECD fixed boundary system set transparency targets for oligotrophic lakes of $\geq 6\text{m}$ annual mean Secchi disk depth, and $\geq 3\text{m}$ annual minimum Secchi disk depth. Free et al. (2009) found high isoetid abundance in lakes with Secchi depths of more than 3m
Water quality: nutrients	$\mu\text{g/l P}$; mg/l N	Maintain the concentration of nutrients in the water column at sufficiently low levels to support the habitat and its typical species	As a nutrient-poor habitat, oligotrophic and Water Framework Directive (WFD) 'high' status targets apply. Where a lake has nutrient concentrations that are lower than these targets, there should be no decline within class, i.e. no upward trend in nutrient concentrations. For lake habitat 3110, annual average total phosphorus (TP) concentration should be $\leq 10\mu\text{g/l TP}$, average annual total ammonia concentration should be $\leq 0.040\text{mg/l N}$ and annual 95th percentile for total ammonia should be $\leq 0.090\text{mg/l N}$. See also The European Communities Environmental Objectives (Surface Waters) Regulations 2009
Water quality: phytoplankton biomass	$\mu\text{g/l Chlorophyll } a$	Maintain appropriate water quality to support the habitat, including high chlorophyll <i>a</i> status	Oligotrophic and WFD 'high' status targets apply to lake habitat 3110. Where a lake has a chlorophyll <i>a</i> concentration that is lower than this target, there should be no decline within class, i.e. no upward trend in phytoplankton biomass. The average growing season (March-October) chlorophyll <i>a</i> concentration must be $< 5.8\mu\text{g/l}$. The annual average chlorophyll <i>a</i> concentration should be $< 2.5\mu\text{g/l}$ and the annual peak chlorophyll <i>a</i> concentration should be $\leq 8.0\mu\text{g/l}$. See also The European Communities Environmental Objectives (Surface Waters) Regulations 2009
Water quality: phytoplankton composition	EPA phytoplankton composition metric	Maintain appropriate water quality to support the habitat, including high phytoplankton composition status	The Environmental Protection Agency (EPA) has developed a phytoplankton composition metric for nutrient enrichment of Irish lakes. As for other water quality indicators, lake habitat 3110 requires WFD high status
Water quality: attached algal biomass	Algal cover and EPA phytobenthos metric	Maintain trace/absent attached algal biomass ($< 5\%$ cover) and high phytobenthos status	Nutrient enrichment can favour epiphytic and epipelagic algae that can out-compete the submerged vegetation. The cover abundance of attached algae in lake habitat 3110 should, therefore, be trace/absent ($< 5\%$ cover). EPA phytobenthos can be used as an indicator of changes in attached algal biomass. As for other water quality indicators, habitat 3110 requires high phytobenthos status
Water quality: macrophyte status	EPA macrophyte metric (The Free Index)	Maintain high macrophyte status	Nutrient enrichment can favour more competitive submerged macrophyte species that out-compete the typical and characteristic species for the lake habitat. The EPA monitors macrophyte status for WFD purposes using the 'Free Index'. The target for lake habitat 3110 is high status or an Ecological Quality Ratio (EQR) for lake macrophytes of ≥ 0.90 , as defined in Schedule Five of the European Communities Environmental Objectives (Surface Waters) Regulations 2009

Acidification status	pH units; mg/l	Maintain appropriate water and sediment pH, alkalinity and cation concentrations to support the habitat, subject to natural processes	Acidification can impact on species abundance and composition in soft water lake habitats. In Europe, acidification of isoetid lakes can lead to loss of isoetids and dominance by submerged <i>Sphagnum</i> mosses and <i>Juncus bulbosus</i> (Arts, 2002). The specific requirements of lake habitat 3110, in terms of water and sediment pH, alkalinity and cation concentration, have not been determined. For lake habitat 3110, and adopting a precautionary approach based on Arts (2002), minimum pH should not be <5.5 pH units. Maximum pH should be <9.0 pH units, in line with the surface water standards established for soft waters (where water hardness is ≤100mg/l calcium carbonate). See Schedule Five of the European Communities Environmental Objectives (Surface Waters) Regulations 2009
Water colour	mg/l PtCo	Maintain appropriate water colour to support the habitat	Increased water colour and turbidity decrease light penetration and can reduce the area of available habitat for lake macrophytes, particularly at the lower euphotic depths. The primary source of increased water colour in Ireland is disturbance to peatland. No habitat-specific or national standards for water colour currently exist. Studies have shown median colour concentrations in Irish lakes of 38mg/l PtCo (Free et al., 2000) and 33mg/l PtCo (Free et al., 2006). It is likely that the water colour in all Irish lake habitats would naturally be <50mg/l PtCo. Water colour can be very low (<20mg/l PtCo or even <10mg/l PtCo) in lake habitat 3110, where the peatland in the lake's catchment is intact
Dissolved organic carbon (DOC)	mg/l	Maintain appropriate organic carbon levels to support the habitat	Dissolved (and particulate) organic carbon (OC) in the water column is linked to water colour and acidification (organic acids). Increasing DOC in water has been documented across the Northern Hemisphere, including afforested peatland catchments in Ireland. Damage and degradation of peatland, leading to decomposition of peat is likely to be the predominant source of OC in Ireland. OC in water promotes decomposition by fungi and bacteria that, in turn, releases dissolved nutrients. The increased biomass of decomposers can also impact directly on the characteristic lake communities through shading, competition, etc.
Turbidity	Nephelometric turbidity units/ mg/l SS/ other appropriate units	Maintain appropriate turbidity to support the habitat	Turbidity can significantly affect the quantity and quality of light reaching rooted and attached vegetation and can, therefore, impact on lake habitats. The settlement of higher loads of inorganic or organic material on lake vegetation communities may also have impacts on sensitive, delicate species. Turbidity can increase as a result of re-suspension of material within the lake, higher loads entering the lake, or eutrophication. Turbidity measurement and interpretation is challenging. As a result, it is likely to be difficult to set habitat-specific targets for turbidity in lakes
Fringing habitat: area and condition	Hectares	Maintain the area and condition of fringing habitats necessary to support the natural structure and functioning of habitat 3110	Most lake shorelines have fringing habitats of reedswamp, other swamp, fen, marsh or wet woodland that intergrade with and support the structure and functions of the lake habitat. In Lough Hoe Bog SAC, active blanket bog and heath communities dominate upland lake shorelines. Transition mire, fen, flush and grassland may also occur. Fringing habitats are dependent on the lake, particularly its water levels, and support wetland communities and species of conservation concern. Many of the fringing wetland habitats support higher invertebrate and plant species richness than the lake habitats themselves

Conservation Objectives for : Lough Hoe Bog SAC [000633]

7130 Blanket bogs (* if active bog)

To restore the favourable conservation condition of Blanket bogs (* if active bog) in Lough Hoe Bog SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Blanket bog has not been mapped in detail for Lough Hoe Bog SAC, but from current available data the total area of the qualifying habitat is estimated to be approximately 1,176ha, covering 37% of the SAC (NPWS internal files). Further details on this and the following attributes can be found in the Lough Hoe Bog SAC conservation objectives supporting document for blanket bogs and associated habitats
Habitat distribution	Occurrence	No decline, subject to natural processes	The habitat mostly occurs in the Co. Sligo section of the SAC, with large expanses present here. In the Co. Mayo section, the habitat occurs at the edge of the SAC at Bunnyconnellan East, south of Fossea Lough, on the flatter ground in the vicinity of Loughalacka, and to the north and west of the lake at Derrynabaunshy (Douglas et al., 1989; NPWS, 2005; NPWS internal files). Further information can be found within Douglas et al. (1989), NPWS (2005), NPWS internal files and the blanket bogs and associated habitats supporting document
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil nutrient status within natural range	See the blanket bogs and associated habitats supporting document for further details
Ecosystem function: peat formation	Active blanket bog as a proportion of the total area of Annex I blanket bog habitat	At least 99% of the total Annex I blanket bog area is active	See the blanket bogs and associated habitats supporting document for further details
Ecosystem function: hydrology	Flow direction, water levels, occurrence of drains and erosion gullies	Natural hydrology unaffected by drains and erosion	Further details and a brief discussion of restoration potential is presented in the blanket bogs and associated habitats supporting document
Community diversity	Abundance of variety of vegetation communities	Maintain variety of vegetation communities, subject to natural processes	A variety of blanket bog vegetation communities have been recorded in this SAC (Douglas et al., 1989; NPWS internal files), two of which correspond to communities recorded in the National Survey of Upland Habitats and listed in the provisional list of vegetation communities described in Perrin et al. (2014). Further information on vegetation communities associated with this habitat is presented in Perrin et al. (2014)
Vegetation composition: positive indicator species	Number of species at a representative number of 2m x 2m monitoring stops	Number of positive indicator species present at each monitoring stop is at least seven	Attribute and target based on Perrin et al. (2014), where the list of positive indicator species for this habitat is also presented
Vegetation composition: lichens and bryophytes	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of bryophytes or lichens, excluding <i>Sphagnum fallax</i> , at least 10%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: negative indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of negative indicator species less than 1%	Attribute and target based on Perrin et al. (2014), where the list of negative indicator species for this habitat is also presented
Vegetation composition: non-native species	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of non-native species less than 1%	Attribute and target based on Perrin et al. (2014). The non-native moss <i>Campylopus introflexus</i> was recorded from the SAC (Douglas et al., 1989), but this species cannot be assigned specifically to blanket bog

Vegetation composition: native trees and shrubs	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of scattered native trees and shrubs less than 10%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: potential dominant species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of each of the potential dominant species less than 75%	Attribute and target based on Perrin et al. (2014). See the blanket bogs and associated habitats supporting document for the list of potential dominant species
Vegetation structure: signs of browsing	Percentage of shoots browsed at a representative number of 2m x 2m monitoring stops	Last complete growing season's shoots of ericoids, crowberry (<i>Empetrum nigrum</i>) and bog-myrtle (<i>Myrica gale</i>) showing signs of browsing collectively less than 33%	Attribute and target based on Perrin et al. (2014)
Vegetation structure: burning	Occurrence in local vicinity of a representative number of monitoring stops	No signs of burning in sensitive areas, into the moss, liverwort or lichen layer or exposure of peat surface due to burning	Attribute and target based on Perrin et al. (2014), where the list of sensitive areas for this habitat is also presented
Vegetation structure: <i>Sphagnum</i> condition	Condition at a representative number of 2m x 2m monitoring stops	Less than 10% of the <i>Sphagnum</i> cover is crushed, broken and/or pulled up	Attribute and target based on Perrin et al. (2014)
Physical structure: disturbed bare ground	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of disturbed bare ground less than 10%	Attribute and target based on Perrin et al. (2014)
Physical structure: drainage	Percentage area in local vicinity of a representative number of monitoring stops	Area showing signs of drainage from heavy trampling, tracking or ditches less than 10%	Attribute and target based on Perrin et al. (2014)
Physical structure: erosion	Percentage area in local vicinity of a representative number of monitoring stops	Less than 5% of the greater bog mosaic comprises erosion gullies and eroded areas	Attribute and target based on Perrin et al. (2014)
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat	This includes species listed in the Flora (Protection) Order, 2015 and/or the red data lists (Lockhart et al., 2012; Wyse Jackson et al., 2016)

Conservation Objectives for : Lough Hoe Bog SAC [000633]

1013 Geyer's Whorl Snail *Vertigo geyeri*

To restore the favourable conservation condition of Geyer's Whorl Snail in Lough Hoe Bog SAC, which is defined by the following list of attributes and targets:


Attribute	Measure	Target	Notes
Distribution: occupied sites	Number of occupied 1km grid squares	Restore at least one sub-population	Geyer's whorl snail (<i>Vertigo geyeri</i>) has been recorded in two separate areas on the shore of Lough Talt in Lough Hoe Bog SAC within a single 1km square, G3915 (Cawley, 2006; site code VgCAM7 in Moorkens and Killeen, 2011). See map 4. The last record from the eastern side was in 2005. The current status of the population on the western shore is uncertain. The habitats occupied by Geyer's whorl snail (<i>V. geyeri</i>) in the SAC are areas of fen and flush close to the shore of Lough Talt
Occurrence in suitable habitat	Number of positive records in a representative number of samples	No decline, subject to natural processes	Positive samples mean the confirmed presence of snails (living or recently dead adults and/or juveniles). See Moorkens and Killeen (2011)
Habitat area	Hectares	Area of suitable habitat stable or increasing, subject to natural processes; at least 1ha of suitable habitat in at least sub-optimal condition	Apparently suitable conditions for the species are present at several places, with the largest area on the east shore of Lough Talt. Two less extensive areas are found on the west shore. Optimal habitat in the SAC is defined (by Moorkens and Killeen, 2011) as flushed fen grassland with sedge/moss lawns 5-15cm tall, containing species such as <i>Carex lepidocarpa</i> , <i>Pinguicula vulgaris</i> , <i>Briza media</i> , <i>Equisetum palustre</i> , <i>Juncus articulatus</i> and the mosses <i>Drepanocladus revolvens</i> and <i>Campyllum stellatum</i> , with scattered tussocks of <i>Schoenus nigricans</i> no more than 80cm tall. During sampling, the water table should be between 0-5cm of the soil surface, but not above ground level. Sub-optimal habitat is defined as above, but vegetation height is less than 5 or more than 15cm tall, or the water table is below 5cm, or ground is flooded at time of sampling
Habitat quality: soil wetness	Percentage of a representative number of sampling stops	At least 67% of a representative number of sampling stops in areas of optimal habitat should be classified as optimal wetness as defined by Moorkens and Killeen (2011); at least 25% should be optimal wetness in areas of sub-optimal habitat	The soil wetness should be assessed using the criteria described in Moorkens and Killeen (2011)

Conservation Objectives for : Lough Hoe Bog SAC [000633]


1092 White-clawed Crayfish *Austropotamobius pallipes*



To maintain the favourable conservation condition of White-clawed Crayfish in Lough Hoe Bog SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Distribution	Number of occupied 1km grid squares	No decline. See map 5	The records for white-clawed crayfish (<i>Austropotamobius pallipes</i>) in Lough Hoe SAC all come from Lough Talt. This lake overlaps five 1km grid squares and the species has been recorded from three of these squares, G3815, G3914, G3915. See map 5. There is no reason to suppose it is not present in G4014 and G4015, but this needs confirmation
Population structure: recruitment	Occurrence of juveniles and females with eggs	Juveniles and/or females with eggs should be present in all occupied 1km squares, subject to natural processes and availability of suitable habitat	See Reynolds et al. (2010) for further details
Negative indicator species	Occurrence	No non-indigenous crayfish species	Non-indigenous crayfish species (NICS) are identified as a major direct threat to the white-clawed crayfish (<i>Austropotamobius pallipes</i>) and as a disease vector, in particular crayfish plague (<i>Aphanomyces astaci</i>), which is fatal to white-clawed crayfish. Ireland is currently free of NICS. See Reynolds (1998) for further details
Disease	Occurrence	No instances of disease	There have been outbreaks of crayfish plague (<i>Aphanomyces astaci</i>) in Ireland since 2015 and it is thought that human activity, especially the carrying of disease vectors on contaminated equipment, has introduced and spread the disease. Strict biosecurity is required
Water quality	EPA Q value	At least Q3-4 at all sites sampled by EPA	Target taken from Demers and Reynolds (2002). Q values based on triennial water quality surveys carried out by the Environmental Protection Agency (EPA)
Habitat quality: heterogeneity	Occurrence of positive habitat features	No decline in habitat quality	Crayfish need high habitat heterogeneity. Larger crayfish must have stones to hide under, or an earthen bank in which to burrow. Hatchlings shelter in vegetation, gravel and among fine tree-roots. Smaller crayfish are typically found among weed and debris in shallow water. Larger juveniles in particular may also be found among cobbles and detritus such as leaf litter. These conditions must be available throughout the area of occupied habitat



Legend

 Lough Hoe Bog SAC 000633





An Roinn
Cultúir, Oidhreachta agus Gaeltachta
Department of
Culture, Heritage and the Gaeltacht

**MAP 1:
LOUGH HOE BOG SAC
CONSERVATION OBJECTIVES
SAC DESIGNATION**

Map to be read in conjunction with the NPWS Conservation Objectives Document.

**SITE CODE:
SAC 000633; version 3. CO. SLIGO / CO. MAYO**




The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
Ordnance Survey of Ireland Licence No EN 0059216. © Ordnance Survey of Ireland Government of Ireland.

Níl sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbheirithníthe a déanamh ar theorainneacha na gceantar comharthaíthe. Suirbhéarachta Ordonáis na hÉireann Ceadúnas Uimh EN 0059216. © Suirbhéarachta Ordonáis na hÉireann Rialtas na hÉireann.



**Map Version 1
Date: Aug 2017**



Legend

- Lough Hoe Bog SAC 000633
- River Moy SAC 002298
- OSi Discovery Series County Boundary

An Roinn
Cultúir, Oidhreachta agus Gaeltachta
Department of
Culture, Heritage and the Gaeltacht

**MAP 2:
LOUGH HOE BOG SAC
CONSERVATION OBJECTIVES
ADJACENT, ADJOINING AND
OVERLAPPING DESIGNATIONS**

Map to be read in conjunction with the NPWS Conservation Objectives Document.


**SITE CODE:
SAC 000633; version 3. SAC 002298; version 3.01
CO. SLIGO / CO. MAYO**

0 3 6 9 12 km

The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
Ordnance Survey of Ireland Licence No EN 0059216. © Ordnance Survey of Ireland Government of Ireland.

Níl sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbheithnithe a déanamh ar theorainneacha na gceantar comharthaíthe. Suirbhéarachta Ordonáis na hÉireann Ceadúnas Uimh EN 0059216. © Suirbhéarachta Ordonáis na hÉireann Rialtas na hÉireann.

**Map Version 1
Date: Aug 2017**



Legend

- Lough Hoe Bog SAC 000633
- OSi Discovery Series County Boundary

Indicative Lake Habitats




Qualifying Interest



- Potential 3110 Potential oligotrophic waters containing very few minerals of sandy plains: *Littorelletalia uniflorae*

Non Qualifying Interest




- Potential 3140 Potential hard oligo-mesotrophic waters with benthic vegetation of *Chara* spp.



Legend

-  1013 Geyer's whorl snail *Vertigo geyeri*
-  Lough Hoe Bog SAC 000633
-  OSi Discovery Series County Boundary



Legend

-  1092 White-clawed Crayfish *Austropotamobius pallipes*
-  Lough Hoe Bog SAC 000633
-  OSi Discovery Series County Boundary



National Parks and Wildlife Service

Conservation Objectives Series

Lough Nabrickkeagh Bog SAC 000634



An Roinn Cultúir,
Oidhreacht agus Gaeltachta
Department of Culture,
Heritage and the Gaeltacht

**National Parks and Wildlife Service,
Department of Culture, Heritage and the Gaeltacht,
90 King Street North, Dublin 7, D07 N7CV, Ireland.**

**Web: www.npws.ie
E-mail: nature.conservation@chg.gov.ie**

Citation:

**NPWS (2019) Conservation Objectives: Lough Nabrickkeagh Bog SAC 000634.
Version 1. National Parks and Wildlife Service, Department of Culture, Heritage
and the Gaeltacht.**

**Series Editor: Rebecca Jeffrey
ISSN 2009-4086**

Introduction

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

** indicates a priority habitat under the Habitats Directive*

000634 Lough Nabrickkeagh Bog SAC

7130 Blanket bogs (* if active bog)

Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

NPWS Documents

Year :	1989
Title :	A survey to locate blanket bogs of scientific interest in County Kerry and County Sligo
Author :	Douglas, C.; Garvey, L.; Kelly, L.; O'Sullivan, A.
Series :	Unpublished report to NPWS
Year :	2005
Title :	Conservation Plan for 2005-2010. Lough Nabrickkeagh Bog cSAC Site Code 000634 Co. Sligo
Author :	NPWS
Series :	Conservation Plan
Year :	2010
Title :	Ireland Red List No. 4: Butterflies
Author :	Regan, E.C.; Nelson, B.; Aldwell, B.; Bertrand, C.; Bond, K.; Harding, J.; Nash, D.; Nixon, D.; Wilson, C.J.
Series :	Ireland Red List series, NPWS
Year :	2011
Title :	Ireland Red List No. 6: Damselflies and Dragonflies (Odonata)
Author :	Nelson, B.; Ronayne, C.; Thompson, R.
Series :	Ireland Red Lists series, NPWS
Year :	2012
Title :	Ireland Red List No. 8: Bryophytes
Author :	Lockhart, N.; Hodgetts, N.; Holyoak, D.
Series :	Ireland Red List series, NPWS
Year :	2014
Title :	Guidelines for a national survey and conservation assessment of upland vegetation and habitats in Ireland, Version 2.0
Author :	Perrin, P.M.; Barron, S.J.; Roche, J.R.; O'Hanrahan, B.
Series :	Irish Wildlife Manuals, No. 79
Year :	2016
Title :	Ireland Red List No. 10: Vascular Plants
Author :	Wyse Jackson, M.; FitzPatrick, Ú.; Cole, E.; Jebb, M.; McFerran, D.; Sheehy Skeffington, M.; Wright, M.
Series :	Ireland Red List Series, NPWS
Year :	2019
Title :	Lough Nabrickkeagh Bog SAC (000634) Conservation objectives supporting document-blanket bog and supporting habitats V1
Author :	NPWS
Series :	Conservation objectives supporting document

Conservation Objectives for : Lough Nabrickkeagh Bog SAC [000634]

7130 Blanket bogs (* if active bog)



To restore the favourable conservation condition of Blanket bogs (* if active bog) in Lough Nabrickkeagh Bog SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Blanket bog has not been mapped in detail for Lough Nabrickkeagh Bog SAC, but from current available data the total area of the qualifying habitat is estimated to be approximately 193ha, covering 71% of the SAC (NPWS internal files). Further details on this and the following attributes can be found in the Lough Nabrickkeagh Bog SAC conservation objectives supporting document for blanket bogs and associated habitats
Habitat distribution	Occurrence	No decline, subject to natural processes	The habitat is documented to occur throughout the SAC (Douglas et al., 1989; NPWS, 2005). Further information can be found within Douglas et al. (1989), NPWS (2005) and the blanket bogs and associated habitats supporting document
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil pH and nutrient status within natural ranges	See the blanket bogs and associated habitats supporting document for further details
Ecosystem function: peat formation	Active blanket bog as a proportion of the total area of Annex I blanket bog habitat	At least 99% of the total Annex I blanket bog area is active	See the blanket bogs and associated habitats supporting document for further details
Ecosystem function: hydrology	Flow direction, water levels, occurrence of drains and erosion gullies	Natural hydrology unaffected by drains and erosion	Further details and a brief discussion of restoration potential is presented in the blanket bogs and associated habitats supporting document
Community diversity	Abundance of variety of vegetation communities	Maintain variety of vegetation communities, subject to natural processes	A variety of blanket bog vegetation communities have been recorded in this SAC (Douglas et al., 1989; NPWS internal files), three of which correspond to communities recorded in the National Survey of Upland Habitats and listed in the provisional list of vegetation communities described in Perrin et al. (2014). Further information on vegetation communities associated with this habitat is presented in Perrin et al. (2014)
Vegetation composition: positive indicator species	Number of species at a representative number of 2m x 2m monitoring stop	Number of positive indicator species present at each monitoring stop is at least seven	Attribute and target based on Perrin et al. (2014), where the list of positive indicator species for this habitat is also presented
Vegetation composition: lichens and bryophytes	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of bryophytes or lichens, excluding <i>Sphagnum fallax</i> , at least 10%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: negative indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of negative indicator species less than 1%	Attribute and target based on Perrin et al. (2014), where the list of negative indicator species for this habitat is also presented
Vegetation composition: non-native species	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of non-native species less than 1%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: native trees and shrubs	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of scattered native trees and shrubs less than 10%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: potential dominant species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of each of the potential dominant species less than 75%	Attribute and target based on Perrin et al. (2014). See the blanket bogs and associated habitats supporting document for the list of potential dominant species

Vegetation structure: signs of browsing	Percentage of shoots browsed at a representative number of 2m x 2m monitoring stops	Last complete growing season's shoots of ericoids, crowberry (<i>Empetrum nigrum</i>) and bog-myrtle (<i>Myrica gale</i>) showing signs of browsing collectively less than 33%	Attribute and target based on Perrin et al. (2014)
Vegetation structure: burning	Occurrence in local vicinity of a representative number of monitoring stops	No signs of burning in sensitive areas, into the moss, liverwort or lichen layer or exposure of peat surface due to burning	Attribute and target based on Perrin et al. (2014), where the list of sensitive areas for this habitat is also presented
Vegetation structure: <i>Sphagnum</i> condition	Condition at a representative number of 2m x 2m monitoring stops	Less than 10% of the <i>Sphagnum</i> cover is crushed, broken and/or pulled up	Attribute and target based on Perrin et al. (2014)
Physical structure: disturbed bare ground	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of disturbed bare ground less than 10%	Attribute and target based on Perrin et al. (2014)
Physical structure: drainage	Percentage area in local vicinity of a representative number of monitoring stops	Area showing signs of drainage from heavy trampling, tracking or ditches less than 10%	Attribute and target based on Perrin et al. (2014)
Physical structure: erosion	Percentage area in local vicinity of a representative number of monitoring stops	Less than 5% of the greater bog mosaic comprises erosion gullies and eroded areas	Attribute and target based on Perrin et al. (2014)
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat	This includes species on the Flora (Protection) Order, 2015 and/or Red Lists (Regan et al., 2010; Nelson et al., 2011; Lockhart et al., 2012; Wyse Jackson et al., 2016, etc.)

Legend

 Lough Nabrickkeagh Bog SAC 000634



© Ordnance Survey Ireland




An Roinn Cultúir,
Oidhreacht agus Gaeltachta
Department of Culture,
Heritage and the Gaeltacht

MAP 1:
LOUGH NABRICKKEAGH BOG SAC
CONSERVATION OBJECTIVES
SAC DESIGNATION

Map to be read in conjunction with the NPWS Conservation Objectives Document.

SITE CODE:
SAC 000634; version 3.02
CO. SLIGO



The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
Ordnance Survey of Ireland Licence No EN 0059216. © Ordnance Survey of Ireland Government of Ireland.

Níl sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbhreithnithe a déanamh ar theorainneacha na gceantar comharthaithe. Suirbhéarachta Ordonáis na hÉireann Ceadúnas Uimh EN 0059216. © Suirbhéarachta Ordonáis na hÉireann Rialtas na hÉireann.



Map Version 1
Date: Jan 2019



Conservation objectives for Templehouse and Cloonacleigh Loughs SAC [000636]

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Objective: To maintain or restore the favourable conservation condition of the Annex I habitat(s) and/or the Annex II species for which the SAC has been selected:

Code	Description
------	-------------

3140	Hard oligo-mesotrophic waters with benthic vegetation of <i>Chara</i> spp.
------	--

3260	Water courses of plain to montane levels with the <i>Ranunculion fluitantis</i> and <i>Callitriche-Batrachion</i> vegetation
------	--

* denotes a priority habitat



Citation: NPWS (2020) Conservation objectives for Templehouse and Cloonacleigha Loughs SAC [000636]. Generic Version 7.0. Department of Culture, Heritage and the Gaeltacht.



Conservation objectives for Turloughmore (Sligo) SAC [000637]

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Objective: To maintain or restore the favourable conservation condition of the Annex I habitat(s) and/or the Annex II species for which the SAC has been selected:

Code	Description
------	-------------

3180	Turloughs*
------	------------

* denotes a priority habitat



Citation: NPWS (2020) Conservation objectives for Turloughmore (Sligo) SAC [000637]. Generic Version 7.0. Department of Culture, Heritage and the Gaeltacht.



Conservation objectives for Rusheenduff Lough SAC [001311]

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Objective: To maintain or restore the favourable conservation condition of the Annex I habitat(s) and/or the Annex II species for which the SAC has been selected:

Code Description

3130 Oligotrophic to mesotrophic standing waters with vegetation of the Littorelletea uniflorae and/or Isoetes-Nanojuncetea

* denotes a priority habitat

Code Common Name Scientific Name

1833 Slender Naiad *Najas flexilis*



Citation: NPWS (2020) Conservation objectives for Rusheenduff Lough SAC [001311]. Generic Version 7.0. Department of Culture, Heritage and the Gaeltacht.

National Parks and Wildlife Service

Conservation Objectives

Clew Bay Complex SAC 001482



*An Roinn
Ealaíon, Oidhreachta agus Gaeltachta*
*Department of
Arts, Heritage and the Gaeltacht*

Introduction

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

** indicates a priority habitat under the Habitats Directive*

001482 Clew Bay Complex SAC

QI	Description
1013	Geyer's whorl snail <i>Vertigo geyeri</i>
1140	Mudflats and sandflats not covered by seawater at low tide
1150	* Coastal lagoons
1160	Large shallow inlets and bays
1210	Annual vegetation of drift lines
1220	Perennial vegetation of stony banks
1330	Atlantic salt meadows (<i>Glauco-Puccinellietalia maritimae</i>)
1355	Otter <i>Lutra lutra</i>
1365	Common seal <i>Phoca vitulina</i>
2110	Embryonic shifting dunes
2120	Shifting dunes along the shoreline with <i>Ammophila arenaria</i> ("white dunes")

Supporting documents, relevant reports & publications (listed by date)

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

Title: Monitoring and Assessment of Irish Lagoons for the purpose of the EU Water Framework Directive
Year: in prep
Author: Roden, C.M.; Oliver, G.
Series: Unpublished report to the EPA

Title: Clew Bay Complex SAC (001482): Conservation objectives supporting document - marine habitats and species [Version 1]
Year: 2011
Author: NPWS
Series: Unpublished Report to NPWS

Title: Clew Bay Complex SAC (001482): Conservation objectives supporting document - coastal habitats [Version 1]
Year: 2011
Author: NPWS
Series: Unpublished Report to NPWS

Title: Otter tracking study of Roaringwater Bay
Year: 2010
Author: De Jongh, A.; O'Neill, L.
Series: Unpublished Draft Report to NPWS

Title: Subtidal benthic surveys (Clew Bay)
Year: 2009
Author: Aquafact
Series: Unpublished Report to NPWS

Title: Saltmarsh Monitoring Report 2007-2008
Year: 2009
Author: McCorry, M.; Ryle, T.
Series: Unpublished Report to NPWS

Title: Clew Bay baseline intertidal survey
Year: 2009
Author: RPS
Series: Unpublished Report to NPWS

Title: Coastal Monitoring Project 2004-2006
Year: 2009
Author: Ryle, T.; Murray, A.; Connolly, C.; Swann, M.
Series: Unpublished Report to NPWS

Title: The phytosociology and conservation value of Irish sand dunes
Year: 2008
Author: Gaynor, K.
Series: Unpublished PhD thesis, National University of Ireland, Dublin

Title: Saltmarsh Monitoring Report 2006
Year: 2007
Author: McCorry, M.
Series: Unpublished Report to NPWS

Title: Inventory of Irish coastal lagoons
Year: 2007
Author: Oliver, G.
Series: Unpublished Report to NPWS

Title: A Survey of Intertidal Mudflats and Sandflats in Ireland
Year: 2006
Author: Aquafact
Series: Unpublished Report to NPWS

Title: Otter Survey of Ireland 2004/2005
Year: 2006
Author: Bailey, M.; Rochford, J.
Series: Irish Wildlife Manuals No. 23

Title: Otters - ecology, behaviour and conservation
Year: 2006
Author: Kruuk, H.
Series: Oxford University Press

Title: Survey of sensitive subtidal benthic marine communities
Year: 2006
Author: MERC
Series: Unpublished Report to NPWS

Title: Harbour seal population assessment in the Republic of Ireland: August 2003
Year: 2004
Author: Cronin, M.; Duck, C.; Ó Cadhla, O.; Nairn, R.; Strong, D.; O'Keeffe, C.
Series: Irish Wildlife Manuals No. 11

Title: Summary of National Parks & Wildlife Service surveys for common (harbour) seals (*Phoca vitulina*) and grey seals (*Halichoerus grypus*), 1978 to 2003
Year: 2004
Author: Lyons, D.O.
Series: Irish Wildlife Manuals No. 13

Title: Broadscale mapping of candidate marine Special Area of Conservation. Clew Bay Complex, cSAC (001482)
Year: 2003
Author: SSI; Aquafact
Series: Unpublished Report to NPWS

Title: A Survey of selected littoral and sublittoral sites in Clew Bay, Co. Mayo
Year: 1999
Author: Aquafact
Series: Unpublished Report to NPWS

-
- Title:** National Shingle Beach Survey of Ireland 1999
Year: 1999
Author: Moore, D.; Wilson, F.
Series: Unpublished Report to NPWS
-
- Title:** Aquatic vegetation of Irish coastal lagoons
Year: 1998
Author: Hatch, P.; Healy, B.
Series: Bulletin of the Irish Biogeographical Society. 21: 2-21
-
- Title:** A survey of the vegetation of Irish coastal lagoons
Year: 1996
Author: Hatch, P.
Series: Unpublished Report to NPWS
-
- Title:** The spatial organization of otters (*Lutra lutra*) in Shetland
Year: 1991
Author: Kruuk, H.; Moorhouse, A.
Series: J. Zool, 224: 41-57
-
- Title:** Otter survey of Ireland
Year: 1982
Author: Chapman, P.J.; Chapman, L.L.
Series: Unpublished Report to Vincent Wildlife Trust
-
- Title:** Lough Furnace, County Mayo; physical and chemical studies of an Irish saline lake, with reference to the biology of *Neomysis integer*
Year: 1977
Author: Parker, M.M.
Series: Unpublished PhD thesis, University of Dublin, Trinity College.
-

Spatial data sources

Year:	Interpolated 2011
Title:	Intertidal and subtidal surveys 1999, 2006, 2009; broadscale mapping 2003
GIS operations:	Polygon feature classes from marine community types base data sub-divided based on interpolation of marine survey data; expert opinion used as necessary to resolve any issues arising
Used for:	Marine community types, 1140 (maps 2 & 4)
Year:	2005
Title:	OSi Discovery series vector data
GIS operations:	High Water Mark (HWM) polyline feature class converted into polygon feature class; clipped to SAC boundary
Used for:	1160, 1365 (maps 3 & 9)
Year:	2005
Title:	OSi Discovery series vector data
GIS operations:	High water mark (HWM) and low water mark (LWM) polyline feature classes converted into polygon feature classes and combined; Saltmarsh and Sand Dune CO datasets erased out if applicable
Used for:	Marine community types base data (map 4)
Year:	Revision 2011
Title:	Inventory of Irish Coastal Lagoons. Version 3
GIS operations:	Clipped to SAC boundary
Used for:	1150 (map 5)
Year:	Revision 2010
Title:	Saltmarsh Monitoring Project 2007-2008. Version 1
GIS operations:	QIs selected; clipped to SAC boundary; overlapping regions with Sand Dune CO data investigated and resolved with expert opinion used
Used for:	1330 (map 6)
Year:	2009
Title:	Coastal Monitoring Project 2004-2006. Version 1
GIS operations:	QIs selected; clipped to SAC boundary; overlapping regions with Saltmarsh CO data investigated and resolved with expert opinion used
Used for:	1210, 2110, 2120 (map 7)
Year:	2005
Title:	OSi Discovery series vector data
GIS operations:	Creation of an 80m buffer on the marine side of the high water mark (HWM); creation of a 10m buffer on the terrestrial side of the HWM; combination of 80m and 10m HWM buffer datasets; creation of a 10m buffer on the landward side of the river banks data; creation of a 20m buffer applied to river centerline and stream data; combination of 10m river banks and 20m river and stream centerline buffer datasets; combined river and stream buffer dataset clipped to HWM; combination of HWM buffer dataset with river and stream buffer dataset; overlapping regions investigated and resolved; resulting dataset clipped to SAC boundary; expert opinion used as necessary to resolve any issues arising
Used for:	1355 (map 8)

Year: 2011
Title: NPWS rare and threatened species database
GIS operations: Dataset created from spatial references in database records; expert opinion used as necessary to resolve any issues arising
Used for: 1365 (map 9)

1013 Geyer's whorl snail *Vertigo geyeri*

The status of Geyer's whorl snail as a qualifying Annex II species for Clew Bay Complex SAC is currently under review. The outcome of this review will determine whether a site-specific conservation objective is set for this species.

1140 Mudflats and sandflats not covered by seawater at low tide

To maintain the favourable conservation condition of Mudflats and sandflats not covered by seawater at low tide in Clew Bay Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	The permanent habitat area is stable or increasing, subject to natural processes. See map 2	Habitat area was estimated using OSI data as 1277ha. See marine supporting document for further details
Community distribution	Hectares	The following sediment communities should be maintained in a natural condition: Intertidal sandy mud with <i>Tubificoides benedii</i> and <i>Pygospio elegans</i> community complex; Sandy mud with polychaetes and bivalves community complex; and Fine sand dominated by <i>Nephtys cirrosa</i> community. See map 4	The likely area of sediment communities was derived from a combination of intertidal and subtidal surveys undertaken in 1999, 2006 and 2009. See marine supporting document for further details

1150 * Coastal lagoons

To maintain the favourable conservation condition of Lagoons in Clew Bay Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat distribution	Occurrence	No decline, subject to natural processes. See map 5 for mapped lagoons	The main lagoon is Furnace Lough. Claggan Lagoon has also been mapped, however, further information is required on this lagoon. NB there maybe other lagoons within the SAC. The following targets and notes concentrate on the largest lagoon, Furnace Lough
Habitat area	Hectares	Area stable, subject to slight natural variation. Favourable reference area of surveyed lagoons is 163.3ha. Furnace Lough- 162.1ha; Claggan Lagoon- 1.2ha. See map 5	Areas calculated from spatial data derived from Oliver, 2007. NB there maybe other lagoons within the SAC
Salinity regime	Practical salinity units (psu)	Maintain current spatial and temporal variation in salinity regime	Furnace Lough is a natural, deep (up to 21m), stratified lagoon with natural periodic overturns and anoxia. It has permanent open connection to the sea through which seawater enters when tides exceed MHWN though this connection is somewhat constricted by weirs. There are major freshwater inputs at the northern end from the large Lough Feeagh/Burrishoole catchment area. The surface layer is oligohaline to mesohaline (0.5-12.0 psu) for most of the time but salinity varies from north (fresh water) to south (high salinity) and summer to winter. The waters are sharply stratified, a permanent halocline runs from 1-3m down to 8m, below which the water is of constant salinity (approx. 20psu), anaerobic and stagnant (Parker, 1977). See Oliver (2007) and Roden and Oliver (in prep.) for further information
Hydrological regime	Metres	Maintain current annual water level fluctuations	This is to ensure maintenance of the current communities of the lagoon margins and the current hydrological functioning of the lagoon itself, especially the salinity regime
Hydrological regime	Discharge (m ³ /second)	Maintain/restore freshwater discharge regime	There is evidence that the original hydrological regime in the Burrishoole catchment has been impacted due to overgrazing and afforestation resulting in changes to run-off regimes with associated increased siltation and eutrophication. The extent to which these changes have impacted on Lough Furnace is unclear but needs further study

1150 * Coastal lagoons

To maintain the favourable conservation condition of Lagoons in Clew Bay Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Barrier	Weir function	Maintain current weir structure at Furnace Lough to ensure maintenance of the current salinity regime	In Furnace Lough, input to and output of saline water is affected to an unknown degree by two weirs. The effect of the weirs needs to be quantified to determine their effect on the salinity regime of the lagoon. These weirs or some similar type structures are shown on the first edition of the 6" OS maps and therefore have been in place for over 170 years
Water quality: chlorophyll a	µg/L	Maintain annual median chlorophyll in Furnace Lough at less than 2.5µg/L	These limits are needed to ensure that excessive shading from phytoplankton does not reduce submergent macrophytes colonisation of the littoral zone the lagoon (J. Ryan, pers comm). The current median levels are less than the target but summer levels are elevated (Roden and Oliver, in prep.) and should be closely monitored
Water quality: Molybdate Reactive Phosphorus (MRP)	mg/L	Maintain annual median MRP in Furnace Lough at less than 0.01mg/L	These limits are needed to ensure that excessive shading from phytoplankton does not reduce submergent macrophytes colonisation of the littoral zone areas of the lagoon (J. Ryan, pers comm). The current median levels in Furnace Lough are 0.005mg/L (Roden and Oliver, in prep). It is possible that the target may be exceeded during periods of overturn. Collection of data on nutrient levels close to the halocline would be useful for the assessment of this possibility
Water quality: Dissolved Inorganic Nitrogen (DIN)	mg/L	Maintain annual median DIN (Dissolved inorganic nitrogen) in Furnace Lough at less than 0.15mg/L	These limits are needed to ensure that excessive shading from phytoplankton does not reduce submergent macrophytes colonisation of the littoral zone of the lagoon (J. Ryan, pers comm). The current median levels of DIN in Furnace Lough are less than 0.1mg/L (Roden and Oliver, in prep)
Water quality: Biological Oxygen Demand (BOD)	mg/L	Maintain annual median BOD (Biological Oxygen Demand) in Furnace Lough at less than 2.0mg/L	These limits are needed to ensure that excessive shading from phytoplankton does not reduce submergent macrophytes colonisation of the littoral zone of the lagoon (J. Ryan, pers comm). The current annual median levels of BOD in Furnace Lough are just below the target (Roden and Oliver, in prep) and should be closely monitored. The relationship between organic matter, mainly peat silt, input from L. Feeagh and BOD in the surface waters and anoxia in the deeper waters warrants further investigation

1150 * Coastal lagoons

To maintain the favourable conservation condition of Lagoons in Clew Bay Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Depth of submergent macrophyte colonisation	Metres	Maintain/increase the depth of submergent macrophyte colonisation of the lagoon	Increased depth of colonisation increases both the extent and diversity of submergent macrophytes. In comparison with similar lagoons the extent of submergent macrophyte colonisation in Furnace Lough appears to be restricted probably due to high water colour. However data on the depth of colonisation and water colour and the relationship between them is lacking. It is also possible that anoxia may be a problem, at least in some areas. These issues need to be investigated
Typical plant species	Number and m ²	Maintain number and extent of listed lagoonal specialists, subject to natural variation	Species in Furnace Lough listed in Oliver (2007), Hatch (1996) and Hatch and Healy (1998). A very limited number of plant species are currently listed for the site based on a series of shallow water transects. A snorkelling survey of this complex lagoon is required establish if that list is fully representative of the flora of the lagoon
Typical animal species	Number	Maintain listed lagoon specialists, subject to natural variation	Species in Furnace Lough listed in Oliver (2007), which rated the aquatic fauna as of moderate-high conservation value based on its high diversity and the presence of rare and unexpected crustaceans
Negative indicator species	Number and % cover	Negative indicator species absent or under control	Eutrophication would favour phytoplankton blooms at the expense of submerged macrophytes

1160 Large shallow inlets and bays

To maintain the favourable conservation condition of Large shallow inlets and bays in Clew Bay Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	The permanent habitat area is stable or increasing, subject to natural processes. See map 3	Habitat area was estimated using OSI data as 10189ha. See marine supporting document for further details.
Community extent	Hectares	Maintain the natural extent of the <i>Zostera</i> dominated and maërl dominated communities. See map 4	The likely extent of the <i>Zostera</i> dominated and maërl dominated communities was derived from the acoustic survey and the dive survey undertaken in 2006. See marine supporting document for further details
Shoot density	Shoots per m ²	Maintain the high quality of <i>Zostera</i> dominated community	2006 diver observation and underwater viewer. See marine supporting document for further details
Community structure	Biological composition	Maintain the high quality of maërl dominated communities	Area established from an acoustic mapping survey 2003 and a 2006 diver observation and underwater viewer. See marine supporting document for further details
Community distribution	Hectares	The following communities should be maintained in a natural condition: Sandy mud with polychaetes and bivalves community complex; Fine sand dominated by <i>Nephtys cirrosa</i> community; Intertidal sandy mud with <i>Tubificoides benedii</i> and <i>Pygospio elegans</i> community complex; Shingle; and Reef. See map 4	The likely area of sediment communities was derived from a combination of acoustic mapping survey in 2003, intertidal data from 1999, 2006 and 2009 and subtidal data obtained in 1999 and 2009. See marine supporting document for further details

1210 Annual vegetation of drift lines

To maintain the favourable conservation condition of Annual vegetation of driftlines in Clew Bay Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes, including erosion and succession. For sub-sites mapped: Bartraw - 0.04ha and Rosmurrevagh - 0.08ha. See map 7	Current area unknown. Two sub-sites (Bartraw and Rosmurrevagh) were mapped during the Coastal Monitoring Project (Ryle et al., 2009), giving a total estimated area of 0.12ha. NB further unsurveyed areas maybe present in the site. Habitat is very difficult to measure in view of its dynamic nature which means that it can appear and disappear within a site from year to year. See coastal habitats supporting document for further details
Habitat distribution	Occurrence	No decline, subject to natural processes	Current distribution unknown. Majority of habitat found at Bartraw and Rosmurrevagh, although there may be additional patches distributed throughout the site. See coastal habitats supporting document for further details
Physical structure: functionality and sediment supply	Presence/absence of physical barriers	Maintain the natural circulation of sediment and organic matter, without any physical obstructions	Dunes are naturally dynamic systems that require continuous supply and circulation of sand. Accumulation of organic matter in tidal litter is essential for trapping sand and initiating dune formation. Physical barriers can lead to fossilisation or over-stabilisation of dunes, as well as beach starvation resulting in increased rates of erosion. See coastal habitats supporting document for further details
Vegetation structure: zonation	Occurrence	Maintain the range of coastal habitats including transitional zones, subject to natural processes including erosion and succession	Based on data from Ryle et al. (2009). See coastal habitats supporting document for further details
Vegetation composition: typical species and sub-communities	Percentage cover at a representative number of monitoring stops	Maintain the presence of species-poor communities with typical species: <i>Cakile maritima</i> , <i>Honckenya peploides</i> , <i>Salsola kali</i> and <i>Atriplex</i> spp.	Based on data from Ryle et al. (2009) . See coastal habitats supporting document for further details
Vegetation composition: negative indicator species	Percentage cover	Negative indicator species (including non-natives) to represent less than 5% cover	Negative indicators include non-native species, species indicative of changes in nutrient status and species not considered characteristic of the habitat. Based on data from Ryle et al. (2009). See coastal habitats supporting document for further details

1220 Perennial vegetation of stony banks

To maintain the favourable conservation condition of Perennial vegetation of stony banks in Clew Bay Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes, including erosion and succession	Current area unknown, but Clew Bay is considered to have the largest shingle reserves in the country. It was recorded from Clew Bay Complex, Bartraw and Rosmurrevagh during the National Shingle Beach Survey (Moore and Wilson, 1999), but the extent was not mapped. The Coastal Monitoring Project mapped 0.48ha of this habitat at Bartraw and 0.01ha at Rosmurrevagh (Ryle et al., 2009). The extent is considerably greater than this figure, as substantial shingle deposits are known to occur in association with many of the drumlins in Clew Bay. See coastal habitats supporting document for further details
Habitat distribution	Occurrence	No decline, subject to natural processes	Distribution unknown at present, although the habitat has been recorded at Clew Bay Complex (Moore and Wilson, 1999), as well as Bartraw and Rosmurrevagh (Moore and Wilson, 1999; Ryle et al., 2009). See coastal habitats supporting document for further details
Physical structure: Functionality and sediment supply	Presence/absence of physical barriers	Maintain the natural circulation of sediment and organic matter, without any physical obstructions	Site represents the only known example of incipient gravel barrier formation in the country. See coastal habitats supporting document for further details
Vegetation structure: zonation	Occurrence	Maintain the range of coastal habitats including transitional zones, subject to natural processes including erosion and succession	Based on data from Moore and Wilson (1999) and Ryle et al. (2009). See coastal habitats supporting document for further details
Vegetation composition: typical species and sub-communities	Percentage cover at a representative sample of monitoring stops	Maintain the presence of species-poor communities with typical species: <i>Honckenya peploides</i> , <i>Beta vulgaris</i> ssp. <i>maritima</i> , <i>Crithmum maritimum</i> , <i>Tripleurospermum maritimum</i> , <i>Glaucium flavum</i> and <i>Silene uniflora</i>	Based on data from Moore and Wilson (1999) and Ryle et al. (2009). See coastal habitats supporting document for further details
Vegetation composition: negative indicator species	Percentage cover	Negative indicator species (including non-natives) to represent less than 5% cover	Based on data from Moore and Wilson (1999) and Ryle et al. (2009). Negative indicators include non-native species, species indicative of changes in nutrient status and species not considered characteristic of the habitat. See coastal habitats supporting document for further details

1330 Atlantic salt meadows (*Glauco-Puccinellietalia maritimae*)

To restore the favourable conservation condition of Atlantic salt meadows in Clew Bay Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes, including erosion and succession. For sub-sites mapped: Mallaranny - 19.76ha, Tooreen - 1.06ha, Rosmurrevagh - 6.40ha, Tierna - 0.39ha, Rockfleet Castle - 0.37ha, Rosharnagh East - 0.03ha, Caraholly - 0.36ha, Kiladangan - 0.96ha, Annagh Island - 5.23ha, Bartraw - 0.38ha. See map 6	Based on data from the Saltmarsh Monitoring Project (McCorry, 2007). Ten sub-sites were mapped (34.94ha) and additional areas of potential saltmarsh (3.92ha) were identified for an examination of aerial photographs, giving a total estimated area of 38.86ha. NB further unsurveyed areas maybe present within the site. See coastal habitats supporting document for further details
Habitat distribution	Occurrence	No decline, subject to natural processes. See map 6 for known distribution	Based on data from McCorry (2007). See coastal habitats supporting document for further details
Physical structure: sediment supply	Presence/absence of physical barriers	Maintain/restore natural circulation of sediments and organic matter, without any physical obstructions	See coastal habitats backing document for further details
Physical structure: creeks and pans	Occurrence	Maintain creek and pan structure, subject to natural processes, including erosion and succession	Based on data from McCorry (2007). The efficiency of sediment circulation throughout a saltmarsh depends on the creek pattern. See coastal habitats supporting document for further details
Physical structure: flooding regime	Hectares flooded; frequency	Maintain natural tidal regime	See coastal habitats supporting document for further details
Vegetation structure: zonation	Occurrence	Maintain the range of coastal habitats including transitional zones, subject to natural processes including erosion and succession	Based on data from McCorry and Ryle (2009). See coastal habitats supporting document for further details
Vegetation structure: vegetation height	Centimetres	Maintain structural variation within sward	Based on data from McCorry (2007). See coastal habitats supporting document for further details
Vegetation structure: vegetation cover	Percentage cover at a representative sample of monitoring stops	Maintain more than 90% area outside creeks vegetated.	Based on data from McCorry (2007). See coastal habitats supporting document for further details
Vegetation composition: typical species and sub-communities	Percentage cover at a representative sample of monitoring stops	Maintain range of sub-communities with typical species listed in Saltmarsh Monitoring Project (McCorry & Ryle, 2009)	Based on data from McCorry (2007). See coastal habitats supporting document for further details

1330 Atlantic salt meadows (*Glauco-Puccinellietalia maritimae*)

To restore the favourable conservation condition of Atlantic salt meadows in Clew Bay Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Vegetation structure: negative indicator species - <i>Spartina anglica</i>	Hectares	No significant expansion of <i>Spartina</i> . No new sites for this species and an annual spread of less than 1% where it is already known to occur	Based on data from McCorry (2007). See coastal habitats supporting document for further details

1355 Otter *Lutra lutra*

To restore the favourable conservation condition of Otter in Clew Bay Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Distribution	Percentage positive survey sites	No significant decline	Measure based on standard otter survey technique. FCS target, based on 1980/81 survey findings, is 88% in SACs. Current range in west estimated at 70% (Bailey and Rochford, 2006)
Extent of terrestrial habitat	Hectares	No significant decline. Area mapped and calculated as 233.1ha above high water mark (HWM); 47.3ha along river banks/ around ponds	No field survey. Areas mapped to include 10m terrestrial buffer along shoreline (above HWM and along river banks) identified as critical for otters (NPWS, 2007)
Extent of marine habitat	Hectares	No significant decline. Area mapped and calculated as 2426.7ha	No field survey. Area mapped based on evidence that otters tend to forage within 80m of the shoreline (HWM) (NPWS, 2007; Kruuk, 2006)
Extent of freshwater (river) habitat	Kilometres	No significant decline. Length mapped and calculated as 10.2km	No field survey. River length calculated on the basis that otters will utilise freshwater habitats from estuary to headwaters (Chapman and Chapman, 1982)
Extent of freshwater (lake/lagoon) habitat	Hectares	No significant decline. Area mapped and calculated as 141.3ha	No field survey. Area mapped based on evidence that otters tend to forage within 80m of the shoreline (NPWS, 2007)
Couching sites and holts	Number	No significant decline	Otters need lying up areas throughout their territory where they are secure from disturbance (Kruuk, 2006; Kruuk and Moorhouse, 1991)
Fish biomass available	Kilograms	No significant decline	Broad diet that varies locally and seasonally, but dominated by fish, in particular salmonids, eels and sticklebacks in freshwater (Bailey and Rochford, 2006) and wrasse and rockling in coastal waters (Kingston et al., 1999)
Barriers to connectivity	Number	No significant increase. For guidance, see map 8	Otters will regularly commute across stretches of open water up to 500m. e.g. between the mainland and an island; between two islands; across an estuary (De Jongh and O'Neill, 2010). It is important that such commuting routes are not obstructed

1365 Common seal *Phoca vitulina*

To maintain the favourable conservation condition of Harbour seal in Clew Bay Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Access to suitable habitat	Number of artificial barriers	Species range within the site should not be restricted by artificial barriers to site use	See marine supporting document for further details
Breeding behaviour	Breeding sites	The breeding sites should be maintained in a natural condition. See map 9	Attribute and target based on background knowledge of Irish breeding populations, review of data from Lyons (2004) and unpublished National Parks and Wildlife Service records. See marine supporting document for further details
Moulting behaviour	Moult haul-out sites	The moult haul-out sites should be maintained in a natural condition. See map 9	Attribute and target based on background knowledge of Irish populations, review of data from Lyons (2004), Cronin et al. (2004) and unpublished National Parks and Wildlife Service records. See marine supporting document for further details
Resting behaviour	Resting haul-out sites	The resting haul-out sites should be maintained in a natural condition. See map 9	Attribute and target based on background knowledge of Irish populations, review of data from Lyons (2004) and unpublished National Parks and Wildlife Service records. See marine supporting document for further details
Disturbance	Level of impact	Human activities should occur at levels that do not adversely affect the harbour seal population at the site	See marine supporting document for further details

2110 Embryonic shifting dunes

To restore the favourable conservation condition of Embryonic shifting dunes in Clew Bay Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes, including erosion and succession. For sub-sites mapped: Bartraw - 0.02ha and Rosmurrevagh - 1.38ha. See map 7	Current area unknown. Two sub-sites (Bartraw and Rosmurrevagh) were mapped during the Coastal Monitoring Project (Ryle et al., 2009), giving a total estimated area of 1.40ha. NB further unsurveyed areas maybe present in the site. Habitat is very difficult to measure in view of its dynamic nature. See coastal habitats supporting document for further details
Habitat distribution	Occurrence	No decline, subject to natural processes. See map 7 for known distribution	Mobile dunes are well developed at Rosmurrevagh, while those at Bartraw have been compromised by the installation of coastal protection works. See coastal habitats supporting document for further details
Physical structure: functionality and sediment supply	Presence/absence of physical barriers	Maintain the natural circulation of sediment and organic matter, without any physical obstructions	Dunes are naturally dynamic systems that require continuous supply and circulation of sand. Physical barriers can lead to fossilisation or over-stabilisation of dunes, as well as beach starvation resulting in increased rates of erosion. See coastal habitats supporting document for further details
Vegetation structure: zonation	Occurrence	Maintain the range of coastal habitats including transitional zones, subject to natural processes including erosion and succession	Based on data from Ryle et al. (2009). See coastal habitats supporting document for further details
Vegetation composition: plant health of foredune grasses	Percentage cover	More than 95% of <i>Elytrigia</i> and/or <i>Leymus</i> should be healthy (i.e. green plant parts above ground and flowering heads present)	Based on data from Ryle et al. (2009). See coastal habitats supporting document for further details
Vegetation composition: typical species and sub-communities	Percentage cover	Maintain the presence of species-poor communities with typical species: <i>Elytrigia juncea</i> and/or <i>Leymus arenarius</i>	Based on data from Ryle et al. (2009). See coastal habitats supporting document for further details
Vegetation composition: negative indicator species	Percentage cover	Negative indicator species (including non-natives) to represent less than 5% cover	Based on data from Ryle et al. (2009). Negative indicators include non-native species, species indicative of changes in nutrient status and species not considered characteristic of the habitat. Sea-buckthorn (<i>Hippophae rhamnoides</i>) should be absent or effectively controlled. See coastal habitats supporting document for further details

2120 Shifting dunes along the shoreline with *Ammophila arenaria* ("white dunes")

To restore the favourable conservation condition of Shifting dunes along the shoreline with *Ammophila arenaria* in Clew Bay Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes including erosion and succession. For sub-sites mapped: Bartraw - 0.18ha and Rosmurrevagh - 0.36ha. See map 7	Current area unknown. Two sub-sites (Bartraw and Rosmurrevagh) were mapped during the Coastal Monitoring Project (Ryle et al., 2009), giving a total estimated area of 0.54ha. NB further unsurveyed areas may be present in the site. Habitat is very difficult to measure in view of its dynamic nature. See coastal habitats supporting document for further details
Habitat distribution	Occurrence	No decline, subject to natural processes. See map 7 for known distribution	Mobile dunes are well developed at Rosmurrevagh, while those at Bartraw have been compromised by the installation of coastal protection works. See coastal habitats supporting document for further details
Physical structure: functionality and sediment supply	Presence/absence of physical barriers	Maintain the natural circulation of sediment and organic matter, without any physical obstructions	Dunes are naturally dynamic systems that require continuous supply and circulation of sand. <i>Ammophila</i> reproduces vegetatively and requires constant accretion of fresh sand to maintain active growth encouraging further accretion. Physical barriers can lead to fossilisation or over-stabilisation of dunes, as well as beach starvation resulting in increased rates of erosion. See coastal habitats supporting document for further details
Vegetation structure: zonation	Occurrence	Maintain the range of coastal habitats including transitional zones, subject to natural processes including erosion and succession	Based on data from Gaynor (2008) and Ryle et al. (2009). See coastal habitats supporting document for further details
Vegetation composition: plant health of dune grasses	Percentage cover	More than 95% of <i>Ammophila</i> and/or <i>Leymus</i> should be healthy (i.e. green plant parts above ground and flowering heads present)	Based on data from Ryle et al. (2009). See coastal habitats supporting document for further details
Vegetation composition: typical species and sub-communities	Percentage cover at a representative number of monitoring stops	Maintain the presence of species-poor communities dominated by <i>Ammophila arenaria</i> and/or <i>Leymus arenarius</i>	Based on data from Gaynor (2008) and Ryle et al. (2009). See coastal habitats supporting document for further details

2120 Shifting dunes along the shoreline with *Ammophila arenaria* ("white dunes")


To restore the favourable conservation condition of Shifting dunes along the shoreline with *Ammophila arenaria* in Clew Bay Complex SAC, which is defined by the following list of attributes and targets:


Attribute	Measure	Target	Notes
Vegetation composition: negative indicator species	Percentage cover	Negative indicator species (including non-natives) to represent less than 5% cover	Based on data from Ryle et al. (2009). Negative indicators include non-native species, species indicative of changes in nutrient status and species not considered characteristic of the habitat. Sea-buckthorn (<i>Hippophae rhamnoides</i>) should be absent or effectively controlled. See coastal habitats supporting document for further details



Legend


SAC 001482





Legend

- SAC 001482
- 1140 Mudflats and sandflats not covered by sea water at low tide
- OSi Discovery Series County Boundary



Legend

- SAC 001482
- 1160 Large shallow inlets and bays
- OSi Discovery Series County Boundary

MAP 3:
CLEW BAY COMPLEX
CONSERVATION OBJECTIVES
LARGE SHALLOW INLETS AND BAYS

Map to be read in conjunction with the NPWS Conservation Objectives Document.

COUNTY MAYO

0 1 2 3 km


The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision. Reproduced from Ordnance Survey material by permission of the Government (Permit number EN 0059208).
 Níl sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbhreithniú a déanamh ar theorainneacha na gceantar comharthaíthe. Macasamhail d'ábhar na Suirbhéarachta Ordonáis le chead ón Rialtas (Ceadúnas Uimh. EN 0059208)

SITE CODE
SAC 001482
Version 1.05

Map Version 1
Date: June 2011

SITE CODE
SAC 001482
Version 1.05

Map Version 1
Date: June 2011





Legend

- SAC 001482
- OSi Discovery Series County Boundary

Marine Community Types


- Fine sand dominated by *Nephtys cirrosa* community
- Intertidal sandy mud with *Tubificoides benedii* and *Pygospio elegans* community complex
- Maerl dominated community
- Reef
- Sandy mud with polychaetes and bivalves community complex
- Shingle
- Zostera* community








Legend

- SAC 001482
- 1150 *Coastal Lagoons
- OSi Discovery Series County Boundary



Area A

Area B



Legend

- SAC 001482
- OSi Discovery Series County Boundary

Coastal Habitats

- 1210 Annual vegetation of drift lines (Qualifying Interest)
- 1220 Perennial vegetation of stony banks (Qualifying Interest)
- 2110 Embryonic shifting dunes (Qualifying Interest)
- 2120 Shifting dunes along the shoreline with *Ammophila arenaria* ('white dunes') (Qualifying Interest)
- 21AO Machairs (*in Ireland)
- 2130 *Fixed coastal dunes with herbaceous vegetation ('grey dunes')



**MAP 7:
CLEW BAY COMPLEX
CONSERVATION OBJECTIVES
COASTAL HABITATS**

Map to be read in conjunction with the NPWS Conservation Objectives Document.


COUNTY MAYO

0 0.25 0.5 0.75 1 km

The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision. Reproduced from Ordnance Survey material by permission of the Government (Permit number EN 0059208).
 Níl sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbheithníthe a déanamh ar theorainneacha na gceantar comharthaíthe. Macasamhail d'ábhar na Suirbhéarachta Ordonáis le chead ón Rialtas (Ceadunas Uimh. EN 0059208)


SITE CODE
SAC 001482
Version 1.05

Map Version 1
Date: June 2011




Legend

- SAC 001482
- OSi Discovery Series County Boundary
- 1355 Otter - *Lutra lutra***
- Freshwater Aquatic 80m
- Freshwater Terrestrial 10m
- Marine Aquatic 80m
- Marine Terrestrial 10m
- Freshwater Aquatic Linear Habitat
- Otter Commuting 250m buffer



Legend

- SAC 001482
- ★ 1365 Common (Harbour) Seal - *Phoca vitulina* Breeding Sites
- ★ 1365 Common (Harbour) Seal - *Phoca vitulina* Moulting Sites
- ★ 1365 Common (Harbour) Seal - *Phoca vitulina* Resting Sites
- 1365 Common (Harbour) Seal - *Phoca vitulina* Habitat
- OSi Discovery Series County Boundary





An Roinn
Ealaíon, Oidhreacht agus Gaeltachta

Department of
Arts, Heritage and the Gaeltacht

**Produced by: National Parks and Wildlife Service,
Department of Arts, Heritage and the Gaeltacht,
7 Ely Place, Dublin 2, Ireland.
Web: www.npws.ie
E-mail: natureconservation@environ.ie**

Citation:

NPWS (2011) Conservation Objectives: Clew Bay Complex SAC 001482. Version 1.0. National Parks and Wildlife Service, Department of Arts, Heritage and the Gaeltacht.

Series Editors: Rebecca Jeffrey & Naomi Kingston

ISSN 2009-4086

National Parks and Wildlife Service

Conservation Objectives Series

Doogort Machair/Lough Doo SAC 001497



An Roinn Ealaíon, Oidhreachta,
Gnóthaí Réigiúnacha, Tuaithe agus Gaeltachta

Department of Arts, Heritage,
Regional, Rural and Gaeltacht Affairs



**National Parks and Wildlife Service,
Department of Arts, Heritage, Regional, Rural and Gaeltacht Affairs,
7 Ely Place, Dublin 2, Ireland.
Web: www.npws.ie
E-mail: nature.conservation@ahg.gov.ie**

Citation:

**NPWS (2017) Conservation Objectives: Doogort Machair/Lough Doo SAC
001497. Version 1. National Parks and Wildlife Service, Department of Arts,
Heritage, Regional, Rural and Gaeltacht Affairs.**

**Series Editor: Rebecca Jeffrey
ISSN 2009-4086**

Introduction

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

** indicates a priority habitat under the Habitats Directive*

001497 Doogort Machair/Lough Doo SAC

1395 Petalwort *Petalophyllum ralfsii*

21A0 Machairs (* in Ireland)

Please note that this SAC overlaps with Doogort Machair SPA (004235). See map 2. The conservation objectives for this site should be used in conjunction with those for the overlapping site as appropriate.

Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

NPWS Documents

Year :	1998
Title :	Biomar survey of Irish machair sites 1996
Author :	Crawford, I.; Bleasdale, A.; Conaghan, J.
Series :	Irish Wildlife Manual No. 3
<hr/>	
Year :	1998
Title :	Biomar Survey of Irish machair sites, 1996. Vol. 2: plant communities
Author :	Crawford, I.; Bleasdale, A.; Conaghan, J.
Series :	Irish Wildlife Manual No. 4
<hr/>	
Year :	2009
Title :	Coastal Monitoring Project 2004-2006
Author :	Ryle, T.; Murray, A.; Connolly, K.; Swann, M.
Series :	Unpublished report to NPWS
<hr/>	
Year :	2013
Title :	Monitoring survey of Annex I sand dune habitats in Ireland
Author :	Delaney, A.; Devaney, F.M.; Martin, J.M.; Barron, S.J.
Series :	Irish Wildlife Manual No. 75
<hr/>	
Year :	2015
Title :	Monitoring methods for <i>Petalophyllum ralfsii</i> (Wils.) Nees & Gottsche (Petalwort) in the Republic of Ireland
Author :	Campbell, C.; Hodgetts, N.; Lockhart, N.
Series :	Irish Wildlife Manual No. 90
<hr/>	
Year :	2017
Title :	Doogort Machair/Lough Doo SAC (site code: 1497) Conservation objectives supporting document- coastal habitats V1
Author :	NPWS
Series :	Conservation objectives supporting document

Other References

Year :	2006
Title :	The vegetation of Irish machair
Author :	Gaynor, K.
Series :	Biology and Environment: Proceedings of the Royal Irish Academy, vol 106B, No. 3: 311-321
<hr/>	
Year :	2013
Title :	Conservation of selected legally protected and Red Listed bryophytes in Ireland
Author :	Campbell, C.
Series :	Unpublished Ph.D. Thesis, Trinity College Dublin

Spatial data sources

Year : 2009
Title : Coastal Monitoring Project 2004-2006. Version 1
GIS Operations : QI selected; clipped to SAC boundary; Expert opinion used as necessary to resolve any issues arising
Used For : 21A0 (map 3)

Year : 2016
Title : NPWS rare and threatened species database
GIS Operations : Dataset created from spatial references in database records. Expert opinion used as necessary to resolve any issues arising
Used For : 1395 (map 4)

Conservation Objectives for : Doogort Machair/Lough Doo SAC [001497]

21A0 Machairs (* in Ireland)

To restore the favourable conservation condition of Machairs (* in Ireland) in Doogort Machair/Lough Doo SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes including erosion and succession. For the sub-site mapped: Lough Doo - 88.2ha. See map 3	Based on data from the Coastal Monitoring Project (CMP) (Ryle et al., 2009). Machair habitat was recorded and mapped at the sub-site Lough Doo (CMP site ID: 114) to give a total estimated area of 88.2ha within Doogort Machair/Lough Doo SAC. See the Doogort Machair/Lough Doo SAC conservation objectives supporting document for coastal habitats for further details
Habitat distribution	Occurrence	No decline or change in habitat distribution, subject to natural processes. See map 3 for known distribution	Based on data from Ryle et al. (2009). The machair can be divided into three distinct areas. The first consists of an area of flat, relatively dry machair, which is located between Lough Nambrack and Barnynagappul Strand. The second area consists of wet machair and is found on slopes reaching down towards both Lough Doo and Lough Nambrack. A third area of machair, located in a wide belt at the back of Gubnahardia Strand stretching from Caraun Point to Ridge Point, is composed of a mosaic of wet and dry areas, with low hummocks throughout. See the coastal habitats supporting document for further details
Physical structure: functionality and sediment supply	Presence/absence of physical barriers	Maintain the natural circulation of sediment and organic matter, without any physical obstructions	Based on data from Ryle et al. (2009). Physical barriers can lead to fossilisation or over-stabilisation of dunes, as well as beach starvation resulting in increased rates of erosion. See the coastal habitats supporting document for further details
Physical structure: hydrological and flooding regime	Water table levels; groundwater fluctuations (metres)	Maintain natural hydrological regime	Based on data from Crawford et al. (1998), Gaynor (2006), Ryle et al. (2009) and Delaney et al. (2013). Two small lakes lie at the back of the machair, Lough Doo and Lough Nambrack, with associated freshwater marshes. The flushed slopes adjacent to the lakes are very calcareous and tufa encrustation is evident in places. See the coastal habitats supporting document for further details
Vegetation structure: zonation	Occurrence	Maintain the range of coastal habitats including transitional zones, subject to natural processes including erosion and succession	Based on data from Ryle et al. (2009). This coastal SAC contains a good diversity of habitats in a small area. The inland machair plain grades into blanket bog and is adjoined by two small lakes with freshwater marsh habitat. See the coastal habitats supporting document for further details
Vegetation structure: bare ground	Percentage cover	Bare ground should not exceed 10% of machair habitat, subject to natural processes	Based on data from Ryle et al. (2009). The machair area includes an area of bare sand of approximately 6ha. See the coastal habitats supporting document for further details
Vegetation structure: sward height	Centimetres	Maintain structural variation within sward	Based on data from Ryle et al. (2009). The machair sward is generally close-cropped and overgrazed. It is part of an unenclosed commonage used for grazing sheep and cattle. See the coastal habitats supporting document for further details
Vegetation composition: typical species and sub-communities	Percentage cover at a representative number of monitoring stops	Maintain range of sub-communities with typical species listed in Delaney et al. (2013)	Based on data from Gaynor (2006), Ryle et al. (2009) and Delaney et al. (2013). See the coastal habitats supporting document for further details


Vegetation composition: negative indicator species	Percentage cover	Negative indicator species (including non-native species) to represent less than 5% cover	Based on data from Ryle et al. (2009). Negative indicators include non-native species, species indicative of changes in nutrient status and species not considered characteristic of the habitat. Negative indicators recorded in this SAC include cock's-foot (<i>Dactylis glomerata</i>), common nettle (<i>Urtica dioica</i>) and common ragwort (<i>Senecio jacobaea</i>). See the coastal habitats supporting document for further details
Vegetation composition: scrub/trees	Percentage cover	No more than 5% cover or under control	Based on data from Ryle et al. (2009). See the coastal habitats supporting document for further details
Vegetation composition: bryophytes	Percentage cover	Should always be at least an occasional component of the vegetation	Based on data from Ryle et al. (2009). Doogort Machair/Lough Doo SAC is relatively species-rich and bryophytes are abundant. There is frequently a carpet of mosses and liverworts covering the machair surface. Petalwort (<i>Petalophyllum ralfsii</i>), a liverwort species listed on Annex II of the EU Habitats Directive, has also been recorded on the machair in this SAC. See the conservation objective for petalwort (1395) and the coastal habitats supporting document for further details

Conservation Objectives for : Doogort Machair/Lough Doo SAC [001497]


1395 Petalwort *Petalophyllum ralfsii*


To maintain the favourable conservation condition of Petalwort in Doogort Machair/Lough Doo SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Distribution	Number and geographical spread of populations	No decline. See map 4 for recorded locations	The known population of <i>Petalophyllum ralfsii</i> in Doogort Machair/Lough Doo SAC is at Doogort Machair. It has been recorded from three locations, where it occurs on compact sandy soil on tightly grazed low sandhills and hummocks on the machair. Data from NPWS surveys and Campbell et al. (2015)
Population size	Number of individuals	No decline. The population at Doogort Machair is estimated to be at least 258 thalli	Campbell and Lockhart recorded 43 <i>Petalophyllum ralfsii</i> thalli (in October 2010) in an area of 25cm x 50cm at one location. Therefore the population is estimated to be a minimum of 258 thalli in 0.75m ² . Numbers of thalli can vary from year to year. See Campbell et al. (2015) for further details
Area of suitable habitat	Hectares	No decline. Area of suitable habitat at Doogort Machair currently unknown, but is estimated to be at least c.0.00008ha	The extent of suitable habitat at Doogort Machair has not been accurately measured using GPS, but is estimated to be at least of 0.75m ² (c.0.00008ha), i.e. a minimum area of 50cm x 50cm at each of the three mapped points. This is an underestimate and suitable habitat within the SAC is likely to be more widespread. See Campbell et al. (2015) for further details
Hydrological conditions: soil moisture	Occurrence of damp soil conditions	Maintain hydrological conditions so that substrate is kept moist and damp throughout the year, but is not subject to prolonged inundation by flooding in winter	<i>Petalophyllum ralfsii</i> grows on damp sandy substrate. Based on Campbell (2013) and Campbell et al. (2015)
Vegetation: open structure	Height and percentage cover of vegetation	Maintain open, low vegetation, with a high percentage cover of bryophytes (small acrocarps and liverwort turf) and bare ground	At Doogort Machair, <i>Petalophyllum ralfsii</i> grows in compacted, sandy ground, maintained by heavy sheep grazing and trampling (by sheep). See Campbell et al. (2015) for further details



Legend

 Doogort Machair/Lough Doo SAC 001497

 An Roinn Ealaíon, Oidhreacht, Gnóthaí Réigiúnacha, Tuaithe agus Gaeltachta
 Department of Arts, Heritage, Regional, Rural and Gaeltacht Affairs

MAP 1:
DOOGORT MACHAIR / LOUGH DOO SAC
CONSERVATION OBJECTIVES
SAC DESIGNATION

Map to be read in conjunction with the NPWS Conservation Objectives Document.


SITE CODE:
SAC 001497; version 3.01.
CO. MAYO

0 0.15 0.3 0.45 0.6 0.75 km

The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
 Ordnance Survey of Ireland Licence No EN 0059216. © Ordnance Survey of Ireland Government of Ireland


Níl sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbhreithniú a déanamh ar theorainneacha na gceantar comharthaite. Suirbhéarachta Ordonáis na hÉireann Ceadúnas


Map Version 1
Date: Oct 2016




Legend


 Doogort Machair/Lough Doo SAC 001497

 Coastal Monitoring Project Survey Area


Annex I Qualifying Interest

 21AO Machair

Annex I Non qualifying Interest

 2120 Shifting dunes along the shoreline with *Ammophila arenaria* ('white dunes')

 OSi Discovery Series County Boundary



CMP Site Code: 114




An Roinn Ealaíon, Oidhreachta,
Gnóthaí Réigiúnacha, Tuaithe agus Gaeltachta
Department of Arts, Heritage,
Regional, Rural and Gaeltacht Affairs

**MAP 3:
DOOGORT MACHAIR / LOUGH DOO SAC
CONSERVATION OBJECTIVES
MACHAIR**

Map to be read in conjunction with the NPWS Conservation Objectives Document.

**SITE CODE:
SAC 001497; version 3.01.
CO. MAYO**




The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
Ordnance Survey of Ireland Licence No EN 0059216. © Ordnance Survey of Ireland Government of Ireland

Níl sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbheithnithe a déanamh ar theorainneacha na gceantar comharthaite. Suirbhéarachta Ordonáis na hÉireann Ceadúnas



**Map Version 1
Date: Oct 2016**



Legend

- Doogort Machair/Lough Doo SAC 001497
- 1395 Petalwort *Petalophyllum ralfsii*
- OSi Discovery Series County Boundary

National Parks and Wildlife Service

Conservation Objectives Series

Erris Head SAC 001501



An Roinn Ealaíon, Oidhreachta,
Gnóthaí Réigiúnacha, Tuaithe agus Gaeltachta

Department of Arts, Heritage,
Regional, Rural and Gaeltacht Affairs



**National Parks and Wildlife Service,
Department of Arts, Heritage, Regional, Rural and Gaeltacht Affairs,
7 Ely Place, Dublin 2, Ireland.
Web: www.npws.ie
E-mail: nature.conservation@ahg.gov.ie**

Citation:

**NPWS (2016) Conservation Objectives: Erris Head SAC 001501. Version 1.
National Parks and Wildlife Service, Department of Arts, Heritage, Regional,
Rural and Gaeltacht Affairs.**

**Series Editor: Rebecca Jeffrey
ISSN 2009-4086**

Introduction

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

** indicates a priority habitat under the Habitats Directive*

001501	Erris Head SAC
1230	Vegetated sea cliffs of the Atlantic and Baltic coasts
4060	Alpine and Boreal heaths

Please note that this SAC overlaps with Mullet Peninsula SPA (004227) and Termoncarragh Lake and Annagh Machair SPA (004093) and adjoins Broadhaven Bay SAC (000472) and West Connaght Coast SAC (002998). See map 2. The conservation objectives for this site should be used in conjunction with those for the overlapping and adjacent sites as appropriate.

Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

NPWS Documents

Year :	2011
Title :	National survey and assessment of the conservation status of Irish sea cliffs
Author :	Barron, S.J.; Delaney, A.; Perrin, P.M.; Martin, J.; O'Neill, F.
Series :	Irish Wildlife Manual No. 53
Year :	2012
Title :	Ireland Red List no. 8: Bryophytes
Author :	Lockhart, N.; Hodgetts, N.; Holyoak, D.
Series :	Ireland Red List series, NPWS
Year :	2013
Title :	The status of EU protected habitats and species in Ireland. Volume 2. Habitats assessments
Author :	NPWS
Series :	Conservation assessments
Year :	2014
Title :	Guidelines for a national survey and conservation assessment of upland vegetation and habitats in Ireland, Version 2.0
Author :	Perrin, P.M.; Barron, S.J.; Roche, J.R.; O'Hanrahan, B.
Series :	Irish Wildlife Manual No. 79
Year :	2016
Title :	Erris Head SAC (site code: 1501) Conservation objectives supporting document- coastal habitats V1
Author :	NPWS
Series :	Conservation objectives supporting document

Other References

Year :	1986
Title :	An outline of the flora of Mayo
Author :	Synnott, D.
Series :	Glasra, 9: 13-117
Year :	1988
Title :	The Irish red data book 1. Vascular plants
Author :	Curtis, T.G.F.; McGough, H.N.
Series :	Wildlife Service, Dublin
Year :	2005
Title :	National inventory of sea cliffs and coastal heaths
Author :	Browne, A.
Series :	Unpublished Report to NPWS

Spatial data sources

Year : 2011
Title : National survey and assessment of the conservation status of Irish sea cliffs
GIS Operations : Clipped to SAC boundary
Used For : 1230 (map 3)

1230 Vegetated sea cliffs of the Atlantic and Baltic coasts

To maintain the favourable conservation condition of Vegetated sea cliffs of the Atlantic and Baltic coasts in Erris Head SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat length	Kilometres	Area stable, subject to natural processes, including erosion. For the sub-site mapped (Glenlara), total length of cliff sections: 24.12km. See map 3	Based on data from the Irish Sea Cliff Survey (ISCS) (Barron et al., 2011). Cliffs are linear features and are therefore measured in kilometres. The sub-site Glenlara (site ID: 08007) was identified using a combination of aerial photos and the DCENR helicopter viewer. The length of cliff was measured (in sections) to give a total estimated area of 24.12km within the SAC. The length of cliff is likely to be underestimated. See the Erris Head SAC conservation objectives supporting document for coastal habitats for further details
Habitat distribution	Occurrence	No decline, subject to natural processes. See map 3	Sea cliffs are known to occur along the coastline from Illaunbaun to Termoncarragh. Hard cliffs have been noted in this SAC and it is thought that all of the cliffs are of the hard type (Browne, 2005; Barron et al., 2011). See the coastal habitats supporting document for further details
Physical structure: functionality and hydrological regime	Occurrence of artificial barriers	No alteration to natural functioning of geomorphological and hydrological processes, including groundwater quality, due to artificial structures	Based on data from Barron et al. (2011). Maintaining natural geomorphological processes, including natural erosion, is important for the health of vegetated sea cliffs. Hydrological processes maintain flushes, and in some cases tufa formations, that can be associated with sea cliffs. Hydrological features such as flushes were noted at Erris Head SAC. See the coastal habitats supporting document for further details
Vegetation structure: zonation	Occurrence	Maintain range of sea cliff habitat zonations including transitional zones, subject to natural processes including erosion and succession	Based on data from Barron et al. (2011). Alpine and Boreal heath and coastal grassland occur adjacent to sea cliff vegetation at Erris Head SAC. See the coastal habitats supporting document for further details
Vegetation structure: vegetation height	Centimetres	Maintain structural variation within sward	Based on data from Barron et al. (2011). Land use in the SAC consists mainly of sheep grazing. See the coastal habitats supporting document for further details
Vegetation composition: typical species and sub-communities	Percentage cover at a representative number of monitoring stops	Maintain range of sub-communities with typical species listed in the Irish Sea Cliff Survey (Barron et al., 2011)	The presence of species such as roseroot (<i>Rhodiola rosea</i>) and rock sea-spurrey (<i>Spergularia rupicola</i>) suggest that there is typical cliff vegetation for such an exposed site. See the coastal habitats supporting document for further details
Vegetation composition: negative indicator species	Percentage	Negative indicator species (including non-native species) to represent less than 5% cover	Based on data from Barron et al. (2011). See the coastal habitats supporting document for further details
Vegetation composition: bracken and woody species	Percentage	Cover of bracken (<i>Pteridium aquilinum</i>) on grassland and/or heath less than 10%. Cover of woody species on grassland and/or heath less than 20%	Based on data from Barron et al. (2011). See the coastal habitats supporting document for further details



Conservation Objectives for : Erris Head SAC [001501]

4060 Alpine and Boreal heaths


To maintain the favourable conservation condition of Alpine and Boreal heaths in Erris Head SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Alpine and Boreal heath has not been mapped in detail for Erris Head SAC and thus the total area of qualifying habitat is unknown. It occurs inland from the tops of the sea cliffs and grades into coastal grassland in places (NPWS internal files)
Habitat distribution	Occurrence	No decline, subject to natural processes	See note on area above
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil nutrient status within natural range	Relevant nutrients and their natural ranges are yet to be defined. However, nitrogen deposition is noted as being relevant to this habitat (NPWS, 2013)
Community diversity	Abundance of variety of vegetation communities	Maintain variety of vegetation communities, subject to natural processes	The diversity of Alpine and Boreal heath communities within this SAC is unknown. Information on vegetation communities associated with this habitat is presented in Perrin et al. (2014)
Vegetation composition: lichens and bryophytes	Number of species at a representative number of 2m x 2m monitoring stops	Number of bryophyte or non-crustose lichen species present at each monitoring stop is at least three	Attribute and target based on Perrin et al. (2014). Alpine and Boreal heath is not necessarily rich in lichen and bryophyte species, but a minimum amount should still be present
Vegetation composition: positive indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of positive indicator species at least 66%	Attribute and target based on Perrin et al. (2014), where the list of positive indicator species for this habitat is also presented. Bearberry (<i>Arctostaphylos uva-ursi</i>), crowberry (<i>Empetrum nigrum</i>) and juniper (<i>Juniperus communis</i>) are listed as being present in the Alpine and Boreal heath in this SAC (NPWS internal files; Synnott, 1986)
Vegetation composition: dwarf shrub species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of dwarf shrub species at least 10%	Attribute and target based on Perrin et al. (2014). A lower cover of dwarf shrubs could indicate that the habitat is transitioning to another vegetation type such as grassland
Vegetation composition: negative indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of negative indicator species less than 10%	Attribute and target based on Perrin et al. (2014), where the list of negative indicator species for this habitat is also presented
Vegetation composition: non-native species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of non-native species less than 1%	Attribute and target based on Perrin et al. (2014). Non-native species can be invasive and have deleterious effects on native vegetation. A low target is set as non-native species can spread rapidly and are most easily dealt with when still at lower abundances
Vegetation structure: signs of grazing	Percentage of leaves browsed at a representative number of 2m x 2m monitoring stops	Less than 10% collectively of the live leaves of specific graminoids showing signs of grazing	Attribute and target based on Perrin et al. (2014). The specific graminoids are stiff sedge (<i>Carex bigelowii</i>), wavy hair-grass (<i>Deschampsia flexuosa</i>), sheep's-fescue (<i>Festuca ovina</i>) and viviparous sheep's-fescue (<i>Festuca vivipara</i>). High levels of grazing of these species would be undesirable as grazing is not required to maintain this habitat
Vegetation structure: signs of browsing	Percentage of shoots browsed at a representative number of 2m x 2m monitoring stops	Less than 33% collectively of the last complete growing season's shoots of ericoids and crowberry (<i>Empetrum nigrum</i>) showing signs of browsing	Attribute and target based on Perrin et al. (2014)
Vegetation structure: burning	Occurrence in local vicinity of a representative number of monitoring stops	No signs of burning within the habitat	Attribute and target based on Perrin et al. (2014). Alpine and Boreal heath does not require burning for the maintenance of the habitat







Physical structure: disturbed bare ground	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of disturbed bare ground less than 10%	Attribute and target based on Perrin et al. (2014). Disturbance can include hoof marks, wallows, human foot prints and vehicle and machinery tracks. Excessive disturbance can result in loss of characteristic species and presage erosion for heaths and peatlands
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat	This includes species listed in the Flora (Protection) Order, 2015 and/or the red data lists (Curtis and McGough, 1988; Lockhart et al., 2012)






Legend

 Erris Head SAC 001501

Legend

-  Termoncarragh Lake and Annagh Machair SPA 004093
-  Mullet Peninsula SPA 004227
-  Erris Head SAC 001501
-  West Connacht Coast SAC 002998
-  Broadhaven Bay SAC 000472
-  OSi Discovery Series County Boundary





Legend

- 1230 Vegetated sea cliffs of the Atlantic and Baltic coasts
- Erris Head SAC 001501
- OSi Discovery Series County Boundary

National Parks and Wildlife Service

Conservation Objectives Series

Keel Machair/Menaun Cliffs SAC 001513



An Roinn
Cultúir, Oidhreachta agus Gaeltachta
Department of
Culture, Heritage and the Gaeltacht



**National Parks and Wildlife Service,
Department of Culture, Heritage and the Gaeltacht,
90 King Street North, Dublin 7, D07 N7CV, Ireland.**

**Web: www.npws.ie
E-mail: nature.conservation@chg.gov.ie**

Citation:

**NPWS (2018) Conservation Objectives: Keel Machair/Menaun Cliffs SAC
001513. Version 1. National Parks and Wildlife Service, Department of Culture,
Heritage and the Gaeltacht.**

**Series Editor: Rebecca Jeffrey
ISSN 2009-4086**

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

** indicates a priority habitat under the Habitats Directive*

001513 Keel Machair/Menaun Cliffs SAC

1220 Perennial vegetation of stony banks

1395 Petalwort *Petalophyllum ralfsii*

21A0 Machairs (* in Ireland)

4060 Alpine and Boreal heaths

**Please note that this SAC is adjacent to Achill Head SAC (002268).
See map 2. The conservation objectives for this site should be used in
conjunction with those for the adjacent site as appropriate.**

Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

NPWS Documents

Year :	1998
Title :	Biomar survey of Irish machair sites 1996
Author :	Crawford, I.; Bleasdale, A.; Conaghan, J.
Series :	Irish Wildlife Manual No. 3
Year :	2009
Title :	Coastal Monitoring Project 2004-2006
Author :	Ryle, T.; Murray, A.; Connolly, K.; Swann, M.
Series :	Unpublished report to NPWS
Year :	2012
Title :	Ireland Red List No. 8: Bryophytes
Author :	Lockhart, N.; Hodgetts, N.; Holyoak, D.
Series :	Ireland Red List series, NPWS
Year :	2013
Title :	Monitoring survey of Annex I sand dune habitats in Ireland
Author :	Delaney, A.; Devaney, F.M.; Martin, J.M.; Barron, S.J.
Series :	Irish Wildlife Manual No. 75
Year :	2013
Title :	The status of EU protected habitats and species in Ireland. Volume 2. Habitats assessments
Author :	NPWS
Series :	Conservation assessments
Year :	2014
Title :	Guidelines for a national survey and conservation assessment of upland vegetation and habitats in Ireland, Version 2.0
Author :	Perrin, P.M.; Barron, S.J.; Roche, J.R.; O'Hanrahan, B.
Series :	Irish Wildlife Manual No. 79
Year :	2015
Title :	Monitoring methods for <i>Petalophyllum ralfsii</i> (Wils.) Nees & Gottsche (Petalwort) in the Republic of Ireland
Author :	Campbell, C.; Hodgetts, N.; Lockhart, N.
Series :	Irish Wildlife Manual No. 90
Year :	2016
Title :	Ireland Red List No. 10: Vascular Plants
Author :	Wyse Jackson, M.; FitzPatrick, Ú.; Cole, E.; Jebb, M.; McFerran, D.; Sheehy Skeffington, M.; Wright, M.
Series :	Ireland Red Lists series, NPWS
Year :	2017
Title :	Survey and assessment of vegetated shingle and associated habitats at 30 coastal sites in Ireland
Author :	Martin, J.R.; Daly, O.H.; Devaney F.M.
Series :	Irish Wildlife Manual No. 98
Year :	2018
Title :	Keel Machair/Menaun Cliffs SAC (site code: 1513) Conservation objectives supporting document: coastal habitats V1
Author :	NPWS
Series :	Conservation objectives supporting document

Other References

- Year :** 2006
Title : The vegetation of Irish machair
Author : Gaynor, K.
Series : Biology and Environment: Proceedings of the Royal Irish Academy, 106B(3): 311-321
-
- Year :** 2008
Title : The phytosociology and conservation value of Irish sand dunes
Author : Gaynor, K.
Series : Unpublished Ph.D. Thesis, National University of Ireland, Dublin
-
- Year :** 2013
Title : Conservation of selected legally protected and Red Listed bryophytes in Ireland
Author : Campbell, C.
Series : Unpublished Ph.D. Thesis, Trinity College Dublin
-

Spatial data sources

Year : 2017
Title : Vegetated Shingle Monitoring Project
GIS Operations : QIs selected; clipped to SAC boundary. Expert opinion used as necessary to resolve any issues arising
Used For : 1220, 21A0 (map 3)

Year : 2017
Title : NPWS rare and threatened species database
GIS Operations : Dataset created from spatial references in database records. Expert opinion used as necessary to resolve any issues arising
Used For : 1395 (map 4)

1220 Perennial vegetation of stony banks

To maintain the favourable conservation condition of Perennial vegetation of stony banks in Keel Machair/Menaun Cliffs SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes, including erosion and succession	Based on data from the Vegetated Shingle Monitoring Project (VSM) (Martin et al., 2017). Perennial vegetation of stony banks was not recorded in the sub-site Trawmore, Keel (VSM site code 006) during the VSM and thus the total area of the qualifying habitat within Keel Machair/Menaun Cliffs SAC is unknown. Martin et al. (2017) did note extensive areas of shingle beach in the sub-site during the VSM; however, no areas of the shingle were vegetated. NB further unsurveyed areas may be present within the SAC. See the Keel Machair/Menaun Cliffs SAC conservation objectives supporting document for coastal habitats for further details
Habitat distribution	Occurrence	No decline, subject to natural processes, including erosion and succession	The full distribution of the habitat in the SAC is unknown at present. It is possible that the habitat on the shingle beach in the Trawmore, Keel sub-site has been temporarily lost due to natural erosion and will re-establish in the future (Martin et al., 2017). See the coastal habitats supporting document for further details. NB further unsurveyed areas may be present within the SAC
Physical structure: functionality and sediment supply	Presence/absence of physical barriers	Maintain, or where necessary restore, the natural circulation of sediment and organic matter, without any physical obstructions	Rock armour is present within the Trawmore, Keel sub-site, but its impact on the formation of the habitat is considered to be minimal (Martin et al., 2017). See the coastal habitats supporting document for further details
Physical structure: disturbance	Percentage	No more than 20% of the habitat affected by disturbance	Based on data from Martin et al. (2017). Disturbance can include damage from heavy trampling, vehicle damage and removal of substrate. See the coastal habitats supporting document for further details
Vegetation structure: zonation	Occurrence	Maintain the range of coastal habitats, including transitional zones, subject to natural processes, including erosion and succession	Based on data from Martin et al. (2017). See the coastal habitats supporting document for further details
Vegetation composition: communities and typical species	Occurrence	Maintain the typical species within the range of vegetated shingle communities	Based on data from Martin et al. (2017) where information on the vegetated shingle communities and associated typical species lists are presented. See the coastal habitats supporting document for further details
Vegetation composition: negative indicator species	Percentage	Negative indicator species cover in any individual monitoring stop should not be more than 25%; no negative indicator species should be present in more than 60% of monitoring stops	Based on data from Martin et al. (2017) where the list of negative indicator species for the habitat is also presented. Negative indicators include species indicative of changes in nutrient status and species not considered characteristic of the habitat. See the coastal habitats supporting document for further details

Vegetation composition: non-native species	Percentage	Non-native species cover in any individual monitoring stop should not be more than 1%; non-native species should not be present in more than 20% of monitoring stops; cover of non-native species across the whole site should not be more than 1%	Based on data from Martin et al. (2017). See the coastal habitats supporting document for further details
--	------------	--	---

Conservation Objectives for : Keel Machair/Menaun Cliffs SAC [001513]

21A0 Machairs (* in Ireland)

To restore the favourable conservation condition of Machairs* in Keel Machair/Menaun Cliffs SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes, including erosion and succession. For the sub-site mapped: Trawmore, Keel - 79.52ha. See map 3	Based on data from the Vegetated Shingle Monitoring Project (VSM) (Martin et al., 2017). Machair habitat was surveyed and mapped at the sub-site Trawmore, Keel (VSM site code 006) to give a total estimated area of 79.52ha within Keel Machair/Menaun Cliffs SAC. See the Keel Machair/Menaun Cliffs SAC conservation objectives supporting document for coastal habitats for further details
Habitat distribution	Occurrence	No decline or change in habitat distribution, subject to natural processes. See map 3 for recorded distribution	Based on data from Martin et al. (2017). The habitat extends along the coast from Keel village in the west to just short of the foothills of Menaun cliffs in the south-east, and landwards to the shores of Keel Lough. See the coastal habitats supporting document for further details
Physical structure: functionality and sediment supply	Presence/absence of physical barriers	Maintain the natural circulation of sediment and organic matter, without any physical obstructions	Dunes are naturally dynamic systems that require continuous supply and circulation of sand. See the coastal habitats supporting document for further details
Physical structure: hydrological and flooding regime	Water table levels; groundwater fluctuations (metres)	Maintain natural hydrological regime	Based on Crawford et al. (1998), Gaynor (2006, 2008), Ryle et al. (2009) and Martin et al. (2017). See the coastal habitats supporting document for further details
Vegetation structure: zonation	Occurrence	Maintain the range of coastal habitats including transitional zones, subject to natural processes, including erosion and succession	Based on Ryle et al. (2009), Delaney et al. (2013) and Martin et al. (2017). See the coastal habitats supporting document for further details
Vegetation structure: bare ground	Percentage cover	Bare ground should not exceed 5% of machair habitat, subject to natural processes	Based on Delaney et al. (2013) and Martin et al. (2017). See the coastal habitats supporting document for further details
Vegetation structure: sward height	Centimetres	Maintain structural variation within sward. The mean sward height should be at least 8cm in July/August	Based on Delaney et al. (2013) and Martin et al. (2017). A mean vegetation height of 2.2cm was recorded by the VSM in the habitat in the Trawmore, Keel sub-site indicating that grazing levels are too high. See the coastal habitats supporting document for further details
Vegetation composition: flowering/fruitleting	Percentage	Positive indicator species should be flowering or fruiting in more than 40% of monitoring stops	Based on Delaney et al. (2013) and Martin et al. (2017). The list of positive indicator species is presented in Delaney et al. (2013). See the coastal habitats supporting document for further details
Vegetation composition: typical species and sub-communities	Occurrence	Maintain range of sub-communities with typical species listed in Delaney et al. (2013)	See the coastal habitats supporting document for further details
Vegetation composition: negative indicator species	Percentage	Negative indicator species cover in any individual monitoring stop should not be more than 25%; no negative indicator species should be present in more than 60% of monitoring stops; cover of negative indicator species across the whole site should not be more than 5%	Based on Delaney et al. (2013) and Martin et al. (2017). Negative indicators include species indicative of changes in nutrient status and species not considered characteristic of the habitat. See the coastal habitats supporting document for further details

Vegetation composition: non-native species	Percentage	Non-native species should not be present in more than 20% of monitoring stops	Based on Delaney et al. (2013) and Martin et al. (2017). The non-native species New Zealand willowherb (<i>Epilobium brunnescens</i>) was recorded by the VSM within the machair habitat in the Trawmore, Keel sub-site. See coastal habitats supporting document for further details
Vegetation composition: scrub/trees	Percentage cover	No more than 5% cover or under control	Based on Delaney et al. (2013) and Martin et al. (2017). See the coastal habitats supporting document for further details
Vegetation composition: bryophytes	Percentage cover	Should always be at least an occasional component of the vegetation, with a minimum cover of 1% within each monitoring stop	Based on Ryle et al. (2009), Delaney et al. (2013) and Martin et al. (2017). The Annex II and Flora (Protection) Order, 2015 listed liverwort petalwort (<i>Petalophyllum ralfsii</i>) has been recorded from the machair in the SAC (see Campbell et al., 2015). See the coastal habitats supporting document for further details. See also the conservation objective for petalwort (1395) in this volume

Conservation Objectives for : Keel Machair/Menaun Cliffs SAC [001513]

4060 Alpine and Boreal heaths

To maintain the favourable conservation condition of Alpine and Boreal heaths in Keel Machair/Menaun Cliffs SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Alpine and Boreal heath has not been mapped in detail for Keel Machair/Menaun Cliffs SAC and thus the total area of qualifying habitat is unknown. It is documented that the habitat occurs around the summit of Menaun in a mosaic with blanket bog with numerous rock outcrops. At lower altitudes, the habitat merges with extensive areas of dry heath dominated by ling (<i>Calluna vulgaris</i>) (NPWS internal files)
Habitat distribution	Occurrence	No decline, subject to natural processes	See the notes on Habitat area above
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil pH and nutrient status within natural ranges	Relevant nutrients and their natural ranges are yet to be defined. However, nitrogen deposition is noted as being relevant to this habitat (NPWS, 2013)
Community diversity	Abundance of variety of vegetation communities	Maintain variety of vegetation communities, subject to natural processes	The entire diversity of Alpine and Boreal heath communities within this SAC is unknown. Information on vegetation communities associated with this habitat is presented in Perrin et al. (2014)
Vegetation composition: lichens and bryophytes	Number of species at a representative number of 2m x 2m monitoring stops	Number of bryophyte or non-crustose lichen species present at each monitoring stop is at least three	Attribute and target based on Perrin et al. (2014). Alpine and Boreal heath is not necessarily rich in lichen and bryophyte species, but a minimum amount should still be present
Vegetation composition: positive indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of positive indicator species at least 66%	Attribute and target based on Perrin et al. (2014), where the list of positive indicator species for this habitat is also presented. Bearberry (<i>Arctostaphylos uva-ursi</i>), crowberry (<i>Empetrum nigrum</i>), juniper (<i>Juniperus communis</i> subsp. <i>nana</i>), bilberry (<i>Vaccinium myrtillus</i>) and the Near Threatened dwarf willow (<i>Salix herbacea</i>) (Wyse Jackson et al., 2016) have been reported from the Alpine and Boreal heath in this SAC (NPWS internal files)
Vegetation composition: dwarf shrub species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of dwarf shrub species at least 10%	Attribute and target based on Perrin et al. (2014). A lower cover of dwarf shrubs could indicate that the habitat is transitioning to another vegetation type such as grassland
Vegetation composition: negative indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of negative indicator species less than 10%	Attribute and target based on Perrin et al. (2014), where the list of negative indicator species for this habitat is also presented
Vegetation composition: non-native species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of non-native species less than 1%	Attribute and target based on Perrin et al. (2014). Non-native species can be invasive and have deleterious effects on native vegetation. A low target is set as non-native species can spread rapidly and are most easily dealt with when still at lower abundances
Vegetation structure: signs of grazing	Percentage of leaves grazed at a representative number of 2m x 2m monitoring stops	Less than 10% collectively of the live leaves of specific graminoids showing signs of grazing	Attribute and target based on Perrin et al. (2014). The specific graminoids are stiff sedge (<i>Carex bigelowii</i>), wavy hair-grass (<i>Deschampsia flexuosa</i>), sheep's-fescue (<i>Festuca ovina</i>) and viviparous sheep's-fescue (<i>Festuca vivipara</i>). High levels of grazing of these species would be undesirable as grazing is not required to maintain this habitat
Vegetation structure: signs of browsing	Percentage of shoots browsed at a representative number of 2m x 2m monitoring stops	Less than 33% collectively of the last complete growing season's shoots of ericoids and crowberry (<i>Empetrum nigrum</i>) showing signs of browsing	Attribute and target based on Perrin et al. (2014)


Vegetation structure: burning	Occurrence in local vicinity of a representative number of monitoring stops	No signs of burning within the habitat	Attribute and target based on Perrin et al. (2014). Alpine and Boreal heath does not require burning for the maintenance of the habitat
Physical structure: disturbed bare ground	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of disturbed bare ground less than 10%	Attribute and target based on Perrin et al. (2014). Disturbance can include hoof marks, wallows, human footprints and vehicle and machinery tracks. Excessive disturbance can result in loss of characteristic species and presage erosion for heaths and peatlands
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat and no decline in status of hepatic mats associated with this habitat	This includes species listed in the Flora (Protection) Order, 2015 and/or the red data lists (Lockhart et al., 2012; Wyse Jackson et al., 2016). The Near Threatened dwarf willow (<i>Salix herbacea</i>) (Wyse Jackson et al., 2016) has been reported from the habitat in the SAC (NPWS internal files). Part of the heath vegetation on Menaun contains Northern Atlantic hepatic mat communities (NPWS internal files)

Conservation Objectives for : Keel Machair/Menaun Cliffs SAC [001513]


1395 Petalwort *Petalophyllum ralfsii*


To maintain the favourable conservation condition of Petalwort in Keel Machair/Menaun Cliffs SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Distribution	Number and geographical spread of populations	No decline, subject to natural processes. See map 4 for recorded locations	The known population of petalwort (<i>Petalophyllum ralfsii</i>) in Keel Machair/Menaun Cliffs SAC occurs at Keel Machair on tightly sheep-grazed turf on the edges of channelised and semi-natural water tracks and on small, partly bare patches of damp unshaded sand on the western and more calcareous side of the machair plain. Data from NPWS surveys (NPWS internal files). See Campbell et al. (2015) for further details
Area of suitable habitat	Hectares	No decline, subject to natural processes	The extent of suitable habitat at Keel Machair has not yet been accurately measured using GPS, but is estimated to be c.10,300m ² (1.03ha) based on NPWS surveys (NPWS internal files). See Campbell et al. (2015) for further details
Hydrological conditions: soil moisture	Occurrence of damp soil conditions	Maintain hydrological conditions so that the substrate is kept moist and damp throughout the year, but is not subject to prolonged inundation by flooding in winter	Petalwort (<i>Petalophyllum ralfsii</i>) grows on damp sandy substrate. Based on Campbell (2013) and Campbell et al. (2015)
Hydrological conditions: water table level	Centimetres in a representative number of 1m x 1m monitoring plots	Mean groundwater level should not be more than 80cm from ground surface	See Campbell et al. (2015) for further details
Physical structure: bare soil	Percentage cover in a representative number of 1m x 1m monitoring plots	Mean percentage cover of bare soil should be more than 5%	At Keel Machair, petalwort (<i>Petalophyllum ralfsii</i>) grows in compacted, sandy ground. There is some threat from potential over-use by vehicles, dumping and mowing at this location (NPWS internal files). See Campbell et al. (2015) for further details
Vegetation structure: vegetation height	Centimetres in a representative number of 1m x 1m monitoring plots	Mean vegetation height should be less than 6cm	Very short vegetation, heavily grazed by sheep, has been reported in petalwort (<i>Petalophyllum ralfsii</i>) habitat at Keel Machair (NPWS internal files). See Campbell et al. (2015) for further details
Vegetation composition: shrub cover	Percentage cover in a representative number of 1m x 1m monitoring plots	Mean percentage shrub cover should be less than 25%	See Campbell et al. (2015) for further details
Vegetation composition: grass cover	Percentage cover in a representative number of 1m x 1m monitoring plots	Mean percentage grass species cover should be less than 60%	Part of this site is managed as a 9-hole pitch and putt course; low intensity management, with only the greens and tees re-seeded, has enabled petalwort (<i>Petalophyllum ralfsii</i>) to survive. Any intensification or expansion should be discouraged (NPWS internal files). See Campbell et al. (2015) for further details



Legend

 Keel Machair/Menaun Cliffs SAC 001513



An Roinn
Cultúir, Oidhreacht agus Gaeltachta
Department of
Culture, Heritage and the Gaeltacht

MAP 1:
KEEL MACHAIR / MENAUN CLIFFS SAC
CONSERVATION OBJECTIVES
SAC DESIGNATION

Map to be read in conjunction with the NPWS Conservation Objectives Document.

SITE CODE:
SAC 001513; version 3.02
CO. MAYO

0 0.5 1 1.5 2 km


The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
Ordnance Survey of Ireland Licence No EN 0059216. © Ordnance Survey of Ireland Government of Ireland.

Níl sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbhreithnithe a déanamh ar theorainneacha na gceantar comharthaithe. Suirbhéarachtá Ordonáis na hÉireann Ceadúnas Uimh EN 0059216. © Suirbhéarachtá Ordonáis na hÉireann Rialtas na hÉireann.

N




Map Version 1
Date: Aug 2017




Legend

- Keel Machair/Menaun Cliffs SAC 001513
- Achill Head SAC 002268

Legend


 Keel Machair/Menaun Cliffs SAC 001513




Annex I Qualifying Interest

 21A0* Machair

Annex I Non Qualifying Interest

 2120 Shifting dunes along the shoreline with *Ammophila arenaria* ("white dunes")

 2130* Fixed dunes with herbaceous vegetation ("grey dunes")




An Roinn
Cultúir, Oidhreacht agus Gaeltachta
Department of
Culture, Heritage and the Gaeltacht

**MAP 3:
KEEL MACHAIR/MENAUN CLIFFS SAC
CONSERVATION OBJECTIVES
COASTAL HABITATS**

Map to be read in conjunction with the NPWS Conservation Objectives Document.

SITE CODE:
SAC 001513; version 3.02,
CO. MAYO



The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
Ordnance Survey of Ireland Licence No EN 0059216. © Ordnance Survey of Ireland Government of Ireland.


Níl sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbheithnithe a déanamh ar theorainneacha na gceantar comharthaíthe. Suirbhéarachta Ordonáis na hÉireann Ceadúnas Uimh EN 0059216. © Suirbhéarachta Ordonáis na hÉireann Riailtas na hÉireann.



Map Version 1
Date: Aug 2017

Legend

- Keel Machair/Menaun Cliffs SAC 001513
- ★ 1395 Petalwort *Petalophyllum ralfsii*



National Parks and Wildlife Service

Conservation Objectives Series

Lough Cahasy, Lough Baun and Roonah Lough SAC
001529



An Roinn Ealaíon, Oidhreachta,
Gnóthaí Réigiúnacha, Tuaithe agus Gaeltachta

Department of Arts, Heritage,
Regional, Rural and Gaeltacht Affairs



**National Parks and Wildlife Service,
Department of Arts, Heritage, Regional, Rural and Gaeltacht Affairs,
7 Ely Place, Dublin 2, Ireland.
Web: www.npws.ie
E-mail: nature.conservation@ahg.gov.ie**

Citation:

**NPWS (2017) Conservation Objectives: Lough Cahasy, Lough Baun and Roonah
Lough SAC 001529. Version 1. National Parks and Wildlife Service, Department
of Arts, Heritage, Regional, Rural and Gaeltacht Affairs.**

**Series Editor: Rebecca Jeffrey
ISSN 2009-4086**

Introduction

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

** indicates a priority habitat under the Habitats Directive*

001529	Lough Cahasy, Lough Baun and Roonah Lough SAC
1150	Coastal lagoonsE
1220	Perennial vegetation of stony banks
2120	Shifting dunes along the shoreline with Cl { [] @ ^ } ^ ^ (white dunes)

Please note that this SAC adjoins West Connacht Coast SAC (002998). See map 2. The conservation objectives for this site should be used in conjunction with those for the adjacent site as appropriate.

Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

NPWS Documents

Year :	1999
Title :	National Shingle Beach Survey of Ireland 1999
Author :	Moore, D.; Wilson, F.
Series :	Unpublished Report to NPWS
<hr/>	
Year :	2007
Title :	Inventory of Irish coastal lagoons (version 2)
Author :	Oliver, G.
Series :	Unpublished report to NPWS
<hr/>	
Year :	2009
Title :	Coastal Monitoring Project 2004-2006
Author :	Ryle, T.; Murray, A.; Connolly, K.; Swann, M.
Series :	Unpublished report to NPWS
<hr/>	
Year :	2013
Title :	Monitoring survey of Annex I sand dune habitats in Ireland
Author :	Delaney, A.; Devaney, F.M.; Martin, J.M.; Barron, S.J.
Series :	Irish Wildlife Manual No. 75
<hr/>	
Year :	2017
Title :	Lough Cahasy, Lough Baun and Roonah Lough SAC (site code: 1529) Conservation objectives supporting document- coastal lagoons V1
Author :	NPWS
Series :	Conservation objectives supporting document
<hr/>	
Year :	2017
Title :	Lough Cahasy, Lough Baun and Roonah Lough SAC (site code: 1529) Conservation objectives supporting document- coastal habitats V1
Author :	NPWS
Series :	Conservation objectives supporting document

Other References

Year :	2008
Title :	The phytosociology and conservation value of Irish sand dunes
Author :	Gaynor, K.
Series :	Unpublished PhD thesis, National University of Ireland, Dublin
<hr/>	
Year :	2013
Title :	Monitoring and assessment of Irish lagoons for the purposes of the EU Water Framework Directive, 2009-2011. Parts 1 and 2
Author :	Roden, C.M; Oliver, G.A.
Series :	Unpublished report to the Environmental Protection Agency

Spatial data sources

Year :	Revision 2011
Title :	Inventory of Irish Coastal Lagoons. Version 3
GIS Operations :	Clipped to SAC boundary
Used For :	1150 (map 3)
<hr/>	
Year :	Revision 2012
Title :	National Shingle Beach Survey
GIS Operations :	Clipped to SAC boundary. Expert opinion used as necessary to resolve any issues arising
Used For :	1220 (map 4)
<hr/>	
Year :	2009
Title :	Coastal Monitoring Project 2004-2006. Version 1
GIS Operations :	QIs selected; clipped to SAC boundary. Expert opinion used as necessary to resolve any issues arising
Used For :	1220, 2120 (map 4)
<hr/>	

1150 Coastal lagoons

To restore the favourable conservation condition of Coastal lagoons* in Lough Cahasy, Lough Baun and Roonah Lough SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable, subject to slight natural variation. Favourable reference area: 40.2ha. See map 3	Area calculated from spatial data derived from Oliver (2007) for Roonah Lough (site code IL072). See the Lough Cahasy, Lough Baun and Roonah Lough SAC conservation objectives supporting document for coastal lagoons for further details
Habitat distribution	Occurrence	No decline, subject to natural processes. See map 3 for mapped lagoon	Site Roonah Lough (site code IL072) in Oliver (2007). See the lagoon supporting document for further details
Salinity regime	Practical salinity units (psu)	Median annual salinity and temporal variation within natural ranges	Roonah Lough is recorded as an oligohaline lagoon. See the lagoon supporting document for further details
Hydrological regime	Metres	Annual water level fluctuations and minima within natural ranges	The maximum depth of Roonah Lough lagoon is recorded as less than 1m. See the lagoon supporting document for further details
Barrier: connectivity between lagoon and sea	Permeability	Appropriate hydrological connections between lagoons and sea, including where necessary, appropriate management	Roonah Lough is described as a sedimentary lagoon with a cobble barrier. See the lagoon supporting document for further details
Water quality: Chlorophyll <i>a</i>	µg/L	Annual median chlorophyll <i>a</i> within natural ranges and less than 5µg/L	Target based on Roden and Oliver (2013). See the lagoon supporting document for further details
Water quality: Molybdate Reactive Phosphorus (MRP)	mg/L	Annual median MRP within natural ranges and less than 0.1mg/L	Target based on Roden and Oliver (2013). See the lagoon supporting document for further details
Water quality: Dissolved Inorganic Nitrogen (DIN)	mg/L	Annual median DIN within natural ranges and less than 0.15mg/L	Target based on Roden and Oliver (2013). See the lagoon supporting document for further details
Depth of macrophyte colonisation	Metres	Macrophyte colonisation to maximum depth of lagoon	Where the lagoon is less than 2m deep, it is expected that macrophyte colonisation would extend to the full depth. See the lagoon supporting document for further details
Typical plant species	Number and m ²	Maintain number and extent of listed lagoonal specialists, subject to natural variation	Species listed in Oliver (2007). See the lagoon supporting document for further details
Typical animal species	Number	Maintain listed lagoon specialists, subject to natural variation	Species listed in Oliver (2007). See the lagoon supporting document for further details
Negative indicator species	Number and percentage cover	Negative indicator species absent or under control	Low salinity, shallow water and elevated nutrient levels increase the threat of unnatural encroachment by reedbeds. See the lagoon supporting document for further details

Conservation Objectives for : Lough Cahasy, Lough Baun and Roonah Lough SAC [001529]

1220 Perennial vegetation of stony banks

To maintain the favourable conservation condition of Perennial vegetation of stony banks in Lough Cahasy, Lough Baun and Roonah Lough SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes, including erosion and succession	During the National Shingle Beach Survey (NSBS) (Moore and Wilson, 1999) perennial vegetation of stony banks was recorded as being present, but its extent was not mapped, from two sub-sites, White Strand (NSBS site ID: 0054) and Sruhir Strand (NSBS Site ID: 0055). Thus, the current area of the qualifying habitat within Lough Cahasy, Lough Baun and Roonah Lough SAC is unknown. During the Coastal Monitoring Project (CMP) (Ryle et al., 2009), an area of vegetated shingle was noted within the sub-site Lough Cahasy (CMP site ID: 109). NB further unsurveyed areas may be present within the SAC. See the Lough Cahasy, Lough Baun and Roonah Lough SAC conservation objectives supporting document for coastal habitats for further details
Habitat distribution	Occurrence	No decline or change in habitat distribution, subject to natural processes. See map 4 for recorded locations	The full distribution of perennial vegetation of stony banks within the SAC is unknown at present, although the habitat has been recorded at White Strand and Sruhir Strand by Moore and Wilson (1999) and within the Lough Cahasy sub-site by Ryle et al. (2009). See map 4 for the NSBS point locations. NB further unsurveyed areas may be present within the SAC. See the coastal habitats supporting document for further details
Physical structure: functionality and sediment supply	Presence/absence of physical barriers	Maintain the natural circulation of sediment and organic matter, without any physical obstructions	Based on data from Moore and Wilson (1999) and Ryle et al. (2009). The shingle beaches at White Strand and Sruhir Strand consist of an extensive and high boulder/shingle ridge along much of the shoreline, most of which is largely unvegetated. See the coastal habitats supporting document for further details
Vegetation structure: zonation	Occurrence	Maintain the range of coastal habitats including transitional zones, subject to natural processes, including erosion and succession	Based on data from Moore and Wilson (1999). See the coastal habitats supporting document for further details
Vegetation composition: typical species and sub-communities	Percentage cover at a representative number of monitoring stops	Maintain the typical vegetated shingle flora including the range of sub-communities within the different zones	Based on data from Moore and Wilson (1999) and Ryle et al. (2009). The shingle beaches at White Strand and Sruhir Strand have sparse vegetation due to the exposed location and dynamic nature. The occurrence of typical species was infrequent, only occurring as scattered individuals. See the coastal habitats supporting document for further details
Vegetation composition: negative indicator species	Percentage cover	Negative indicator species (including non-native species) to represent less than 5% cover	Based on data from Moore and Wilson (1999). Negative indicators include non-native species indicative of changes in nutrient status and species not considered characteristic of the habitat. See the coastal habitats supporting document for further details



Conservation Objectives for : Lough Cahasy, Lough Baun and Roonah Lough SAC [001529]

2120 Shifting dunes along the shoreline with *Ammophila arenaria* (white dunes)

To maintain the favourable conservation condition of Shifting dunes along the shoreline with *Ammophila arenaria* (white dunes) in Lough Cahasy, Lough Baun and Roonah Lough SAC, which is defined by the following list of attributes and targets:


Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes, including erosion and succession. For the sub-site mapped: Lough Cahasy - 0.9ha. See map 4	Based on data from the Coastal Monitoring Project (CMP) (Ryle et al., 2009). Shifting dunes along the shoreline with <i>Ammophila arenaria</i> was mapped at the sub-site Lough Cahasy (CMP site ID: 109) to give a total estimated area of 0.9ha within Lough Cahasy, Lough Baun and Roonah Lough SAC. The habitat is very difficult to measure in view of its dynamic nature. See the Lough Cahasy, Lough Baun and Roonah Lough SAC conservation objectives supporting document for coastal habitats for further details
Habitat distribution	Occurrence	No decline or change in habitat distribution, subject to natural processes. See map 4 for known distribution	Based on data from Ryle et al. (2009). The mobile dunes are not extensive and are mainly confined to a narrow band along Gortnagarryan Strand. See the coastal habitats supporting document for further details
Physical structure: functionality and sediment supply	Presence/absence of physical barriers	Maintain the natural circulation of sediment and organic matter, without any physical obstructions	Based on data from Ryle et al. (2009). Dunes are naturally dynamic systems that require continuous supply and circulation of sand. Marram grass (<i>Ammophila arenaria</i>) reproduces vegetatively and requires constant accretion of fresh sand to maintain active growth encouraging further accretion. The CMP considers that the mobile dunes in the SAC will persist owing to the continuous input of fresh sand from longshore drift. See the coastal habitats supporting document for further details
Vegetation structure: zonation	Occurrence	Maintain the range of coastal habitats including transitional zones, subject to natural processes including erosion and succession	Based on data from Gaynor (2008) and Ryle et al. (2009). See the coastal habitats supporting document for further details
Vegetation composition: plant health of dune grasses	Percentage cover	More than 95% of marram grass (<i>Ammophila arenaria</i>) and/or lyme-grass (<i>Leymus arenarius</i>) should be healthy (i.e. green plant parts above ground and flowering heads present)	Based on data from Ryle et al. (2009). See the coastal habitats supporting document for further details
Vegetation composition: typical species and sub-communities	Percentage cover at a representative number of monitoring stops	Maintain the presence of species-poor communities dominated by marram grass (<i>Ammophila arenaria</i>) and/or lyme-grass (<i>Leymus arenarius</i>)	Based on data from Ryle et al. (2009). See the coastal habitats supporting document for further details
Vegetation composition: negative indicator species	Percentage cover	Negative indicator species (including non-native species) to represent less than 5% cover	Based on data from Gaynor (2008), Ryle et al. (2009) and Delaney et al. (2013). Negative indicators include non-native species, species indicative of changes in nutrient status and species not considered characteristic of the habitat. Sea buckthorn (<i>Hippophae rhamnoides</i>) should be absent or effectively controlled. The occurrence of the negative indicator species common ragwort (<i>Senecio jacobea</i>) in the mobile dunes in the SAC was negligible. See the coastal habitats supporting document for further details





Legend


- Lough Cahasy, Lough Baun and Roonah Lough SAC 001529
- West Connacht Coast SAC 002998




Irish Lagoon Inventory, SITE CODE: IL072

Legend

- Lough Cahasy, Lough Baun and Roonah Lough SAC 001529
- 1150 *Coastal lagoons





Legend

- Lough Cahasy, Lough Baun and Roonah Lough SAC 001529
- Coastal Monitoring Project Survey Area

Annex I Qualifying Interests

- ★ 1220 Perennial vegetation of stony banks
- 2120 Shifting dunes along the shoreline with *Ammophila arenaria* (white dunes)

Annex I Non-qualifying Interests

- 1210 Annual vegetation of drift lines
- 2110 Embryonic shifting dunes
- 2130 *Fixed dunes with herbaceous vegetation ('grey dunes')
- 21AO *Machair
- OSi Discovery Series County Boundary

National Parks and Wildlife Service

Conservation Objectives Series

Mocorha Lough SAC 001536



An Roinn Cultúir,
Oidhreacht agus Gaeltachta
Department of Culture,
Heritage and the Gaeltacht

**National Parks and Wildlife Service,
Department of Culture, Heritage and the Gaeltacht,
90 King Street North, Dublin 7, D07 N7CV, Ireland.**

**Web: www.npws.ie
E-mail: nature.conservation@chg.gov.ie**

Citation:

**NPWS (2019) Conservation Objectives: Mocarha Lough SAC 001536. Version 1.
National Parks and Wildlife Service, Department of Culture, Heritage and the
Gaeltacht.**

**Series Editor: Rebecca Jeffrey
ISSN 2009-4086**

Introduction

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

** indicates a priority habitat under the Habitats Directive*

001536 Mocarha Lough SAC

7210 Calcareous fens with *Cladium mariscus* and species of the Caricion davallianae*

Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

NPWS Documents

Year :	1979
Title :	A Preliminary Report on Areas of Scientific Interest in County Mayo
Author :	Goodwillie, R.N.
Series :	Unpublished Report
Year :	2009
Title :	Ireland Red List No. 2: Non-marine molluscs
Author :	Byrne, A.; Moorkens, E.A.; Anderson, R.; Killeen, I.J.; Regan, E.C.
Series :	Ireland Red List series, NPWS
Year :	2010
Title :	Ireland Red List No. 4: Butterflies
Author :	Regan, E.C.; Nelson, B.; Aldwell, B.; Bertrand, C.; Bond, K.; Harding, J.; Nash, D.; Nixon, D.; Wilson, C.J.
Series :	Ireland Red List series, NPWS
Year :	2012
Title :	Ireland Red List No. 8: Bryophytes
Author :	Lockhart, N.; Hodgetts, N.; Holyoak, D.
Series :	Ireland Red List series, NPWS
Year :	2013
Title :	Conservation status assessments for three fen habitat types - 7230 – Alkaline fens, 7210 – Calcareous fens with <i>Cladium mariscus</i> and species of <i>Caricion davallianae</i> and 7140 – Transition mires and quaking bogs
Author :	Kimberley, S.
Series :	Unpublished report to NPWS
Year :	2014
Title :	Guidelines for a national survey and conservation assessment of upland vegetation and habitats in Ireland, Version 2.0
Author :	Perrin, P.M.; Barron, S.J.; Roche, J.R.; O'Hanrahan, B.
Series :	Irish Wildlife Manuals, No. 79
Year :	2016
Title :	Ireland Red List No. 10: Vascular Plants
Author :	Wyse Jackson, M.; FitzPatrick, Ú.; Cole, E.; Jebb, M.; McFerran, D.; Sheehy Skeffington, M.; Wright, M.
Series :	Ireland Red List Series, NPWS
Year :	2018
Title :	Backing document – National Conservation Status Assessments (NCAs) for three fen habitat types: 7140 – Transition mires and quaking bogs, 7210 – Calcareous fens with <i>Cladium mariscus</i> and species of <i>Caricion davallianae</i> , 7230 – Alkaline fens
Author :	Long, M.P.; Crowe, O.; Kimberley, S.; Denyer, J.
Series :	Unpublished report to NPWS
Year :	2019
Title :	The Status of EU Protected Habitats and Species in Ireland. Volume 2: Habitat Assessments
Author :	NPWS
Series :	Conservation assessments

Other References

Year : 2004
Title : Common Standards Monitoring guidance for lowland wetland habitats
Author : JNCC
Series : Joint Nature Conservation Committee, Peterborough


Conservation Objectives for : Macorha Lough SAC [001536]

7210 Calcareous fens with *Cladium mariscus* and species of the *Caricion davallianae**

To maintain the favourable conservation condition of Calcareous fens with *Cladium mariscus* and species of the *Caricion davallianae in Macorha Lough SAC, which is defined by the following list of attributes and targets:**

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Calcareous fens with <i>Cladium mariscus</i> and species of the <i>Caricion davallianae</i> * has not been mapped in detail for Macorha Lough SAC and thus the total current area of the priority qualifying habitat in the SAC is unknown. The SAC supports a good example of <i>Cladium</i> fen in a calcareous lake basin which is considered one of the largest stands in the west of Ireland. It occurs in association with common reed (<i>Phragmites australis</i>) swamp, black bog-rush (<i>Schoenus nigricans</i>) dominated fen and other wetland vegetation. Wet grassland, which floods at times, is found adjacent to the wetland habitats. The lake basin is mostly overgrown with the swamp and fen vegetation, with very little open water remaining (Goodwillie, 1979; NPWS internal files)
Habitat distribution	Occurrence	No decline, subject to natural processes	See the notes for Habitat area above
Ecosystem function: peat formation	Percentage cover of peat-forming vegetation and water table levels	Maintain active peat formation, where appropriate	In order for peat to form, water levels need to be slightly below or above the soil surface for c.90% of the time
Ecosystem function: hydrology - groundwater levels	Water levels (centimetres); duration of levels; hydraulic gradients	Maintain, or where necessary restore, appropriate natural hydrological regimes necessary to support the natural structure and functioning of the habitat	Fen habitats require high groundwater levels (i.e. water levels at or above the ground surface) for a large proportion of the calendar year (i.e. duration of mean groundwater level). Fen groundwater levels are controlled by regional groundwater levels in the contributing catchment area (which sustain the hydraulic gradients of the fen groundwater table). Regional abstraction of groundwater may affect fen groundwater levels
Ecosystem function: hydrology - surface water flow	Drain density and form	Maintain, or where necessary restore, as close as possible to natural or semi-natural, drainage conditions	Drainage, either within or surrounding the fen habitat, can result in the drawdown of the alkaline fen groundwater table. The depth, geometry and density of drainage (hydromorphology) will indicate the scale and impact on fen hydrology. Drainage can result in loss of characteristic species and transition to drier habitats
Ecosystem function: water quality	Water chemistry measures	Maintain appropriate water quality, particularly pH and nutrient levels, to support the natural structure and functioning of the habitat	Fens receive natural levels of nutrients (e.g. iron, magnesium and calcium) from water sources. However, they are generally poor in nitrogen and phosphorus, with the latter tending to be the limiting nutrient under natural conditions. Water supply should be also relatively calcium-rich
Vegetation composition: typical species	Percentage cover at a representative number of 2m x 2m monitoring stops	Maintain adequate cover of typical species including brown mosses and vascular plants	For lists of typical vascular plant and bryophyte species, including high quality indicator species, see the Article 17 conservation status assessment for <i>Cladium</i> fens (NPWS, 2019) and the fen habitats supporting document (Long et al., 2018)
Vegetation composition: native negative indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of native negative indicator species at insignificant levels	Negative indicators include species not characteristic of the habitat and species indicative of undesirable activities such as overgrazing, undergrazing, nutrient enrichment, agricultural improvement or impacts on hydrology. See JNCC (2004) and Kimberley (2013)
Vegetation composition: non-native species	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of non-native species less than 1%	Attribute and target based on Perrin et al. (2014). Non-native species can be invasive and have deleterious effects on native vegetation. A low target is set as non-native species can spread rapidly and are most easily dealt with when still at lower abundances

Vegetation composition: trees and shrubs	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of scattered native trees and shrubs less than 10%	Attribute and target based on Perrin et al. (2014). Scrub and trees will tend to invade if fen conditions become drier
Physical structure: disturbed bare ground	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of disturbed bare ground not more than 10%. Where tufa is present, disturbed bare ground not more than 1%	Attribute and target based on Perrin et al. (2014). While grazing may be appropriate in this habitat, excessive areas of disturbed bare ground may develop due to unsuitable grazing regimes. Disturbance can include hoof marks, wallows, human footprints, vehicle and machinery tracks. Excessive disturbance can result in loss of characteristic species and presage erosion for peatlands
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat; maintain features of local distinctiveness, subject to natural processes	This includes species on the Flora (Protection) Order, 2015 and/or Red Lists (Byrne et al., 2009; Regan et al., 2010; Lockhart et al., 2012; Wyse Jackson et al., 2016, etc.)



Legend
 Macorha Lough SAC 001536

An Roinn Cultúir,
 Oidhreacht agus Gaeltachta
 Department of Culture,
 Heritage and the Gaeltacht

**MAP 1:
 MOCORHA LOUGH SAC
 CONSERVATION OBJECTIVES
 SAC DESIGNATION**
 Map to be read in conjunction with the NPWS Conservation Objectives Document.

**SITE CODE:
 SAC 001536; version 3.01. CO. MAYO**

0 110 220 330 440 Meters

The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
 Ordnance Survey Ireland Licence No OSI-NMA-014. © Ordnance Survey Ireland Government of Ireland

Níl sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbheithnithe a déanamh ar theorainneacha na gceantar comharthaíthe. Suirbhéarachtá Ordonáis na hÉireann Ceadúnas Uimh OSI-NMA-014. © Suirbhéarachtá Ordonáis na hÉireann Rialtas na hÉireann

N
Map Version 1
Date: Feb 2019

National Parks and Wildlife Service

Conservation Objectives Series

Urlaur Lakes SAC 001571



An Roinn
Cultúir, Oidhreachta agus Gaeltachta

Department of
Culture, Heritage and the Gaeltacht



**National Parks and Wildlife Service,
Department of Culture, Heritage and the Gaeltacht,
7 Ely Place, Dublin 2, Ireland.
Web: www.npws.ie
E-mail: nature.conservation@ahg.gov.ie**

Citation:

**NPWS (2017) Conservation Objectives: Urlaur Lakes SAC 001571. Version 1.
National Parks and Wildlife Service, Department of Culture, Heritage and the
Gaeltacht.**

**Series Editor: Rebecca Jeffrey
ISSN 2009-4086**

Introduction

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

** indicates a priority habitat under the Habitats Directive*

001571 Urlaur Lakes SAC

3140 Hard oligo-mesotrophic waters with benthic vegetation of *O⁺l⁺ac* spp.

Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

NPWS Documents

Year :	2013
Title :	A survey of the benthic macrophytes of three hard-water lakes: Lough Bunny, Lough Carra and Lough Owel
Author :	Roden, C.; Murphy, P.
Series :	Irish Wildlife Manual No. 70
Year :	2013
Title :	The status of EU protected habitats and species in Ireland. Volume 2. Habitats assessments
Author :	NPWS
Series :	Conservation assessments
Year :	2015
Title :	Habitats Directive Annex I lake habitats: a working interpretation for the purposes of site-specific conservation objectives and Article 17 reporting
Author :	O Connor, Á.
Series :	Unpublished document by NPWS

Other References

Year :	1982
Title :	Eutrophication of waters. Monitoring assessment and control
Author :	OECD
Series :	OECD, Paris
Year :	2000
Title :	Colour in Irish lakes
Author :	Free, G.; Allott, N.; Mills, P.; Kennelly, C.; Day, S.
Series :	Verhandlungen Internationale Vereinigung für theoretische und angewandte Limnologie, 27: 2620-2623
Year :	2006
Title :	A reference-based typology and ecological assessment system for Irish lakes. Preliminary investigations. Final report. Project 2000-FS-1-M1 Ecological assessment of lakes pilot study to establish monitoring methodologies EU (WFD)
Author :	Free, G.; Little, R.; Tierney, D.; Donnelly, K.; Coroni, R.
Series :	EPA, Wexford
Year :	2015
Title :	Water quality in Ireland 2010-2012
Author :	Bradley, C.; Byrne, C.; Craig, M.; Free, G.; Gallagher, T.; Kennedy, B.; Little, R.; Lucey, J.; Mannix, A.; McCreesh, P.; McDermott, G.; McGarrigle, M.; Ní Longphuirt, S.; O'Boyle, S.; Plant, C.; Tierney, D.; Trodd, W.; Webster, P.; Wilkes, R.; Wynne, C.
Series :	EPA, Wexford
Year :	in prep.
Title :	Monitoring of hard-water lakes in Ireland using charophytes and other macrophytes
Author :	Roden, C.; Murphy, P.
Series :	Unpublished report to NPWS

Spatial data sources

Year : 2008

Title : OSi 1:5000 IG vector dataset

GIS Operations : WaterPolygons feature class clipped to the SAC boundary. Expert opinion used to identify Annex I habitat and to resolve any issues arising

Used For : 3140 (map 2)

Conservation Objectives for : Urlaur Lakes SAC [001571]

3140 Hard oligo-mesotrophic waters with benthic vegetation of Chara spp.


To restore the favourable conservation condition of Hard oligo-mesotrophic waters with benthic vegetation of *Chara* spp. in Urlaur Lakes SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Urlaur Lakes SAC contains three marl lakes with habitat 3140 (Urlaur Lough, Lough Nanoge and Lough Roe). The vegetation of Urlaur Lough was surveyed in 2012 and assessed as being in poor conservation condition (Roden and Murphy, in prep.). Urlaur Lough is on the Water Framework Directive (WFD) monitoring programme and regular macrophyte surveys are conducted by the Environmental Protection Agency (EPA). Two measures of extent should be used: 1. the area of the lake itself and; 2. the extent of the vegetation communities/zones that typify the habitat. Further information relating to all attributes is provided in the lake habitats supporting document for the purposes of site-specific conservation objectives and Article 17 reporting (O Connor, 2015)
Habitat distribution	Occurrence	No decline, subject to natural processes	As noted above, lake habitat 3140 occurs in Urlaur Lough, Lough Nanoge and Lough Roe in the SAC. See map 2
Vegetation composition: typical species	Occurrence	Typical species present, in good condition, and demonstrating typical abundances and distribution	For lists of 3140 typical species (cyanobacteria, algae, higher plants and water beetles), see the Article 17 habitat assessment for lake habitat 3140 (NPWS, 2013) and the lake habitats supporting document (O Connor, 2015). Roden and Murphy (in prep.) recorded krustenstein, <i>Chara aspera</i> , <i>C. contraria</i> , <i>C. curta</i> , <i>C. virgata</i> , <i>Ophrydium versatile</i> , <i>Baldellia ranunculoides</i> , <i>Callitriche hermaphroditica</i> , <i>Eleocharis palustris</i> , <i>Elodea canadensis</i> , <i>Fontinalis antipyretica</i> , <i>Hippuris vulgaris</i> , <i>Lemna trisulca</i> , <i>Littorella uniflora</i> , <i>Potamogeton berchtoldii</i> , <i>P. perfoliatus</i> , <i>Ranunculus flammula</i> , <i>Utricularia minor</i> and <i>U. vulgaris</i> in Urlaur Lough. NPWS site files also note <i>Cladium mariscus</i> , <i>Equisetum fluviatile</i> , <i>Nuphar lutea</i> , <i>Nymphaea alba</i> , <i>Phragmites australis</i> and <i>Schoenoplectus lacustris</i> in the SAC (NPWS internal files)
Vegetation composition: characteristic zonation	Occurrence	All characteristic zones should be present, correctly distributed and in good condition	The characteristic zonation of lake habitat 3140 is described in Roden and Murphy (2013). Urlaur Lough had few charophyte bands in 2012: <i>Chara contraria</i> or <i>C. curta</i> and <i>C. virgata</i> extending to c.2m; common <i>Elodea canadensis</i> at the base of the euphotic zone and well-developed krustenstein on occasional boulders (Roden and Murphy, in prep.)
Vegetation distribution: maximum depth	Metres	Restore maximum depth of vegetation, subject to natural processes	Maximum vegetation depth is expected to be deep in clear, hard water lakes, and extremely clear marl lakes can have charophyte vegetation to more than 9m (e.g. Lough Rea has charophytes to 10-11m, Coolorta >9m) (Roden and Murphy, in prep.). The indicative target of >6m for lake habitat 3140 may need to be modified based on the habitat sub-type/form and/or the specific lake in question (Roden and Murphy, 2013, in prep.). In this SAC, the maximum depth of vegetation at Urlaur Lough was very shallow at 2-2.1m in 2012 (Roden and Murphy, in prep.). The water is highly coloured in Urlaur Lough and this may contribute to the limited vegetation development. Areas of drained peatland (for turf-cutting, conifer forest, agricultural use) in the catchments of Urlaur Lough, Lough Nanoge and Lough Roe are likely to artificially increase the lakes' water colour


Hydrological regime: water level fluctuations	Metres	Maintain appropriate hydrological regime necessary to support the habitat	The hydrological regime of lakes with habitat 3140 is driven by groundwater flows. Groundwater can discharge directly to the lake, via springs or seepages, or to in-flowing rivers. Fluctuations in lake water level are typical in Ireland, but can be amplified by activities such as abstraction and drainage. Increased water level fluctuations can increase wave action and turbidity, up-root vegetation, alter the substratum and lead to nutrient release from sediment. The hydrological regime, particularly the groundwater contribution, must be maintained so that the area, distribution and depth of the lake habitat and its constituent/characteristic vegetation zones and communities are not reduced
Lake substratum quality	Various	Maintain appropriate substratum type, extent and chemistry to support the vegetation	The hard water lake habitat (3140) is associated with a range of base-rich substratum types, from marl and limestone bedrock, through rocks, cobbles, gravel, muds and even peat. Further research into substratum quality (notably calcium, iron and nutrient concentrations) in the hard water lake habitat would be beneficial. Roden and Murphy (in prep.) recorded sand, mud and occasional shoreline boulders at Urlaur Lough. The EPA have recorded cobble and gravel on the shoreline and silt at depth in Urlaur Lough. NPWS site files note stony, gravelly and sandy shorelines at Lough Nanoge (NPWS internal files)
Water quality: transparency	Metres	Restore appropriate Secchi transparency. There should be no decline in Secchi depth/transparency	Transparency relates to light penetration and, hence, to the depth of colonisation of vegetation. It can be affected by phytoplankton blooms, water colour and turbidity. A target of >6m has been set for hard water lakes (3140) (Roden and Murphy, in prep.). The OECD fixed boundary system set transparency targets for oligotrophic lakes of $\geq 6\text{m}$ annual mean Secchi disk depth and $\geq 3\text{m}$ annual minimum Secchi disk depth. Hard water lakes typically have high transparency, particularly in the very clear and typical marl forms (Roden and Murphy, in prep.). Secchi depth at Urlaur Lough was 1.5m in 2001 (Free et al., 2006) and 1.54m in 2012 (Roden and Murphy, in prep.)
Water quality: nutrients	$\mu\text{g/l P}$; mg/l N	Restore the concentration of nutrients in the water column to sufficiently low levels to support the habitat and its typical species	Lake habitat 3140 is typically associated with high water quality, as demonstrated by naturally low dissolved nutrients. The target for Loughs Urlaur, Nanoge and Roe is WFD High Status or oligotrophic (OECD, 1982). Annual average total phosphorus (TP) concentration should be $\leq 10\mu\text{g/l TP}$, average annual total ammonia concentration should be $\leq 0.04\text{mg/l N}$ and annual 95th percentile for total ammonia should be $\leq 0.09\text{mg/l N}$. Where nutrient concentrations are lower than the targets, there should be no upward trend in concentrations. See also The European Communities Environmental Objectives (Surface Waters) Regulations 2009. Free et al. (2006) measured $< 10\mu\text{g/l TP}$ in Urlaur Lough in April 2001. Urlaur Lough failed to reach the target in 2010-12, however, having good nutrient status (Bradley et al., 2015)


Water quality: phytoplankton biomass	µg/l Chlorophyll <i>a</i>	Maintain/restore appropriate water quality to support the habitat, including high chlorophyll <i>a</i> status	Lake habitat 3140 is associated with high water quality, as demonstrated by naturally low algal growth. As for nutrients, the default target is WFD High Status or oligotrophic (OECD, 1982). Average growing season (March-October) chlorophyll <i>a</i> concentration must be <5.8µg/l. Annual average chlorophyll <i>a</i> concentration should be <2.5µg/l and the annual peak should be <8.0µg/l. Where chlorophyll <i>a</i> concentrations are lower than the targets, there should be no upward trend in phytoplankton biomass. See also The European Communities Environmental Objectives (Surface Waters) Regulations 2009. In Urlaur Lough, Free et al. (2006) measured 7.7µg/l chlorophyll <i>a</i> in April 2001 and chlorophyll <i>a</i> status was high in 2010-12 (Bradley et al., 2015)
Water quality: phytoplankton composition	EPA phytoplankton composition metric	Restore appropriate water quality to support the habitat, including high phytoplankton composition status	The EPA has developed a phytoplankton composition metric for nutrient enrichment of Irish lakes. As for other water quality indicators, the default target for lake habitat 3140 is WFD high status. Urlaur Lough failed to reach the target in 2010-12, having good phytoplankton composition status (Bradley et al., 2015)
Water quality: attached algal biomass	Algal cover and EPA phytobenthos metric	Maintain/restore trace/absent attached algal biomass (<5% cover) and high phytobenthos status	Nutrient enrichment can favour epiphytic and epipelic algae that can out-compete the submerged vegetation. The cover abundance of attached algae in hard water lakes (3140) should, therefore, be trace/absent (<5% cover). EPA phytobenthos status can be used as an indicator of changes in attached algal biomass. As for other water quality indicators, the default target for lake habitat 3140 is high phytobenthos status. Phytobenthos status was high in Urlaur Lough in 2010-12 (Bradley et al., 2015); however, filamentous algae were recorded by the EPA in the lake in 2001 and 2011
Water quality: macrophyte status	EPA macrophyte metric (The Free Index)	Restore high macrophyte status	Nutrient enrichment can favour more competitive submerged macrophyte species that out-compete the typical and characteristic species for hard water lakes (3140). The EPA monitors macrophyte status for WFD purposes using the 'Free Index'. The target for lake habitat 3140 is high status or an Ecological Quality Ratio (EQR) for lake macrophytes of ≥0.90, as defined in Schedule Five of the European Communities Environmental Objectives (Surface Waters) Regulations 2009. Urlaur Lough failed to reach the target in 2010-12, having good macrophyte status (Bradley et al., 2015)
Acidification status	pH units; mg/l	Maintain appropriate water and sediment pH, alkalinity and cation concentrations to support the habitat, subject to natural processes	The specific requirements of lake habitat 3140, in terms of water and sediment pH, alkalinity and cation concentration, have not been fully determined. Acidification is not considered a threat to lake habitat 3140; however, eutrophication can lead to at least temporary increases in pH to toxic levels (>9/9.5 pH units). Maximum pH should be <9.0 pH units, in line with the surface water standards. See Schedule Five of the European Communities Environmental Objectives (Surface Waters) Regulations 2009
Water colour	mg/l PtCo	Maintain/restore appropriate water colour to support the habitat	Increased colour decreases light penetration and reduces the area of macrophyte habitat, particularly at the lower euphotic depths. Higher colour also appears to favour angiosperms over charophytes in hard water lakes (Roden and Murphy, in prep.). The primary source of increased colour in Ireland is peatland disturbance. No habitat-specific or national standards for water colour exist. Studies have shown median colour concentrations in Irish lakes of 38mg/l PtCo (Free et al., 2000) and 33mg/l PtCo (Free et al., 2006). Lake habitat 3140 is typically associated with very clear waters and expected colour would be <10mg/l PtCo or, more likely, <5mg/l PtCo. Free et al. (2006) recorded colour of 33mg/l PtCo in Urlaur Lough

Dissolved organic carbon (DOC)	mg/l	Maintain/restore appropriate organic carbon levels to support the habitat	Dissolved (and particulate) organic carbon (OC) in the water column is linked to water colour and acidification (organic acids). Increasing DOC in water has been documented across the Northern Hemisphere, including afforested peatland catchments in Ireland. Damage and degradation of peatland, leading to decomposition of peat is likely to be the predominant source of OC in Ireland. OC in water promotes decomposition by fungi and bacteria that, in turn, releases dissolved nutrients. The increased biomass of decomposers can also impact directly on the characteristic lake communities through shading, competition, etc. As noted above, increased water colour, low transparency and shallow vegetation zones at Urlaur Lough may be linked to peatland disturbance
Turbidity	Nephelometric turbidity units/ mg/l SS/ other appropriate unit	Maintain appropriate turbidity to support the habitat	Turbidity can significantly affect the quantity and quality of light reaching rooted and attached vegetation and can, therefore, impact on lake habitats. The settlement of higher loads of inorganic or organic material on lake vegetation communities may also have impacts on sensitive, delicate species. Turbidity can increase as a result of re-suspension of material within the lake, higher loads entering the lake, or eutrophication. Turbidity measurement and interpretation is challenging. As a result, it is likely to be difficult to set habitat-specific targets for turbidity in lakes
Fringing habitat: area and condition	Hectares	Maintain the area and condition of fringing habitats necessary to support the natural structure and functioning of habitat 3140	Most lake shorelines have fringing habitats of reedswamp, other swamp, fen, marsh or wet woodland that intergrade with and support the structure and functions of the lake habitat. Equally, fringing habitats are dependent on the lake, particularly its water levels, and support wetland communities and species of conservation concern. Many of the fringing wetland habitats support higher invertebrate and plant species richness than the lake habitats themselves. Fringing fen habitats can be particularly important around hard water lakes, notably the Annex I habitats alkaline fen, <i>Cladium</i> fen and petrifying springs (habitat codes 7230, 7210 and 7220). Reedbeds (<i>Schoenoplectus lacustris</i> , <i>Phragmites australis</i>), swamp (<i>Carex rostrata</i> , <i>Cladium mariscus</i> , <i>Typha latifolia</i> , <i>Equisetum fluviatile</i>), fen/flush, heath, revegetated cutaway bog, scrub and calcareous grassland have been recorded on the shores of Urlaur, Nanoge and Roe. Transition mire/quaking bog may also occur



Legend

 Urlaur Lakes SAC 001571



An Roinn
Cultúir, Oidhreacht agus Gaeltachta
Department of
Culture, Heritage and the Gaeltacht

**MAP 1:
URLAUR LAKES SAC
CONSERVATION OBJECTIVES
SAC DESIGNATION**

Map to be read in conjunction with the NPWS Conservation Objectives Document.

**SITE CODE:
SAC 001571; version 3. CO. MAYO**


0 200 400 600 800 Meters

The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
Ordnance Survey of Ireland Licence No EN 0059216. © Ordnance Survey of Ireland Government of Ireland

Níl sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbheithnithe a déanamh ar theorainneacha na gceantar comharthaíthe. Suirbhéarachta Ordonáis na hÉireann Ceadúnas Uimh EN 0059216. © Suirbhéarachta Ordonáis na hÉireann Rialtas na hÉireann




**Map Version 1
Date: Nov 2017**



Legend

- Urlaur Lakes SAC 001571
- 3140 Hard oligo-mesotrophic waters with benthic vegetation of Chara spp.
- OSi Discovery Series County Boundary





Conservation objectives for Bricklieve Mountains and Keishcorran SAC [001656]

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Objective: To maintain or restore the favourable conservation condition of the Annex I habitat(s) and/or the Annex II species for which the SAC has been selected:

Code	Description
3180	Turloughs*
6210	Semi-natural dry grasslands and scrubland facies on calcareous substrates (<i>Festuco Brometalia</i>) (* important orchid sites)*
6510	Lowland hay meadows (<i>Alopecurus pratensis</i> , <i>Sanguisorba officinalis</i>)
8120	Calcareous and calchist screes of the montane to alpine levels (<i>Thlaspietea rotundifolii</i>)

* denotes a priority habitat

Code	Common Name	Scientific Name
------	-------------	-----------------

For more information please go to: www.npws.ie/protected-sites/conservation-management-planning



1065	Marsh Fritillary	<i>Euphydryas aurinia</i>
1092	White-clawed Crayfish	<i>Austropotamobius pallipes</i>

Citation: NPWS (2020) Conservation objectives for Bricklieve Mountains and Keishcorran SAC [001656]. Generic Version 7.0. Department of Culture, Heritage and the Gaeltacht.



Conservation objectives for Knockalongy and Knockachree Cliffs SAC [001669]

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Objective: To maintain or restore the favourable conservation condition of the Annex I habitat(s) and/or the Annex II species for which the SAC has been selected:

ADD HABITATS

Code	Common Name	Scientific Name
1421	Killarney Fern	<i>Trichomanes speciosum</i>



Citation: NPWS (2020) Conservation objectives for Knockalongy and Knockachree Cliffs SAC [001669].
Generic Version 7.0. Department of Culture, Heritage and the Gaeltacht.



Conservation objectives for Lough Carra/Mask Complex SAC [001774]

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Objective: To maintain or restore the favourable conservation condition of the Annex I habitat(s) and/or the Annex II species for which the SAC has been selected:

Code	Description
3110	Oligotrophic waters containing very few minerals of sandy plains (<i>Littorelletalia uniflorae</i>)
3130	Oligotrophic to mesotrophic standing waters with vegetation of the <i>Littorelletea uniflorae</i> and/or <i>Isoeto-Nanojuncetea</i>
3140	Hard oligo-mesotrophic waters with benthic vegetation of <i>Chara</i> spp.
4030	European dry heaths
6210	Semi-natural dry grasslands and scrubland facies on calcareous substrates (<i>Festuco Brometalia</i>) (* important orchid sites)*
7210	Calcareous fens with <i>Cladium mariscus</i> and species of the <i>Caricion davallianae</i> *
7230	Alkaline fens
8240	Limestone pavements*



91E0 Alluvial forests with *Alnus glutinosa* and *Fraxinus excelsior* (Alno-Padion, Alnion incanae, Salicion albae)*

* denotes a priority habitat

Code	Common Name	Scientific Name
1303	Lesser Horseshoe Bat	<i>Rhinolophus hipposideros</i>
1355	Otter	<i>Lutra lutra</i>
1393	Slender Green Feather-moss	<i>Drepanocladus vernicosus</i>

Citation: NPWS (2020) Conservation objectives for Lough Carra/Mask Complex SAC [001774]. Generic Version 7.0. Department of Culture, Heritage and the Gaeltacht.



Conservation objectives for Unshin River SAC [001898]

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Objective: To maintain or restore the favourable conservation condition of the Annex I habitat(s) and/or the Annex II species for which the SAC has been selected:

Code	Description
3260	Water courses of plain to montane levels with the <i>Ranunculus fluitans</i> and <i>Callitriche-Batrachion</i> vegetation
6210	Semi-natural dry grasslands and scrubland facies on calcareous substrates (<i>Festuco Brometalia</i>) (* important orchid sites)*
6410	<i>Molinia</i> meadows on calcareous, peaty or clayey-silt-laden soils (<i>Molinion caeruleae</i>)
91E0	Alluvial forests with <i>Alnus glutinosa</i> and <i>Fraxinus excelsior</i> (<i>Alno-Padion</i> , <i>Alnion incanae</i> , <i>Salicion albae</i>)*

* denotes a priority habitat



Code	Common Name	Scientific Name
1106	Salmon	<i>Salmo salar</i>
1355	Otter	<i>Lutra lutra</i>

Citation: NPWS (2020) Conservation objectives for Unshin River SAC [001898]. Generic Version 7.0.
Department of Culture, Heritage and the Gaeltacht.

National Parks and Wildlife Service

Conservation Objectives Series

Cloonakillina Lough SAC 001899



An Roinn Cultúir,
Oidhreacht agus Gaeltachta
Department of Culture,
Heritage and the Gaeltacht

**National Parks and Wildlife Service,
Department of Culture, Heritage and the Gaeltacht,
90 King Street North, Dublin 7, D07 N7CV, Ireland.**

**Web: www.npws.ie
E-mail: nature.conservation@chg.gov.ie**

Citation:

NPWS (2019) Conservation Objectives: Cloonakillina Lough SAC 001899. Version 1. National Parks and Wildlife Service, Department of Culture, Heritage and the Gaeltacht.

**Series Editor: Rebecca Jeffrey
ISSN 2009-4086**

Introduction

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

** indicates a priority habitat under the Habitats Directive*

001899 Cloonakillina Lough SAC

7140 Transition mires and quaking bogs

Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

NPWS Documents

Year :	1992
Title :	Owenmore River Catchment. Proposed Arterial Drainage Environmental Impact Assessment - Botanical and Ornithological Surveys.
Author :	Goodwillie, R.N.; Buckley, P.; Douglas, C.
Series :	Unpublished report
Year :	2009
Title :	Ireland Red List No. 2: Non-marine molluscs
Author :	Byrne, A.; Moorkens, E.A.; Anderson, R.; Killeen, I.J.; Regan, E.C.
Series :	Ireland Red List series, NPWS
Year :	2010
Title :	Ireland Red List No. 4: Butterflies
Author :	Regan, E.C.; Nelson, B.; Aldwell, B.; Bertrand, C.; Bond, K.; Harding, J.; Nash, D.; Nixon, D.; Wilson, C.J.
Series :	Ireland Red List series, NPWS
Year :	2012
Title :	Ireland Red List No. 8: Bryophytes
Author :	Lockhart, N.; Hodgetts, N.; Holyoak, D.
Series :	Ireland Red List series, NPWS
Year :	2013
Title :	The status of EU protected habitats and species in Ireland. Volume 2. Habitats assessments
Author :	NPWS
Series :	Conservation assessments
Year :	2013
Title :	Conservation status assessments for three fen habitat types - 7230 – Alkaline fens, 7210 – Calcareous fens with <i>Cladium mariscus</i> and species of <i>Caricion davallianae</i> and 7140 – Transition mires and quaking bogs
Author :	Kimberley, S.
Series :	Unpublished report to NPWS
Year :	2014
Title :	Guidelines for a national survey and conservation assessment of upland vegetation and habitats in Ireland, Version 2.0
Author :	Perrin, P.M.; Barron, S.J.; Roche, J.R.; O'Hanrahan, B.
Series :	Irish Wildlife Manuals, No. 79
Year :	2016
Title :	Ireland Red List No. 10: Vascular Plants
Author :	Wyse Jackson, M.; FitzPatrick, Ú.; Cole, E.; Jebb, M.; McFerran, D.; Sheehy Skeffington, M.; Wright, M.
Series :	Ireland Red List Series, NPWS
Year :	2018
Title :	Backing document – National Conservation Status Assessments (NCAs) for three fen habitat types: 7140 – Transition mires and quaking bogs, 7210 – Calcareous fens with <i>Cladium mariscus</i> and species of <i>Caricion davallianae</i> , 7230 – Alkaline fens
Author :	Long, M.P.; Crowe, O.; Kimberley, S.; Denyer, J.
Series :	Unpublished report to NPWS

Year : in prep.
Title : The Status of EU Protected Habitats and Species in Ireland (2013-2018). Habitat Assessments
Author : NPWS
Series : Conservation assessments

Other References

Year : 1993
Title : Notes on the flora of the Owenmore Catchment Cos Sligo (H28) and East Mayo (H26)
Author : Douglas, C.; Goodwillie, R.; Mooney, E.
Series : Irish Naturalists' Journal, 24(5): 218-220

Year : 2004
Title : Common Standards Monitoring guidance for lowland wetland habitats
Author : JNCC
Series : Joint Nature Conservation Committee, Peterborough

Year : 2011
Title : Review and revision of empirical critical loads and dose-response relationships. Proceedings of an expert workshop, Noordwijkerhout, 23-25 June 2010
Author : Bobbink, R.; Hettelingh, J.P.
Series : RIVM report 680359002, Coordination Centre for Effects, National Institute for Public Health and the Environment (RIVM)

Year : 2018
Title : Irish Vegetation Classification: Technical Progress Report No. 4
Author : Perrin, P.
Series : Report submitted to National Biodiversity Data Centre


Conservation Objectives for : Cloonakillina Lough SAC [001899]


7140 Transition mires and quaking bogs


To maintain the favourable conservation condition of Transition mires and quaking bogs in Cloonakillina Lough SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Transition mires and quaking bogs have not been mapped in detail for Cloonakillina Lough SAC and thus the total current area of the qualifying habitat in the SAC is unknown. The habitat occurs as a scraw (floating vegetation), with some open pools, covering the western half of the Cloonakillina Lough basin. Cloonakillina Lough has undergone rapid succession from open water to transition mire since it was mapped in 1915, most likely due to drainage and lowering of the water table in the area (NPWS internal files). The SAC contains an interesting variety of stages in mire development, and transition mire occurs in association with areas of open water, fen grassland, reed swamp and wet woodland (NPWS internal files). See also Goodwillie et al. (1992)
Habitat distribution	Occurrence	No decline, subject to natural processes	See the notes for Habitat area above
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil pH and nutrient status within natural ranges	Relevant nutrients and their natural ranges are yet to be defined. However, nitrogen deposition is noted as being relevant to this habitat in NPWS (2013). See also Bobbink and Hettelingh (2011)
Ecosystem function: peat formation	Percentage cover of peat-forming vegetation and water table levels	Maintain active peat formation, where appropriate	In order for peat to form, water levels need to be slightly below or above the soil surface for c.90% of the time
Ecosystem function: hydrology - water levels	Centimetres; duration of water levels	Maintain appropriate water levels necessary to support the natural structure and functioning of the habitat	Maintenance of a permanently high water level, remaining close to the peat surface all year, with water level fluctuations within natural ranges, is required for this wetland habitat. See Kimberley (2013) and Long et al. (2018)
Ecosystem function: hydrology - flow patterns	Flow direction	Maintain appropriate topography and water movement regime necessary to support the natural structure and functioning of the habitat	Maintenance, both within and surrounding the habitat, of topography and flow patterns within natural ranges is essential in order to ensure the hydrological integrity of this wetland habitat
Ecosystem function: water quality	Water chemistry measures	Maintain appropriate water quality to support the natural structure and functioning of the habitat	The surface water conditions necessary to maintain transition mires range from acidic to slightly base-rich. The vegetation typically has intimate mixtures of species considered to be acidophile and others considered calciphile. In other cases, these intermediate properties may reflect the actual process of succession, as peat accumulates in groundwater-fed fen or open water to produce rainwater-fed bog isolated from groundwater influence
Community diversity	Abundance of variety of vegetation communities	Maintain variety of vegetation communities, subject to natural processes	The entire diversity of transition mire vegetation communities present in the SAC is currently unknown. Information on vegetation communities associated with this habitat in the uplands is presented in Perrin et al. (2014). See also the Irish Vegetation Classification (Perrin, 2018; www.biodiversityireland.ie/projects/national-vegetation-database/irish-vegetation-classification)

Vegetation composition: typical vascular plants and bryophytes	Percentage cover at a representative number of 2m x 2m monitoring stops	Maintain adequate cover of typical vascular plant and bryophyte species	For lists of typical vascular plant and bryophyte species, including high quality indicators, see the Article 17 conservation status assessment for transition mires and quaking bogs (NPWS, in prep.) and the fen habitats supporting document (Long et al., 2018). See also Perrin et al. (2014) and JNCC (2004). In this SAC, the habitat is species-rich with typical species recorded including bog myrtle (<i>Myrica gale</i>), bogbean (<i>Menyanthes trifoliata</i>), marsh pennywort (<i>Hydrocotyle vulgaris</i>), marsh lousewort (<i>Pedicularis palustris</i>) marsh cinquefoil (<i>Comarum palustre</i>), <i>Sphagnum contortum</i> and <i>S. squarrosum</i> and a diverse range of sedges including slender sedge (<i>Carex lasiocarpa</i>), lesser tussock-sedge (<i>C. diandra</i>), bottle sedge (<i>C. rostrata</i>), common sedge (<i>C. nigra</i>), bog sedge (<i>C. limosa</i>) and bog cotton (<i>Eriophorum angustifolium</i>) (Goodwillie et al., 1992; NPWS internal files)
Vegetation composition: native negative indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Native negative indicator species at insignificant levels	Negative indicators include species not characteristic of the habitat and species indicative of undesirable activities such as overgrazing, undergrazing, nutrient enrichment, agricultural improvement or impacts on hydrology. Native negative indicator species that could suggest drying out include ling (<i>Calluna vulgaris</i>) and birch (<i>Betula pubescens</i>)
Vegetation composition: non-native species	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of non-native species less than 1%	Attribute and target based on Perrin et al. (2014). Non-native species can be invasive and have deleterious effects on native vegetation. A low target is set as non-native species can spread rapidly and are most easily dealt with when still at lower abundances
Physical structure: drainage	Percentage area in local vicinity of a representative number of monitoring stops	Area showing signs of drainage from heavy trampling, tracking or ditches less than 10%	Attribute and target based on Perrin et al. (2014). Drainage can result in loss of characteristic species and transition to drier habitats. Cloonakillina Lough SAC contains a dynamic hydrological and ecological system susceptible to drainage (NPWS internal files)
Physical structure: disturbed bare ground	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of disturbed bare ground less than 10%	Attribute and target based on Perrin et al. (2014). Disturbance can include hoof marks, wallows, human footprints, vehicle and machinery tracks. Excessive disturbance can result in loss of characteristic species and presage erosion for peatlands
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat; maintain features of local distinctiveness, subject to natural processes	This includes species on the Flora (Protection) Order, 2015 and/or Red Lists (Byrne et al., 2009; Regan et al., 2010; Lockhart et al., 2012; Wyse Jackson et al., 2016, etc.). The Near Threatened slender tufted-sedge (<i>Carex acuta</i>) (Wyse Jackson et al., 2016) has been recorded in the habitat in the SAC (Goodwillie et al., 1992; Douglas et al., 1993)



Legend
 Cloonakillina Lough SAC 001899

 An Roinn Cultúir,
 Oidhreacht agus Gaeltachta
 Department of Culture,
 Heritage and the Gaeltacht

**MAP 1:
 CLOONAKILLINA LOUGH SAC
 CONSERVATION OBJECTIVES
 SAC DESIGNATION**
 Map to be read in conjunction with the NPWS Conservation Objectives Document.

**SITE CODE:
 SAC 001899; version 3.01. CO. MAYO**

0 100 200 300 400 Meters

The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
 Ordnance Survey of Ireland Licence No OSI-NMA-014. © Ordnance Survey of Ireland Government of Ireland

Níl sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbhreithnithe a déanamh ar theorainneacha na gceantar comharthaithe. Suirbhéarachta Ordonáis na hÉireann Ceadúnas Uimh OSI-NMA-014. © Suirbhéarachta Ordonáis na hÉireann Rialtas na hÉireann


**Map Version 1
 Date: Feb 2019**

National Parks and Wildlife Service

Conservation Objectives Series

Bellacorick Bog Complex SAC 001922



An Roinn
Cultúir, Oidhreachta agus Gaeltachta

Department of
Culture, Heritage and the Gaeltacht



**National Parks and Wildlife Service,
Department of Culture, Heritage and the Gaeltacht,
7 Ely Place, Dublin 2, Ireland.
Web: www.npws.ie
E-mail: nature.conservation@ahg.gov.ie**

Citation:

**NPWS (2017) Conservation Objectives: Bellacorick Bog Complex SAC 001922.
Version 1. National Parks and Wildlife Service, Department of Culture, Heritage
and the Gaeltacht.**

**Series Editor: Rebecca Jeffrey
ISSN 2009-4086**

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

** indicates a priority habitat under the Habitats Directive*

001922	Bellacorick Bog Complex SAC
1013	Geyer's Whorl Snail <i>Vertigo geyeri</i>
1528	Marsh Saxifrage <i>Saxifraga hirculus</i>
3160	Natural dystrophic lakes and ponds
4010	Northern Atlantic wet heaths with <i>Cladonia</i> <i>Cladonia</i>
7130	Blanket bogs (* if active bog)
7150	Depressions on peat substrates of the Rhynchosporion
7230	Alkaline fens

Please note that this SAC is adjacent to River Moy SAC (002298). See map 2. The conservation objectives for this site should be used in conjunction with those for the adjacent site as appropriate.

Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

NPWS Documents

Year :	1987
Title :	A survey to locate blanket bogs of scientific interest in County Mayo. Part I
Author :	Foss, P.; McGee, E.
Series :	A report commissioned by the Wildlife Service
Year :	1989
Title :	Survey to locate blanket bogs of scientific interest in Mayo. Part II
Author :	Douglas, C.; Garvey, L.; Kelly, L.; O'Sullivan, A.; Van Doorsleer, L.
Series :	A report commissioned by the Wildlife Service
Year :	2011
Title :	Monitoring and condition assessment of populations of <i>Vertigo geyeri</i> , <i>Vertigo angustior</i> and <i>Vertigo moulinsiana</i> in Ireland
Author :	Moorkens, E.; Killeen, I.
Series :	Irish Wildlife Manual No. 55
Year :	2011
Title :	Implementation of a <i>Vertigo</i> monitoring programme: <i>Vertigo angustior</i> monitoring at Fermoy
Author :	Moorkens, E.A.; Killeen, I.J.
Series :	Unpublished report to NPWS
Year :	2011
Title :	Implementation of a <i>Vertigo</i> monitoring programme: <i>Vertigo geyeri</i> monitoring at Brackloon
Author :	Moorkens, E.A.; Killeen, I.J.
Series :	Unpublished report to NPWS
Year :	2012
Title :	Ireland Red List No. 8: Bryophytes
Author :	Lockhart, N.; Hodgetts, N.; Holyoak, D.
Series :	Ireland Red List series, NPWS
Year :	2013
Title :	The status of EU protected habitats and species in Ireland. Volume 2. Habitats assessments
Author :	NPWS
Series :	Conservation assessments
Year :	2013
Title :	The status of EU protected habitats and species in Ireland. Volume 3. Species assessments
Author :	NPWS
Series :	Conservation assessments
Year :	2014
Title :	Guidelines for a national survey and conservation assessment of upland vegetation and habitats in Ireland, Version 2.0
Author :	Perrin, P.M.; Barron, S.J.; Roche, J.R.; O'Hanrahan, B.
Series :	Irish Wildlife Manual No. 79
Year :	2015
Title :	Monitoring recommendations for Marsh Saxifrage (<i>Saxifraga hirculus</i> L.) in the Republic of Ireland
Author :	Muldoon, C.S.; Waldren, S.; Lynn, D.
Series :	Irish Wildlife Manual No. 88

Year : 2015
Title : Habitats Directive Annex I lake habitats: a working interpretation for the purposes of site-specific conservation objectives and Article 17 reporting
Author : O Connor, Á.
Series : Unpublished document by NPWS

Year : 2016
Title : Ireland Red List No. 10: Vascular Plants
Author : Wyse Jackson, M.; FitzPatrick, Ú.; Cole, E.; Jebb, M.; McFerran, D.; Sheehy Skeffington, M.; Wright, M.
Series : Ireland Red Lists series, NPWS

Year : 2017
Title : Bellacorick Bog Complex SAC (site code: 1922) Conservation objectives supporting document-blanket bogs and associated habitats V1
Author : NPWS
Series : Conservation objectives supporting document

Other References

Year : 1982
Title : Eutrophication of waters. Monitoring assessment and control
Author : OECD
Series : OECD, Paris

Year : 2000
Title : Colour in Irish lakes
Author : Free, G.; Allott, N.; Mills, P.; Kennelly, C.; Day, S.
Series : Verhandlungen Internationale Vereinigung für theoretische und angewandte Limnologie, 27: 2620-2623

Year : 2005
Title : Widespread occurrence of *Vertigo geyeri* (Gastropoda: Vertiginidae) in north and west Ireland
Author : Holyoak, G.A.
Series : Irish Naturalists' Journal, 28: 141-150

Year : 2006
Title : A reference-based typology and ecological assessment system for Irish lakes. Preliminary investigations. Final report. Project 2000-FS-1-M1 Ecological assessment of lakes pilot study to establish monitoring methodologies EU (WFD)
Author : Free, G.; Little, R.; Tierney, D.; Donnelly, K.; Coroni, R.
Series : EPA, Wexford

Year : 2008
Title : Water Quality in Ireland 2004-2006
Author : Clabby, K.J.; Bradley, C.; Craig, M.; Daly, D.; Lucey, J.; McGarrigle, M.; O'Boyle, S.; Tierney, D.; Bowman, J.
Series : EPA, Wexford

Year : 2010
Title : Water quality in Ireland 2007-2009
Author : McGarrigle, M.; Lucey, J.; Ó Cinnéide, M.
Series : EPA, Wexford

Year : 2011
Title : Conservation biology of *Saxifraga hirculus* L. in Ireland
Author : Muldoon, C.S.
Series : Unpublished Ph.D. Thesis, Trinity College Dublin

Year : 2012
Title : The impact of conifer plantation forestry on the ecology of peatland lakes
Author : Drinan, T.J.
Series : Unpublished Ph.D. thesis, University College Cork

Year : 2013
Title : Interpretation manual of European Union habitats- Eur 28
Author : European Commission- DG Environment
Series : European Commission

Year : 2015
Title : Water quality in Ireland 2010-2012
Author : Bradley, C.; Byrne, C.; Craig, M.; Free, G.; Gallagher, T.; Kennedy, B.; Little, R.; Lucey, J.; Mannix, A.; McCreesh, P.; McDermott, G.; McGarrigle, M.; Ní Longphuirt, S.; O'Boyle, S.; Plant, C.; Tierney, D.; Trodd, W.; Webster, P.; Wilkes, R.; Wynne, C.
Series : EPA, Wexford

Spatial data sources

Year : 2008
Title : OSi 1:5000 IG vector dataset
GIS Operations : WaterPolygons feature class clipped to the SAC boundary. Expert opinion used to identify Annex I habitat and to resolve any issues arising
Used For : 3160 (map 3)

Year : 2017
Title : NPWS rare and threatened species database
GIS Operations : Dataset created from spatial references in database records. Expert opinion used as necessary to resolve any issues arising
Used For : 1013 (map 4)

Conservation Objectives for : Bellacorick Bog Complex SAC [001922]

3160 Natural dystrophic lakes and ponds

To maintain the favourable conservation condition of Natural dystrophic lakes and ponds in Bellacorick Bog Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Bellacorick Bog Complex SAC has some of the most extensive extant areas of lowland blanket bog pool systems. Habitat 3160 is likely to occur in all pools and lakes and all are mapped as potential 3160 (map 3). Although there are more than 5,700 lake/pool polygons, many pools are not mapped in the 1:5,000 OSi data (see map 3). The habitat is of high conservation value in the SAC, owing to the area, extent and morphological diversity of pools. For further information on the distribution, vegetation and morphology of the habitat in the SAC, see Foss and McGee (1987) and Douglas et al. (1989). Two measures of extent should be used: 1. the area of the lake itself and; 2. the extent of the vegetation communities/zones that typify the habitat. Further information relating to all attributes is provided in the lake habitats supporting document for the purposes of site-specific conservation objectives and Article 17 reporting (O Connor, 2015)
Habitat distribution	Occurrence	No decline, subject to natural processes	As noted above, the habitat is widespread in the SAC (see map 3). All lake/pond polygons have been mapped as potential 3160
Typical species	Occurrence	Typical species present, in good condition, and demonstrating typical abundances and distribution	For lists of typical plant and invertebrate species, see the Article 17 habitat assessment for 3160 (NPWS, 2013) and O Connor (2015). Douglas et al. (1989) state that pools and lakes typically have many-stalked spike-rush (<i>Eleocharis multicaulis</i>), bogbean (<i>Menyanthes trifoliata</i>), the bog mosses <i>Sphagnum denticulatum</i> and <i>S. cuspidatum</i> , with great sundew (<i>Drosera anglica</i>) and round-leaved sundew (<i>D. rotundifolia</i>) in shallower interconnecting pools. Pipewort (<i>Eriocaulon aquaticum</i>), bulbous rush (<i>Juncus bulbosus</i>) and water lobelia (<i>Lobelia dortmanna</i>) also occur in pools and lakes, while bladderworts (<i>Utricularia</i> spp.), broad-leaved pondweed (<i>Potamogeton natans</i>), common reed (<i>Phragmites australis</i>) and bottle sedge (<i>Carex rostrata</i>) were also recorded
Vegetation composition: characteristic zonation	Occurrence	All characteristic zones should be present, correctly distributed and in good condition	Further work is necessary to describe the characteristic zonation and other spatial patterns in lake habitat 3160 (see O Connor, 2015). Spatial patterns are likely to be relatively simple in 3160 lakes and ponds, with limited zonation
Vegetation distribution: maximum depth	Metres	Maintain maximum depth of vegetation, subject to natural processes	The maximum depth of vegetation is likely to be specific to the lake shoreline in question. Further work is necessary to develop indicative targets for lake habitat 3160. 3160 lakes and pools naturally have very clear water and, therefore, maximum depth can be large

Hydrological regime: water level fluctuations	Metres	Maintain appropriate natural hydrological regime necessary to support the habitat	Fluctuations in lake water level are typical in Ireland, but can be amplified by activities such as abstraction and drainage. Increased water level fluctuations can increase wave action, up-root vegetation, increase turbidity, alter the substratum and lead to release of nutrients from the sediment. The hydrological regime of the lakes and pools must be maintained so that the area, distribution and depth of the lake habitat and its constituent/characteristic vegetation zones and communities are not reduced. The hydrological regime of 3160 lakes and pools is integrally linked to that of the surrounding blanket bog, transition mire/quaking bog and other peatland habitats. Owing to their size and the sensitivity of peatland, 3160 lakes and pools can easily be damaged or destroyed by drainage
Lake substratum quality	Various	Maintain appropriate substratum type, extent and chemistry to support the vegetation	Research is required to further characterise the substratum types (particle size and origin) and substratum quality (notably pH, calcium, iron and nutrient concentrations) favoured by each of the five Annex I lake habitats in Ireland. It is likely that lake habitat 3160 is associated with nutrient-poor peat and silt substrates
Water quality: transparency	Metres	Maintain appropriate Secchi transparency. There should be no decline in Secchi depth/transparency	Transparency relates to light penetration and, hence, to the depth of colonisation of vegetation. It can be affected by phytoplankton blooms, water colour and turbidity. A specific target has yet to be established for this Annex I lake habitat. Habitat 3160 is associated with very clear water. The OECD fixed boundary system set transparency targets for ultra-oligotrophic lakes of $\geq 12\text{m}$ annual mean Secchi disk depth, and $\geq 6\text{m}$ annual minimum Secchi disk depth
Water quality: nutrients	$\mu\text{g/l P}$; mg/l N	Maintain the concentration of nutrients in the water column at sufficiently low levels to support the habitat and its typical species	As a nutrient-poor habitat, oligotrophic and Water Framework Directive (WFD) 'high' status targets apply. Where a lake has nutrient concentrations that are lower than these targets, there should be no decline within class, i.e. no upward trend in nutrient concentrations. For 3160 lakes and pools, annual average total phosphorus (TP) concentration should be $\leq 5\mu\text{g/l TP}$, average annual total ammonia concentration should be $\leq 0.040\text{mg/l N}$ and annual 95th percentile for total ammonia should be $\leq 0.090\text{mg/l N}$. See also The European Communities Environmental Objectives (Surface Waters) Regulations 2009
Water quality: phytoplankton biomass	$\mu\text{g/l Chlorophyll } a$	Maintain appropriate water quality to support the habitat, including high chlorophyll <i>a</i> status	Oligotrophic and WFD 'high' status targets apply to lake habitat 3160. The average growing season (March-October) chlorophyll <i>a</i> concentration must be $< 5.8\mu\text{g/l}$ (The European Communities Environmental Objectives (Surface Waters) Regulations 2009). Where a lake has a chlorophyll <i>a</i> concentration that is lower than this target, there should be no decline within class, i.e. no upward trend in phytoplankton biomass. The OECD targets may be more appropriate for habitat 3160: annual average chlorophyll <i>a</i> concentration $< 1\mu\text{g/l}$ and annual peak chlorophyll <i>a</i> concentration $\leq 2.5\mu\text{g/l}$
Water quality: phytoplankton composition	EPA phytoplankton composition metric	Maintain appropriate water quality to support the habitat, including high phytoplankton composition status	The Environmental Protection Agency (EPA) has developed a phytoplankton composition metric for nutrient enrichment of Irish lakes. As for other water quality indicators, lake habitat 3160 requires WFD high status
Water quality: attached algal biomass	Algal cover and EPA phytobenthos metric	Maintain trace/absent attached algal biomass ($< 5\%$ cover) and high phytobenthos status	Nutrient enrichment can favour epiphytic and epipelagic algae that can out-compete the submerged vegetation. The cover abundance of attached algae in the oligotrophic soft water habitat should, therefore, be trace/absent ($< 5\%$ cover). EPA phytobenthos can be used as an indicator of changes in attached algal biomass. As for other water quality indicators, lake habitat 3160 requires high phytobenthos status

Water quality: macrophyte status	EPA macrophyte metric (The Free Index)	Maintain high macrophyte status	Nutrient enrichment can favour more competitive submerged macrophyte species that out-compete the typical and characteristic species for the lake habitat. The EPA monitors macrophyte status for WFD purposes using the 'Free Index'. The target for 3160 lakes and pools is high status or an Ecological Quality Ratio (EQR) for lake macrophytes of ≥ 0.90 , as defined in Schedule Five of the European Communities Environmental Objectives (Surface Waters) Regulations 2009
Acidification status	pH units; mg/l	Maintain appropriate water and sediment pH, alkalinity and cation concentrations to support the habitat, subject to natural processes	Acidification can impact on species abundance and composition in soft water lake habitats. Although European Commission (2013) describes lake habitat 3160 as having pH 3-6, Drinan (2012) found mean pHs of 5.16 and 5.62 in upland and lowland 3160 lakes, respectively. The target for habitat 3160 is pH >4.5 and <9.0 , in line with the surface water standards for soft waters (where water hardness is $\leq 100\text{mg/l}$ calcium carbonate). See Schedule Five of the European Communities Environmental Objectives (Surface Waters) Regulations 2009. The specific requirements of lake habitat 3160, in terms of water and sediment pH, alkalinity and cation concentration, have not been determined
Water colour	mg/l PtCo	Maintain appropriate water colour to support the habitat	Increased water colour and turbidity decrease light penetration and can reduce the area of available habitat for lake macrophytes, particularly at the lower euphotic depths. The primary source of increased water colour in Ireland is disturbance to peatland. No habitat-specific or national standards for water colour currently exist. Studies have shown median colour concentrations in Irish lakes of 38mg/l PtCo (Free et al., 2000) and 33mg/l PtCo (Free et al., 2006). It is likely that the water colour in all Irish lake habitats would naturally be $<50\text{mg/l}$ PtCo. Water colour can be very low ($<20\text{mg/l}$ PtCo or even $<10\text{mg/l}$ PtCo) in 3160 lakes and pools where the peatland in the lake's catchment is intact
Dissolved organic carbon (DOC)	mg/l	Maintain appropriate organic carbon levels to support the habitat	Dissolved (and particulate) organic carbon (OC) in the water column is linked to water colour and acidification (organic acids). Increasing DOC in water has been documented across the Northern Hemisphere, including afforested peatland catchments in Ireland. Damage and degradation of peatland, leading to decomposition of peat is likely to be the predominant source of OC in Ireland. OC in water promotes decomposition by fungi and bacteria that, in turn, releases dissolved nutrients. The increased biomass of decomposers can also impact directly on the characteristic lake communities through shading, competition, etc.
Turbidity	Nephelometric turbidity units/ mg/l SS/ other appropriate units	Maintain appropriate turbidity to support the habitat	Turbidity can significantly affect the quantity and quality of light reaching rooted and attached vegetation and can, therefore, impact on lake habitats. The settlement of higher loads of inorganic or organic material on lake vegetation communities may also have impacts on sensitive, delicate species. Turbidity can increase as a result of re-suspension of material within the lake, higher loads entering the lake, or eutrophication. Turbidity measurement and interpretation is challenging. As a result, it is likely to be difficult to set habitat-specific targets for turbidity in lakes

Fringing habitat: Hectares
area and condition

Maintain the area and condition of fringing habitats necessary to support the natural structure and functioning of habitat 3160

3160 lakes and pool intergrade with blanket bog communities in this SAC. Spring-fed, species-rich flushes are a significant feature, occurring throughout the complex and supporting poor and rich fen and swamp carr communities. Quaking bog is also associated with pool systems in the SAC. These habitats support the structure and functions of the lake habitat. The fringing habitats are also dependent on the lake/pool, particularly its water levels, and can support wetland communities and species of conservation concern

Conservation Objectives for : Bellacorick Bog Complex SAC [001922]

4010 Northern Atlantic wet heaths with *Erica tetralix*

To restore the favourable conservation condition of Northern Atlantic wet heaths with *Erica tetralix* in Bellacorick Bog Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Northern Atlantic wet heaths with <i>Erica tetralix</i> has not been mapped in detail for Bellacorick Bog Complex SAC, but from current available data the total area of the qualifying habitat is estimated to be approximately 187ha, covering 2% of the SAC (NPWS internal files). Further details on this and the following attributes can be found in the Bellacorick Bog Complex SAC conservation objectives supporting document for blanket bogs and associated habitats
Habitat distribution	Occurrence	No decline, subject to natural processes	The habitat occurs occasionally on sloping ground and on elevated mounds of mineral soil that are scattered throughout the lowland blanket bog-covered plains. These are particularly evident in the Owenboy Nature Reserve and along some of the steeper stream valley sides (NPWS internal files). Further information can be found within Foss and McGee (1987), NPWS internal files and the blanket bogs and associated habitats supporting document
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil nutrient status within natural range	See the blanket bogs and associated habitats supporting document for further details
Community diversity	Abundance of variety of vegetation communities	Maintain variety of vegetation communities, subject to natural processes	Wet heath vegetation communities have been recorded in this SAC (Douglas et al., 1989; NPWS internal files), one of which corresponds to a community recorded in the National Survey of Upland Habitats and listed in the provisional list of vegetation communities described in Perrin et al. (2014). Further information on vegetation communities associated with this habitat is presented in Perrin et al. (2014)
Vegetation composition: cross-leaved heath	Occurrence within 20m of a representative number of monitoring stops	Cross-leaved heath (<i>Erica tetralix</i>) present within a 20m radius of each monitoring stop	Attribute and target based on Perrin et al. (2014)
Vegetation composition: positive indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of positive indicator species at least 50%	Attribute and target based on Perrin et al. (2014), where the list of positive indicator species for this habitat is also presented
Vegetation composition: lichens and bryophytes	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of <i>Cladonia</i> and <i>Sphagnum</i> species, <i>Racomitrium lanuginosum</i> and pleurocarpous mosses at least 10%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: ericoid species and crowberry	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of ericoid species and crowberry (<i>Empetrum nigrum</i>) at least 15%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: dwarf shrub species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of dwarf shrubs less than 75%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: negative indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of negative indicator species less than 1%	Attribute and target based on Perrin et al. (2014), where the list of negative indicator species for this habitat is also presented

Vegetation composition: non-native species	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of non-native species less than 1%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: native trees and shrubs	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of scattered native trees and shrubs less than 20%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: bracken	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of bracken (<i>Pteridium aquilinum</i>) less than 10%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: soft rush	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of soft rush (<i>Juncus effusus</i>) less than 10%	Attribute and target based on Perrin et al. (2014)
Vegetation structure: <i>Sphagnum</i> condition	Condition at a representative number of 2m x 2m monitoring stops	Less than 10% of the <i>Sphagnum</i> cover is crushed, broken and/or pulled up	Attribute and target based on Perrin et al. (2014)
Vegetation structure: signs of browsing	Percentage of shoots browsed at a representative number of 2m x 2m monitoring stops	Less than 33% collectively of the last complete growing season's shoots of ericoids, crowberry (<i>Empetrum nigrum</i>) and bog-myrtle (<i>Myrica gale</i>) showing signs of browsing	Attribute and target based on Perrin et al. (2014)
Vegetation structure: burning	Occurrence in local vicinity of a representative number of monitoring stops	No signs of burning in sensitive areas, into the moss, liverwort or lichen layer or exposure of peat surface due to burning	Attribute and target based on Perrin et al. (2014), where the list of sensitive areas for this habitat is also presented
Physical structure: disturbed bare ground	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of disturbed bare ground less than 10%	Attribute and target based on Perrin et al. (2014)
Physical structure: drainage	Percentage area in local vicinity of a representative number of monitoring stops	Area showing signs of drainage from heavy trampling, tracking or ditches less than 10%	Attribute and target based on Perrin et al. (2014)
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat	This includes species listed in the Flora (Protection) Order, 2015 and/or the red data lists (Lockhart et al., 2012; Wyse Jackson et al., 2016)

Conservation Objectives for : Bellacorick Bog Complex SAC [001922]

7130 Blanket bogs (* if active bog)

To restore the favourable conservation condition of Blanket bogs (* if active bog) in Bellacorick Bog Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Blanket bog has not been mapped in detail for Bellacorick Bog Complex SAC, but from current available data the total area of the qualifying habitat is estimated to be approximately 6,286ha, covering 66% of the SAC (NPWS internal files). Further details on this and the following attributes can be found in the Bellacorick Bog Complex SAC conservation objectives supporting document for blanket bogs and associated habitats
Habitat distribution	Occurrence	No decline, subject to natural processes	The habitat is documented to occur throughout the SAC. Further information can be found within Foss and McGee (1987), Douglas et al. (1989), NPWS internal files and the blanket bogs and associated habitats supporting document
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil nutrient status within natural range	See the blanket bogs and associated habitats supporting document for further details
Ecosystem function: peat formation	Active blanket bog as a proportion of the total area of Annex I blanket bog habitat	At least 99% of the total Annex I blanket bog area is active	See the blanket bogs and associated habitats supporting document for further details
Ecosystem function: hydrology	Flow direction, water levels, occurrence of drains and erosion gullies	Natural hydrology unaffected by drains and erosion	Further details and a brief discussion of restoration potential is presented in the blanket bogs and associated habitats supporting document
Community diversity	Abundance of variety of vegetation communities	Maintain variety of vegetation communities, subject to natural processes	A variety of blanket bog vegetation communities have been recorded in this SAC (Foss and McGee, 1987; Douglas et al., 1989; NPWS internal files), five of which correspond to communities recorded in the National Survey of Upland Habitats and listed in the provisional list of vegetation communities described in Perrin et al. (2014). Further information on vegetation communities associated with this habitat is presented in Perrin et al. (2014)
Vegetation composition: positive indicator species	Number of species at a representative number of 2m x 2m monitoring stops	Number of positive indicator species present at each monitoring stop is at least seven	Attribute and target based on Perrin et al. (2014), where the list of positive indicator species for this habitat is also presented
Vegetation composition: lichens and bryophytes	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of bryophytes or lichens, excluding <i>Sphagnum fallax</i> , at least 10%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: potential dominant species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of each of the potential dominant species less than 75%	Attribute and target based on Perrin et al. (2014). See the blanket bogs and associated habitats supporting document for the list of potential dominant species
Vegetation composition: negative indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of negative indicator species less than 1%	Attribute and target based on Perrin et al. (2014), where the list of negative indicator species for this habitat is also presented
Vegetation composition: non-native species	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of non-native species less than 1%	Attribute and target based on Perrin et al. (2014). <i>Rhododendron</i> (<i>Rhododendron ponticum</i>) occurs within this habitat in the SAC (NPWS internal files)

Vegetation composition: native trees and shrubs	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of scattered native trees and shrubs less than 10%	Attribute and target based on Perrin et al. (2014)
Vegetation structure: <i>Sphagnum</i> condition	Condition at a representative number of 2m x 2m monitoring stops	Less than 10% of the <i>Sphagnum</i> cover is crushed, broken and/or pulled up	Attribute and target based on Perrin et al. (2014)
Vegetation structure: signs of browsing	Percentage of shoots browsed at a representative number of 2m x 2m monitoring stops	Last complete growing season's shoots of ericoids, crowberry (<i>Empetrum nigrum</i>) and bog-myrtle (<i>Myrica gale</i>) showing signs of browsing collectively less than 33%	Attribute and target based on Perrin et al. (2014)
Vegetation structure: burning	Occurrence in local vicinity of a representative number of monitoring stops	No signs of burning in sensitive areas, into the moss, liverwort or lichen layer or exposure of peat surface due to burning	Attribute and target based on Perrin et al. (2014), where the list of sensitive areas for this habitat is also presented
Physical structure: disturbed bare ground	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of disturbed bare ground less than 10%	Attribute and target based on Perrin et al. (2014)
Physical structure: drainage	Percentage area in local vicinity of a representative number of monitoring stops	Area showing signs of drainage from heavy trampling, tracking or ditches less than 10%	Attribute and target based on Perrin et al. (2014)
Physical structure: erosion	Percentage area in local vicinity of a representative number of monitoring stops	Less than 5% of the greater bog mosaic comprises erosion gullies and eroded areas	Attribute and target based on Perrin et al. (2014)
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat	This includes species listed in the Flora (Protection) Order, 2015 and/or the red data lists (Lockhart et al., 2012; Wyse Jackson et al., 2016). The Near Threatened brown beak-sedge (<i>Rhynchospora fusca</i>) (Wyse Jackson et al., 2016) was recorded from the Bellacorick Bog area (Douglas et al., 1989), but this species cannot be assigned specifically to blanket bog

Conservation Objectives for : Bellacorick Bog Complex SAC [001922]

7150 Depressions on peat substrates of the Rhynchosporion

To restore the favourable conservation condition of Depressions on peat substrates of the Rhynchosporion in Bellacorick Bog Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Depressions on peat substrates of the Rhynchosporion has not been mapped in detail for Bellacorick Bog Complex SAC and thus the total area of the qualifying habitat in the SAC is unknown. Further details on this and the following attributes can be found in the Bellacorick Bog Complex SAC conservation objectives supporting document for blanket bogs and associated habitats
Habitat distribution	Occurrence	No decline, subject to natural processes	The habitat occurs in locations supporting pools and wet quaking peat (NPWS internal files). Further information can be found within the blanket bogs and associated habitats supporting document
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil nutrient status within natural range	See the blanket bogs and associated habitats supporting document for further details
Vegetation composition: positive indicator species	Number of species at a representative number of 2m x 2m monitoring stops	Number of positive indicator species at each monitoring stop is at least five	Attribute and target based on Perrin et al. (2014), where the list of positive indicator species for this habitat is also presented
Vegetation composition: <i>Rhynchospora</i> spp.	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of white beaked sedge (<i>Rhynchospora alba</i>) and brown beaked sedge (<i>R. fusca</i>) at least 10%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: potential dominant species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of each of the potential dominant species individually less than 35%	Attribute and target based on Perrin et al. (2014). See the blanket bogs and associated habitats supporting document for the list of potential dominant species
Vegetation composition: negative indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of negative indicator species less than 1%	Attribute and target based on Perrin et al. (2014), where the list of negative indicator species for this habitat is also presented
Vegetation composition: non-native species	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of non-native species less than 1%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: native trees and shrubs	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of scattered native trees and shrubs less than 10%	Attribute and target based on Perrin et al. (2014)
Vegetation structure: <i>Sphagnum</i> condition	Condition at a representative number of 2m x 2m monitoring stops	Less than 10% of the <i>Sphagnum</i> cover is crushed, broken and/or pulled up	Attribute and target based on Perrin et al. (2014)
Vegetation structure: signs of browsing	Percentage of shoots browsed at a representative number of 2m x 2m monitoring stops	Last complete growing season's shoots of ericoids, crowberry (<i>Empetrum nigrum</i>) and bog-myrtle (<i>Myrica gale</i>) showing signs of browsing collectively less than 33%	Attribute and target based on Perrin et al. (2014)
Vegetation structure: burning	Occurrence in local vicinity of a representative number of monitoring stops	No signs of burning in sensitive areas, into the moss, liverwort or lichen layer or exposure of peat surface due to burning	Attribute and target based on Perrin et al. (2014), where the list of sensitive areas for this habitat is also presented

Physical structure: disturbed bare ground	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of disturbed bare ground less than 10%	Attribute and target based on Perrin et al. (2014)
Physical structure: drainage	Percentage area in local vicinity of a representative number of monitoring stops	Area showing signs of drainage from heavy trampling, tracking or ditches less than 10%	Attribute and target based on Perrin et al. (2014)
Physical structure: erosion	Percentage area in local vicinity of a representative number of monitoring stops	Less than 5% of the greater bog mosaic comprises erosion gullies and eroded areas	Attribute and target based on Perrin et al. (2014)
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat	This includes species listed in the Flora (Protection) Order, 2015 and/or the red data lists (Lockhart et al., 2012; Wyse Jackson et al., 2016). The Near Threatened brown beak-sedge (<i>Rhynchospora fusca</i>) (Wyse Jackson et al., 2016) was recorded from the Bellacorick Bog area (Douglas et al., 1989), but this species cannot be assigned specifically to this habitat

Conservation Objectives for : Bellacorick Bog Complex SAC [001922]

7230 Alkaline fens

To restore the favourable conservation condition of Alkaline fens in Bellacorick Bog Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Alkaline fens has not been mapped in detail for Bellacorick Bog Complex SAC and thus the total area of the qualifying habitat in the SAC is unknown. Further details on this and the following attributes can be found in the Bellacorick Bog Complex SAC conservation objectives supporting document for blanket bogs and associated habitats
Habitat distribution	Occurrence	No decline, subject to natural processes	The habitat is documented to occur throughout the SAC, but is most well-developed along the eastern margin. Further information can be found within Foss and McGee (1987), Douglas et al. (1989), NPWS internal files and the blanket bogs and associated habitats supporting document
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil nutrient status within natural range	See the blanket bogs and associated habitats supporting document for further details
Ecosystem function: peat formation	Flood duration	Maintain active peat formation, where appropriate	In order for peat to form, water levels need to be slightly below or above the soil surface for c.90% of the time (Jim Ryan, pers. comm.)
Ecosystem function: hydrology	Metres	Maintain appropriate natural hydrological regimes necessary to support the natural structure and functioning of the habitat	Maintenance of groundwater, surface water flows and water table levels within natural ranges is essential for this wetland habitat
Ecosystem function: water quality	Water chemistry measures	Maintain appropriate water quality, particularly nutrient levels, to support the natural structure and functioning of the habitat	Fens receive natural levels of nutrients (e.g. iron, magnesium and calcium) from water sources. However, they are generally poor in nitrogen and phosphorus, with the latter tending to be the limiting nutrient
Community diversity	Abundance of variety of vegetation communities	Maintain variety of vegetation communities, subject to natural processes	The diversity of alkaline fen habitats in this SAC is unknown. Further information on vegetation communities associated with this habitat is presented in Perrin et al. (2014)
Vegetation composition: number of positive indicator species (brown mosses)	Number of species at a representative number of 2m x 2m monitoring stops	Number of brown moss species present at each monitoring stop is at least one	Attribute and target based on Perrin et al. (2014), where the list of positive indicator species for this habitat is also presented
Vegetation composition: number of positive indicator species (vascular plants)	Number of species at a representative number of 2m x 2m monitoring stops	Number of positive vascular plant indicator species present at each monitoring stop is at least two for small-sedge flushes and at least three for black bog-rush (<i>Schoenus nigricans</i>) flush and bottle sedge (<i>Carex rostrata</i>) fen	Attribute and target based on Perrin et al. (2014), where the list of positive indicator species for this habitat is also presented
Vegetation composition: cover of positive indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of brown moss species and positive vascular plant indicator species at least 20% for small-sedge flushes and at least 75% cover for black bog-rush (<i>Schoenus nigricans</i>) flush and bottle sedge (<i>Carex rostrata</i>) fen	Attribute and target based on Perrin et al. (2014), where the list of positive indicator species for this habitat is also presented

Vegetation composition: negative indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of negative indicator species less than 1%	Attribute and target based on Perrin et al. (2014), where the list of negative indicator species for this habitat is also presented
Vegetation composition: non-native species	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of non-native species less than 1%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: native trees and shrubs	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of scattered native trees and shrubs less than 10%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: soft rush and common reed cover	Percentage cover in local vicinity of a representative number of monitoring stops	Total cover of soft rush (<i>Juncus effusus</i>) and common reed (<i>Phragmites australis</i>) less than 10%	Attribute and target based on Perrin et al. (2014)
Vegetation structure: height	Percentage of leaves/shoots at a representative number of 2m x 2m monitoring stops	Proportion of live leaves and/or flowering shoots of vascular plants that are more than 5cm above the ground surface should be at least 50%	Attribute and target based on Perrin et al. (2014)
Physical structure: disturbed bare ground	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of disturbed bare ground less than 10%	Attribute and target based on Perrin et al. (2014)
Physical structure: drainage	Percentage area in local vicinity of a representative number of monitoring stops	Area showing signs of drainage from heavy trampling, tracking or ditches less than 10%	Attribute and target based on Perrin et al. (2014)
Physical structure: tufa formations	Percentage cover in local vicinity of a representative number of 2m x 2m monitoring stops	Disturbed proportion of vegetation cover where tufa is present is less than 1%	Attribute and target based on Perrin et al. (2014)
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat	This includes species listed in the Flora (Protection) Order, 2015 (FPO) and/or the red data lists (Lockhart et al., 2012; Wyse Jackson et al., 2016). The FPO listed and Near Threatened marsh saxifrage (<i>Saxifraga hirculus</i>) (Wyse Jackson et al., 2016) is present within the SAC (Muldoon et al., 2015), but this species cannot be assigned specifically to alkaline fens. Bryophytes present include the FPO and Critically Endangered moss <i>Paludella squarrosa</i> , the FPO and Endangered liverwort <i>Leiocolea rutheana</i> , the Vulnerable mosses <i>Tomentypnum nitens</i> and <i>Sphagnum warnstorffii</i> , and the Near Threatened moss <i>Sphagnum teres</i> (Lockhart et al., 2012), but again these species cannot be assigned specifically to alkaline fens. Further information can be found within Foss and McGee (1987), Douglas et al. (1989) and NPWS internal files

Conservation Objectives for : Bellacorick Bog Complex SAC [001922]

1013 Geyer's Whorl Snail *Vertigo geyeri*

To maintain the favourable conservation condition of Geyer's Whorl Snail in Bellacorick Bog Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Distribution: occupied sites	Number	No decline. There are three known sites for this species in the SAC within the 1km grid squares G0522, G0718 and G0818. See map 4	There have been records of Geyer's whorl snail (<i>Vertigo geyeri</i>) from three 1km grid squares within the Bellacorick Bog Complex SAC boundary (G0522, G0718 and G0818). This corresponds to the three discrete occupied parts of the SAC at Fermoyale (G0522; site code VgCAM16 in Moorkens and Killeen (2011)); Coolturk (G0718; Holyoak (2005)) and Brackloon (G0818; site code VgCAM3 in Moorkens and Killeen (2011))
Occurrence in suitable habitat	Percentage positive records in a representative number of samples	No decline. A minimum of 50% positive samples in optimum habitat areas; 25% in areas described as sub-optimal	Positive samples mean the confirmed presence of snails (living or recently dead adults and/or juveniles). 50% of samples taken from optimal habitat polygons and 25% of sample from sub-optimal polygons should be positive for the species. Samples should be spread throughout the sub-sites with a minimum of eight in Fermoyale and four in Brackloon. Note that as Coolturk has not been surveyed in detail, no minimum sampling level has been specified. The habitats occupied by Geyer's whorl snail (<i>Vertigo geyeri</i>) in this SAC are areas of fen and flush. See the notes on habitat area below for definitions of optimal and sub-optimal habitats at Fermoyale and at Brackloon. See also Moorkens and Killeen (2011)
Habitat area - Fermoyale	Hectares	Stable or increasing, subject to natural processes. No less than 25.0ha of at least sub-optimal habitat	Suitable conditions for the species can be found throughout the southern portion of the SAC, with the largest known extent in the fen at Fermoyale. Moorkens and Killeen (2011) give a target figure of 25.0ha of habitat in at least sub-optimal condition, and describe the methodology that should be used to assess the target. Optimal habitat is defined for Fermoyale as flush areas of fen and bog with <i>Schoenus nigricans</i> , <i>Carex viridula</i> , <i>C. pulicaria</i> , other low carices, <i>C. rostrata</i> , <i>Menyanthes trifoliata</i> , <i>Campyllum stellatum</i> , <i>Juncus articulatus</i> , <i>Eriophorum angustifolium</i> and orchids. Vegetation height mostly 10-30cm with higher <i>S. nigricans</i> tussocks. During sampling the water table should be between 0-5cm of the soil surface, but not above ground level. Sub-optimal habitat is defined as above but with more <i>S. nigricans</i> tussocks with mosses between them, or overall sward height is >30cm, or the water table is <5cm or ground is flooded at the time of sampling
Habitat area - Brackloon	Hectares	Stable or increasing, subject to natural processes. No less than 0.6ha of at least sub-optimal habitat	The amount of suitable habitat at Brackloon is not precisely known, but Moorkens and Killeen (2011) give a target figure of 0.6ha of habitat for this sub-site in at least sub-optimal condition, and describe the methodology that should be used to assess the target. Optimal habitat is defined for Brackloon as a mosaic of spring flush vegetation 5-55cm tall, containing species such as <i>Carex viridula</i> , <i>Equisetum</i> spp., <i>Menyanthes trifoliata</i> , <i>Platanthera bifolia</i> and mosses, with scattered tussocks of <i>Schoenus nigricans</i> no greater than 50cm tall. During sampling the water table should be between 0-5cm of the soil surface, but not above ground level. Sub-optimal habitat is defined as above (or with <i>Myrica gale</i> , <i>Hydrocotyle vulgaris</i> , <i>Eriophorum angustifolium</i> , <i>Drosera rotundifolia</i> and <i>D. intermedia</i>), but either vegetation height is greater than 50cm, or the water table is below 5cm or ground is flooded at the time of sampling


Habitat area - Coolturk	Hectares	Stable or increasing, subject to natural processes	The amount of suitable habitat at Coolturk is currently unknown
Habitat quality: soil wetness	Percentage of a representative number of monitoring stops	At each sub-site, at least 67% of a representative number of sampling stops in areas of optimal habitat should be classified as optimal wetness as defined by Moorkens and Killeen (2011); at least 25% in areas of sub-optimal habitat	The soil wetness should be assessed using the methodology described in Moorkens and Killeen (2011). In optimal areas, 67% of stops should meet the soil wetness criterion of Moorkens and Killeen (2011); in sub-optimal habitat at least 25% should be optimal wetness


Conservation Objectives for : Bellacorick Bog Complex SAC [001922]

1528 Marsh Saxifrage *Saxifraga hirculus*

To maintain the favourable conservation condition of Marsh Saxifrage in Bellacorick Bog Complex SAC, which is defined by the following list of attributes and targets:


Attribute	Measure	Target	Notes
Distribution	Number and geographical spread of populations	No loss in geographical spread and number of populations, subject to natural processes	The known populations of marsh saxifrage (<i>Saxifraga hirculus</i>) in Bellacorick Bog Complex SAC occur in five flushes: Formoyle, Sheskin A, Sheskin B, Sheskin C and Croaghaun East. See Muldoon (2011) and Muldoon et al. (2015) for further details. The exact locations are not mapped here on account of the presence of sensitive associated species
Population size: number of rosettes	Number	Maintain the size of the known populations, subject to natural processes. The target numbers of rosettes are: at least 136 in Formoyle, at least 288 in Sheskin A, at least 93 in Sheskin B, at least 800 in Sheskin C and at least 1,600 in Croaghaun East	The number of rosettes recorded were: 170 in Formoyle, 360 in Sheskin A, 116 in Sheskin B (Muldoon, 2011), 1,000 in Sheskin C and 2,000 in Croaghaun East (NPWS internal files). The target figures are a c.20% reduction of the recorded numbers to allow a margin of error and variability over monitoring seasons. See Muldoon et al. (2015) for further details
Population size: area of occupancy	Hectares	Maintain the areas of occupancy of the known populations, subject to natural processes. The target areas are: at least 0.0066ha in Formoyle, at least 0.0014ha in Sheskin A, at least 0.0071ha in Sheskin B, at least 0.09ha in Sheskin C and at least 0.1093ha in Croaghaun East	The areas of occupancy were estimated as: 73m ² (0.0073ha) in Formoyle, 16m ² (0.0016ha) in Sheskin A, 79.4m ² (0.0079ha) in Sheskin B (Muldoon, 2011), 1,000m ² (0.1ha) in Sheskin C and 1,214m ² (0.1214ha) in Croaghaun East (NPWS internal files). The target area figures are a 10% reduction of the recorded areas to allow for a margin of error. See Muldoon et al. (2015) for further details
Hydrological conditions: water level	Occurrence of high or fluctuating water levels	Maintain the appropriate natural hydrological regime necessary to support the habitat for the species	In Ireland, marsh saxifrage (<i>Saxifraga hirculus</i>) is now restricted to mineral flushes in blanket bog where rising groundwater forms small streams and seepage areas suitable for the species. Based on Muldoon (2011) and Muldoon et al. (2015)
Vegetation composition: positive indicator species	Occurrence in a representative number of 1m x 1m monitoring stops	Knotted pearlwort (<i>Sagina nodosa</i>) should be present in at least two of five 1m x 1m monitoring stops	The presence of the positive indicator species knotted pearlwort (<i>Sagina nodosa</i>) should be maintained (Muldoon, 2011; Muldoon et al., 2015)
Vegetation composition: negative indicator species	Mean percentage cover in five 1m x 1m monitoring stops	Mean percentage cover of purple moor-grass (<i>Molinia caerulea</i>) should not exceed 5%; mean percentage cover of Yorkshire fog (<i>Holcus lanatus</i>) should not exceed 15%	Low cover of the negative indicator species purple moor-grass (<i>Molinia caerulea</i>) and Yorkshire fog (<i>Holcus lanatus</i>) should be maintained. See Muldoon (2011) and Muldoon et al. (2015) for further details
Vegetation structure: grazing level	Evidence of grazing	Maintain grazing at light to moderate levels to ensure an open vegetation structure and to allow flowering to occur	Undergrazing is a minor issue at Sheskin A and Sheskin B. See Muldoon (2011) and Muldoon et al. (2015) for further details





Legend


- Bellacorick Bog Complex SAC 001922
- River Moy SAC SAC 002298
- OSi Discovery Series County Boundary



**MAP 2:
BELLACORICK BOG COMPLEX SAC
CONSERVATION OBJECTIVES
ADJACENT, ADJOINING AND
OVERLAPPING DESIGNATIONS**

Map to be read in conjunction with the NPWS Conservation Objectives Document.



**SITE CODE: SAC 001922; version 3.
SAC 002298; version 1.14, 1.09, 1.03
CO. MAYO, CO. SLIGO, CO. ROSCOMMON**





The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
Ordnance Survey of Ireland Licence No EN 0059216. © Ordnance Survey of Ireland Government of Ireland.

Níl sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbheithnithe a déanamh ar theorainneacha na gceantar comharthaithe. Suirbhéarachta Ordonáis na hÉireann Ceadúnas Uimh EN 0059216. © Suirbhéarachta Ordonáis na hÉireann Rialtas na hÉireann.







Legend

-  Bellacorick Bog Complex SAC 001922
-  OSi Discovery Series County Boundary

Indicative Lake Habitats

-  Potential 3160 Potential natural dystrophic lakes and ponds



**MAP 3:
BELLACORICK BOG COMPLEX SAC
CONSERVATION OBJECTIVES
INDICATIVE LAKE HABITATS**

Map to be read in conjunction with the NPWS Conservation Objectives Document.


**SITE CODE:
SAC 001922; version 3
CO. MAYO**






The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
Ordnance Survey of Ireland Licence No EN 0059216. © Ordnance Survey of Ireland Government of Ireland.
Níl sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbheithníthe a déanamh ar theorainneacha na gceantar comharthaithe. Suirbhéarachta Ordonáis na hÉireann Ceadúnas Uimh EN 0059216. © Suirbhéarachta Ordonáis na hÉireann Rialtas na hÉireann.



**Map Version 1
Date: Sept 2017**



Legend

-  1013 Geyer's whorl snail *Vertigo geyeri*
-  Bellacorick Bog Complex SAC 001922
-  OSi Discovery Series County Boundary




An Roinn
Cultúir, Oidhreacht agus Gaeltachta
Department of
Culture, Heritage and the Gaeltacht

**MAP 4:
BELLACORICK BOG COMPLEX SAC
CONSERVATION OBJECTIVES
GEYER'S WHORL SNAIL**

Map to be read in conjunction with the NPWS Conservation Objectives Document.

**SITE CODE:
SAC 001922; version 3.
CO. MAYO**



The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
Ordnance Survey of Ireland Licence No EN 0059216. © Ordnance Survey of Ireland Government of Ireland.

Níl sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbheithníthe a déanamh ar theorainneacha na gceantar comharthaithe. Suirbhéarachta Ordonáis na hÉireann Ceadúnas Uimh EN 0059216. © Suirbhéarachta Ordonáis na hÉireann Rialtas na hÉireann.



**Map Version 1
Date: Sept 2017**

National Parks and Wildlife Service

Conservation Objectives Series

Mweelrea/Sheeffry/Erriff Complex SAC 001932



An Roinn
Cultúir, Oidhreachta agus Gaeltachta

Department of
Culture, Heritage and the Gaeltacht



**National Parks and Wildlife Service,
Department of Culture, Heritage and the Gaeltacht,
7 Ely Place, Dublin 2, Ireland.
Web: www.npws.ie
E-mail: nature.conservation@ahg.gov.ie**

Citation:

**NPWS (2017) Conservation Objectives: Mweelrea/Sheeffry/Erriff Complex SAC
001932. Version 1. National Parks and Wildlife Service, Department of Culture,
Heritage and the Gaeltacht.**

**Series Editor: Rebecca Jeffrey
ISSN 2009-4086**

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

* indicates a priority habitat under the Habitats Directive

001932 Mweelrea/Sheeffry/Erriff Complex SAC

- 1013 Geyer's Whorl Snail *Vertigo geyeri*
- 1014 Narrow-mouthed Whorl Snail *Vertigo angustior*
- 1029 Freshwater Pearl Mussel *Margaritifera margaritifera*
- 1106 Salmon *Salmo salar*
- 1150 Coastal lagoonsE
- 1210 Annual vegetation of drift lines
- 1330 Atlantic salt meadows (*Glauco-Puccinellietalia maritimae*)
- 1355 Otter *Lutra lutra*
- 1395 Petalwort *Petalophyllum ralfsii*
- 1410 Mediterranean salt meadows (*Juncetalia maritimi*)
- 1833 Slender Naiad *Najas flexilis*
- 2110 Embryonic shifting dunes
- 2120 Shifting dunes along the shoreline with *Cladonia* (white dunes)
- 2150 Atlantic decalcified fixed dunes (*Calluno-Ulicetea*)E
- 2170 Dunes with *Salicornia* ssp. *arenariae* (*Salicion arenariae*)
- 21A0 Machairs (* in Ireland)
- 3110 Oligotrophic waters containing very few minerals of sandy plains (*Littorelletalia uniflorae*)
- 3130 Oligotrophic to mesotrophic standing waters with vegetation of the *Littorelletea uniflorae* and/or *Isoeto-Nanojuncetea*
- 3160 Natural dystrophic lakes and ponds
- 3260 Water courses of plain to montane levels with the *Ranunculion fluitantis* and *Callitriche-Batrachion* vegetation
- 4010 Northern Atlantic wet heaths with *Calluna*
- 4030 European dry heaths
- 4060 Alpine and Boreal heaths
- 5130 *Racomitrium lanuginosum* formations on heaths or calcareous grasslands
- 6430 Hydrophilous tall herb fringe communities of plains and of the montane to alpine levels
- 7130 Blanket bogs (* if active bog)
- 7140 Transition mires and quaking bogs
- 7150 Depressions on peat substrates of the *Rhynchosporion*
- 7220 Petrifying springs with tufa formation (*Cratoneurion*)E
- 7230 Alkaline fens
- 8110 Siliceous scree of the montane to snow levels (*Androsacetalia alpinae* and *Galeopsietalia ladani*)
- 8210 Calcareous rocky slopes with chasmophytic vegetation
- 8220 Siliceous rocky slopes with chasmophytic vegetation

Please note that this SAC is adjacent to West Connaght Coast SAC (002998). See map 2. The conservation objectives for this site should be used in conjunction with those for the adjacent site as appropriate.

Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

NPWS Documents

Year :	1987
Title :	The vegetation of Irish rivers
Author :	Heuff, H.
Series :	Unpublished report to NPWS
Year :	1987
Title :	A survey to locate blanket bogs of scientific interest in County Mayo. Part I
Author :	Foss, P.; McGee, E.
Series :	A report commissioned by the Wildlife Service
Year :	1989
Title :	Survey to locate blanket bogs of scientific interest in Mayo. Part II
Author :	Douglas, C.; Garvey, L.; Kelly, L.; O'Sullivan, A.; Van Doorsleer, L.
Series :	A report commissioned by the Wildlife Service
Year :	1995
Title :	Mapping of proposed SAC rivers for <i>Margaritifera margaritifera</i> . A report for the National Parks and Wildlife Service on work carried out from August to October 1995 (in two volumes) Volume 1
Author :	Moorkens, E.
Series :	Unpublished report to NPWS
Year :	1998
Title :	Biomar survey of Irish machair sites 1996
Author :	Crawford, I.; Bleasdale, A.; Conaghan, J.
Series :	Irish Wildlife Manual No. 3
Year :	1998
Title :	Biomar survey of Irish machair sites, 1996. Vol. 2: plant communities
Author :	Crawford, I.; Bleasdale, A.; Conaghan, J.
Series :	Irish Wildlife Manual No. 4
Year :	2002
Title :	<i>Najas flexilis</i> in Donegal
Author :	Roden, C.M.
Series :	Unpublished report to NPWS
Year :	2004
Title :	The distribution of <i>Najas flexilis</i> in Ireland 2002-2004
Author :	Roden, C.M.
Series :	Unpublished report to NPWS
Year :	2005
Title :	Monitoring populations of the freshwater pearl mussel <i>Margaritifera margaritifera</i> . Baseline survey of the Bundorragha River cSAC, County Mayo
Author :	Moorkens, E.
Series :	Unpublished report to NPWS
Year :	2006
Title :	Otter survey of Ireland 2004/2005
Author :	Bailey, M.; Rochford, J.
Series :	Irish Wildlife Manual No. 23

Year :	2007
Title :	Saltmarsh Monitoring Project 2006
Author :	McCorry, M.
Series :	Unpublished report to NPWS
Year :	2007
Title :	Supporting documentation for the Habitats Directive Conservation Status Assessment - backing documents. Article 17 forms and supporting maps
Author :	NPWS
Series :	Unpublished report to NPWS
Year :	2007
Title :	Inventory of Irish coastal lagoons (version 2)
Author :	Oliver, G.
Series :	Unpublished report to NPWS
Year :	2007
Title :	Management prescriptions for <i>Vertigo angustior</i> at cSAC sites for the species in the Republic of Ireland
Author :	Moorkens, E.
Series :	Unpublished report to NPWS
Year :	2009
Title :	Coastal Monitoring Project 2004-2006
Author :	Ryle, T.; Murray, A.; Connolly, K.; Swann, M.
Series :	Unpublished report to NPWS
Year :	2009
Title :	Saltmarsh monitoring project 2007-2008
Author :	McCorry, M.; Ryle, T.
Series :	Unpublished report to NPWS
Year :	2009
Title :	NS II Freshwater pearl mussel sub-basin management plans: fisheries survey. Stage 1 report
Author :	Paul Johnston Associates
Series :	Unpublished report to NPWS
Year :	2009
Title :	NS II Freshwater pearl mussel sub-basin management plans: Monitoring of the freshwater pearl mussel in the Bundorragha
Author :	Moorkens, E.A.
Series :	Unpublished report to NPWS
Year :	2009
Title :	Lichen searches of rivers, loughs, mountains: Cos Wicklow, Sligo, Mayo and Galway. September 2009
Author :	Giavarini, V.
Series :	Unpublished report to NPWS
Year :	2009
Title :	NS II Freshwater pearl mussel sub-basin management plans: Report on biological monitoring of surface water quality in the Bundorragha catchment
Author :	Williams, L.
Series :	Unpublished report to NPWS

Year :	2010
Title :	Second draft Bundorragha freshwater pearl mussel sub-basin management plan (2009-2015). March 2010
Author :	NPWS
Series :	Unpublished document to the Department of the Environment, Heritage and Local Government
Year :	2011
Title :	Monitoring and condition assessment of populations of <i>Vertigo geyeri</i> , <i>Vertigo angustior</i> and <i>Vertigo moulinsiana</i> in Ireland
Author :	Moorkens, E.; Killeen, I.
Series :	Irish Wildlife Manual No. 55
Year :	2011
Title :	Implementation of a <i>Vertigo</i> monitoring programme: <i>Vertigo angustior</i> monitoring at Dooaghtry
Author :	Moorkens, E.A.; Killeen, I.J.
Series :	Unpublished report to NPWS
Year :	2011
Title :	Implementation of a <i>Vertigo</i> monitoring programme: <i>Vertigo geyeri</i> monitoring at Dooaghtry
Author :	Moorkens, E.A.; Killeen, I.J.
Series :	Unpublished report to NPWS
Year :	2012
Title :	The conservation status of juniper formations in Ireland
Author :	Cooper, F.; Stone, R.E.; McEvoy, P.; Wilkins, T.; Reid, N.
Series :	Irish Wildlife Manual No. 63
Year :	2012
Title :	Monitoring populations of the freshwater pearl mussel <i>Margaritifera margaritifera</i> . A condition assessment survey of the freshwater pearl mussel in the Bundorragha River, Co. Mayo
Author :	Moorkens, E.
Series :	Unpublished report to NPWS
Year :	2012
Title :	Ireland Red List No. 8: Bryophytes
Author :	Lockhart, N.; Hodgetts, N.; Holyoak, D.
Series :	Ireland Red List series, NPWS
Year :	2013
Title :	Conservation status assessment for petrifying springs
Author :	Lyons, M.D.; Kelly, D.L.
Series :	Unpublished report to NPWS
Year :	2013
Title :	National otter survey of Ireland 2010/12
Author :	Reid, N.; Hayden, B.; Lundy, M.G.; Pietravalle, S.; McDonald, R.A.; Montgomery, W.I.
Series :	Irish Wildlife Manual No. 76
Year :	2013
Title :	Monitoring survey of Annex I sand dune habitats in Ireland
Author :	Delaney, A.; Devaney, F.M.; Martin, J.M.; Barron, S.J.
Series :	Irish Wildlife Manual No. 75
Year :	2013
Title :	The status of EU protected habitats and species in Ireland. Volume 2. Habitats assessments
Author :	NPWS
Series :	Conservation assessments

Year :	2013
Title :	The status of EU protected habitats and species in Ireland. Volume 3. Species assessments
Author :	NPWS
Series :	Conservation assessments
Year :	2014
Title :	Guidelines for a national survey and conservation assessment of upland vegetation and habitats in Ireland, Version 2.0
Author :	Perrin, P.M.; Barron, S.J.; Roche, J.R.; O'Hanrahan, B.
Series :	Irish Wildlife Manual No. 79
Year :	2014
Title :	Interim Bundorragha condition assessment. Preliminary results from Bundorragha River monitoring work 2014
Author :	Moorkens, E.A.
Series :	Unpublished report to NPWS
Year :	2014
Title :	Targeted survey of <i>Najas flexilis</i>
Author :	Roden, C.; Murphy, P.
Series :	Unpublished report to NPWS
Year :	2014
Title :	National Survey of Upland Habitats (Phase 1, 2010-2011) Site report no 1: Mweelrea/Sheeffry/Erriff Complex cSAC (001932) Co. Mayo (Revision)
Author :	Roche, J.R.; Perrin, P.M.; Barron, S.J.; Daly, O.H.
Series :	Unpublished report to NPWS
Year :	2015
Title :	Habitats Directive Annex I lake habitats: a working interpretation for the purposes of site-specific conservation objectives and Article 17 reporting
Author :	O Connor, Á.
Series :	Unpublished document by NPWS
Year :	2015
Title :	Monitoring methods for <i>Petalophyllum ralfsii</i> (Wils.) Nees & Gottsche (Petalwort) in the Republic of Ireland
Author :	Campbell, C.; Hodgetts, N.; Lockhart, N.
Series :	Irish Wildlife Manual No. 90
Year :	2016
Title :	Monitoring guidelines for the assessment of petrifying springs in Ireland
Author :	Lyons, M.D.; Kelly, D.L.
Series :	Irish Wildlife Manual No. 94
Year :	2016
Title :	Ireland Red List No. 10: Vascular Plants
Author :	Wyse Jackson, M.; FitzPatrick, Ú.; Cole, E.; Jebb, M.; McFerran, D.; Sheehy Skeffington, M.; Wright, M.
Series :	Ireland Red Lists series, NPWS
Year :	2017
Title :	Mweelrea/Sheeffry/Erriff Complex SAC (site code: 1932) Conservation objectives supporting objectives supporting document- coastal lagoons V1
Author :	NPWS
Series :	Conservation objectives supporting document

Year : 2017
Title : Survey and condition assessment of the population of the freshwater mussel *Margaritifera margaritifera* in the Bundorragha River, County Mayo
Author : Moorkens, E.
Series : Unpublished report to NPWS

Year : 2017
Title : Mweelrea/Sheeffry/Erriff Complex SAC (site code: 1932) Conservation objectives supporting document- blanket bogs and associated habitats V1
Author : NPWS
Series : Conservation objectives supporting document

Year : 2017
Title : Mweelrea/Sheeffry/Erriff Complex SAC (site code: 1932) Conservation objectives supporting document- coastal habitats V1
Author : NPWS
Series : Conservation objectives supporting document

Year : 2017
Title : Mweelrea/Sheeffry/Erriff Complex SAC (site code: 1932) Conservation objectives supporting document- *Najas flexilis* V1
Author : NPWS
Series : Conservation objectives supporting document

Other References

Year : 1982
Title : Otter survey of Ireland
Author : Chapman, P.J.; Chapman, L.L.
Series : Unpublished report to Vincent Wildlife Trust

Year : 1982
Title : Eutrophication of waters. Monitoring assessment and control
Author : OECD
Series : OECD, Paris

Year : 1984
Title : Studies on the biology of freshwater mussels (Lamellibranchia: Unionacea) in Ireland
Author : Ross, E.D.
Series : Unpublished MSc Thesis. National University of Ireland, Galway

Year : 1988
Title : The reproductive biology of freshwater mussels in Ireland, with observations on their distribution and demography
Author : Ross, E.D.
Series : Unpublished Ph.D. Thesis, National University of Ireland, Galway

Year : 1991
Title : The spatial organization of otters (*Lutra lutra*) in Shetland
Author : Kruuk, H.; Moorhouse, A.
Series : Journal of Zoology, 224: 41-57

Year : 1993
Title : The non-marine Mollusca of Dooaghtry, Co. Mayo: changes in the fauna since 1910
Author : Tattersfield, P.
Series : Irish Naturalists' Journal, 24: 183-192

Year :	1996
Title :	Studies on the biology and ecology of <i>Margaritifera</i> in Ireland
Author :	Moorkens, E.
Series :	Unpublished Ph.D. thesis, University of Dublin, Trinity College.
Year :	1999
Title :	Diet of otters (<i>Lutra lutra</i>) on Inishmore, Aran Islands, west coast of Ireland
Author :	Kingston, S.; O'Connell, M.; Fairley, J.S.
Series :	Biology and Environment: Proceedings of the Royal Irish Academy, 99B: 173-182
Year :	2000
Title :	Colour in Irish lakes
Author :	Free, G.; Allott, N.; Mills, P.; Kennelly, C.; Day, S.
Series :	Verhandlungen Internationale Vereinigung für theoretische und angewandte Limnologie, 27: 2620-2623
Year :	2001
Title :	Aquatic plants in Britain and Ireland
Author :	Preston, C.D.; Croft, J.M.
Series :	Harley Books, Colchester
Year :	2002
Title :	Deterioration of Atlantic soft water macrophyte communities by acidification, eutrophication and alkalinisation
Author :	Arts, G.H.P.
Series :	Aquatic Botany, 73: 373-393
Year :	2003
Title :	Ecology of watercourses characterised by Ranunculion fluitantis and Callitriche-Batrachion vegetation
Author :	Hatton-Ellis, T.W.; Grieve, N.
Series :	Conserving Natura 2000 Rivers Ecology Series No. 11. English Nature, Peterborough
Year :	2004
Title :	The ecology of <i>Najas flexilis</i>
Author :	Wingfield, R.A.; Murphy, K.J.; Hollingsworth, P.; Gaywood, M.J.
Series :	Scottish Natural Heritage Commissioned Report No. 017 (ROAME No. F98PA02)
Year :	2004
Title :	Non-marine Mollusca: New and notable records for Ireland
Author :	Moorkens, E.A.
Series :	Bulletin of the Irish Biogeographical Society, 28: 189-198
Year :	2005
Title :	Widespread occurrence of <i>Vertigo geyeri</i> (Gastropoda: Vertiginidae) in north and west Ireland
Author :	Holyoak, G.A.
Series :	Irish Naturalists' Journal, 28: 141-150
Year :	2006
Title :	Otters - ecology, behaviour and conservation
Author :	Kruuk, H.
Series :	Oxford University Press
Year :	2006
Title :	The status of host fish populations and fish species richness in European freshwater pearl mussel (<i>Margaritifera margaritifera</i>) streams
Author :	Geist, J.; Porkka, M.; Kuehn, R.
Series :	Aquatic Conservation: Marine and Freshwater Ecosystems, 16: 251-266

Year :	2006
Title :	The vegetation of Irish machair
Author :	Gaynor, K.
Series :	Biology and Environment: Proceedings of the Royal Irish Academy, vol 106B, No. 3: 311-321
Year :	2006
Title :	A reference-based typology and ecological assessment system for Irish lakes. Preliminary investigations. Final report. Project 2000-FS-1-M1 Ecological assessment of lakes pilot study to establish monitoring methodologies EU (WFD)
Author :	Free, G.; Little, R.; Tierney, D.; Donnelly, K.; Coroni, R.
Series :	EPA, Wexford
Year :	2008
Title :	The phytosociology and conservation value of Irish sand dunes
Author :	Gaynor, K.
Series :	Unpublished Ph.D. Thesis, National University of Ireland, Dublin
Year :	2008
Title :	Water Quality in Ireland 2004-2006
Author :	Clabby, K.J.; Bradley, C.; Craig, M.; Daly, D.; Lucey, J.; McGarrigle, M.; O'Boyle, S.; Tierney, D.; Bowman, J.
Series :	EPA, Wexford
Year :	2009
Title :	The identification, characterization and conservation value of isoetid lakes in Ireland
Author :	Free, G.; Bowman, J.; McGarrigle, M.; Little, R.; Coroni, R.; Donnelly, K.; Tierney, D.; Trodd, W.
Series :	Aquatic Conservation: Marine and Freshwater Ecosystems, 19(3): 264–273
Year :	2010
Title :	Otter tracking study of Roaringwater Bay
Author :	De Jongh, A.; O'Neill, L.
Series :	Unpublished draft report to NPWS
Year :	2010
Title :	Addressing the conservation and rehabilitation of <i>Margaritifera margaritifera</i> populations in the Republic of Ireland within the framework of the habitats and species directive
Author :	Moorkens, E.
Series :	Journal of Conchology, 40: 339
Year :	2010
Title :	Water quality in Ireland 2007-2009
Author :	McGarrigle, M.; Lucey, J.; Ó Cinnéide, M.
Series :	EPA, Wexford
Year :	2012
Title :	The impact of conifer plantation forestry on the ecology of peatland lakes
Author :	Drinan, T.J.
Series :	Unpublished Ph.D. thesis, University College Cork
Year :	2013
Title :	Monitoring and Assessment of Irish Lagoons for the purpose of the EU Water Framework Directive
Author :	Roden, C.M.; Oliver, G.
Series :	Unpublished Report to the EPA

Year :	2013
Title :	Conservation of selected legally protected and Red Listed bryophytes in Ireland
Author :	Campbell, C.
Series :	Unpublished Ph.D. Thesis, Trinity College Dublin
Year :	2013
Title :	Management strategies for the protection of high status water bodies
Author :	Ní Chatháin, B.; Moorkens, E.; Irvine, K.
Series :	Strive Report Series No. 99. EPA, Wexford
Year :	2013
Title :	Interpretation manual of European Union habitats- Eur 28
Author :	European Commission- DG Environment
Series :	European Commission
Year :	2014
Title :	Assessing near-bed velocity in a recruiting population of the endangered freshwater pearl mussel (<i>Margaritifera margaritifera</i>) in Ireland
Author :	Moorkens, E.; Killeen, I.
Series :	Aquatic Conservation: Marine and Freshwater Ecosystems, 24(6): 853-862
Year :	2014
Title :	The impact of rural land management changes on soil hydraulic properties and runoff processes: results from experimental plots in upland UK
Author :	Marshall, M.R.; Ballard, C.E.; Frogbrook, Z.L.; Solloway, I.; McIntyre, N.; Reynolds, B.; Wheater, H.S.
Series :	Hydrological Processes, 28: 2617-2629
Year :	2015
Title :	Water quality in Ireland 2010-2012
Author :	Bradley, C.; Byrne, C.; Craig, M.; Free, G.; Gallagher, T.; Kennedy, B.; Little, R.; Lucey, J.; Mannix, A.; McCreesh, P.; McDermott, G.; McGarrigle, M.; Ní Longphuirt, S.; O'Boyle, S.; Plant, C.; Tierney, D.; Trodd, W.; Webster, P.; Wilkes, R.; Wynne, C.
Series :	EPA, Wexford
Year :	2015
Title :	The flora and conservation status of petrifying springs in Ireland
Author :	Lyons, M.D.
Series :	Unpublished Ph.D. Thesis, Trinity College Dublin
Year :	2016
Title :	A narrative for conserving freshwater and wetland habitats in England
Author :	Mainstone, C.; Hall, R.; Diack, I.
Series :	Natural England Research Reports Number 064
Year :	2016
Title :	The Status of Irish Salmon Stocks in 2015 with Precautionary Catch Advice for 2016
Author :	SSCS (Standing Scientific Committee on Salmon)
Series :	Independent Scientific Report to Inland Fisheries Ireland

Spatial data sources

Year :	Revision 2011
Title :	Inventory of Irish Coastal Lagoons. Version 3
GIS Operations :	Clipped to SAC boundary
Used For :	1150 (map 3)
Year :	Revision 2010
Title :	Saltmarsh Monitoring Project 2007-2008. Version 1
GIS Operations :	QIs selected; clipped to SAC boundary; overlapping regions with Coastal CO data investigated and resolved with expert opinion used
Used For :	1330, 1410 (map 4)
Year :	2013
Title :	Sand Dune Monitoring Project 2011. Version 1
GIS Operations :	QIs selected; clipped to SAC boundary; overlapping regions with Saltmarsh CO data investigated and resolved with expert opinion used
Used For :	1210, 2110, 2120, 2150, 2170, 21A0 (map 5)
Year :	2008
Title :	OSi 1:5000 IG vector dataset
GIS Operations :	WaterPolygons feature class clipped to the SAC boundary. Expert opinion used to identify Annex I habitat and to resolve any issues arising
Used For :	3110, 3130, 3160 (map 6)
Year :	2011
Title :	National Survey of Upland Habitats
GIS Operations :	Habitat dataset for site clipped to SAC boundary. Relevant QI selected and exported to new dataset. Expert opinion used as necessary to resolve any issues arising
Used For :	4010, 4030, 4060, 6430, 7130, 7140, 7150, 7230, 8110, 8210, 8220 (maps 7, 8, 9, 10, 11, 12, 13, 15, 16, 17, 18)
Year :	Derived 2012
Title :	Internal NPWS files
GIS Operations :	Dataset created from spatial reference contained in files
Used For :	7220 (map 14)
Year :	2017
Title :	NPWS rare and threatened species database
GIS Operations :	Dataset created from spatial references in database records. Expert opinion used as necessary to resolve any issues arising
Used For :	1013, 1014, 1029, 1395 (maps 19, 20 and 22)
Year :	Revision 2012
Title :	Margaritifera Sensitive Areas data
GIS Operations :	Relevant catchment boundary identified. Expert opinion used as necessary to resolve any issues arising
Used For :	1029 (map 20)
Year :	2005
Title :	OSi Discovery series vector data
GIS Operations :	Creation of 80m buffer on marine side of high water mark (HWM); creation of 10m buffer on terrestrial side of HWM; combination of 80m and 10m HWM buffer datasets; creation of 10m buffer on terrestrial side of river banks data; creation of 20m buffer applied to canal centreline data. These datasets combined with derived EPA WFD Waterbodies data and Coastal Lagoon data for the 1355 CO. Overlapping regions investigated and resolved; resulting dataset clipped to SAC boundary. Expert opinion used as necessary to resolve any issues arising. Creation of 250m buffer on marine side of HWM to highlight potential commuting points
Used For :	1355 (map 21)

Year :	2010
Title :	EPA WFD Waterbodies data
GIS Operations :	Creation of 20m buffer applied to river and stream centreline data; creation of 80m buffer on aquatic side of lake data; creation of 10m buffer on terrestrial side of lake data. These datasets combined with derived OSi data and coastal lagoon data for the 1355 CO. Overlapping regions investigated and resolved; resulting dataset clipped to SAC boundary. Expert opinion used as necessary to resolve any issues arising. Creation of 250m buffer on aquatic side of the lake boundary to highlight potential commuting points
Used For :	1355 (map 21)
<hr/>	
Year :	Revision 2011
Title :	Inventory of Irish Coastal Lagoons. Version 3
GIS Operations :	Creation of 80m buffer on the aquatic side of lagoon data; creation of 10m buffer on the terrestrial side of lagoon data. These datasets combined with derived OSi data and EPA WFD Waterbodies data for the 1355 CO. Overlapping regions investigated and resolved; resulting dataset clipped to SAC boundary. Expert opinion used as necessary to resolve any issues arising
Used For :	1355 (map 21)
<hr/>	
Year :	2013
Title :	Najas flexilis data
GIS Operations :	Lake habitat for species clipped to SAC boundary
Used For :	1833 (map 23)
<hr/>	

1150 Coastal lagoons

To restore the favourable conservation condition of Coastal lagoons* in Mweelrea/Sheeffry/Erriff Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable, subject to slight natural variation. Favourable reference area: 7.9ha. See map 3	Area calculated from spatial data derived from Oliver (2007) for Corragaun Lough (site code IL071). See the Mweelrea/Sheeffry/Erriff Complex SAC conservation objectives supporting document for coastal lagoons for further details
Habitat distribution	Occurrence	No decline, subject to natural processes. See map 3 for mapped lagoon	Site code IL071 (Corragaun Lough) in Oliver (2007). See the lagoon supporting document for further details
Salinity regime	Practical salinity units (psu)	Annual median salinity and temporal variation within natural ranges	Corragaun Lough is recorded as a mesohaline to euhaline lagoon. See the lagoon supporting document for further details
Hydrological regime	Metres	Annual water level fluctuations and minima within natural ranges	The maximum depth of Corragaun Lough is recorded as less than 1m. See the lagoon supporting document for further details
Barrier: connectivity between lagoon and sea	Permeability	Appropriate hydrological connections between lagoon and sea, including where necessary, appropriate management	Corragaun Lough is described as a natural sedimentary lagoon. See the lagoon supporting document for further details
Water quality: Chlorophyll <i>a</i>	µg/L	Annual median chlorophyll <i>a</i> within natural ranges and less than 5µg/L	Target based on Roden and Oliver (2013). See the lagoon supporting document for further details
Water quality: Molybdate Reactive Phosphorus (MRP)	mg/L	Annual median MRP within natural ranges and less than 0.1mg/L	Target based on Roden and Oliver (2013). See the lagoon supporting document for further details
Water quality: Dissolved Inorganic Nitrogen (DIN)	mg/L	Annual median DIN within natural ranges and less than 0.15mg/L	Target based on Roden and Oliver (2013). See the lagoon supporting document for further details
Depth of macrophyte colonisation	Metres	Macrophyte colonisation to maximum depth of lagoon	As the lagoon is less than 2m deep, it is expected that macrophyte colonisation would extend to the full depth. See the lagoon supporting document for further details
Typical plant species	Number and m ²	Maintain number and extent of listed lagoonal specialists, subject to natural variation	Species listed in Oliver (2007). See the lagoon supporting document for further details
Typical animal species	Number	Maintain listed lagoonal specialists, subject to natural variation	Species listed in Oliver (2007). See the lagoon supporting document for further details
Negative indicator species	Number and percentage cover	Negative indicator species absent or under control	Low salinity, shallow water and elevated nutrient levels increase the threat of accelerated encroachment by reedbeds. See the lagoon supporting document for further details

1210 Annual vegetation of drift lines

To maintain the favourable conservation condition of Annual vegetation of drift lines in Mweelrea/Sheeffry/Erriff Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes including erosion and succession. For the sub-site mapped: Dooaghtry - 0.18ha. See map 5	Based on data from the Sand Dunes Monitoring Project (SDM) (Delaney et al., 2013). Annual vegetation of drift lines was mapped at the sub-site Dooaghtry (SDM site ID: 108) to give a total estimated area of 0.18ha within Mweelrea/Sheeffry/Erriff Complex SAC. The habitat is very difficult to measure in view of its dynamic nature which means that it can appear and disappear within a site from year to year. See the Mweelrea/Sheeffry/Erriff Complex SAC conservation objectives supporting document for coastal habitats for further details
Habitat distribution	Occurrence	No decline or change in habitat distribution, subject to natural processes. See map 5 for known distribution	Based on data from Delaney et al. (2013). The greatest area of the annual vegetation of drift lines habitat is found in the southern part of the Dooaghtry sub-site. See the coastal habitats supporting document for further details
Physical structure: functionality and sediment supply	Presence/absence of physical barriers	Maintain the natural circulation of sediment and organic matter, without any physical obstructions	Based on data from Delaney et al. (2013). Dunes are naturally dynamic systems that require continuous supply and circulation of sand. Accumulation of organic matter in tidal litter is essential for trapping sand and initiating dune formation. Physical barriers will effect sediment supply at these sites. Coastal defences are present close to the outflow of the Owenadornaun river, but these appear to be longstanding structures. See the coastal habitats supporting document for further details
Vegetation structure: zonation	Occurrence	Maintain the range of coastal habitats including transitional zones, subject to natural processes including erosion and succession	Based on data from Ryle et al. (2009) and Delaney et al. (2013). There are transitional communities between a range of sand dune habitats in the SAC. See the coastal habitats supporting document for further details
Vegetation composition: typical species and sub-communities	Percentage cover at a representative number of monitoring stops	Maintain the presence of species-poor communities with typical species: sea rocket (<i>Cakile maritima</i>), sea sandwort (<i>Honckenya peploides</i>), prickly saltwort (<i>Salsola kali</i>) and oraches (<i>Atriplex</i> spp.)	Based on data from Ryle et al. (2009) and Delaney et al. (2013). Typical species of annual vegetation of drift lines occurring at Dooaghtry includes prickly saltwort (<i>Salsola kali</i>). See the coastal habitats supporting document for further details
Vegetation composition: negative indicator species	Percentage cover	Negative indicator species (including non-native species) to represent less than 5% cover	Based on data from Gaynor (2008), Ryle et al. (2009) and Delaney et al. (2013). Negative indicators include non-native species, species indicative of changes in nutrient status and species not considered characteristic of the habitat. See the coastal habitats supporting document for further details

1330 Atlantic salt meadows (*Glauco-Puccinellietalia maritimae*)

To restore the favourable conservation condition of Atlantic salt meadows (*Glauco-Puccinellietalia maritimae*) in Mweelrea/Sheeffry/Erriff Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes including erosion and succession. For the sub-sites (Dooaghtry and Aasleagh Falls) and potential areas mapped: 21.2ha. See map 4	Based on data from the Saltmarsh Monitoring Project (SMP) (McCorry, 2007; McCorry and Ryle, 2009). Two sub-sites that support Atlantic salt meadows (ASM) were mapped, Dooaghtry (SMP site ID: SMP0010) - 16.23ha and Aasleagh Falls (SMP site ID: SMP0106) - 0.31ha, and additional areas of potential ASM habitat (4.66ha) were identified from an examination of aerial photographs, giving a total estimated area of 21.2ha within Mweelrea/Sheeffry/Erriff Complex SAC. NB further unsurveyed areas may be present within the SAC. See the Mweelrea/Sheeffry/Erriff Complex SAC conservation objectives supporting document for coastal habitats for further details
Habitat distribution	Occurrence	No decline or change in habitat distribution, subject to natural processes. See map 4 for mapped and potential distribution	Based on data from McCorry (2007) and McCorry and Ryle (2009). In the northern section of the Dooaghtry sub-site, which is located on the seaward side of Corragaun Lough, ASM is present on both sides of the entrance/outflow from Corragaun Lough. The southern section of the Dooaghtry sub-site occurs to the south-east of Dooaghtry machair at Trawleckachoolia Bay. ASM occupies only a small proportion of the Aasleagh Falls sub-site and is generally poorly developed with only small fragments present. See the coastal habitats supporting document for further details. NB further unsurveyed areas may be present within the SAC
Physical structure: sediment supply	Presence/absence of physical barriers	Maintain natural circulation of sediments and organic matter, without any physical obstructions	Based on data from McCorry (2007) and McCorry and Ryle (2009). There have been significant changes to the seaward side of Corragaun Lough and the channel connecting it to the sea. Up to half of the lough (classified as a lagoon) has in-filled by sand accretion. See the coastal habitats supporting document and the conservation objective for coastal lagoons (1150) for further details
Physical structure: creeks and pans	Occurrence	Maintain creek and pan structure, subject to natural processes including erosion and succession	Based on data from McCorry (2007) and McCorry and Ryle (2009). In the northern section of the Dooaghtry sub-site, the saltmarsh topography is poorly developed, with few salt pans and creeks. The ASM saltmarsh in the main part of the southern section of the Dooaghtry sub-site has an excellent creek and salt pan structure. The saltmarsh topography of the ASM in the Aasleagh Falls sub-site is poorly developed, which is typical of these small fragments of saltmarsh. See the coastal habitats supporting document for further details
Physical structure: flooding regime	Hectares flooded; frequency	Maintain natural tidal regime	Based on data from McCorry (2007) and McCorry and Ryle (2009). See the coastal habitats supporting document for further details
Vegetation structure: zonation	Occurrence	Maintain the range of coastal habitats including transitional zones, subject to natural processes including erosion and succession	Based on data from McCorry (2007) and McCorry and Ryle (2009). The largest area of saltmarsh in the northern section of the Dooaghtry sub-site is part of a large flat coastal plain and there is a gradual transition to machair vegetation communities towards the west. In the ASM habitat in the Aasleagh Falls sub-site, typical zonation of the shoreline is poorly developed. There are natural transitions to other coastal habitats at both the lower and upper ASM boundaries. See the coastal habitats supporting document for further details

Vegetation structure: vegetation height	Centimetres	Maintain structural variation within sward	Based on data from McCorry (2007) and McCorry and Ryle (2009). The northern section of the Dooaghtry sub-site around the seaward side of Corragaun Lough is badly affected by overgrazing from sheep. The ASM sward height in the Aasleagh Falls sub-site varies between 1-10cm high and is light to moderately grazed overall. See the coastal habitats supporting document for further details
Vegetation structure: vegetation cover	Percentage cover at a representative number of monitoring stops	Maintain more than 90% of the area outside of creeks vegetated	Based on data from McCorry (2007) and McCorry and Ryle (2009). The northern section of the Dooaghtry sub-site has undergone damage caused by heavy overgrazing by sheep and by vehicle wheel ruts. See the coastal habitats supporting document for further details
Vegetation composition: typical species and sub-communities	Percentage cover at a representative number of monitoring stops	Maintain range of sub-communities with typical species listed in McCorry and Ryle (2009)	Based on data from McCorry (2007) and McCorry and Ryle (2009). In the Dooaghtry sub-site, ASM saltmarsh vegetation is dominated by saltmarsh rush (<i>Juncus gerardii</i>) and sea milkwort (<i>Glaux maritima</i>), with sea plantain (<i>Plantago maritima</i>), buck's-horn plantain (<i>P. coronopus</i>), common saltmarsh-grass (<i>Puccinellia maritima</i>), red fescue (<i>Festuca rubra</i>), thrift (<i>Armeria maritima</i>), sea arrowgrass (<i>Triglochin maritimum</i>), creeping bent (<i>Agrostis stolonifera</i>), brookweed (<i>Samolus valerandi</i>) and sea arrowgrass (<i>Triglochin maritimum</i>). Species diversity in the ASM was typical of the habitat in the Aasleagh Falls sub-site. Turf fucoids, an indicator of local distinctiveness, were recorded at several locations in the ASM habitat; this is fairly typical of fringe type saltmarshes along the west coast of Ireland. See the coastal habitats supporting document for further details
Vegetation composition: negative indicator species - <i>Spartina anglica</i>	Hectares	There is no record of common cordgrass (<i>Spartina anglica</i>) in the SAC and its establishment should be prevented	Based on data from McCorry (2007) and McCorry and Ryle (2009). The SMP did not record common cordgrass (<i>Spartina anglica</i>) in the Dooaghtry or Aasleagh Falls sub-sites. See the coastal habitats supporting document for further details

1410 Mediterranean salt meadows (*Juncetalia maritimi*)

To restore the favourable conservation condition of Mediterranean salt meadows (*Juncetalia maritimi*) in Mweelrea/Sheeffry/Erriff Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes including erosion and succession. For the sub-sites (Dooaghtry and Aasleagh Falls) and potential areas mapped: 3.36ha. See map 4	Based on data from the Saltmarsh Monitoring Project (SMP) (McCorry, 2007; McCorry and Ryle, 2009). Two sub-sites that support Mediterranean salt meadows (MSM) were mapped, Dooaghtry (SMP site ID: SMP0010) - 1.13ha and Aasleagh Falls (SMP site ID: SMP0106) - 2.2ha, and additional areas of potential MSM habitat (0.03ha) were identified from an examination of aerial photographs, giving a total estimated area of 3.36ha within Mweelrea/Sheeffry/Erriff Complex SAC. NB further unsurveyed areas may be present within the SAC. See the Mweelrea/Sheeffry/Erriff Complex SAC conservation objectives supporting document for coastal habitats for further details
Habitat distribution	Occurrence	No decline or change in habitat distribution, subject to natural processes. See map 4 for mapped and potential distribution	Based on data from McCorry (2007) and McCorry and Ryle (2009). In the Dooaghtry sub-site, MSM habitat is predominantly present in the southern section and is mainly located along the landward boundary. MSM dominates the saltmarsh in the Aasleagh Falls sub-site. See the coastal habitats supporting document for further details. NB further unsurveyed areas may be present within the SAC
Physical structure: sediment supply	Presence/absence of physical barriers	Maintain natural circulation of sediments and organic matter, without any physical obstructions	Based on data from McCorry (2007) and McCorry and Ryle (2009). See the coastal habitats supporting document for further details
Physical structure: creeks and pans	Occurrence	Maintain creek and pan structure, subject to natural processes including erosion and succession	Based on data from McCorry (2007) and McCorry and Ryle (2009). There are few creeks and pans within the MSM habitat in the Dooaghtry sub-site. The saltmarsh topography of the MSM in the Aasleagh Falls sub-site is also poorly developed, although there are some small creeks present in the most extensive area. See the coastal habitats supporting document for further details
Physical structure: flooding regime	Hectares flooded; frequency	Maintain natural tidal regime	Based on data from McCorry (2007) and McCorry and Ryle (2009). Mediterranean salt meadow is found high up in the saltmarsh but requires occasional tidal inundation. See the coastal habitats supporting document for further details
Vegetation structure: zonation	Occurrence	Maintain the range of coastal habitats including transitional zones, subject to natural processes including erosion and succession	Based on data from McCorry (2007) and McCorry and Ryle (2009). In the Dooaghtry sub-site, the MSM is generally located at the back of the southern saltmarsh area and there are only narrow patches of transitional (brackish and freshwater marsh) habitats before the development of terrestrial habitats as the slope is relatively steep at the back of the saltmarsh. Shoreline zonation is relatively poorly developed in the MSM habitat in the Aasleagh Falls sub-site. There are natural transitions to other coastal habitats at both the lower and upper MSM boundaries. See the coastal habitats supporting document for further details
Vegetation structure: vegetation height	Centimetres	Maintain structural variation in the sward	Based on data from McCorry (2007) and McCorry and Ryle (2009). In the Dooaghtry sub-site, the sward height is varied in the sea rush (<i>Juncus maritimus</i>) dominated MSM areas. The sward height of the MSM in the Aasleagh Falls sub-site is typical of the habitat and varies between 5-10cm. See the coastal habitats supporting document for further details

Vegetation structure: vegetation cover	Percentage cover at a representative number of monitoring stops	Maintain more than 90% of the area outside of creeks vegetated	Based on data from McCorry (2007) and McCorry and Ryle (2009). See the coastal habitats supporting document for further details
Vegetation composition: typical species and sub-communities	Percentage cover at a representative number of monitoring stops	Maintain range of sub-communities with typical species listed in McCorry and Ryle (2009)	Based on data from McCorry (2007) and McCorry and Ryle (2009). In the MSM habitat in the southern section of the Dooaghtry sub-site, species diversity within the clumps of sea rush (<i>Juncus maritimus</i>) is relatively high. Other species that occur frequently amongst the sea rush include red fescue (<i>Festuca rubra</i>), thrift (<i>Armeria maritima</i>), autumn hawkbit (<i>Leontodon autumnalis</i>), sea plantain (<i>Plantago maritima</i>) and white clover (<i>Trifolium repens</i>). The MSM habitat in the Aasleagh Falls sub-site contained a typical species assemblage with the vegetation generally being dominated by sea rush and frequent creeping bent (<i>Agrostis stolonifera</i>). Turf fucoids were also recorded in the MSM habitat. See the coastal habitats supporting document for further details
Vegetation composition: negative indicator species - <i>Spartina anglica</i>	Hectares	There is no record of common cordgrass (<i>Spartina anglica</i>) in the SAC and its establishment should be prevented	Based on data from McCorry (2007) and McCorry and Ryle (2009). The SMP did not record common cordgrass (<i>Spartina anglica</i>) in the Dooaghtry or Aasleagh Falls sub-sites. See the coastal habitats supporting document for further details

2110 Embryonic shifting dunes

To maintain the favourable conservation condition of Embryonic shifting dunes in Mweelrea/Sheeffry/Erriff Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes including erosion and succession. For the sub-site mapped: Dooaghtry - 0.53ha. See map 5	Based on data from the Sand Dunes Monitoring Project (SDM) (Delaney et al., 2013). Embryonic shifting dunes habitat was mapped at the sub-site Dooaghtry (SDM site ID: 108) to give a total estimated area of 0.53ha within Mweelrea/Sheeffry/Erriff Complex SAC. The habitat is very difficult to measure in view of its dynamic nature. See the Mweelrea/Sheeffry/Erriff Complex SAC conservation objectives supporting document for coastal habitats for further details
Habitat distribution	Occurrence	No decline or change in habitat distribution, subject to natural processes. See map 5 for recorded distribution	Based on data from Delaney et al. (2013). Embryonic shifting dunes were recorded in two small areas at Dooaghtry. See the coastal habitats supporting document for further details
Physical structure: functionality and sediment supply	Presence/absence of physical barriers	Maintain the natural circulation of sediment and organic matter, without any physical obstructions	Based on data from Delaney et al. (2013). Dunes are naturally dynamic systems that require continuous supply and circulation of sand. Physical barriers can lead to fossilisation or over-stabilisation of dunes, as well as beach starvation, resulting in increased rates of erosion. Coastal defences are present close to the outflow of the Owenadornaun River, but these appear to be longstanding structures. See the coastal habitats supporting document for further details
Vegetation structure: zonation	Occurrence	Maintain the range of coastal habitats including transitional zones, subject to natural processes including erosion and succession	Based on data from Ryle et al. (2009) and Delaney et al. (2013). See the coastal habitats supporting document for further details
Vegetation composition: plant health of foredune grasses	Percentage cover	More than 95% of sand couch grass (<i>Elytrigia juncea</i>) and/or lyme-grass (<i>Leymus arenarius</i>) should be healthy (i.e. green plant parts above ground and flowering heads present)	Based on data from Ryle et al. (2009) and Delaney et al. (2013). All of the shoots of sand couch grass (<i>Elytrigia juncea</i>) were healthy in the habitat in the SAC at the time of the SDM survey. See the coastal habitats supporting document for further details
Vegetation composition: typical species and sub-communities	Percentage cover at a representative number of monitoring stops	Maintain the presence of species-poor communities with typical species: sand couch grass (<i>Elytrigia juncea</i>) and/or lyme-grass (<i>Leymus arenarius</i>)	Based on data from Ryle et al. (2009) and Delaney et al. (2013). See the coastal habitats supporting document for further details
Vegetation composition: negative indicator species	Percentage cover	Negative indicator species (including non-native species) to represent less than 5% cover	Based on data from Gaynor (2008), Ryle et al. (2009) and Delaney et al. (2013). Negative indicators include non-native species, species indicative of changes in nutrient status and species not considered characteristic of the habitat. Sea buckthorn (<i>Hippophae rhamnoides</i>) should be absent or effectively controlled. See the coastal habitats supporting document for further details

2120 Shifting dunes along the shoreline with *Ammophila arenaria* (white dunes)

To maintain the favourable conservation condition of Shifting dunes along the shoreline with *Ammophila arenaria* (white dunes) in Mweelrea/Sheeffry/Erriff Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes including erosion and succession. For the sub-site mapped: Dooaghtry - 12.43ha. See map 5	Based on data from the Sand Dunes Monitoring Project (SDM) (Delaney et al., 2013). Shifting dunes along the shoreline with <i>Ammophila arenaria</i> was mapped at the sub-site Dooaghtry (SDM site ID: 108) to give a total estimated area of 12.43ha within Mweelrea/Sheeffry/Erriff Complex SAC. The habitat is very difficult to measure in view of its dynamic nature. See the Mweelrea/Sheeffry/Erriff Complex SAC conservation objectives supporting document for coastal habitats for further details
Habitat distribution	Occurrence	No decline or change in habitat distribution, subject to natural processes. See map 5 for recorded distribution	Based on data from Delaney et al. (2013). There is a wide band of marram (<i>Ammophila arenaria</i>) dunes along the western side of the dunes at Dooaghtry and for much of the sub-site the habitat fronts directly onto the beach. See the coastal habitats supporting document for further details
Physical structure: functionality and sediment supply	Presence/absence of physical barriers	Maintain the natural circulation of sediment and organic matter, without any physical obstructions	Based on data from Ryle et al. (2009) and Delaney et al. (2013). Dunes are naturally dynamic systems that require continuous supply and circulation of sand. Marram grass (<i>Ammophila arenaria</i>) reproduces vegetatively and requires constant accretion of fresh sand to maintain active growth encouraging further accretion. See the coastal habitats supporting document for further details
Vegetation structure: zonation	Occurrence	Maintain the range of coastal habitats including transitional zones, subject to natural processes including erosion and succession	Based on data from Ryle et al. (2009) and Delaney et al. (2013). See the coastal habitats supporting document for further details
Vegetation composition: plant health of dune grasses	Percentage cover	More than 95% of marram grass (<i>Ammophila arenaria</i>) and/or lyme-grass (<i>Leymus arenarius</i>) should be healthy (i.e. green plant parts above ground and flowering heads present)	Based on data from Ryle et al. (2009) and Delaney et al. (2013). See the coastal habitats supporting document for further details
Vegetation composition: typical species and sub-communities	Percentage cover at a representative number of monitoring stops	Maintain the presence of species-poor communities dominated by marram grass (<i>Ammophila arenaria</i>) and/or lyme-grass (<i>Leymus arenarius</i>)	Based on data from Ryle et al. (2009) and Delaney et al. (2013). See the coastal habitats supporting document for further details
Vegetation composition: negative indicator species	Percentage cover	Negative indicator species (including non-native species) to represent less than 5% cover	Based on data from Gaynor (2008), Ryle et al. (2009) and Delaney et al. (2013). Negative indicators include non-native species, species indicative of changes in nutrient status and species not considered characteristic of the habitat. Sea buckthorn (<i>Hippophae rhamnoides</i>) should be absent or effectively controlled. See the coastal habitats supporting document for further details

2150 Atlantic decalcified fixed dunes (Calluno-Ulicetea)

To maintain the favourable conservation condition of Atlantic decalcified fixed dunes (Calluno-Ulicetea)* in Mweelrea/Sheeffry/Erriff Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes including erosion and succession	Based on data from the Sand Dunes Monitoring Project (SDM) (Delaney et al., 2013). Atlantic decalcified fixed dunes habitat was not recorded by the SDM in the Dooaghtry sub-site. The current status of this habitat in Ireland is under review. See the Mweelrea/Sheeffry/Erriff Complex SAC conservation objectives supporting document for coastal habitats for further details
Habitat distribution	Occurrence	No decline or change in habitat distribution, subject to natural processes	The current distribution and status of the habitat in the SAC is unknown. See the coastal habitats supporting document for further details
Physical structure: functionality and sediment supply	Presence/absence of physical barriers	Maintain the natural circulation of sediment and organic matter, without any physical obstructions	Physical barriers can lead to fossilisation or over-stabilisation of dunes, as well as beach starvation, resulting in increased rates of erosion. See the coastal habitats supporting document for further details
Vegetation structure: zonation	Occurrence	Maintain the range of coastal habitats including transitional zones, subject to natural processes including erosion and succession	Based on data from Ryle et al. (2009) and Delaney et al. (2013). See the coastal habitats supporting document for further details
Vegetation structure: bare ground	Percentage cover	Bare ground should not exceed 10%, subject to natural processes	Based on data from Gaynor (2008), Ryle et al. (2009) and Delaney et al. (2013). See the coastal habitats supporting document for further details
Vegetation composition: sward height	Centimetres	Maintain structural variation within sward	Based on data from Gaynor (2008), Ryle et al. (2009) and Delaney et al. (2013). See the coastal habitats supporting document for further details
Vegetation composition: typical species and sub-communities	Percentage cover at a representative number of monitoring stops	Maintain range of sub-communities with typical species listed in Delaney et al. (2013)	Based on data from Gaynor (2008), Ryle et al. (2009) and Delaney et al. (2013). See the coastal habitats supporting document for further details
Vegetation composition: negative indicator species	Percentage cover	Negative indicator species (including non-native species) to represent less than 5% cover	Based on data from Gaynor (2008), Ryle et al. (2009) and Delaney et al. (2013). Negative indicators include non-native species, species indicative of changes in nutrient status and species not considered characteristic of the habitat. Sea buckthorn (<i>Hippophae rhamnoides</i>) should be absent or effectively controlled. See the coastal habitats supporting document for further details
Vegetation composition: scrub/trees	Percentage cover	No more than 5% cover or under control	Based on data from Ryle et al. (2009) and Delaney et al. (2013). See the coastal habitats supporting document for further details

2170 Dunes with *Salix repens* ssp. *argentea* (Salicion arenariae)

To maintain the favourable conservation condition of Dunes with *Salix repens* ssp. *argentea* (Salicion arenariae) in Mweelrea/Sheeffry/Erriff Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes including erosion and succession	Based on data from the Sand Dunes Monitoring Project (SDM) (Delaney et al., 2013). Dunes with <i>Salix repens</i> ssp. <i>argentea</i> (Salix arenariae) was not recorded by the SDM in the Dooaghtry sub-site. The current area and status of this habitat in the SAC is unknown. The habitat can be difficult to distinguish from 2190 Humid dune slacks. See the Mweelrea/Sheeffry/Erriff Complex SAC conservation objectives supporting document for coastal habitats for further details
Habitat distribution	Occurrence	No decline or change in habitat distribution, subject to natural processes	The current distribution and status of the habitat in the SAC is unknown. See the coastal habitats supporting document for further details
Physical structure: functionality and sediment supply	Presence/absence of physical barriers	Maintain the natural circulation of sediment and organic matter, without any physical obstructions	Physical barriers can lead to fossilisation or over-stabilisation of dunes, as well as beach starvation, resulting in increased rates of erosion. See the coastal habitats supporting document for further details
Vegetation structure: zonation	Occurrence	Maintain the range of coastal habitats including transitional zones, subject to natural processes including erosion and succession	Based on data from Ryle et al. (2009) and Delaney et al. (2013). See the coastal habitats supporting document for further details
Vegetation structure: bare ground	Percentage cover	Bare ground should not exceed 10% cover, subject to natural processes	Based on data from Gaynor (2008), Ryle et al. (2009) and Delaney et al. (2013). See the coastal habitats supporting document for further details
Vegetation structure: vegetation height	Centimetres	Maintain structural variation within sward	Based on data from Gaynor (2008), Ryle et al. (2009) and Delaney et al. (2013). See the coastal habitats supporting document for further details
Vegetation composition: typical species and sub-communities	Percentage cover at a representative number of monitoring stops	Maintain range of sub-communities with typical species listed in Delaney et al. (2013)	See the coastal habitats supporting document for further details
Vegetation composition: cover and height of <i>Salix repens</i>	Percentage cover; centimetres	Maintain more than 10% cover of creeping willow (<i>Salix repens</i>); vegetation height should be in the average range of 5-20cm	Based on data from Gaynor (2008), Ryle et al. (2009) and Delaney et al. (2013). Cover of creeping willow (<i>Salix repens</i>) needs to be maintained (e.g. through an appropriate grazing regime) to prevent the development of a coarse, rank vegetation cover. See the coastal habitats supporting document for further details
Vegetation composition: negative indicator species	Percentage cover at a representative number of monitoring stops	Negative indicator species (including non-native species) to represent less than 5% cover	Based on data from Ryle et al. (2009) and Delaney et al. (2013). Negative indicators include non-native species, species indicative of changes in nutrient status and species not considered characteristic of the habitat. Sea buckthorn (<i>Hippophae rhamnoides</i>) should be absent or effectively controlled. See the coastal habitats supporting document for further details
Vegetation composition: scrub/trees	Percentage cover	For trees and scrub other than creeping willow (<i>Salix repens</i>), there should be no more than 5% cover or their presence should be under control	Based on data from Ryle et al. (2009) and Delaney et al. (2013). See the coastal habitats supporting document for further details

Conservation Objectives for : Mweelrea/Sheeffry/Erriff Complex SAC [001932]

21A0 Machairs (* in Ireland)

To restore the favourable conservation condition of Machairs (* in Ireland) in Mweelrea/Sheeffry/Erriff Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes including erosion and succession. For the sub-site mapped: Dooaghtry - 137.01ha. See map 5	Based on data from the Sand Dunes Monitoring Project (SDM) (Delaney et al., 2013). Machair was recorded and mapped from the sub-site Dooaghtry (SDM site ID: 108) to give a total estimated area of 137.01ha within Mweelrea/Sheeffry/Erriff Complex SAC. See the Mweelrea/Sheeffry/Erriff Complex SAC conservation objectives supporting document for coastal habitats for further details
Habitat distribution	Occurrence	No decline or change in habitat distribution, subject to natural processes. See map 5 for recorded distribution	Based on data from Delaney et al. (2013). Machair is the most extensive coastal Annex I habitat at Dooaghtry. See the coastal habitats supporting document for further details
Physical structure: functionality and sediment supply	Presence/absence of physical barriers	Maintain the natural circulation of sediment and organic matter, without any physical obstructions	Physical barriers can lead to fossilisation or over-stabilisation of dunes, as well as beach starvation, resulting in increased rates of erosion. See the coastal habitats supporting document for further details
Physical structure: hydrological and flooding regime	Water table levels; groundwater fluctuations (metres)	Maintain natural hydrological regime	Based on data from Crawford et al. (1998), Gaynor (2006), Ryle et al. (2009) and Delaney et al. (2013). See the coastal habitats supporting document for further details
Vegetation structure: zonation	Occurrence	Maintain the range of coastal habitats including transitional zones, subject to natural processes including erosion and succession	Based on data from Ryle et al. (2009) and Delaney et al. (2013). In this SAC, machair grades into saltmarsh towards the tidal inlet and to different fen types as it approaches the open water of the lough at Carrickskeewaun. See the coastal habitats supporting document for further details
Vegetation structure: bare ground	Percentage cover	Bare ground should not exceed 10% of the machair habitat, subject to natural processes	Based on data from Ryle et al. (2009) and Delaney et al. (2013). See the coastal habitats supporting document for further details
Vegetation structure: sward height	Centimetres	Maintain structural variation within sward	Based on data from Ryle et al. (2009) and Delaney et al. (2013). Rabbits, cattle and sheep graze the machair at Dooaghtry. See the coastal habitats supporting document for further details
Vegetation composition: typical species and sub-communities	Percentage cover at a representative number of monitoring stops	Maintain range of sub-communities with typical species listed in Delaney et al. (2013)	Based on data from Ryle et al. (2009) and Delaney et al. (2013). Typical machair species present include creeping bent (<i>Agrostis stolonifera</i>), daisy (<i>Bellis perennis</i>), sand sedge (<i>Carex arenaria</i>), glaucous sedge (<i>C. flacca</i>), common mouse-ear (<i>Cerastium fontanum</i>), lady's bedstraw (<i>Galium verum</i>), fairy flax (<i>Linum catharticum</i>), common bird's-foot trefoil (<i>Lotus corniculatus</i>), red bartsia (<i>Odontites vernus</i>), ribwort plantain (<i>Plantago lanceolata</i>), selfheal (<i>Prunella vulgaris</i>) and white clover (<i>Trifolium repens</i>). See the coastal habitats supporting document for further details
Vegetation composition: negative indicator species	Percentage cover	Negative indicator species (including non-native species) to represent less than 5% cover	Based on data from Gaynor (2008), Ryle et al. (2009) and Delaney et al. (2013). Negative indicators include non-native species, species indicative of changes in nutrient status and species not considered characteristic of the habitat. The presence of perennial rye grass (<i>Lolium perenne</i>) in places throughout the machair in the SAC is indicative of agricultural improvement. See the coastal habitats supporting document for further details

Vegetation composition: scrub/trees	Percentage cover	No more than 5% cover or under control	Based on data from Ryle et al. (2009) and Delaney et al. (2013). See the coastal habitats supporting document for further details
Vegetation composition: bryophytes	Percentage cover	Should always be at least an occasional component of the vegetation	Based on data from Ryle et al. (2009) and Delaney et al. (2013). The mosses <i>Climacium dendroides</i> , <i>Homalothecium lutescens</i> , <i>Rhytidiadelphus triquetrus</i> , <i>Thuidium tamariscinum</i> and <i>Syntrichia ruralis</i> subsp. <i>ruraliformis</i> have been recorded from the machair. Petalwort (<i>Petalophyllum ralfsii</i>), a liverwort species listed on Annex II of the EU Habitats Directive and on the Flora (Protection) Order, 2015, has also been recorded on the machair in this SAC (Campbell et al., 2015). See the conservation objective for petalwort (1395) and the coastal habitats supporting document for further details

3110 Oligotrophic waters containing very few minerals of sandy plains (Littorelletalia uniflorae)

To maintain the favourable conservation condition of Oligotrophic waters containing very few minerals of sandy plains (Littorelletalia uniflorae) in Mweelrea/Sheeffry/Erriff Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Lake habitat 3110 is considered likely to occur in most lakes in Mweelrea/Sheeffry/Erriff Complex SAC, from upland corrie lakes such as Loughs Glenawough, Lugacollivee and Bellawaum (where it may co-occur with habitat 3160) to lowland lakes such as Doo, Glencullin and Tawnyard Loughs. It may also co-occur with habitat 3160 in lakes in blanket bog, and with habitat 3130 in Lough Nahaltora and Fin Lough. In line with Article 17 reporting (NPWS, 2013), all lakes larger than 1ha were mapped as potential 3110 (see map 6). It is likely, however, that the habitat also occurs in smaller lakes and ponds. Two measures of extent should be used: 1. the area of the lake itself and; 2. the extent of the vegetation communities/zones that typify the habitat. Further information relating to all attributes is provided in the lake habitats supporting document for the purposes of site-specific conservation objectives and Article 17 reporting (O Connor, 2015)
Habitat distribution	Occurrence	No decline, subject to natural processes	As noted above, all lakes larger than 1ha have been mapped as potential 3110, but the habitat is likely to be even more widespread in the SAC (see map 6)
Typical species	Occurrence	Typical species present, in good condition, and demonstrating typical abundances and distribution	For lists of typical plant species, see the Article 17 habitat assessment for 3110 (NPWS, 2013) and the lake habitats supporting document (O Connor, 2015). See the Mweelrea/Sheeffry/Erriff Complex SAC conservation objectives supporting document for <i>Najas flexilis</i> for information on Lough Nahaltora in the SAC. Information on Fin Lough can be found in Roden and Murphy (2014). Glencullin and Doo Loughs are Water Framework Directive (WFD) monitoring lakes and regular macrophyte surveys are conducted by the Environmental Protection Agency (EPA)
Vegetation composition: characteristic zonation	Occurrence	All characteristic zones should be present, correctly distributed and in good condition	Further work is necessary to describe the characteristic zonation and other spatial patterns in lake habitat 3110 (see O Connor, 2015). Lake-specific information on vegetation zonation may be available from sources such as Roden (2004, 2007 in NPWS, 2007), Roden and Murphy (2014), EPA surveys and others
Vegetation distribution: maximum depth	Metres	Maintain maximum depth of vegetation, subject to natural processes	The maximum depth of vegetation is likely to be specific to the lake shoreline in question. Further work is necessary to develop indicative targets for lake habitat 3110. Maximum depth should be large in lakes in the SAC within undisturbed peatland and uplands; however, pressures such as overgrazing, forestry and peat-cutting may have reduced vegetation depth in some lakes

Hydrological regime: water level fluctuations	Metres	Maintain appropriate natural hydrological regime necessary to support the habitat	Fluctuations in lake water level are typical in Ireland, but can be amplified by activities such as abstraction, drainage and overgrazing. Increased water level fluctuations can increase wave action, up-root vegetation, increase turbidity, alter the substratum and lead to release of nutrients from the sediment. The hydrological regime of the lakes must be maintained so that the area, distribution and depth of the lake habitat and its constituent/characteristic vegetation zones and communities are not reduced. It is possible that the hydrological regimes of some of the lakes in this SAC have been altered by historic overgrazing (faster run-off, higher flood peaks, lower base flows, etc.; see Marshall et al., 2014)
Lake substratum quality	Various	Maintain appropriate substratum type, extent and chemistry to support the vegetation	Research is required to further characterise the substratum types (particle size and origin) and substratum quality (notably pH, calcium, iron and nutrient concentrations) favoured by each of the five Annex I lake habitats in Ireland. It is likely that lake habitat 3110 is associated with a range of nutrient-poor substrates, from stones, cobble and gravel, through sands, silt, clay and peat. Substratum particle size is likely to vary with depth and along the shoreline within a single lake
Water quality: transparency	Metres	Maintain/restore appropriate Secchi transparency. There should be no decline in Secchi depth/transparency	Transparency relates to light penetration and, hence, to the depth of colonisation of vegetation. It can be affected by phytoplankton blooms, water colour and turbidity. Specific targets have yet to be established for lake habitat 3110 (O Connor, 2015). Lake habitat 3110 is associated with very clear water. The OECD fixed boundary system set transparency targets for oligotrophic lakes of $\geq 6\text{m}$ annual mean Secchi disk depth, and $\geq 3\text{m}$ annual minimum Secchi disk depth. Free et al. (2009) found high isoetid abundance in lakes with Secchi depths of more than 3m. Free et al. (2006) give Secchi depth of 2.9m in Glencullin, 5.6m in Doo and 5.8m in Fin Loughs in this SAC
Water quality: nutrients	$\mu\text{g/l P}$; mg/l N	Maintain/restore the concentration of nutrients in the water column to sufficiently low levels to support the habitat and its typical species	As a nutrient-poor habitat, oligotrophic and WFD 'high' status targets apply. Where a lake has nutrient concentrations that are lower than these targets, there should be no decline within class, i.e. no upward trend in nutrient concentrations. For lake habitat 3110, annual average total phosphorus (TP) concentration should be $\leq 10\mu\text{g/l TP}$, average annual total ammonia concentration should be $\leq 0.040\text{mg/l N}$ and annual 95th percentile for total ammonia should be $\leq 0.090\text{mg/l N}$. See also The European Communities Environmental Objectives (Surface Waters) Regulations 2009. Glencullin and Doo Loughs had high nutrient status in 2010-12 (Bradley et al., 2015). Both lakes had failed in 2007-09, however, having good nutrient status and TP $> 10\mu\text{g/l}$ (McGarrigle et al., 2010)
Water quality: phytoplankton biomass	$\mu\text{g/l Chlorophyll } a$	Maintain appropriate water quality to support the habitat, including high chlorophyll <i>a</i> status	Oligotrophic and WFD 'high' status targets apply to lake habitat 3110. Where a lake has a chlorophyll <i>a</i> concentration that is lower than this target, there should be no decline within class, i.e. no upward trend in phytoplankton biomass. The average growing season (March-October) chlorophyll <i>a</i> concentration must be $< 5.8\mu\text{g/l}$. The annual average chlorophyll <i>a</i> concentration should be $< 2.5\mu\text{g/l}$ and the annual peak chlorophyll <i>a</i> concentration should be $\leq 8.0\mu\text{g/l}$. See also The European Communities Environmental Objectives (Surface Waters) Regulations 2009. Glencullin and Doo Loughs had high status in 2007-2009 and 2010-2012 (McGarrigle et al., 2010; Bradley et al., 2015)

Water quality: phytoplankton composition	EPA phytoplankton composition metric	Maintain appropriate water quality to support the habitat, including high phytoplankton composition status	The EPA has developed a phytoplankton composition metric for nutrient enrichment of Irish lakes. As for other water quality indicators, lake habitat 3110 requires WFD high status. Glencullin and Doo Loughs had high phytoplankton composition status in 2010-12 (Bradley et al., 2015)
Water quality: attached algal biomass	Algal cover and EPA phytobenthos metric	Maintain trace/absent attached algal biomass (<5% cover) and high phytobenthos status	Nutrient enrichment can favour epiphytic and epipelic algae that can out-compete the submerged vegetation. The cover abundance of attached algae in lake habitat 3110 should, therefore, be trace/absent (<5% cover). EPA phytobenthos can be used as an indicator of changes in attached algal biomass. As for other water quality indicators, lake habitat 3110 requires high phytobenthos status. Glencullin and Doo Loughs had high phytobenthos status in 2010-12 (Bradley et al., 2015)
Water quality: macrophyte status	EPA macrophyte metric (The Free Index)	Maintain high macrophyte status	Nutrient enrichment can favour more competitive submerged macrophyte species that out-compete the typical and characteristic species for the lake habitat. The EPA monitors macrophyte status for WFD purposes using the 'Free Index'. The target for lake habitat 3110 is high status or an Ecological Quality Ratio (EQR) for lake macrophytes of ≥ 0.90 , as defined in Schedule Five of the European Communities Environmental Objectives (Surface Waters) Regulations 2009. Glencullin and Doo Loughs had high macrophyte status in 2007-2009 and 2010-2012 (McGarrigle et al., 2010; Bradley et al., 2015)
Acidification status	pH units; mg/l	Maintain appropriate water and sediment pH, alkalinity and cation concentrations to support the habitat, subject to natural processes	Acidification can impact on species abundance and composition in soft water lake habitats. In Europe, acidification of isoetid lakes can lead to loss of isoetids and dominance by submerged <i>Sphagnum</i> mosses and <i>Juncus bulbosus</i> (Arts, 2002). The specific requirements of lake habitat 3110, in terms of water and sediment pH, alkalinity and cation concentration, have not been determined. For lake habitat 3110, and adopting a precautionary approach based on Arts (2002), minimum pH should not be <5.5 pH units. Maximum pH should be <9.0 pH units, in line with the surface water standards established for soft waters (where water hardness is ≤ 100 mg/l calcium carbonate). See Schedule Five of the European Communities Environmental Objectives (Surface Waters) Regulations 2009. Glencullin and Doo Loughs passed for acidification status in the 2007-09 and 2010-12 periods (McGarrigle et al., 2010; Bradley et al., 2015)
Water colour	mg/l PtCo	Maintain appropriate water colour to support the habitat	Increased water colour and turbidity decrease light penetration and can reduce the area of available habitat for lake macrophytes, particularly at the lower euphotic depths. The primary source of increased water colour in Ireland is disturbance to peatland. No habitat-specific or national standards for water colour currently exist. Studies have shown median colour concentrations in Irish lakes of 38mg/l PtCo (Free et al., 2000) and 33mg/l PtCo (Free et al., 2006). It is likely that the water colour in all Irish lake habitats would naturally be <50mg/l PtCo. Water colour can be very low (<20mg/l or even <10mg/l PtCo) in lake habitat 3110, where the peatland in the lake's catchment is intact. Free et al. (2006) reported colour of 17mg/l, 8mg/l and 10mg/l PtCo in Glencullin, Doo and Fin Loughs, respectively. Roden (2004) stated the water in Lough Nahaltora was coloured/peat-stained. Overgrazing, peat-cutting or other peatland damage may have increased colour in some lakes in the SAC

Dissolved organic carbon (DOC)	mg/l	Maintain appropriate organic carbon levels to support the habitat	Dissolved (and particulate) organic carbon (OC) in the water column is linked to water colour and acidification (organic acids). Increasing DOC in water has been documented across the Northern Hemisphere, including afforested peatland catchments in Ireland. Damage and degradation of peatland, leading to decomposition of peat is likely to be the predominant source of OC in Ireland. OC in water promotes decomposition by fungi and bacteria that, in turn, releases dissolved nutrients. The increased biomass of decomposers can also impact directly on the characteristic lake communities through shading, competition, etc. Overgrazing and other peatland degradation is also likely to have increased DOC in some lakes in Mweelrea/Sheeffry/Erriff Complex SAC
Turbidity	Nephelometric turbidity units/ mg/l SS/ other appropriate units	Maintain appropriate turbidity to support the habitat	Turbidity can significantly affect the quantity and quality of light reaching rooted and attached vegetation and can, therefore, impact on lake habitats. The settlement of higher loads of inorganic or organic material on lake vegetation communities may also have impacts on sensitive, delicate species. Turbidity can increase as a result of re-suspension of material within the lake, higher loads entering the lake, or eutrophication. Turbidity measurement and interpretation is challenging. As a result, it is likely to be difficult to set habitat-specific targets for turbidity in lakes. Increased loads of fine organic and inorganic particles from overgrazing may have increased turbidity in lakes in the SAC
Fringing habitat: area	Hectares	Maintain the area and condition of fringing habitats necessary to support the natural structure and functioning of habitat 3110	Most lake shorelines have fringing habitats of reedswamp, other swamp, fen, marsh or wet woodland that intergrade with and support the structure and functions of the lake habitat. In this SAC, lake shorelines are likely to have acid grassland, swamp, heath, blanket bog and rock communities. Fringing habitats are dependent on the lake, particularly its water levels, and support wetland communities and species of conservation concern. Many of the fringing wetland habitats support higher invertebrate and plant species richness than the lake habitats themselves

3130 Oligotrophic to mesotrophic standing waters with vegetation of the Littorelletea uniflorae and/or Isoeto-Nanojuncetea

To maintain the favourable conservation condition of Oligotrophic to mesotrophic standing waters with vegetation of the Littorelletea uniflorae and/or of the Isoeto-Nanojuncetea in Mweelrea/Sheeffry/Erriff Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Lake habitat 3130 is considered likely to occur in Lough Nahaltora in Mweelrea/Sheeffry/Erriff Complex SAC, which has a population of <i>Najas flexilis</i> (slender naiad). It may also occur in Fin Lough and other lakes in the SAC, particularly those on more base-rich geology or near the coast (see map 6). It is likely to co-occur with lake habitat 3110 in these lakes. Two measures of extent should be used: 1. the area of the lake itself and; 2. the extent of the vegetation communities/zones that typify the habitat. Further information relating to all attributes is provided in the lake habitats supporting document for the purposes of site-specific conservation objectives and Article 17 reporting (O Connor, 2015). See also the Mweelrea/Sheeffry/Erriff Complex SAC conservation objectives supporting document for <i>Najas flexilis</i>
Habitat distribution	Occurrence	No decline, subject to natural processes	The characteristics and distribution of lake habitat 3130 in Ireland are not yet fully understood. The Annex II macrophyte <i>Najas flexilis</i> (slender naiad) is considered to be characteristic of the habitat and occurs in Lough Nahaltora (see map 6). As noted above, the habitat may be more widespread in the SAC. See O Connor (2015) and the <i>Najas flexilis</i> conservation objectives supporting document for further information
Typical species	Occurrence	Typical species present, in good condition, and demonstrating typical abundances and distribution	For lists of typical plant species, see the Article 17 habitat assessment for 3130 (NPWS, 2013), O Connor (2015) and the <i>Najas flexilis</i> conservation objectives supporting document for further information
Vegetation composition: characteristic zonation	Occurrence	All characteristic zones should be present, correctly distributed and in good condition	Further work is necessary to describe the characteristic zonation and other spatial patterns in lake habitat 3130 (see O Connor, 2015). Lake-specific information on vegetation zonation may be available from Roden (2004, 2007 in NPWS 2007), Roden and Murphy (2014), Environmental Protection Agency (EPA) surveys or other sources
Vegetation distribution: maximum depth	Metres	Maintain maximum depth of vegetation, subject to natural processes	The maximum depth of vegetation is likely to be specific to the lake shoreline in question. Further work is necessary to develop indicative targets for lake habitat 3130. Maximum depth should be large in lakes in the SAC within undisturbed peatland and uplands; however, pressures such as overgrazing, forestry and peat-cutting may have reduced vegetation depth in some lakes

Hydrological regime: water level fluctuations	Metres	Maintain appropriate natural hydrological regime necessary to support the habitat	Fluctuations in lake water level are typical in Ireland, but can be amplified by activities such as abstraction, drainage and overgrazing. Increased water level fluctuations can increase wave action, up-root vegetation, increase turbidity, alter the substratum and lead to release of nutrients from the sediment. The hydrological regime of the lakes must be maintained so that the area, distribution and depth of the lake habitat and its constituent/characteristic vegetation zones and communities are not reduced. It is possible that the hydrological regimes of some of the lakes in Mweelrea/Sheeffry/Erriff Complex SAC have been altered by historic overgrazing (faster run-off, higher flood peaks, lower base flows, etc.; see Marshall et al. (2014))
Lake substratum quality	Various	Maintain appropriate substratum type, extent and chemistry to support the vegetation	Research is required to further characterise the substratum types (particle size and origin) and substratum quality (notably pH, calcium, iron and nutrient concentrations) favoured by each of the five Annex I lake habitats in Ireland. It is likely that lake habitat 3130 is associated with a range of substrate types that are more productive/base-rich relative to the substratum of lake habitat 3110. Substratum particle size is likely to vary with depth and along the shoreline within a single lake; however, it should be noted that <i>Najas flexilis</i> is typically found on soft substrata of mud, silt or fine sand (Preston and Croft, 2001; Roden, 2002, 2004). For further information see the lake habitats supporting document (O Connor, 2015) and the <i>Najas flexilis</i> conservation objectives supporting document
Water quality: transparency	Metres	Maintain/restore appropriate Secchi transparency. There should be no decline in Secchi depth/transparency	Transparency relates to light penetration and, hence, to the depth of colonisation of vegetation. It can be affected by phytoplankton blooms, water colour and turbidity. Specific targets have yet to be established for lake habitat 3130 (O Connor, 2015). Habitat 3130 is associated with clear water, as evidenced by the growth of the character species <i>Najas flexilis</i> at depths of up to 10m. There is likely to be some variation in Secchi depth across lakes with habitat 3130 and site-specific conditions should also be considered. The OECD fixed boundary system set transparency targets for oligotrophic lakes of $\geq 6\text{m}$ annual mean Secchi disk depth, and $\geq 3\text{m}$ annual minimum Secchi disk depth. Roden (2004, 2007 in NPWS, 2007) stated Lough Nahaltora had moderate visibility and peat-stained water. Free et al. (2006) recorded Secchi depth of 5.8m in Fin Lough. Roden and Murphy (2014) stated Fin Lough had very clear water
Water quality: nutrients	$\mu\text{g/l P}$; mg/l N	Maintain the concentration of nutrients in the water column at sufficiently low levels to support the habitat and its typical species	Lake habitat 3130 is associated with high water quality, with naturally low dissolved nutrients. It is naturally more productive than 3110, probably reflecting higher concentrations of nutrients such as calcium, rather than P alone. Lake habitat 3130 may reach favourable condition slightly above the oligotrophic boundary for nutrients, but in the absence of habitat-specific targets, the targets are Water Framework Directive (WFD) 'High Status' or oligotrophic (OECD, 1982). The "good-moderate" boundary is too enriched to support the habitat. Annual average total phosphorus (TP) concentration should be $\leq 10\mu\text{g/l TP}$, average annual total ammonia concentration should be $\leq 0.040\text{mg/l N}$ and annual 95th percentile for total ammonia should be $\leq 0.090\text{mg/l N}$. Where nutrient concentrations are lower, there should be no upward trend in nutrient concentration. See also The European Communities Environmental Objectives (Surface Waters) Regulations 2009

Water quality: phytoplankton biomass	µg/l Chlorophyll <i>a</i>	Maintain appropriate water quality to support the habitat, including high chlorophyll <i>a</i> status	Lake habitat 3130 is associated with high water quality, and naturally low algal growth. As for nutrients, the targets are WFD 'High Status' or oligotrophic (OECD, 1982). The "good-moderate" boundary is too enriched to support the habitat. The average growing season (March-October) chlorophyll <i>a</i> concentration must be <5.8µg/l. The annual average chlorophyll <i>a</i> concentration should be <2.5µg/l and the annual peak chlorophyll <i>a</i> concentration should be ≤8.0µg/l. Where a lake has a chlorophyll <i>a</i> concentration that is lower than this target, there should be no decline within class, i.e. no upward trend in phytoplankton biomass. See also The European Communities Environmental Objectives (Surface Waters) Regulations 2009
Water quality: phytoplankton composition	EPA phytoplankton composition metric	Maintain appropriate water quality to support the habitat, including high phytoplankton composition status	The EPA has developed a phytoplankton composition metric for nutrient enrichment of Irish lakes. As for other water quality indicators, lake habitat 3130 requires WFD high status
Water quality: attached algal biomass	Algal cover and EPA phytobenthos metric	Maintain trace/absent attached algal biomass (<5% cover) and high phytobenthos status	Nutrient enrichment can favour epiphytic and epipelic algae that can out-compete the submerged vegetation. The cover abundance of attached algae in habitat 3130 should, therefore, be trace/absent (<5% cover). EPA phytobenthos can be used as an indicator of changes in attached algal biomass. As for other water quality indicators, habitat 3130 requires high phytobenthos status
Water quality: macrophyte status	EPA macrophyte metric (The Free Index)	Maintain high macrophyte status	Nutrient enrichment can favour more competitive submerged macrophyte species that out-compete the typical and characteristic species for the lake habitat. The EPA monitors macrophyte status for WFD purposes using the 'Free Index'. The target for lake habitat 3130 is high status or an Ecological Quality Ratio (EQR) for lake macrophytes of ≥0.90, as defined in Schedule Five of the European Communities Environmental Objectives (Surface Waters) Regulations 2009
Acidification status	pH units; mg/l	Maintain appropriate water and sediment pH, alkalinity and cation concentrations to support the habitat, subject to natural processes	Acidification can impact on species abundance and composition in lake habitat 3130. Acidification reduces the abundance and reproductive capacity of <i>Najas flexilis</i> (Wingfield et al., 2004). The specific requirements of habitat 3130, in terms of water and sediment pH, alkalinity and cation concentration, have not been determined. In line with targets for <i>Najas flexilis</i> , median pH values should be greater than 7 pH units. Water and sediment alkalinity and concentrations of cations (notably calcium) should be appropriate to the habitat. The target for WFD Acidification/Alkalisation status is high. Maximum pH should be <9.0 pH units, in line with the surface water standards. See Schedule Five of the European Communities Environmental Objectives (Surface Waters) Regulations 2009

Water colour	mg/l PtCo	Maintain/restore appropriate water colour to support the habitat	Increased water colour and turbidity decrease light penetration and can reduce the area of available habitat for lake macrophytes, particularly at the lower euphotic depths. The primary source of increased water colour in Ireland is disturbance to peatland. No habitat-specific or national standards for water colour currently exist. Studies have shown median colour concentrations in Irish lakes of 38mg/l PtCo (Free et al., 2000) and 33mg/l PtCo (Free et al., 2006). It is likely that the water colour in all Irish lake habitats would naturally be <50mg/l PtCo. Water colour is generally <30mg/l PtCo or, more naturally, <20mg/l PtCo in lakes with habitat 3130, where the peatland in the lake's catchment is intact. Roden (2004, 2007 in NPWS, 2007) stated the water in Lough Nahaltora was coloured/peat-stained. Free et al. (2006) reported colour of 10mg/l PtCo in Fin Lough. Overgrazing, peat-cutting or other peatland damage may have increased colour in some lakes in the SAC
Dissolved organic carbon (DOC)	mg/l	Maintain appropriate organic carbon levels to support the habitat	Dissolved (and particulate) organic carbon (OC) in the water column is linked to water colour and acidification (organic acids). Increasing DOC in water has been documented across the Northern Hemisphere, including afforested peatland catchments in Ireland. Damage and degradation of peatland, leading to decomposition of peat is likely to be the predominant source of OC in Ireland. OC in water promotes decomposition by fungi and bacteria that, in turn, releases dissolved nutrients. The increased biomass of decomposers can also impact directly on the characteristic lake communities through shading, competition, etc. Overgrazing and other peatland degradation is also likely to have increased DOC in some lakes in Mweelrea/Sheeffry/Erriff Complex SAC
Turbidity	Nephelometric turbidity units/ mg/l SS/ other appropriate units	Maintain appropriate turbidity to support the habitat	Turbidity can significantly affect the quantity and quality of light reaching rooted and attached vegetation and can, therefore, impact on lake habitats. The settlement of higher loads of inorganic or organic material on lake vegetation communities may also have impacts on sensitive, delicate species. Turbidity can increase as a result of re-suspension of material within the lake, higher loads entering the lake, or eutrophication. Turbidity measurement and interpretation is challenging. As a result, it is likely to be difficult to set habitat-specific targets for turbidity in lakes. Increased loads of fine organic and inorganic particles from overgrazing may have increased turbidity in lakes in the SAC
Fringing habitat: area	Hectares	Maintain the area and condition of fringing habitats necessary to support the natural structure and functioning of habitat 3130	Most lake shorelines have fringing habitats of reedswamp, other swamp, fen, marsh or wet woodland that intergrade with and support the structure and functions of the lake habitat. In this SAC, lake shorelines are likely to have acid grassland, swamp, heath, blanket bog and rock communities. Calcareous fen and machair occur around coastal lakes. Fringing habitats are dependent on the lake, particularly its water levels, and support wetland communities and species of conservation concern. Many of the fringing wetland habitats support higher invertebrate and plant species richness than the lake habitats themselves

3160 Natural dystrophic lakes and ponds

To maintain the favourable conservation condition of Natural dystrophic lakes and ponds in Mweelrea/Sheeffry/Erriff Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Mweelrea/Sheeffry/Erriff Complex SAC has both blanket bog pool systems and upland lakes with habitat 3160. The habitat is likely to co-occur with habitat 3110 in lakes. All lakes/pools smaller than 1ha and all upland corrie lakes have been mapped as 3160 (see map 6). Many of the bog pools are not mapped in the 1:5,000 OSi data used. Bog pool systems have been recorded at Srahiroosky, Knockeen, Derrinkee, Derrintin, Derryaun, Derrycraff, Srahlea and North of Lough Glenawough (Foss and McGee, 1987; Douglas et al., 1989; Roche et al., 2014). Two measures of extent should be used: 1. the area of the lake itself and; 2. the extent of the vegetation communities/zones that typify the habitat. Further information relating to all attributes is provided in the lake habitats supporting document for the purposes of site-specific conservation objectives and Article 17 reporting (O Connor, 2015)
Habitat distribution	Occurrence	No decline, subject to natural processes	As noted above, all lakes and pools smaller than 1ha in area and all upland lakes have been mapped as potential 3160 (see map 6). Atlantic blanket bog pools, including interconnecting pool systems, were recorded at Srahiroosky, Knockeen, Derrinkee, Derrintin, Derryaun, Derrycraff, Srahlea and North of Lough Glenawough, and other areas of the SAC (Foss and McGee, 1987; Douglas et al., 1989). The habitat is likely to be even more widespread in the SAC as not all pools on blanket bog are mapped in the 1:5,000 OSi data
Typical species	Occurrence	Typical species present, in good condition, and demonstrating typical abundances and distribution	For lists of typical plant and invertebrate species, see the Article 17 habitat assessment for 3160 (NPWS, 2013) and O Connor (2015). See Foss and McGee (1987) and Douglas et al. (1989) for records of plant species in blanket bog pools in the SAC
Vegetation composition: characteristic zonation	Occurrence	All characteristic zones should be present, correctly distributed and in good condition	Further work is necessary to describe the characteristic zonation and other spatial patterns in lake habitat 3160 (see O Connor, 2015). Spatial patterns are likely to be relatively simple in lake habitat 3160, with limited zonation
Vegetation distribution: maximum depth	Metres	Maintain maximum depth of vegetation, subject to natural processes	The maximum depth of vegetation is likely to be specific to the lake shoreline in question. Further work is necessary to develop indicative targets for lake habitat 3160. 3160 lakes and pools naturally have very clear water and, therefore, maximum depth can be large within undisturbed peatland and uplands. Pressures such as overgrazing, forestry and peat-cutting may have reduced vegetation depth in some lakes in the SAC

Hydrological regime: water level fluctuations	Metres	Maintain/restore appropriate natural hydrological regime necessary to support the habitat	Natural water level fluctuations can be amplified by activities such as abstraction, drainage and overgrazing. Increased fluctuations can increase wave action, up-root vegetation, increase turbidity, alter the substratum and lead to release of nutrients from the sediment. The hydrological regime of the lakes and pools must be maintained so that the area, distribution and depth of the lake habitat and its constituent/characteristic vegetation zones and communities are not reduced. The hydrological regime of 3160 lakes and pools is integrally linked to that of the surrounding blanket bog, transition mire/quaking bog and other peatland habitats. Owing to their size and the sensitivity of peatland, 3160 lakes and pools can easily be damaged or destroyed by drainage. It is likely that the hydrological regimes of lakes and pools may still be altered owing to historic overgrazing (faster run-off, higher flood peaks, lower base flows, etc.; see Marshall et al., 2014)
Lake substratum quality	Various	Maintain appropriate substratum type, extent and chemistry to support the vegetation	Research is required to further characterise the substratum types (particle size and origin) and substratum quality (notably pH, calcium, iron and nutrient concentrations) favoured by each of the five Annex I lake habitats in Ireland. It is likely that habitat 3160 is associated with nutrient-poor peat and silt substrates
Water quality: transparency	Metres	Maintain appropriate Secchi transparency. There should be no decline in Secchi depth/transparency	Transparency relates to light penetration and, hence, to the depth of colonisation of vegetation. It can be affected by phytoplankton blooms, water colour and turbidity. A specific target has yet to be established for this Annex I lake habitat. Lake habitat 3160 is associated with very clear water. The OECD fixed boundary system set transparency targets for ultra-oligotrophic lakes of $\geq 12\text{m}$ annual mean Secchi disk depth, and $\geq 6\text{m}$ annual minimum Secchi disk depth
Water quality: nutrients	$\mu\text{g/l P}$; mg/l N	Maintain the concentration of nutrients in the water column at sufficiently low levels to support the habitat and its typical species	As a nutrient-poor habitat, oligotrophic and Water Framework Directive (WFD) 'high' status targets apply. Where a lake has nutrient concentrations that are lower than these targets, there should be no decline within class, i.e. no upward trend in nutrient concentrations. For 3160 lakes and pools, annual average total phosphorus (TP) concentration should be $\leq 5\mu\text{g/l TP}$, average annual total ammonia concentration should be $\leq 0.040\text{mg/l N}$ and annual 95th percentile for total ammonia should be $\leq 0.090\text{mg/l N}$. See also The European Communities Environmental Objectives (Surface Waters) Regulations 2009
Water quality: phytoplankton biomass	$\mu\text{g/l Chlorophyll } a$	Maintain appropriate water quality to support the habitat, including high chlorophyll <i>a</i> status	Oligotrophic and WFD 'high' status targets apply to habitat 3160. The average growing season (March-October) chlorophyll <i>a</i> concentration must be $< 5.8\mu\text{g/l}$ (The European Communities Environmental Objectives (Surface Waters) Regulations 2009). Where a lake has a chlorophyll <i>a</i> concentration that is lower than this target, there should be no decline within class, i.e. no upward trend in phytoplankton biomass. The OECD targets may be more appropriate for habitat 3160: annual average chlorophyll <i>a</i> concentration $< 1\mu\text{g/l}$ and annual peak chlorophyll <i>a</i> concentration $\leq 2.5\mu\text{g/l}$. See also The European Communities Environmental Objectives (Surface Waters) Regulations 2009
Water quality: phytoplankton composition	EPA phytoplankton composition metric	Maintain appropriate water quality to support the habitat, including high phytoplankton composition status	The Environmental Protection Agency (EPA) has developed a phytoplankton composition metric for nutrient enrichment of Irish lakes. As for other water quality indicators, lake habitat 3160 requires WFD high status

Water quality: attached algal biomass	Algal cover and EPA phytobenthos metric	Maintain trace/absent attached algal biomass (<5% cover) and high phytobenthos status	Nutrient enrichment can favour epiphytic and epipelic algae that can out-compete the submerged vegetation. The cover abundance of attached algae in the oligotrophic soft water habitat should, therefore, be trace/absent (<5% cover). EPA phytobenthos can be used as an indicator of changes in attached algal biomass. As for other water quality indicators, lake habitat 3160 requires high phytobenthos status
Water quality: macrophyte status	EPA macrophyte metric (The Free Index)	Maintain high macrophyte status	Nutrient enrichment can favour more competitive submerged macrophyte species that out-compete the typical and characteristic species for the lake habitat. The EPA monitors macrophyte status for WFD purposes using the 'Free Index'. The target for 3160 lakes and pools is high status or an Ecological Quality Ratio (EQR) for lake macrophytes of ≥ 0.90 , as defined in Schedule Five of the European Communities Environmental Objectives (Surface Waters) Regulations 2009
Acidification status	pH units; mg/l	Maintain appropriate water and sediment pH, alkalinity and cation concentrations to support the habitat, subject to natural processes	Acidification can impact on species abundance and composition in soft water lake habitats. Although European Commission (2013) describes lake habitat 3160 as having pH 3-6, Drinan (2012) found mean pH values of 5.16 and 5.62 in upland and lowland 3160 lakes, respectively. The target for habitat 3160 is pH >4.5 and <9.0, in line with the surface water standards for soft waters (where water hardness is ≤ 100 mg/l calcium carbonate). See Schedule Five of the European Communities Environmental Objectives (Surface Waters) Regulations 2009. The specific requirements of lake habitat 3160, in terms of water and sediment pH, alkalinity and cation concentration, have not been determined
Water colour	mg/l PtCo	Maintain/restore appropriate water colour to support the habitat	Increased water colour and turbidity decrease light penetration and can reduce the area of available habitat for lake macrophytes, particularly at the lower euphotic depths. The primary source of increased water colour in Ireland is disturbance to peatland. No habitat-specific or national standards for water colour currently exist. Studies have shown median colour concentrations in Irish lakes of 38mg/l PtCo (Free et al., 2000) and 33mg/l PtCo (Free et al., 2006). It is likely that the water colour in all Irish lake habitats would naturally be <50mg/l PtCo. Water colour can be very low (<20mg/l PtCo or even <10mg/l PtCo) in 3160 lakes and pools where the peatland in the lake's catchment is intact. Overgrazing and other peatland degradation is likely to have increased colour in some lakes and pools in Mweelrea/Sheeffry/Erriff Complex SAC
Dissolved organic carbon (DOC)	mg/l	Maintain/restore appropriate organic carbon levels to support the habitat	Dissolved (and particulate) organic carbon (OC) in the water column is linked to water colour and acidification (organic acids). Increasing DOC in water has been documented across the Northern Hemisphere, including afforested peatland catchments in Ireland. Damage and degradation of peatland, leading to decomposition of peat is likely to be the predominant source of OC in Ireland. OC in water promotes decomposition by fungi and bacteria that, in turn, releases dissolved nutrients. The increased biomass of decomposers can also impact directly on the characteristic lake communities through shading, competition, etc. Overgrazing and other peatland degradation is likely to have increased DOC in some lakes and pools in the SAC

Turbidity	Nephelometric turbidity units/ mg/l SS/ other appropriate units	Maintain/restore appropriate turbidity to support the habitat	Turbidity can significantly affect the quantity and quality of light reaching rooted and attached vegetation and can, therefore, impact on lake habitats. The settlement of higher loads of inorganic or organic material on lake vegetation communities may also have impacts on sensitive, delicate species. Turbidity can increase as a result of re-suspension of material within the lake, higher loads entering the lake, or eutrophication. Turbidity measurement and interpretation is challenging. As a result, it is likely to be difficult to set habitat-specific targets for turbidity in lakes. Increased loads of fine organic and inorganic particles from overgrazing may have increased turbidity in lakes in the SAC
Fringing habitat: area	Hectares	Maintain the area and condition of fringing habitats necessary to support the natural structure and functioning of habitat 3160	3160 pools intergrade with blanket bog communities in Mweelrea/Sheeffry/Erriff Complex SAC. Spring-fed flushes are also a feature of the SAC, as is quaking bog. Lakes with 3160, particularly in uplands, are likely to be fringed by acid grassland, heath and rock communities. Fringing habitats support the structure and functions of the lake/pool habitat. Equally, fringing habitats are dependent on the lake/pool, particularly its water levels, and can support wetland communities and species of conservation concern

3260 Water courses of plain to montane levels with the Ranunculion fluitantis and Callitricho-Batrachion vegetation

To maintain the favourable conservation condition of Water courses of plain to montane levels with the Ranunculion fluitantis and Callitricho-Batrachion vegetation in Mweelrea/Sheeffry/Erriff Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Kilometres	Area stable or increasing, subject to natural processes	Conservation objectives concentrate on the high conservation value sub-types of the habitat. Selection of the SAC used a broad interpretation and the habitat's full distribution and sub-types are not yet documented. Rivers in the SAC are naturally highly oligotrophic and species-poor; rare and threatened river bryophytes and lichens indicate high conservation value. Upland streams in particular deserve further study. Note: rooted macrophytes should be trace/absent (<5% cover) in freshwater pearl mussel (<i>Margaritifera margaritifera</i>) habitat. The SAC contains the priority freshwater pearl mussel Bundorragha catchment, one of the world's most important pearl mussel rivers. The freshwater pearl mussel (1029) conservation objective takes precedence over the 3260 objective in the Bundorragha in this SAC because the mussel requires environmental conditions closer to natural background levels
Habitat distribution	Occurrence	No decline, subject to natural processes	Further study of Irish rivers is needed to interpret the broad description of habitat 3260 which covers upland bryophyte/macroalgal dominated to lowland depositing rivers with pondweeds and starworts (European Commission, 2013). The rivers in the SAC are naturally very nutrient-poor. The Bundorragha and upland rivers/streams are of particular interest. The upland rivers/streams require further investigation. Heuff (1987) surveyed the Erriff. Some information on the Bundorragha is available from NPWS (2010) and Williams (2009) (see also the conservation objective for the freshwater pearl mussel (<i>Margaritifera margaritifera</i>)). Lockhart et al. (2012) provide information on bryophytes. The SAC contains a number of high status river sites (see McGarrigle et al., 2010; Ní Chatháin et al., 2013; Bradley et al., 2015)
Hydrological regime: river flow	Metres per second	Maintain/restore appropriate hydrological regimes	High conservation value sub-types are associated with natural hydrological regimes. A natural flow regime is required for both plant communities and channel geomorphology to be in favourable condition, exhibiting typical dynamics for the river type (Hatton-Ellis and Grieve, 2003). For many sub-types of the habitat, high flows are required to maintain the substratum necessary for the characteristic species. Flow variation can be particularly important, with high and flood flows being critical to the hydromorphology. Other aspects of hydrology, such as tidal regime, are important for certain sub-types of the habitat. The hydrological regimes of some of the SAC's rivers, particularly in the Erriff system, may have been altered by overgrazing and peat erosion (faster run-off, higher flood peaks, lower base flows, etc.; see Marshall et al. (2014))
Hydrological regime: groundwater discharge	Metres per second	Maintain appropriate hydrological regime	Even small groundwater contributions can significantly alter hydrochemistry, particularly where there is basic bedrock and/or subsoils. Freshwater seepages can be very important in tidal reaches

Substratum composition: particle size range	Millimetres	Maintain/restore appropriate substratum particle size range, quantity and quality, subject to natural processes	Although many of the high conservation value sub-types of the habitat are dominated by coarse substrata and bedrock, certain sub-types, notably those associated with lake inflows/outflows and peatlands, are dominated by fine substrata. The size and distribution of particles is largely determined by the river flow and geology. The chemical composition (particularly minerals and nutrients) of the substratum is also important. The quality of finer sediment particles is a notable driver for rooted plant communities. Increased loads of fine organic and inorganic particles arising from drainage, overgrazing and other peatland degradation is likely to have increased the fine sediment load to rivers in the SAC, particularly the Erriff
Water quality	Various	Maintain/restore appropriate water quality to support the natural structure and functioning of the habitat	The specific targets may vary among sub-types. The rivers within the SAC are naturally very nutrient-poor and require Water Framework Directive high status, in terms of nutrient and oxygenation standards, and EQRs (Ecological Quality Ratios) for macroinvertebrates and phytobenthos. High status targets apply to freshwater pearl mussel (<i>Margaritifera margaritifera</i>) habitat in the Bundorragha (see The European Communities Environmental Objectives (Freshwater Pearl Mussel) Regulations 2009). Overgrazing, peat-cutting or other peatland damage may have increased ammonia, colour, fine sediment, DOC (dissolved organic carbon) and other pollutant loads to rivers such as the Erriff in the SAC. See also The European Communities Environmental Objectives (Surface Waters) Regulations 2009, Environmental Protection Agency (EPA) river water quality reports (e.g. Bradley et al., 2015) and Ní Chatháin et al. (2013)
Typical species	Occurrence	Typical species of the relevant habitat sub-type should be present and in good condition	Typical species have not yet been fully defined. Heuff (1987) recorded typical oligotrophic, base-poor species in the Erriff including <i>Fissidens viridulus</i> , <i>Fontinalis squamosa</i> , <i>Juncus bulbosus</i> , <i>Myriophyllum alterniflorum</i> and <i>Potamogeton natans</i> . <i>Callitriche hermaphroditica</i> , <i>C. stagnalis</i> , <i>J. bulbosus</i> , <i>M. alterniflorum</i> and <i>P. polygonifolius</i> were recorded in the Bundorragha in 2014 (M. Wyse Jackson, pers. comm.). Red-listed bryophytes in/near rivers include <i>Hygrohypnum duriusculum</i> (CR) (fast-flowing upland stream, Mweelrea); <i>Bryum riparium</i> (EN) and <i>Fissidens serrulatus</i> (VU) (Lockhart et al., 2012). Scarce lichens recorded in the Erriff include <i>Dermatocarpon meiophyllizum</i> , <i>Catillaria chalybeia</i> var. <i>chloropoliza</i> , <i>Pertusaria excludens</i> and <i>Porpidia hydrophila</i> (NPWS internal files). Giavarini (2009) recorded <i>Dermatocarpon luridum</i> , <i>Lecidea lithophila</i> and <i>Porpidia hydrophila</i> in the Bundorragha
Floodplain connectivity: area	Hectares	The area of active floodplain at and upstream of the habitat should be maintained	River connectivity with the floodplain is important for the functioning of this habitat. Channels with a naturally functioning floodplain are better able to maintain habitat and water quality (Hatton-Ellis and Grieve, 2003). Floodplain connectivity is particularly important in terms of sediment sorting and nutrient deposition. High conservation value rivers are intimately connected to floodplain habitats and function as important wildlife corridors, connecting otherwise isolated or fragmented habitats in the wider countryside (Hatton-Ellis and Grieve, 2003; Mainstone et al., 2016)

Riparian habitat: Hectares
area and condition

Maintain the area and
condition of fringing
habitats necessary to
support the habitat and its
sub-types

Riparian habitats, including those along lakes, e.g. woodlands and wetlands, are integral to the structure and functioning of rivers, even where not part of a floodplain. Fringing habitats can contribute to the aquatic food web (e.g. allochthonous matter such as leaf fall), provide habitat for certain life-stages of fish, birds and aquatic invertebrates, assist in the settlement of fine sediment, protect banks from erosion and contribute to nutrient cycling. Shade may be important in suppressing algal growth and moderating temperatures. Equally, fringing habitats are dependent on rivers/lakes, particularly their water levels, and support wetland communities and species of conservation concern. See Mainstone et al. (2016). Semi-natural grassland and peatland are a feature of many rivers in the SAC. *Erica erigena* occurs along streams to the N. of Killary. *Carex aquatilis* is found along Erriff. The upland streams are associated with flushes and other important habitats

4010 Northern Atlantic wet heaths with *Erica tetralix*

To restore the favourable conservation condition of Northern Atlantic wet heaths with *Erica tetralix* in Mweelrea/Sheeffry/Erriff Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Mweelrea/Sheeffry/Erriff Complex SAC was surveyed as part of the National Survey of Upland Habitats (NSUH; see Perrin et al., 2014 and Roche et al., 2014). Northern Atlantic wet heaths with <i>Erica tetralix</i> was mapped in detail for the SAC and the total area of the qualifying habitat stated by Roche et al. (2014) is 4,861.9ha, covering 23.2% of the SAC. Roche et al. (2014) report obvious losses of the habitat since 1995 of approximately 1.46ha. Further information can be found in Roche et al. (2014). Further details on this and the following attributes can be found in the Mweelrea/Sheeffry/Erriff Complex SAC conservation objectives supporting document for blanket bogs and associated habitats where a summary of the mapping methodology and a brief discussion of restoration potential are also presented
Habitat distribution	Occurrence	No decline, subject to natural processes. See map 7	Roche et al. (2014) recorded wet heaths throughout the SAC, including to the north-west of the main ridge of the Sheeffry Hills, the northern hill of Tangincartoor and the neighbouring ridge of Laghta Eighter, the slopes of Ben Gorm and Ben Creggan, the northern lip of the Glenawough valley, the lower flanks of the Mweelrea Mountains and the lower lying lands of the Erriff valley and around Cregganbaun. Further information can be found within Roche et al. (2014) and the blanket bogs and associated habitats supporting document
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil nutrient status within natural range	See the blanket bogs and associated habitats supporting document for further details
Community diversity	Abundance of variety of vegetation communities	Maintain variety of vegetation communities, subject to natural processes	Roche et al. (2014) recorded six different wet heath communities within this SAC. Data on the abundance of these communities is reproduced in the blanket bogs and associated habitats supporting document. Further information on the vegetation communities associated with this habitat is presented in Perrin et al. (2014)
Vegetation composition: cross-leaved heath	Occurrence within 20m of a representative number of monitoring stops	Cross-leaved heath (<i>Erica tetralix</i>) present within a 20m radius of each monitoring stop	Attribute and target based on Perrin et al. (2014)
Vegetation composition: positive indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of positive indicator species at least 50%	Attribute and target based on Perrin et al. (2014), where the list of positive indicator species for this habitat is also presented
Vegetation composition: lichens and bryophytes	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of <i>Cladonia</i> and <i>Sphagnum</i> species, <i>Racomitrium lanuginosum</i> and pleurocarpous mosses at least 10%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: ericoid species and crowberry	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of ericoid species and crowberry (<i>Empetrum nigrum</i>) at least 15%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: dwarf shrub species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of dwarf shrubs less than 75%	Attribute and target based on Perrin et al. (2014)

Vegetation composition: negative indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of negative indicator species less than 1%	Attribute and target based on Perrin et al. (2014), where the list of negative indicator species for this habitat is also presented
Vegetation composition: non-native species	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of non-native species less than 1%	Attribute and target based on Perrin et al. (2014). During the NSUH, the non-native moss <i>Campylopus introflexus</i> was recorded in 8.6% of the wet heath monitoring stops, but its cover was not sufficient to cause the stops to fail. Rhododendron (<i>Rhododendron ponticum</i>) has a localised distribution within habitat 4010 in the SAC. Further information can be found in Roche et al. (2014)
Vegetation composition: native trees and shrubs	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of scattered native trees and shrubs less than 20%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: bracken	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of bracken (<i>Pteridium aquilinum</i>) less than 10%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: soft rush	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of soft rush (<i>Juncus effusus</i>) less than 10%	Attribute and target based on Perrin et al. (2014)
Vegetation structure: <i>Sphagnum</i> condition	Condition at a representative number of 2m x 2m monitoring stops	Less than 10% of the <i>Sphagnum</i> cover is crushed, broken and/or pulled up	Attribute and target based on Perrin et al. (2014)
Vegetation structure: signs of browsing	Percentage of shoots browsed at a representative number of 2m x 2m monitoring stops	Less than 33% collectively of the last complete growing season's shoots of ericoids, crowberry (<i>Empetrum nigrum</i>) and bog-myrtle (<i>Myrica gale</i>) showing signs of browsing	Attribute and target based on Perrin et al. (2014)
Vegetation structure: burning	Occurrence in local vicinity of a representative number of monitoring stops	No signs of burning in sensitive areas, into the moss, liverwort or lichen layer or exposure of peat surface due to burning	Attribute and target based on Perrin et al. (2014), where the list of sensitive areas for this habitat is also presented
Physical structure: disturbed bare ground	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of disturbed bare ground less than 10%	Attribute and target based on Perrin et al. (2014)
Physical structure: drainage	Percentage area in local vicinity of a representative number of monitoring stops	Area showing signs of drainage from heavy trampling, tracking or ditches less than 10%	Attribute and target based on Perrin et al. (2014)
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat and no decline in status of hepatic mats associated with this habitat	This includes species listed in the Flora (Protection) Order, 2015 (FPO) and/or the red data lists (Lockhart et al., 2012; Wyse Jackson et al., 2016). Roche et al. (2014) compiled and mapped existing rare and notable plant records for the SAC and added any new records collected during the NSUH survey. The Near Threatened and FPO listed liverwort <i>Mastigophora woodsii</i> and the Near Threatened moss <i>Sphagnum subsecundum</i> (Lockhart et al., 2012) can be attributed specifically to wet heath in the SAC (Roche et al., 2014)

4030 European dry heaths

To restore the favourable conservation condition of European dry heaths in Mweelrea/Sheeffry/Erriff Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Mweelrea/Sheeffry/Erriff Complex SAC was surveyed as part of the National Survey of Upland Habitats (NSUH; see Perrin et al., 2014 and Roche et al., 2014). European dry heaths was mapped in detail for the SAC and the total area of the qualifying habitat stated by Roche et al. (2014) is 359.8ha, covering 1.7% of the SAC. Roche et al. (2014) report obvious losses of the habitat since 1995 of approximately 0.08ha. Further information can be found in Roche et al. (2014). Further details on this and the following attributes can be found in the Mweelrea/Sheeffry/Erriff Complex SAC conservation objectives supporting document for blanket bogs and associated habitats where a summary of the mapping methodology and a brief discussion of restoration potential are also presented
Habitat distribution	Occurrence	No decline, subject to natural processes. See map 8	Roche et al. (2014) recorded dry heaths throughout the SAC, with the main areas being the drier tops of Tangincartoor and the neighbouring ridge at Laghta Eighter. It also occurs on the back wall of the Glenawough corrie and on the slopes of the large broad valley of Laghta Ougher, north of the main Sheeffry Hills ridge. Further information can be found within Roche et al. (2014) and the blanket bogs and associated habitats supporting document
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil nutrient status within natural range	See the blanket bogs and associated habitats supporting document for further details
Community diversity	Abundance of variety of vegetation communities	Maintain variety of vegetation communities, subject to natural processes	Roche et al. (2014) recorded six different dry heath communities within this SAC. Data on the abundance of these communities is reproduced in the blanket bogs and associated habitats supporting document. Further information on the vegetation communities associated with this habitat is presented in Perrin et al. (2014)
Vegetation composition: lichens and bryophytes	Number of species at a representative number of 2m x 2m monitoring stops	Number of bryophyte or non-crustose lichen species present at each monitoring stop is at least three, excluding <i>Campylopus</i> and <i>Polytrichum</i> mosses	Attribute and target based on Perrin et al. (2014)
Vegetation composition: number of positive indicator species	Number of species at a representative number of 2m x 2m monitoring stops	Number of positive indicator species present at each monitoring stop is at least two	Attribute and target based on Perrin et al. (2014), where the list of positive indicator species for this habitat, which is composed of dwarf shrubs, is also presented
Vegetation composition: cover of positive indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of positive indicator species at least 50% for siliceous dry heath and 50-75% for calcareous dry heath	Attribute and target based on Perrin et al. (2014), where the list of positive indicator species for this habitat, which is composed of dwarf shrubs, is also presented
Vegetation composition: dwarf shrub composition	Percentage cover at a representative number of 2m x 2m monitoring stops	Proportion of dwarf shrub cover composed collectively of bog-myrtle (<i>Myrica gale</i>), creeping willow (<i>Salix repens</i>) and western gorse (<i>Ulex gallii</i>) is less than 50%	Attribute and target based on Perrin et al. (2014)

Vegetation composition: negative indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of negative indicator species less than 1%	Attribute and target based on Perrin et al. (2014), where the list of negative indicator species for this habitat is also presented
Vegetation composition: non-native species	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of non-native species less than 1%	Attribute and target based on Perrin et al. (2014). During the NSUH, the non-native moss <i>Campylopus introflexus</i> was recorded within one dry heath monitoring stop, but its cover was not sufficient to cause the stop to fail. Further information can be found in Roche et al. (2014)
Vegetation composition: native trees and shrubs	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of scattered native trees and shrubs less than 20%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: bracken	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of bracken (<i>Pteridium aquilinum</i>) less than 10%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: soft rush	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of soft rush (<i>Juncus effusus</i>) less than 10%	Attribute and target based on Perrin et al. (2014)
Vegetation structure: senescent ling	Percentage cover at a representative number of 2m x 2m monitoring stops	Senescent proportion of ling (<i>Calluna vulgaris</i>) cover less than 50%	Attribute and target based on Perrin et al. (2014)
Vegetation structure: signs of browsing	Percentage of shoots browsed at a representative number of 2m x 2m monitoring stops	Less than 33% collectively of the last complete growing season's shoots of ericoids and crowberry (<i>Empetrum nigrum</i>) showing signs of browsing	Attribute and target based on Perrin et al. (2014)
Vegetation structure: burning	Occurrence in local vicinity of a representative number of monitoring stops	No signs of burning in sensitive areas	Attribute and target based on Perrin et al. (2014), where the list of sensitive areas for this habitat is also presented
Vegetation structure: growth phases of ling	Percentage cover in local vicinity of a representative number of monitoring stops	Outside sensitive areas, all growth phases of ling (<i>Calluna vulgaris</i>) should occur throughout, with at least 10% of cover in the mature phase	Attribute and target based on Perrin et al. (2014), where the list of sensitive areas for this habitat is also presented
Physical structure: disturbed bare ground	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of disturbed bare ground less than 10%	Attribute and target based on Perrin et al. (2014)
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat and no decline in status of hepatic mats associated with this habitat	This includes species listed in the Flora (Protection Order, 2015 (FPO) and/or the red data lists (Lockhart et al., 2012; Wyse Jackson et al., 2016). Roche et al. (2014) compiled and mapped existing rare and notable plant records for the SAC and added any new records collected during the NSUH survey. During the NSUH, the Near Threatened and FPO listed liverwort species <i>Mastigophora woodsii</i> and the Vulnerable and FPO liverworts <i>Bazzania pearsonii</i> and <i>Scapania ornithopodioides</i> (Lockhart et al., 2012) were recorded within a hepatic mat on dry heath. There is a historic record for the Vulnerable and FPO listed species small-white orchid (<i>Pseudorchis albida</i>) (Wyse Jackson et al., 2016) from the SAC, but this species cannot be attributed specifically to dry heaths (Roche et al., 2014)

4060 Alpine and Boreal heaths

To restore the favourable conservation condition of Alpine and Boreal heaths in Mweelrea/Sheeffry/Erriff Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Mweelrea/Sheeffry/Erriff Complex SAC was surveyed as part of the National Survey of Upland Habitats (NSUH; see Perrin et al., 2014 and Roche et al., 2014). Alpine and Boreal heaths was mapped in detail for the SAC and the total area of the qualifying habitat stated by Roche et al. (2014) is 148.7ha, covering 0.7% of the SAC. Roche et al. (2014) report no obvious losses of this habitat since 1995. Further information can be found in Roche et al. (2014). Further details on this and the following attributes can be found in the Mweelrea/Sheeffry/Erriff Complex SAC conservation objectives supporting document for blanket bogs and associated habitats where a summary of the mapping methodology is also presented
Habitat distribution	Occurrence	No decline, subject to natural processes. See map 9	Roche et al. (2014) recorded Alpine and Boreal heaths on high ground across the SAC, including the summit plateau of the Sheeffry Hills, the summits of Tangincartoor and the neighbouring ridge of Laghta Eighter, above the Glenawough corrie, the northern side of the Mweelrea Mountains and close to the summits of Ben Creggan and Ben Gorm. Further information can be found within Roche et al. (2014) and the blanket bogs and associated habitats supporting document
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil nutrient status within natural range	See the blanket bogs and associated habitats supporting document for further details
Community diversity	Abundance of variety of vegetation communities	Maintain variety of vegetation communities, subject to natural processes	Roche et al. (2014) recorded four different Alpine and Boreal heath vegetation communities within this SAC. Data on the abundance of these communities is reproduced in the blanket bogs and associated habitats supporting document. Further information on the vegetation communities associated with this habitat is presented in Perrin et al. (2014)
Vegetation composition: lichens and bryophytes	Number of species at a representative number of 2m x 2m monitoring stops	Number of bryophyte or non-crustose lichen species present at each monitoring stop is at least three	Attribute and target based on Perrin et al. (2014)
Vegetation composition: positive indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of positive indicator species at least 66%	Attribute and target based on Perrin et al. (2014), where the list of positive indicator species for this habitat is also presented
Vegetation composition: dwarf shrub species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of dwarf shrub species at least 10%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: negative indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of negative indicator species less than 10%	Attribute and target based on Perrin et al. (2014), where the list of negative indicator species for this habitat is also presented
Vegetation composition: non-native species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of non-native species less than 1%	Attribute and target based on Perrin et al. (2014). During the NSUH, the non-native moss <i>Campylopus introflexus</i> was recorded within one Alpine and Boreal heath monitoring stop, but its cover was not sufficient to cause the stop to fail. Further information can be found in Roche et al. (2014)

Vegetation structure: signs of grazing	Percentage of leaves grazed at a representative number of 2m x 2m monitoring stops	Less than 10% collectively of the live leaves of specific graminoids showing signs of grazing	Attribute and target based on Perrin et al. (2014). See the blanket bogs and associated habitats supporting document for the list of specific graminoids
Vegetation structure: signs of browsing	Percentage of shoots browsed at a representative number of 2m x 2m monitoring stops	Less than 33% collectively of the last complete growing season's shoots of ericoids and crowberry (<i>Empetrum nigrum</i>) showing signs of browsing	Attribute and target based on Perrin et al. (2014)
Vegetation structure: burning	Occurrence in local vicinity of a representative number of monitoring stops	No signs of burning within the habitat	Attribute and target based on Perrin et al. (2014)
Physical structure: disturbed bare ground	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of disturbed bare ground less than 10%	Attribute and target based on Perrin et al. (2014)
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat and no decline in status of hepatic mats associated with this habitat	This includes species listed in the Flora (Protection) Order, 2015 and/or the red data lists (Lockhart et al., 2012; Wyse Jackson et al., 2016). Roche et al. (2014) compiled and mapped existing rare and notable plant records for the SAC and added any new records collected during the NSUH survey. No relevant species were recorded in this habitat; however, new records should be considered within this attribute

5130 Juniperus communis formations on heaths or calcareous grasslands

To maintain the favourable conservation condition of *Juniperus communis* formations on heaths or calcareous grasslands in Mweelrea/Sheeffry/Erriff Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	<i>Juniperus communis</i> formations on heath or calcareous grasslands has not been mapped in detail in Mweelrea/Sheeffry/Erriff Complex SAC and thus the total area of the qualifying habitat in the SAC is unknown. The habitat was not recorded during the National Survey of Upland Habitats (Roche et al., 2014), although <i>Juniperus communis</i> was recorded as a component of habitat 4060. Douglas et al. (1989) also recorded the species in the middle of Knockeen Bog in the north-west of the SAC, where it occurs both on small islands in pools and in flat areas of blanket bog (7130*) between the pools. The habitat may occur in Creganawoody townland in the north-west of the SAC, where juniper has been recorded on shallow peat, where there is much outcropping rock (NPWS internal files)
Habitat distribution	Occurrence	No decline, subject to natural processes	See the notes for habitat area above
Juniper population size	Number per formation	At least 50 plants per formation	To classify as a juniper (<i>Juniperus communis</i>) formation, at least 50 plants should be present (Cooper et al., 2012)
Vegetation composition: typical species	Number per formation	At least 50% of the listed positive indicator species for the relevant vegetation group present	Attribute and target based on Cooper et al. (2012), where positive indicator species for five vegetation groups are listed
Vegetation composition: negative indicator species	Occurrence per formation	Negative indicator species, particularly non-native invasive species, absent or under control	Attribute and target based on Cooper et al. (2012), where the list of negative indicator species is presented
Vegetation structure: cone-bearing plants	Percentage per formation	At least 10% of juniper plants are bearing cones	Attribute and target based on Cooper et al. (2012)
Vegetation structure: seedling recruitment	Percentage per formation	At least 10% of juniper plants are seedlings	Attribute and target based on Cooper et al. (2012)
Vegetation structure: dead juniper	Percentage per formation	Mean percentage of each juniper plant dead less than 10%	Attribute and target based on Cooper et al. (2012)

Conservation Objectives for : Mweelrea/Sheeffry/Erriff Complex SAC [001932]

6430 Hydrophilous tall herb fringe communities of plains and of the montane to alpine levels

To maintain the favourable conservation condition of Hydrophilous tall herb fringe communities of plains and of the montane to alpine levels in Mweelrea/Sheeffry/Erriff Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Mweelrea/Sheeffry/Erriff Complex SAC was surveyed as part of the National Survey of Upland Habitats (NSUH; see Perrin et al., 2014 and Roche et al., 2014). Hydrophilous tall herb fringe communities of plains and of the montane to alpine levels habitat was recorded in an upland context and was mapped in detail for the SAC by Roche et al. (2014). The total area of the qualifying habitat stated by Roche et al. (2014) is 1.5ha, covering 0.01% of the SAC. Further information can be found in Roche et al. (2014) and Perrin et al. (2014)
Habitat distribution	Occurrence	No decline, subject to natural processes. See map 10	The upland ledge type of the habitat was recorded from several corrie walls in the SAC and particularly among the Sheeffry Hills. See Roche et al. (2014) for further details
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil nutrient status within natural range	Relevant nutrients and their natural ranges are yet to be defined. However, nitrogen deposition is noted as being relevant to this habitat (NPWS, 2013)
Community diversity	Abundance of variety of vegetation communities	Maintain variety of vegetation communities, subject to natural processes	Roche et al. (2014) recorded one tall herb community of rocky upland ledges within this SAC. Further information on the vegetation communities associated with this habitat can be found in Perrin et al. (2014)
Vegetation composition: number of positive indicator species	Number of species at a representative number of monitoring stops	At least one positive indicator species at each monitoring stop	The list of positive indicator species for the upland type community can be found in the Article 17 habitat assessment for 6430 (NPWS, 2013). See Roche et al. (2014) and Perrin et al. (2014) for further details
Vegetation composition: cover of positive indicator species	Percentage cover at a representative number of monitoring stops	Cover of positive indicator species is at least 25%	The list of positive indicator species for the upland type community can be found in the Article 17 habitat assessment for 6430 (NPWS, 2013). See Roche et al. (2014) and Perrin et al. (2014) for further details
Vegetation composition: non-native species	Percentage at a representative number of monitoring stops	Cover of non-native species less than 1%	See NPWS (2013), Roche et al. (2014) and Perrin et al. (2014) for further details
Vegetation structure: height/flowering	Percentage/occurrence at a representative number of monitoring stops	At least 50% of tall herb stems should be greater than 20cm tall or signs of flowering/ability to flower should be present	See NPWS (2013), Roche et al. (2014) and Perrin et al. (2014) for further details
Vegetation structure: grazing	Percentage of flowering tall herb shoots grazed at a representative number of monitoring stops	Live shoots of flowering tall herb shoots showing signs of grazing less than 50%	See NPWS (2013), Roche et al. (2014) and Perrin et al. (2014) for further details
Physical structure: disturbed bare ground	Percentage cover at, and in local vicinity of, a representative number of monitoring stops	Cover of disturbed bare ground in monitoring stop less than 25% and less than 10% in local vicinity of monitoring stop	See NPWS (2013), Roche et al. (2014) and Perrin et al. (2014) for further details

Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat and no decline in status of hepatic mats associated with this habitat	This includes species listed in the Flora (Protection) Order, 2015 and/or the red data lists (Lockhart et al., 2012; Wyse Jackson et al., 2016). Roche et al. (2014) compiled and mapped existing rare and notable plant records for the SAC and added any new records collected during the NSUH survey. No relevant species were recorded in this habitat; however, new records should be considered within this attribute
-------------------------------------	--------------------------------	--	---

7130 Blanket bogs (* if active bog)

To restore the favourable conservation condition of Blanket bogs (* if active bog) in Mweelrea/Sheeffry/Erriff Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Mweelrea/Sheeffry/Erriff Complex SAC was surveyed as part of the National Survey of Upland Habitats (NSUH; see Perrin et al., 2014 and Roche et al., 2014). Blanket bog was mapped in detail and the total area of the qualifying habitat stated by Roche et al. (2014) is 4523ha, covering 21.6% of the SAC. This comprises 4287.7ha of active blanket bog and 236.3ha of inactive blanket bog. Roche et al. (2014) report obvious losses of approximately 2.2ha since 1995. However, this is almost certainly an underestimate as chronic losses due to erosion since 1995 cannot be quantified (495.9ha were mapped as eroding blanket bog by Roche et al. (2014)). It should be noted that further restoration of blanket bog would be required in order to fulfil the targets for peat formation and hydrology presented below. Further details on this and the following attributes can be found in the Mweelrea/Sheeffry/Erriff Complex SAC conservation objectives supporting document for blanket bogs and associated habitats
Habitat distribution	Occurrence	No decline, subject to natural processes. See map 11	Roche et al. (2014) recorded blanket bog throughout the SAC including the large broad valley of Laghta Ougher, the lower ground surrounding Laghta Eighter and Tangincartoor, lower areas around the massif in the vicinity of Ben Gorm and Ben Creggan, within and above the Glenawough valley and in the lowlands of the Erriff valley and around Cregganbaun. Further information can be found within Roche et al. (2014) and the blanket bogs and associated habitats supporting document where a summary of the mapping methodology is presented
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil nutrient status within natural range	See the blanket bogs and associated habitats supporting document for further details
Ecosystem function: peat formation	Active blanket bog as a proportion of the total area of Annex I blanket bog habitat	At least 99% of the total Annex I blanket bog area is active	From the habitat areas given by Roche et al. (2014) above, 94.8% of the Annex I blanket bog habitat is currently actively peat-forming
Ecosystem function: hydrology	Flow direction, water levels, occurrence of drains and erosion gullies	Natural hydrology unaffected by drains and erosion	Further details and a brief discussion of restoration potential is presented in the blanket bogs and associated habitats supporting document
Community diversity	Abundance of variety of vegetation communities	Maintain variety of vegetation communities, subject to natural processes	Roche et al. (2014) recorded seven active blanket bog communities within this SAC. Data on the abundance of the active blanket bog communities is reproduced in the blanket bogs and associated habitats supporting document. Further information on the vegetation communities associated with this habitat is presented in Perrin et al. (2014)
Vegetation composition: positive indicator species	Number of species at a representative number of 2m x 2m monitoring stops	Number of positive indicator species present at each monitoring stop is at least seven	Attribute and target based on Perrin et al. (2014), where the list of positive indicator species for this habitat is also presented
Vegetation composition: lichens and bryophytes	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of bryophytes or lichens, excluding <i>Sphagnum fallax</i> , at least 10%	Attribute and target based on Perrin et al. (2014)

Vegetation composition: potential dominant species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of each of the potential dominant species less than 75%	Attribute and target based on Perrin et al. (2014). See the blanket bogs and associated habitats supporting document for the list of potential dominant species
Vegetation composition: negative indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of negative indicator species less than 1%	Attribute and target based on Perrin et al. (2014), where the list of negative indicator species for this habitat is also presented
Vegetation composition: non-native species	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of non-native species less than 1%	Attribute and target based on Perrin et al. (2014). During the NSUH, the non-native moss <i>Campylopus introflexus</i> formed extensive carpets within this habitat. Rhododendron (<i>Rhododendron ponticum</i>) was also found within the habitat in the SAC. Further information can be found in Roche et al. (2014)
Vegetation composition: native trees and shrubs	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of scattered native trees and shrubs less than 10%	Attribute and target based on Perrin et al. (2014)
Vegetation structure: <i>Sphagnum</i> condition	Condition at a representative number of 2m x 2m monitoring stops	Less than 10% of the <i>Sphagnum</i> cover is crushed, broken and/or pulled up	Attribute and target based on Perrin et al. (2014)
Vegetation structure: signs of browsing	Percentage of shoots browsed at a representative number of 2m x 2m monitoring stops	Last complete growing season's shoots of ericoids, crowberry (<i>Empetrum nigrum</i>) and bog-myrtle (<i>Myrica gale</i>) showing signs of browsing collectively less than 33%	Attribute and target based on Perrin et al. (2014)
Vegetation structure: burning	Occurrence in local vicinity of a representative number of monitoring stops	No signs of burning in sensitive areas, into the moss, liverwort or lichen layer or exposure of peat surface due to burning	Attribute and target based on Perrin et al. (2014), where the list of sensitive areas for this habitat is also presented
Physical structure: disturbed bare ground	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of disturbed bare ground less than 10%	Attribute and target based on Perrin et al. (2014)
Physical structure: drainage	Percentage area in local vicinity of a representative number of monitoring stops	Area showing signs of drainage from heavy trampling, tracking or ditches less than 10%	Attribute and target based on Perrin et al. (2014)
Physical structure: erosion	Percentage area in local vicinity of a representative number of monitoring stops	Less than 5% of the greater bog mosaic comprises erosion gullies and eroded areas	Attribute and target based on Perrin et al. (2014)
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat	This includes species listed in the Flora (Protection) Order, 2015 and/or the red data lists (Lockhart et al., 2012; Wyse Jackson et al., 2016). Roche et al. (2014) compiled and mapped existing rare and notable plant records for the SAC and added any new records collected during the NSUH survey. No relevant species were recorded in this habitat; however, new records should be considered within this attribute

7140 Transition mires and quaking bogs

To restore the favourable conservation condition of Transition mires and quaking bogs in Mweelrea/Sheeffry/Erriff Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Mweelrea/Sheeffry/Erriff Complex SAC was surveyed as part of the National Survey of Upland Habitats (NSUH; see Perrin et al., 2014 and Roche et al., 2014). Transition mires and quaking bogs was mapped in detail for the SAC and the total area of the qualifying habitat stated by Roche et al. (2014) is 96.2ha, covering 0.5% of the SAC. Roche et al. (2014) report no obvious losses of this habitat since 1995. Further information can be found in Roche et al. (2014). Further details on this and the following attributes can be found in the Mweelrea/Sheeffry/Erriff Complex SAC conservation objectives supporting document for blanket bogs and associated habitats where a summary of the mapping methodology is also presented
Habitat distribution	Occurrence	No decline, subject to natural processes. See map 12	Roche et al. (2014) recorded transition mires throughout the SAC, including the lower ground surrounding Laghta Eighter and Tangincartoor, Cregganbaun, Srahrooskey, Drummin West, Derrycraff and Derrintin. Further information can be found within Roche et al. (2014) and the blanket bogs and associated habitats supporting document
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil nutrient status within natural range	See the blanket bogs and associated habitats supporting document for further details
Community diversity	Abundance of variety of vegetation communities	Maintain variety of vegetation communities, subject to natural processes	Roche et al. (2014) recorded three different transition mire and quaking bog vegetation communities within this SAC. Data on the abundance of these communities is reproduced in the blanket bogs and associated habitats supporting document. Further information on vegetation communities associated with this habitat is presented in Perrin et al. (2014)
Vegetation composition: number of positive indicator species	Number of species at a representative number of 2m x 2m monitoring stops	Number of positive indicator species at each monitoring stop is at least three for infilling pools and flushes and at least six for fens	Attribute and target based on Perrin et al. (2014), where the list of positive indicator species for this habitat is also presented
Vegetation composition: number of core positive indicator species	Number of species at a representative number of 2m x 2m monitoring stops	At least one core positive indicator species present	Attribute and target based on Perrin et al. (2014), where the list of positive indicator species for this habitat is also presented
Vegetation composition: cover of positive indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of positive indicator species is at least 25%	Attribute and target based on Perrin et al. (2014), where the list of positive indicator species for this habitat is also presented
Vegetation composition: negative indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of negative indicator species less than 1%	Attribute and target based on Perrin et al. (2014), where the list of negative indicator species for this habitat is also presented
Vegetation composition: non-native species	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of non-native species less than 1%	Attribute and target based on Perrin et al. (2014). No non-native species were recorded within this habitat during the NSUH (Roche et al., 2014)

Vegetation structure: height	Percentage of leaves/shoots at a representative number of 2m x 2m monitoring stops	Proportion of live leaves and/or flowering shoots of vascular plants that are more than 15cm above the ground surface should be at least 50%	Attribute and target based on Perrin et al. (2014). This attribute is only applicable to fen and flush examples of the habitat, not to infilling pool examples
Physical structure: disturbed bare ground	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of disturbed bare ground less than 10%	Attribute and target based on Perrin et al. (2014)
Physical structure: drainage	Percentage area in local vicinity of a representative number of monitoring stops	Area showing signs of drainage from heavy trampling, tracking or ditches less than 10%	Attribute and target based on Perrin et al. (2014)
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat	This includes species listed in the Flora (Protection Order, 2015 and/or the red data lists (Lockhart et al., 2012; Wyse Jackson et al., 2016). Roche et al. (2014) compiled and mapped existing rare and notable plant records for the SAC and added any new records collected during the NSUH survey. No relevant species were recorded in this habitat; however, new records should be considered within this attribute

7150 Depressions on peat substrates of the Rhynchosporion

To restore the favourable conservation condition of Depressions on peat substrates of the Rhynchosporion in Mweelrea/Sheeffry/Erriff Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Mweelrea/Sheeffry/Erriff Complex SAC was surveyed as part of the National Survey of Upland Habitats (NSUH; see Perrin et al., 2014 and Roche et al., 2014). Depressions on peat substrates of the Rhynchosporion habitat was mapped in detail for the SAC and the total area of the qualifying habitat stated by Roche et al. (2014) is 406ha, covering 1.9% of the SAC. Roche et al. (2014) report obvious losses of approximately 0.01ha since 1995. Further information can be found in Roche et al. (2014). Further details on this and the following attributes can be found in the Mweelrea/Sheeffry/Erriff Complex SAC conservation objectives supporting document for blanket bogs and associated habitats where a summary of the mapping methodology is also presented
Habitat distribution	Occurrence	No decline, subject to natural processes. See map 13	Roche et al. (2014) recorded Rhynchosporion depressions throughout the SAC, including the large broad valley of Laghta Ougher, the lower ground surrounding Laghta Eighter and Tangincartoor, the land surrounding Cregganbaun, Srahrooskey and Srahlea and the land adjacent to and west of Tawnyard Lough. Further information can be found within Roche et al. (2014) and the blanket bogs and associated habitats supporting document
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil nutrient status within natural range	See the blanket bogs and associated habitats supporting document for further details
Vegetation composition: positive indicator species	Number of species at a representative number of 2m x 2m monitoring stops	Number of positive indicator species at each monitoring stop is at least five	Attribute and target based on Perrin et al. (2014), where the list of positive indicator species for this habitat is also presented
Vegetation composition: <i>Rhynchospora</i> spp.	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of white beaked sedge (<i>Rhynchospora alba</i>) and brown beaked sedge (<i>R. fusca</i>) at least 10%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: potential dominant species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of each of the potential dominant species individually less than 35%	Attribute and target based on Perrin et al. (2014). See the blanket bogs and associated habitats supporting document for the list of potential dominant species
Vegetation composition: negative indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of negative indicator species less than 1%	Attribute and target based on Perrin et al. (2014), where the list of negative indicator species for this habitat is also presented
Vegetation composition: non-native species	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of non-native species less than 1%	Attribute and target based on Perrin et al. (2014). No non-native species were recorded within this habitat during the NSUH (Roche et al., 2014)
Vegetation composition: native trees and shrubs	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of scattered native trees and shrubs less than 10%	Attribute and target based on Perrin et al. (2014)
Vegetation structure: <i>Sphagnum</i> condition	Condition at a representative number of 2m x 2m monitoring stops	Less than 10% of the <i>Sphagnum</i> cover is crushed, broken and/or pulled up	Attribute and target based on Perrin et al. (2014)

Vegetation structure: signs of browsing	Percentage of shoots browsed at a representative number of 2m x 2m monitoring stops	Last complete growing season's shoots of ericoids, crowberry (<i>Empetrum nigrum</i>) and bog-myrtle (<i>Myrica gale</i>) showing signs of browsing collectively less than 33%	Attribute and target based on Perrin et al. (2014)
Vegetation structure: burning	Occurrence in local vicinity of a representative number of monitoring stops	No signs of burning in sensitive areas, into the moss, liverwort or lichen layer or exposure of peat surface due to burning	Attribute and target based on Perrin et al. (2014), where the list of sensitive areas for this habitat is also presented
Physical structure: disturbed bare ground	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of disturbed bare ground less than 10%	Attribute and target based on Perrin et al. (2014)
Physical structure: drainage	Percentage area in local vicinity of a representative number of monitoring stops	Area showing signs of drainage from heavy trampling, tracking or ditches less than 10%	Attribute and target based on Perrin et al. (2014)
Physical structure: erosion	Percentage area in local vicinity of a representative number of monitoring stops	Less than 5% of the greater bog mosaic comprises erosion gullies and eroded areas	Attribute and target based on Perrin et al. (2014)
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat	This includes species listed in the Flora (Protection) Order, 2015 and/or the red data lists (Lockhart et al., 2012; Wyse Jackson et al., 2016). Roche et al. (2014) compiled and mapped existing rare and notable plant records for the SAC and added any new records collected during the NSUH survey. No relevant species were recorded in this habitat; however, new records should be considered within this attribute

Conservation Objectives for : Mweelrea/Sheeffry/Erriff Complex SAC [001932]

7220 Petrifying springs with tufa formation (Cratoneurion)

To maintain the favourable conservation condition of Petrifying springs with tufa formation (Cratoneurion)* in Mweelrea/Sheeffry/Erriff Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Square metres	Area stable or increasing, subject to natural processes	A total of 3,070m ² (0.307ha) of this habitat was recorded at three locations within Mweelrea/Sheeffry/Erriff Complex SAC, two at Dooaghtry Coast and one at Dooaghtry Flush, by Lyons (2015) (see map 14). The first location at Dooaghtry Coast (site ID: PS027a) was recorded as small tufa-forming springs at the upper limits of the beach with an area of c.20m ² , the second at Dooaghtry Coast (site ID: PS027b) was recorded as an inactive tufa deposit in a spray zone with smaller actively forming tufa deposits with an area of c.50m ² and the third location at Dooaghtry Flush (site ID: PS028) has been described as a large petrifying spring and flush complex with paludal tufa covering an area of c.3,000m ² and is of very high conservation value (Lyons, 2015)
Habitat distribution	Occurrence	No decline, subject to natural processes. See map 14 for point locations	The habitat has been recorded at three locations, two at Dooaghtry Coast and one at Dooaghtry Flush, within Mweelrea/Sheeffry/Erriff Complex SAC by Lyons (2015). Lyons and Kelly (2016) describe eight plant communities of Irish petrifying springs based on relevé data. The Dooaghtry Flush site (PS028) falls into the <i>Palustriella falcata-Carex panicea</i> springs group (Lyons, 2015). It is not known into what groups the Dooaghtry Coast sites fall into as a full survey was not carried out at these locations. Further information on all the vegetation communities associated with this habitat is presented in Lyons and Kelly (2016)
Hydrological regime: height of water table; water flow	Metres; metres per second	Maintain appropriate hydrological regimes	Petrifying springs rely on permanent irrigation, usually from upwelling groundwater sources or seepage sources (Lyons and Kelly, 2013). Water flow should not be altered anthropogenically. See Lyons and Kelly (2016) for further details
Water quality - nitrate level	mg/l	No increase from baseline nitrate level and less than 10mg/l	Target based on data from McGarrigle et al. (2010). See Lyons and Kelly (2016) for further details
Water quality - phosphate level	µg/l	No increase from baseline phosphate level and less than 15µg/l	Based on data from Lyons (2015). See Lyons and Kelly (2016) for further details
Vegetation composition: positive indicator species	Number per spring	At least three positive/high quality indicator species as listed in Lyons and Kelly (2016) and no loss from baseline number	Based on Lyons and Kelly (2016), where the lists of positive and high quality indicator species are presented. The positive indicator moss <i>Didymodon tophaceus</i> was found at both Dooaghtry Coast sites (PS027a, b). The high quality indicator moss <i>Catoscopium nigratum</i> (Near Threatened, Lockhart et al., 2012) was recorded at the Dooaghtry Flush site (PS028), along with the positive indicators bog pimpernel (<i>Anagallis tenella</i>), long-stalked yellow sedge (<i>Carex lepidocarpa</i>), carnation sedge (<i>C. panicea</i>), grass-of-parnassus (<i>Parnassia palustris</i>), common butterwort (<i>Pinguicula vulgaris</i>), brookweed (<i>Samolus valerandi</i>), the bryophytes <i>Aneura pinguis</i> , <i>Bryum pseudotriquetrum</i> , <i>Campylium stellatum</i> , <i>Jungermannia atrovirens</i> , <i>Palustriella falcata</i> , <i>Philonotis calcarea</i> , <i>Scorpidium cossonii</i> and <i>S. scorpioides</i> , the stonewort <i>Chara vulgaris</i> and lesser clubmoss (<i>Selaginella selaginoides</i>) (Lyons, 2015)

Vegetation composition: negative indicator species	Cover (DAFOR scale)	Potentially negative indicator species should not be Dominant or Abundant; invasive species should be absent	Based on Lyons and Kelly (2016), where the lists of potentially negative herbaceous, bryophyte (and alga) and woody species are presented. See Lyons and Kelly (2016) also for details on potentially invasive species, including sycamore (<i>Acer pseudoplatanus</i>) which is invasive in non-wooded springs and a negative indicator species in wooded springs. If two or more potentially negative bryophyte species are present, and if at least two are frequent, or at least one is abundant, then the habitat fails for this attribute. See Lyons and Kelly (2016) for further details. The potentially negative herbaceous species fool's water-cress (<i>Apium nodiflorum</i>) and the potentially negative woody species creeping willow (<i>Salix repens</i>) were recorded at Dooaghtry Flush, but neither species was Dominant or Abundant (Lyons, 2015)
Vegetation structure: sward height	Centimetres	Field layer height between 10cm and 50cm (except for bryophyte-dominated ground <10cm)	See Lyons and Kelly (2016) for further details
Physical structure: trampling/dung	Cover (DAFOR scale)	Cover should not be Dominant or Abundant	See Lyons and Kelly (2016) for further details

7230 Alkaline fens

To restore the favourable conservation condition of Alkaline fens in Mweelrea/Sheeffry/Erriff Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Mweelrea/Sheeffry/Erriff Complex SAC was surveyed as part of the National Survey of Upland Habitats (NSUH; see Perrin et al., 2014 and Roche et al., 2014). Alkaline fens habitat was mapped in detail for the SAC and the total area of the qualifying habitat stated by Roche et al. (2014) is 44.8ha, covering 0.2% of the SAC. Roche et al. (2014) report no obvious losses of this habitat since 1995. Further information can be found in Roche et al. (2014). Further details on this and the following attributes can be found in the Mweelrea/Sheeffry/Erriff Complex SAC conservation objectives supporting document for blanket bogs and associated habitats where a summary of the mapping methodology is also presented
Habitat distribution	Occurrence	No decline, subject to natural processes. See map 15	Roche et al. (2014) recorded alkaline fens throughout the SAC, including the slopes of the Sheeffry Hills, Ben Creggan and Ben Gorm, the Glenlaur valley, Srahrooskey, Glenkeen, Tonatlewa and to the south and south-east of Corragau Lough. Further information can be found within Roche et al. (2014) and the blanket bogs and associated habitats supporting document
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil nutrient status within natural range	See the blanket bogs and associated habitats supporting document for further details
Ecosystem function: peat formation	Flood duration	Maintain active peat formation, where appropriate	In order for peat to form, water levels need to be slightly below or above the soil surface for c.90% of the time (Jim Ryan, pers. comm.)
Ecosystem function: hydrology	Metres	Maintain appropriate natural hydrological regimes necessary to support the natural structure and functioning of the habitat	Maintenance of groundwater, surface water flows and water table levels within natural ranges is essential for this wetland habitat
Ecosystem function: water quality	Water chemistry measures	Maintain appropriate water quality, particularly nutrient levels, to support the natural structure and functioning of the habitat	Fens receive natural levels of nutrients (e.g. iron, magnesium and calcium) from water sources. However, they are generally poor in nitrogen and phosphorus, with the latter tending to be the limiting nutrient
Community diversity	Abundance of variety of vegetation communities	Maintain variety of vegetation communities, subject to natural processes	Roche et al. (2014) recorded three different alkaline fen vegetation communities within this SAC. Data on the abundance of these communities is reproduced in the blanket bogs and associated habitats supporting document. Further information on vegetation communities associated with this habitat is presented in Perrin et al. (2014)
Vegetation composition: number of positive indicator species (brown mosses)	Number of species at a representative number of 2m x 2m monitoring stops	Number of brown moss species present at each monitoring stop is at least one	Attribute and target based on Perrin et al. (2014), where the list of positive indicator species for this habitat is also presented

Vegetation composition: number of positive indicator species (vascular plants)	Number of species at a representative number of 2m x 2m monitoring stops	Number of positive vascular plant indicator species present at each monitoring stop is at least two for small-sedge flushes and at least three for black bog-rush (<i>Schoenus nigricans</i>) flush and bottle sedge (<i>Carex rostrata</i>) fen	Attribute and target based on Perrin et al. (2014), where the list of positive indicator species for this habitat is also presented
Vegetation composition: cover of positive indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of brown moss species and positive vascular plant indicator species at least 20% for small-sedge flushes and at least 75% cover for black bog-rush (<i>Schoenus nigricans</i>) flush and bottle sedge (<i>Carex rostrata</i>) fen	Attribute and target based on Perrin et al. (2014), where the list of positive indicator species for this habitat is also presented
Vegetation composition: negative indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of negative indicator species less than 1%	Attribute and target based on Perrin et al. (2014), where the list of negative indicator species for this habitat is also presented
Vegetation composition: non-native species	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of non-native species less than 1%	Attribute and target based on Perrin et al. (2014). No non-native species were recorded within this habitat during the NSUH (Roche et al., 2014)
Vegetation composition: native trees and shrubs	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of scattered native trees and shrubs less than 10%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: soft rush and common reed cover	Percentage cover in local vicinity of a representative number of monitoring stops	Total cover of soft rush (<i>Juncus effusus</i>) and common reed (<i>Phragmites australis</i>) less than 10%	Attribute and target based on Perrin et al. (2014)
Vegetation structure: height	Percentage of leaves/shoots at a representative number of 2m x 2m monitoring stops	Proportion of live leaves and/or flowering shoots of vascular plants that are more than 5cm above the ground surface should be at least 50%	Attribute and target based on Perrin et al. (2014)
Physical structure: disturbed bare ground	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of disturbed bare ground less than 10%	Attribute and target based on Perrin et al. (2014)
Physical structure: drainage	Percentage area in local vicinity of a representative number of monitoring stops	Area showing signs of drainage from heavy trampling, tracking or ditches less than 10%	Attribute and target based on Perrin et al. (2014)
Physical structure: tufa formations	Percentage cover in local vicinity of a representative number of 2m x 2m monitoring stops	Disturbed proportion of vegetation cover where tufa is present is less than 1%	Attribute and target based on Perrin et al. (2014)
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat	This includes species listed in the Flora (Protection) Order, 2015 (FPO) and/or the red data lists (Lockhart et al., 2012; Wyse Jackson et al., 2016). Roche et al. (2014) compiled and mapped rare and notable plant records for the SAC and added any new records collected during the NSUH survey. During the NSUH, the Near Threatened and FPO listed species bog orchid (<i>Hammarbya paludosa</i>) (Wyse Jackson et al., 2016) and the Near Threatened moss <i>Sphagnum platyphyllum</i> (Lockhart et al., 2012) were recorded within alkaline fen in the SAC by Roche et al. (2014)

Conservation Objectives for : Mweelrea/Sheeffry/Erriff Complex SAC [001932]

8110 Siliceous scree of the montane to snow levels (*Androsacetalia alpinae* and *Galeopsietalia ladani*)

To restore the favourable conservation condition of Siliceous scree of the montane to snow levels (*Androsacetalia alpinae* and *Galeopsietalia ladani*) in Mweelrea/Sheeffry/Erriff Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Mweelrea/Sheeffry/Erriff Complex SAC was surveyed as part of the National Survey of Upland Habitats (NSUH; see Perrin et al., 2014 and Roche et al., 2014). Siliceous scree of the montane to snow levels (<i>Androsacetalia alpinae</i> and <i>Galeopsietalia ladani</i>) was mapped in detail for the SAC and the total area of the qualifying habitat stated by Roche et al. (2014) is 234.7ha, covering 0.1% of the SAC. Roche et al. (2014) report no obvious losses of this habitat since 1995. Further information can be found in Roche et al. (2014). Further details on this and the following attributes can be found in the Mweelrea/Sheeffry/Erriff Complex SAC conservation objectives supporting document for blanket bogs and associated habitats where a summary of the mapping methodology is also presented
Habitat distribution	Occurrence	No decline, subject to natural processes. See map 16	Roche et al. (2014) recorded siliceous screes on several hillsides within the SAC and the habitat was particularly concentrated on the higher slopes and corries of the Sheeffry Hills, on the steeper slopes of Ben Gorm and Ben Creggan and the upper slopes and corries of the Mweelrea Mountains. Further information can be found within Roche et al. (2014) and the blanket bogs and associated habitats supporting document
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil nutrient status within natural range	See the blanket bogs and associated habitats supporting document for further details
Vegetation composition: lichens and bryophytes	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of bryophytes and non-crustose lichen species at least 5%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: negative indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Proportion of vegetation composed of negative indicator species less than 1%	Attribute and target based on Perrin et al. (2014), where the list of negative indicator species for this habitat is also presented
Vegetation composition: non-native species	Percentage cover at a representative number of 2m x 2m monitoring stops	Proportion of vegetation composed of non-native species less than 1%	Attribute and target based on Perrin et al. (2014). During the NSUH, the non-native New Zealand willowherb (<i>Epilobium brunnescens</i>) was recorded within two siliceous screes monitoring stops. One monitoring stop failed due to excessive cover of the species. Pirri-pirri-bur (<i>Acaena novae-zelandiae</i>) was also recorded within siliceous scree habitat during the NSUH (Roche et al., 2014)
Vegetation composition: positive indicator species	Number of species in local vicinity of a representative number of monitoring stops	At least one positive indicator species present in vicinity of each monitoring stop in block scree	Attribute and target based on Perrin et al. (2014). The list of positive indicator species for this habitat is also presented in Perrin et al. (2014) and is the same as for 8220 Siliceous rocky slopes
Vegetation composition: grass species and dwarf shrubs	Percentage cover in local vicinity of a representative number of monitoring stops	Total cover of grass species and dwarf shrubs less than 20%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: bracken, native trees and shrubs	Percentage cover in local vicinity of a representative number of monitoring stops	Total cover of bracken (<i>Pteridium aquilinum</i>), native trees and shrubs less than 25%	Attribute and target based on Perrin et al. (2014)

Vegetation structure: grazing and browsing	Percentage of leaves/shoots grazed/browsed at a representative number of 2m x 2m monitoring stops	Live leaves of forbs and shoots of dwarf shrubs showing signs of grazing or browsing collectively less than 50%	Attribute and target based on Perrin et al. (2014)
Physical structure: disturbance	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Ground disturbed by human and animal paths, scree running, vehicles less than 10%	Attribute and target based on Perrin et al. (2014)
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat and no decline in status of hepatic mats associated with this habitat	This includes species listed in the Flora (Protection Order, 2015 and/or the red data lists (Lockhart et al., 2012; Wyse Jackson et al., 2016). Roche et al. (2014) compiled and mapped existing rare and notable plant records for the SAC and added any new records collected during the NSUH survey. During the NSUH, the Near Threatened and FPO listed liverwort <i>Mastigophora woodsii</i> and the Near Threatened moss <i>Grimmia donniana</i> (Lockhart et al., 2012) were recorded within siliceous scree habitat in the SAC by Roche et al. (2014)

8210 Calcareous rocky slopes with chasmophytic vegetation

To maintain the favourable conservation condition of Calcareous rocky slopes with chasmophytic vegetation in Mweelrea/Sheeffry/Erriff Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Mweelrea/Sheeffry/Erriff Complex SAC was surveyed as part of the National Survey of Upland Habitats (NSUH; see Perrin et al., 2014 and Roche et al., 2014). Calcareous rocky slopes with chasmophytic vegetation was mapped in detail for the SAC and the total area of the qualifying habitat stated by Roche et al. (2014) is 1.7ha, covering 0.01% of the SAC. Roche et al. (2014) report no obvious losses of this habitat since 1995. Further information can be found in Roche et al. (2014). Further details on this and the following attributes can be found in the Mweelrea/Sheeffry/Erriff Complex SAC conservation objectives supporting document for blanket bogs and associated habitats where a summary of the mapping methodology is also presented
Habitat distribution	Occurrence	No decline, subject to natural processes. See map 17	Roche et al. (2014) recorded calcareous rocky slope habitat in the north-eastern corrie of the Mweelrea Mountains and on the slopes of Sheeffry Hills including the Lough Tarriff corrie, north-east of the Lough Brawn corrie, the slopes above Lugaloughaun and on the slopes north-east of the Doo Lough parking area. It is also present on the high slopes west of Drummin village. Further information can be found within Roche et al. (2014) and the blanket bogs and associated habitats supporting document
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil nutrient status within natural range	See the blanket bogs and associated habitats supporting document for further details
Vegetation composition: positive indicator fern and <i>Saxifraga</i> species	Number of species in local vicinity of a representative number of monitoring stops	Number of ferns and <i>Saxifraga</i> indicators at each monitoring stop is at least one	Attribute and target based on Perrin et al. (2014), where the list of positive indicator species for this habitat is also presented
Vegetation composition: positive indicator species	Number of species in local vicinity of a representative number of monitoring stops	Number of positive indicator species at each monitoring stop is at least three	Attribute and target based on Perrin et al. (2014), where the list of positive indicator species for this habitat is also presented
Vegetation composition: non-native species	Percentage cover in local vicinity of a representative number of monitoring stops	Proportion of vegetation composed of non-native species less than 1%	Attribute and target based on Perrin et al. (2014). During the NSUH, the non-native New Zealand willowherb (<i>Epilobium brunnescens</i>) was recorded within the one calcareous rocky slopes monitoring stop, but its cover was not sufficiently high enough to cause the monitoring stop to fail the target (Roche et al., 2014)
Vegetation composition: bracken, native trees and shrubs	Percentage cover in local vicinity of a representative number of monitoring stops	Total cover of bracken (<i>Pteridium aquilinum</i>), native trees and shrubs less than 25%	Attribute and target based on Perrin et al. (2014)
Vegetation structure: grazing and browsing	Percentage of leaves/shoots grazed/browsed in local vicinity of a representative number of monitoring stops	Live leaves of forbs and shoots of dwarf shrubs showing signs of grazing or browsing collectively less than 50%	Attribute and target based on Perrin et al. (2014)

Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat	This includes species listed in the Flora (Protection) Order, 2015 and/or the red data lists (Lockhart et al., 2012; Wyse Jackson et al., 2016). Roche et al. (2014) compiled and mapped existing rare and notable plant records for the SAC and added any new records collected during the NSUH survey. The Vulnerable Alpine saw-wort (<i>Saussurea alpina</i>) (Wyse Jackson et al., 2016) has been recorded on the Sheeffry Hills in the Lough Brawn corrie (historic record) and the adjacent Lough Tarriff corrie (NSUH record), but this species cannot be attributed specifically to calcareous rocky slopes (Roche et al., 2014)
-------------------------------------	--------------------------------	--	---

Conservation Objectives for : Mweelrea/Sheeffry/Erriff Complex SAC [001932]

8220 Siliceous rocky slopes with chasmophytic vegetation

To restore the favourable conservation condition of Siliceous rocky slopes with chasmophytic vegetation in Mweelrea/Sheeffry/Erriff Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Mweelrea/Sheeffry/Erriff Complex SAC was surveyed as part of the National Survey of Upland Habitats (NSUH; see Perrin et al., 2014 and Roche et al., 2014). Siliceous rocky slopes with chasmophytic vegetation was mapped in detail for the SAC and the total area of the qualifying habitat stated by Roche et al. (2014) is 164.9ha, covering 0.8% of the SAC. Roche et al. (2014) report no obvious losses of this habitat since 1995. Further information can be found in Roche et al. (2014). Further details on this and the following attributes can be found in the Mweelrea/Sheeffry/Erriff Complex SAC conservation objectives supporting document for blanket bogs and associated habitats where a summary of the mapping methodology is also presented
Habitat distribution	Occurrence	No decline, subject to natural processes. See map 18	Roche et al. (2014) recorded siliceous rocky slope habitat on hillsides across the SAC including Laghta Eighter, the Sheeffry Hills, the Mweelrea Mountains, Ben Gorm and Ben Creggan and the Glenawough corrie. Further information can be found within Roche et al. (2014) and the blanket bogs and associated habitats supporting document
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil nutrient status within natural range	See the blanket bogs and associated habitats supporting document for further details
Vegetation composition: positive indicator species	Number of species in local vicinity of a representative number of monitoring stops	At least one positive indicator species present in vicinity of each monitoring stop	Attribute and target based on Perrin et al. (2014). The list of positive indicator species for this habitat is also presented in Perrin et al. (2014) and is the same as for 8110 Siliceous screes
Vegetation composition: non-native species	Percentage cover in local vicinity of a representative number of monitoring stops	Proportion of vegetation composed of non-native species less than 1%	Attribute and target based on Perrin et al. (2014). During the NSUH, the non-native New Zealand willowherb (<i>Epilobium brunnescens</i>) was recorded within three siliceous scree monitoring stops, two of which failed due to excessive cover of the species. Cotoneaster (<i>Cotoneaster</i> sp.) was also noted in this habitat during the NSUH (Roche et al., 2014)
Vegetation composition: bracken, native trees and shrubs	Percentage cover in local vicinity of a representative number of monitoring stops	Total cover of bracken (<i>Pteridium aquilinum</i>), native trees and shrubs less than 25%	Attribute and target based on Perrin et al. (2014)
Vegetation structure: grazing and browsing	Percentage of leaves/shoots grazed/browsed in local vicinity of a representative number of monitoring stops	Live leaves of forbs and shoots of dwarf shrubs showing signs of grazing or browsing collectively less than 50%	Attribute and target based on Perrin et al. (2014)
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat and no decline in status of hepatic mats associated with this habitat	This includes species listed in the Flora (Protection) Order, 2015 and/or the red data lists (Lockhart et al., 2012; Wyse Jackson et al., 2016). Roche et al. (2014) compiled and mapped rare and notable plant records for the SAC and added any new records collected during the NSUH survey. No relevant species were recorded in this habitat; however, new records should be considered within this attribute

1013 Geyer's Whorl Snail *Vertigo geyeri*

To maintain the favourable conservation condition of Geyer's Whorl Snail in Mweelrea/Sheeffry/Erriff Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Distribution: occupied sites	Number	No decline. There is one known site for this species in the SAC within five 1km grid squares - L7369, L7468, L7469, L7568 and L7569. See map 19	Geyer's whorl snail (<i>Vertigo geyeri</i>) in Mweelrea/Sheeffry/Erriff Complex SAC is found in wetland habitat at Dooaghtry (Holyoak, 2005; Moorkens, 2004; Moorkens and Killeen, 2011). This locality should be considered as a single site (site code VgCAM5 in Moorkens and Killeen (2011)). There have been records from five 1km grid squares within the site and within the SAC boundary (L7369, L7468, L7469, L7568 and L7569)
Occurrence in suitable habitat	Percentage positive records in a representative number of samples	Number of positive samples at least stable, subject to natural processes	Attribute and target based on Moorkens and Killeen (2011). Positive samples mean the confirmed presence of snails (living or recently dead adults and/or juveniles)
Habitat quality: soil wetness	Metres along monitoring transect	No decline, subject to natural processes	Attribute and target based on Moorkens and Killeen (2011)
Habitat area	Hectares	Stable or increasing subject to natural processes. No less than 30ha of at least sub-optimal habitat	Suitable conditions for the species can be found throughout the site. Moorkens and Killeen (2011) give a target figure of 30ha to be at least in sub-optimal condition. Optimal habitat is defined as flushed fen grassland with sedge/moss lawns 5-30cm tall, containing a high diversity, with species such as <i>Carex lepidocarpa</i> , <i>Parnassia palustris</i> , <i>Pinguicula vulgaris</i> , <i>Juncus articulatus</i> , <i>Eriophorum</i> sp. and the mosses <i>Scorpidium revolvens</i> and <i>Campylium stellatum</i> , with scattered tussocks of <i>Schoenus nigricans</i> ≤80cm tall. During sampling, the water table should be between 0-5cm of the soil surface, but not above ground level. Sub-optimal habitat is defined as above, but more dominated by <i>Schoenus nigricans</i> tussocks with mosses between the tussocks, or overall sward height is more than 30cm, or the water table is below 5cm, or the ground is flooded at the time of sampling

1014 **Narrow-mouthed Whorl Snail *Vertigo angustior***

To maintain the favourable conservation condition of Narrow-mouthed Whorl Snail in Mweelrea/Sheeffry/Erriff Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Distribution: occupied sites	Number	No decline. There is one known site for this species in the SAC, within three 1km squares - L7567, L7568, L7470. See map 19	Narrow-mouthed whorl snail (<i>Vertigo angustior</i>) in Mweelrea/Sheeffry/Erriff Complex SAC is found on the coastal dunes and machair at Dooaghtry (Holyoak, 2005; Moorkens, 2007; Moorkens and Killeen, 2011, where the site code is VaCAM3). This locality, which has been known for over 100 years, should be considered as a single site. There have been records from three 1km grid squares within the site and within the SAC boundary (L7567, L7568, L7470). There has been just a single record from L7567 in 1991, but the species has been recorded from L7568 and L7470 on more than one occasion and since 2000
Occurrence in suitable habitat	Percentage positive records in a representative number of samples	Number of positive samples at least stable, subject to natural processes	Attribute and target based on Moorkens and Killeen (2011). Positive samples mean the confirmed presence of snails (living or recently dead adults and/or juveniles). Optimal and sub-optimal habitats are defined below
Habitat quality: optimal soil wetness	Metres along monitoring transect	No decline, subject to natural processes	Attribute and target based on Moorkens and Killeen (2011)
Habitat extent	Hectares	Stable or increasing, subject to natural processes. No less than 0.23ha of optimal habitat and 0.44ha of sub-optimal habitat	Suitable conditions for the species can be found throughout the site, but the exact locations have clearly changed historically (Tattersfield, 1993; Moorkens, 2007). Moorkens and Killeen (2011) give a figure of 0.23ha of optimal habitat and 0.41ha of sub-optimal habitat. These should be viewed as minimum figures to maintain the population. Optimal habitat is defined as maritime grassland with red fescue (<i>Festuca rubra</i>) and silverweed (<i>Potentilla anserina</i>), 10-25cm in height OR damp grassland dominated by yellow flag (<i>Iris pseudacorus</i>), silverweed and other grasses up to 0.9m high. Soil is damp and friable soil, covered with a layer of humid, open structured thatch and area is lightly grazed by cattle. Sub-optimal habitat is defined as above, but with vegetation height less than 5cm or greater than 15cm, or the yellow flag grassland greater than 0.9m high, or the water table is below 5cm, or the ground is flooded at the time of sampling

1029 Freshwater Pearl Mussel *Margaritifera margaritifera*

To restore the favourable conservation condition of Freshwater Pearl Mussel in Mweelrea/Sheeffry/Erriff Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Distribution	Kilometres	Maintain distribution at 4.34km. Note that the distribution target length includes the perimeter of Fin Lough. See map 20	The conservation objective applies to the Bundorragha freshwater pearl mussel population in Mweelrea/Sheeffry/Erriff Complex SAC, which is of international importance and one of eight Irish populations prioritised for conservation action (Moorkens, 2010; NPWS, 2010). The main distribution and abundance of the freshwater pearl mussel in the Bundorragha catchment was mapped by Moorkens (2005) from downstream of the confluence of the Owennaglogh (just upstream of Delphi Bridge) to the river mouth at Killary Harbour. In 2014, a small number of mussels was found in Fin Lough, near the mouth of the Owengarr River (Roden and Murphy, 2014). Moorkens (2017) found mussels from the cottage near Doo Lough to Fin Lough in the Owengarr River. The target is for the species to be sufficiently widespread to maintain itself on a long-term basis as a viable component of the Bundorragha system. See NPWS (2010) for further information
Population size	Number of adult mussels	Restore populations to at least 2 million adult mussels	The Bundorragha population was estimated as 2,000,000 (NPWS, 2010). Mussels are categorised as abundant along much of the Bundorragha, densities exceed 250/m ² throughout the lower sections and are up to 500 mussels/m ² in places (Moorkens, 2005, 2012, 2017). Mussel density is low in the Owengarr River (Moorkens, 2017). Moorkens (2012) recorded a net decrease of 2.6% in adults between 2005-2012. NPWS (2013) assumed that all priority populations had declined at a rate of 1% per year 2007-2012. A mussel kill and associated decline in adult and juvenile mussels was recorded in 2014 (Moorkens, 2014). In 2016, recruitment was excellent and the conservation condition favourable (Moorkens, 2017); however, a further kill was recorded in 2017, demonstrating that the Bundorragha is close to sustainable, but is subject to on-going significant pressures. The target is for the species to be sufficiently abundant to maintain itself on a long-term basis as a viable component of the Bundorragha system
Population structure: recruitment	Percentage per size class	Restore to at least 20% of population no more than 65mm in length; and at least 5% of population no more than 30mm in length	Mussels of no more than 65mm are considered 'young mussels' and may be found buried in the substratum and/or beneath adult mussels. Mussels of no more than 30mm are 'juvenile mussels' and are always buried in the substratum. See the European Communities Environmental Objectives (Freshwater Pearl Mussel) Regulations 2009. The Sub-basin Management Plan (NPWS, 2010) summarises the demographic work up to 2010 (Ross, 1984; Moorkens, 2005, 2009). The Bundorragha had recovered to favourable condition in 2009, with 21.1% of mussels ≤65mm and 7.5% ≤30mm (Moorkens, 2009; NPWS, 2010). It returned to unfavourable condition in 2012 (6% ≤65mm, 0.4% ≤30mm) and 2014 (12.7% ≤65mm, 3.3% ≤30mm) (Moorkens, 2012, 2014). It was favourable again in 2016 (43.5% ≤65mm, 23.2% ≤30mm) (Moorkens, 2017). The Bundorragha population is almost sustainable. The target is for sufficient juvenile recruitment to allow the species to maintain itself on a long-term basis as a viable component of the Bundorragha system

Population structure: adult mortality	Percentage	No more than 5% decline from previous number of live adults counted; dead shells less than 1% of the adult population and scattered in distribution	5% is considered the cut-off between the combined errors associated with natural fluctuations and sampling methods and evidence of true population decline. 1% of dead shells is considered to be indicative of natural losses. The Bundorragha passed both targets in 2009 (Moorkens, 2009; NPWS, 2010). It failed the targets in 2012 (2.55% decline in live adults and 2.91% dead shells) and again in 2014, when a significant kill was recorded, and passed in 2016 (Moorkens, 2012, 2014, 2017). The target is for sufficient survival of adults to allow the species to maintain itself on a long-term basis as a viable component of the Bundorragha system
Suitable habitat: extent	Kilometres	Maintain suitable habitat extent in 2.67km of the Bundorragha (see map 20) and any additional stretches necessary for salmonid spawning. Note that the suitable habitat target length includes the perimeter of Fin Lough	The extent of the mussel habitat in the Bundorragha has been well-documented: from the early surveys of Ross (1984, 1988) and Moorkens (1995, 1996) to full baseline monitoring in 2005 (Moorkens, 2005), and survey upstream of Fin Lough in 2016 (Moorkens, 2017). Much of the Owengarr river is unsuitable owing to unstable substratum, and mussel numbers are small. Most of the available habitat in the Bundorragha system is occupied by mussels (Moorkens, 2005, 2009, 2012, 2014). Despite the Bundorragha habitat's recovery to favourable condition in 2009 (NPWS, 2010), it returned to unfavourable condition (Moorkens, 2012, 2014). It reached favourable again in 2016, but flow/drought, sedimentation and nutrient impacts are on-going concerns. The target is for sufficient habitat in favourable condition to allow the species to maintain itself on a long-term basis as a viable component of the Bundorragha system
Suitable habitat: condition	Kilometres	Restore condition of suitable habitat	The habitat is a combination of the area of 1) habitat adult and juvenile mussels can occupy; 2) spawning and nursery habitats host fish can occupy. Fish nursery habitat typically overlaps with mussel habitat. Fish spawning habitat is generally adjacent to mussel habitat, but may lie upstream of the generalised mussel distribution. Only spawning areas that can regularly contribute juvenile fish to areas occupied by adult mussels should be considered. Availability of mussel and fish habitat is determined by flow and substratum conditions. It is highly sensitive to hydromorphological changes, sedimentation and nutrient enrichment. Pressures throughout the catchment (map 20) contribute to such impacts. The habitat in the Bundorragha periodically cannot support sufficient juvenile and adult survival (Moorkens, 2012, 2014). The target is for sufficient habitat in favourable condition to allow the species to maintain itself on a long-term basis as a viable component of the Bundorragha system
Water quality: macroinvertebrate and phytobenthos (diatoms)	Ecological quality ratio (EQR)	Restore water quality - macroinvertebrates: EQR greater than 0.90 (Q4-5 or Q5); phytobenthos: EQR greater than 0.93	The EQR targets correspond to high ecological status for these two Water Framework Directive biological quality elements. They represent high water quality with very low nutrient concentrations (oligotrophic conditions). In 2009, the habitat in the Bundorragha system passed both targets (Williams, 2009; NPWS, 2010). See also The European Communities Environmental Objectives (Surface Waters) Regulations 2009. The target is for sufficient habitat in favourable condition to allow the species to maintain itself on a long-term basis as a viable component of the Bundorragha system

Substratum quality: filamentous algae (macroalgae); macrophytes (rooted higher plants)	Percentage	Restore substratum quality - filamentous algae: absent or trace (less than 5%); macrophytes: absent or trace (less than 5%)	The Bundorragha passed both targets in 2009 (borderline pass/fail for macrophytes) (Moorkens, 2009; NPWS, 2010). It failed the macrophyte target in 2012 and <i>Myriophyllum</i> cover reached 80% in parts of the mussel habitat surveyed (Moorkens, 2012). In 2014, it failed both targets, with 75% macrophyte cover in places, and a general increase in filamentous algae reaching cover abundance of up to 90% (Moorkens, 2014). Filamentous algae were present in 2016, but at trace levels, and <i>Myriophyllum</i> was common, but less abundant than in previous surveys (Moorkens, 2017). Sufficient recruitment of juvenile mussels is being prevented by the poor condition of the river substratum. The target is for sufficient habitat in favourable condition to allow the species to maintain itself on a long-term basis as a viable component of the Bundorragha system
Substratum quality: sediment	Occurrence	Restore substratum quality - stable cobble and gravel substrate with very little fine material; no artificially elevated levels of fine sediment	The Bundorragha passed this target in 2009 (Moorkens, 2009; NPWS, 2010). In 2012, sand cover had increased and although heavy in places, it passed the target (Moorkens, 2012). In 2014, sand cover was 'deep and heavy in places': a borderline pass/fail (Moorkens, 2014). Sand was not recorded in 2016 and silt plumes were restricted to margins and other areas of slower flow (Moorkens, 2017). Sand deposition may be acting in combination with bank erosion, flow changes and nutrients to make the habitat unfavourable for mussels. The impacts include increased frequency of drought and emersion of mussels, and increased macrophyte cover (Moorkens, 2014), and temporary declines in oxygen in the substratum may also occur. Sufficient survival of juvenile and adult mussels is being prevented by periodic poor substratum condition. The target is for sufficient habitat in favourable condition to allow the species to maintain itself on a long-term basis as a viable component of the Bundorragha system
Substratum quality: oxygen availability	Redox potential	Restore to no more than 20% decline from water column to 5cm depth in substrate	Differences in redox potential between the water column and the substrate correlate with differences in oxygen levels. Juvenile mussels require full oxygenation while buried in gravel. In suitable habitat, there should be very little loss of redox potential between the water column and underlying gravels. The Bundorragha passed the redox target in 2009, 2012, 2014 and 2016, with average declines of 16%, 18.8%, 14.7% and 9.1% respectively (Moorkens, 2009, 2012, 2014, 2017; NPWS, 2010). As noted above, however, the habitat is periodically impacted, sand deposition is a problem and declines in oxygen in the substratum may be occurring under certain conditions (e.g. at low flows, or when organic material is decaying). The target is for sufficient habitat in favourable condition to allow the species to maintain itself on a long-term basis as a viable component of the Bundorragha system

Hydrological regime: flow variability	Metres per second	Restore appropriate hydrological regime	The availability of suitable habitat is largely determined by flow (catchment geology is the other important factor). In order to restore the species' habitat, flow variability over the annual cycle must be such that: 1) high flows can wash fine sediments from the substratum; 2) high flows are not artificially increased so as to cause excessive scour of mussel habitat; 3) low flows do not exacerbate the deposition of fine sediments or growth of algae/macrophytes and 4) low flows do not cause stress to mussels in terms of exposure, water temperatures, food availability or aspects of the reproductive cycle; see Moorkens and Killeen (2014). Groundwater inflow to the substratum also contributes to water-cycling and favourable habitat condition. The flow in the Bundorragha has been significantly impacted (Moorkens, 2014). The target is for sufficient habitat in favourable condition to allow the species to maintain itself on a long-term basis as a viable component of the Bundorragha system
Host fish	Number	Maintain sufficient juvenile salmonids to host glochidial larvae	Salmonid fish are host to the larval stage of the freshwater pearl mussel and, thus, are essential to completion of the life cycle. 0+ and 1+ fish are typically used, both because of habitat overlaps and the development of immunity with age in fish. Fish presence is sufficient as higher fish density and biomass is indicative of enriched conditions in mussel rivers. Geist et al. (2006) found that higher densities of host fish coincided with eutrophication, poor substrate quality for mussels and a lack of mussel recruitment, while significantly lower host fish density and biomass were associated with high juvenile mussel numbers. Fish movement patterns must be such that 0+ fish remain in the mussel habitat until their 1+ summer. No fish stocking should occur within the mussel habitat, nor any works that may change the salmonid balance or residency time. In 2009, glochidia were found on salmon, but not trout, in the Bundorragha (Johnston, 2009; NPWS, 2010)
Fringing habitat: area and condition	Hectares	Maintain the area and condition of fringing habitats necessary to support the population	Riparian habitats, including those along lake fringes, particularly natural/semi-natural woodlands and wetlands, even where they do not form part of a natural floodplain, are a vital part of the structure and functioning of river systems. Fringing habitats aid in the settlement of fine suspended matter, protect banks from erosion, contribute to nutrient cycling and the aquatic food web (e.g. allochthonous matter) and provide habitat for certain life-stages of fish, birds and invertebrates. Shade may also be important in suppressing algal and macrophyte growth in enriched rivers and moderating temperatures. Fringing habitats are dependent on rivers/lakes, particularly their water levels, and support wetland communities and species of conservation concern. Riverbank erosion is impacting on mussels in the Bundorragha. The target is for sufficient habitat in favourable condition to allow the species to maintain itself on a long-term basis as a viable component of the Bundorragha system

Conservation Objectives for : Mweelrea/Sheeffry/Erriff Complex SAC [001932]

1106 Salmon *Salmo salar*

To restore the favourable conservation condition of Atlantic Salmon in Mweelrea/Sheeffry/Erriff Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Distribution: extent of anadromy	Percentage of river accessible	100% of river channels down to second order accessible from estuary	Artificial barriers block salmon's upstream migration, thereby limiting species to lower stretches and restricting access to spawning areas
Adult spawning fish	Number	Conservation limit (CL) for each system consistently exceeded	A conservation limit (CL) is defined by the North Atlantic Salmon Conservation Organisation (NASCO) as "the spawning stock level that produces long-term average maximum sustainable yield as derived from the adult to adult stock and recruitment relationship". The target is based on the Standing Scientific Committee on Salmon (SSCS) annual model output of CL attainment levels. See SSCS (2016). Attainment of CL estimates are derived from direct counts of adults (rod catch, fish counter) or indirectly by fry abundance counts. The Carrownisky River is currently below CL, meeting 55% of CL. The other river systems overlapping the SAC, namely the Bundorragha, Erriff and Bunowen systems, are currently reaching CL targets
Salmon fry abundance	Number of fry/5 minutes electrofishing	Maintain or exceed 0+ fry mean catchment-wide abundance threshold value. Currently set at 17 salmon fry/5 minutes sampling	The target is the threshold value for rivers currently exceeding their conservation limit (CL)
Out-migrating smolt abundance	Number	No significant decline	Smolt abundance can be negatively affected by a number of impacts such as estuarine pollution, predation and sea lice (<i>Lepeophtheirus salmonis</i>)
Number and distribution of redds	Number and occurrence	No decline in number and distribution of spawning redds due to anthropogenic causes	Salmon spawn in clean gravels
Water quality	EPA Q value	At least Q4 at all sites sampled by EPA	Q values based on triennial water quality surveys carried out by the Environmental Protection Agency (EPA)

Conservation Objectives for : Mweelrea/Sheeffry/Erriff Complex SAC [001932]

1355 **Otter *Lutra lutra***

To maintain the favourable conservation condition of Otter in Mweelrea/Sheeffry/Erriff Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Distribution	Percentage positive survey sites	No significant decline	Measure based on standard otter survey technique. Favourable Conservation Status (FCS) target, based on 1980/81 survey findings, is 88% in SACs. Current range is estimated at 93.6% (Reid et al., 2013)
Extent of terrestrial habitat	Hectares	No significant decline. Area mapped and calculated as 826.0ha along river banks/lake shoreline/ around ponds and 26.3ha above high water mark (HWM)	No field survey. Areas mapped to include 10m terrestrial buffer, identified as critical for otters (NPWS, 2007), along shoreline (above HWM) and along rivers and around water bodies
Extent of marine habitat	Hectares	No significant decline. Area mapped and calculated as 118.0ha	No field survey. Area mapped based on evidence that otters tend to forage within 80m of the shoreline (HWM) (Kruuk, 2006; NPWS, 2007)
Extent of freshwater (river) habitat	Kilometres	No significant decline. Length mapped and calculated as 397.9km	No field survey. River length calculated on the basis that otters will utilise freshwater habitats from estuary to headwaters (Chapman and Chapman, 1982)
Extent of freshwater (lake) habitat	Hectares	No significant decline. Area mapped and calculated as 244.2ha	No field survey. Area mapped based on evidence that otters tend to forage within 80m of the shoreline (NPWS, 2007)
Couching sites and holts	Number	No significant decline	Otters need lying up areas throughout their territory where they are secure from disturbance (Kruuk and Moorhouse, 1991; Kruuk, 2006)
Fish biomass available	Kilograms	No significant decline	Broad diet that varies locally and seasonally, but dominated by fish, in particular salmonids, eels and sticklebacks in freshwater (Bailey and Rochford, 2006) and wrasse and rockling in coastal waters (Kingston et al., 1999)
Barriers to connectivity	Number	No significant increase. For guidance, see map 21	Otters will regularly commute across stretches of open water up to 500m e.g. between the mainland and an island; between two islands; across an estuary (De Jongh and O'Neill, 2010). It is important that such commuting routes are not obstructed

1395 Petalwort *Petalophyllum ralfsii*

To maintain the favourable conservation condition of Petalwort in Mweelrea/Sheeffry/Erriff Complex SAC, which is defined by the following list of attributes and targets:


Attribute	Measure	Target	Notes
Distribution	Number and geographical spread of populations	No decline, subject to natural processes. See map 22 for recorded locations	The known population of petalwort (<i>Petalophyllum ralfsii</i>) in Mweelrea/Sheeffry/Erriff Complex SAC is at Dooaghtry, where it occurs on tightly sheep-grazed turf on flat machair plain, on the sides of low sandy hummocks and in flushed machair. Data from NPWS surveys and Campbell (2013). See also Campbell et al. (2015) for further details
Area of suitable habitat	Hectares	No decline, subject to natural processes	The area of occupancy of petalwort (<i>Petalophyllum ralfsii</i>) at Dooaghtry, estimated from polygons drawn around GPS co-ordinates taken from NPWS surveys and Campbell (2013), is c.159,648m ² . However, only c.60% of this area is suitable for petalwort, i.e. c.95,790m ² (9.58ha). See also Campbell et al. (2015)
Hydrological conditions: soil moisture	Occurrence of damp soil conditions	Maintain hydrological conditions so that substrate is kept moist and damp throughout the year, but is not subject to prolonged inundation by flooding in winter	Petalwort (<i>Petalophyllum ralfsii</i>) grows on damp sandy substrate. Based on Campbell (2013) and Campbell et al. (2015)
Hydrological conditions: water table level	Centimetres in a representative number of 1m x 1m monitoring plots	Mean groundwater level should not be more than 80cm from ground surface	See Campbell et al. (2015) for further details
Physical structure: bare soil	Percentage cover in a representative number of 1m x 1m monitoring plots	Mean percentage cover of bare soil should be more than 5%	At Dooaghtry, petalwort (<i>Petalophyllum ralfsii</i>) grows in compacted, sandy ground. Campbell (2013) recorded bryophyte cover of 34-90% in four plots at Dooaghtry. See Campbell et al. (2015) for further details
Vegetation structure: vegetation height	Centimetres in a representative number of 1m x 1m monitoring plots	Mean vegetation height should be less than 6cm	At Dooaghtry, petalwort (<i>Petalophyllum ralfsii</i>) habitat is maintained by rabbit (<i>Oryctolagus cuniculus</i>) and sheep grazing and some trampling (by walkers and sheep). Campbell (2013) recorded a mean height of vegetation of 3.4cm in four plots at Dooaghtry. See Campbell et al. (2015) for further details
Vegetation composition: shrub cover	Percentage cover in a representative number of 1m x 1m monitoring plots	Mean percentage shrub cover should be less than 25%	See Campbell et al. (2015) for further details

Conservation Objectives for : Mweelrea/Sheeffry/Erriff Complex SAC [001932]

1833 Slender Naiad *Najas flexilis*


To maintain the favourable conservation condition of Slender Naiad in Mweelrea/Sheeffry/Erriff Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Population extent	Hectares; distribution	No change to the spatial extent of <i>Najas flexilis</i> within Lough Nahaltora, subject to natural processes. See map 23 for known location	A population of <i>Najas flexilis</i> (slender naiad) occurs in Lough Nahaltora in Mweelrea/Sheeffry/Erriff Complex SAC. A record for <i>Najas flexilis</i> in Fin Lough has been rejected (Roden and Murphy, 2014). It is nevertheless possible that the species is more widespread within the SAC. See the Mweelrea/Sheeffry/Erriff Complex SAC conservation objectives supporting document for <i>Najas flexilis</i> for further details
Population depth	Metres	No change to the depth range of <i>Najas flexilis</i> within Lough Nahaltora, subject to natural processes	See the <i>Najas flexilis</i> supporting document for further details
Population viability	Plant traits	No decline in plant fitness, subject to natural processes	See the <i>Najas flexilis</i> supporting document for further details
Population abundance	Square metres	No change to the cover abundance of <i>Najas flexilis</i> , subject to natural processes	See the <i>Najas flexilis</i> supporting document for further details
Species distribution	Occurrence	No decline, subject to natural processes	See the <i>Najas flexilis</i> supporting document for further details
Habitat extent	Hectares	No decline, subject to natural processes	See the <i>Najas flexilis</i> supporting document for further details
Hydrological regime: water level fluctuations	Metres	Maintain appropriate natural hydrological regime necessary to support the habitat for the species	See the <i>Najas flexilis</i> supporting document for further details
Lake substratum quality	Various	Maintain appropriate substratum type, extent and chemistry to support the population of the species	See the <i>Najas flexilis</i> supporting document for further details
Water quality	Various	Maintain appropriate water quality to support the population of the species	See the <i>Najas flexilis</i> supporting document for further details
Acidification status	pH units; mg/l	Maintain appropriate water and sediment pH, alkalinity and cation concentrations to support the population of <i>Najas flexilis</i> , subject to natural processes	See the <i>Najas flexilis</i> supporting document for further details
Water colour	mg/l PtCo	Maintain/restore appropriate water colour to support the population of <i>Najas flexilis</i>	See the <i>Najas flexilis</i> supporting document for further details
Associated species	Species composition and abundance	Maintain appropriate associated species and vegetation communities to support the population of <i>Najas flexilis</i>	See the <i>Najas flexilis</i> supporting document for further details
Fringing habitat: area and condition	Hectares	Maintain the area and condition of fringing habitats necessary to support the population of <i>Najas flexilis</i>	See the <i>Najas flexilis</i> supporting document for further details




Legend


 Mweelrea/Sheefry/Erriff Complex SAC 001932



Legend

- Mweelrea / Sheeffry / Erriff Complex SAC 001932
- West Connacht Coast SAC 002998
- OSi Discovery Series County Boundaries





An Roinn
Cultúir, Oidhreachta agus Gaeltachta
Department of
Culture, Heritage and the Gaeltacht

**MAP 4:
MWEELREA/SHEEFFRY/ERRIFF SAC
CONSERVATION OBJECTIVES
SALTMARSH HABITATS**

Map to be read in conjunction with the NPWS Conservation Objectives Document.

**SITE CODE:
SAC 001932; version 3.01
CO. MAYO**

0 2 4 6 8 10 km


The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
Ordnance Survey of Ireland Licence No EN 0059216. © Ordnance Survey of Ireland Government of Ireland
Níl sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbhreithniú a déanamh ar theorainneacha na gceantar comharthaithe. Suirbhéarachta Ordonáis na hÉireann Ceadúnas




**Map Version 1
Date: Aug 2017**


Legend


 Mweelrea/Sheeffry/Erriff Complex SAC 001932


 Sand Dune Monitoring Project Boundary

Qualifying Interests


 1210 Annual vegetation of drift lines

 2110 Embryonic shifting dunes

 2120 Shifting dunes along the shoreline with *Ammophila arenaria* (white dunes)




 21A0 Machairs (* in Ireland)


Non Qualifying Interests

 2130 Fixed dunes with herbaceous vegetation ("grey dunes")

 2190 Humid dune slacks

 OSi Discovery Series County Boundary





Legend

- Mweelrea/Sheeffry/Erriff Complex SAC 001932
- OSi Discovery Series County Boundary

Indicative Lake Habitats

- 3110 Oligotrophic waters containing very few minerals of sandy plains: *Littorelletalia uniflorae*
- 3130 Oligotrophic to mesotrophic standing waters with vegetation of the *Littorelletea uniflorae* and/or of the *Isoëto-Nanojuncetea*
- 3160 Natural dystrophic lakes and ponds
- 3110 / 3160 Oligotrophic waters containing very few minerals of sandy plains: *Littorelletalia uniflorae* / Natural dystrophic lakes and ponds
- 3110 / 3130 Oligotrophic waters containing very few minerals of sandy plains: *Littorelletalia uniflorae* / Oligotrophic to mesotrophic standing waters with vegetation of the *Littorelletea uniflorae* and/or of the *Isoëto-Nanojuncetea*

An Roinn
Cultúir, Oidhreacht agus Gaeltachta
Department of
Culture, Heritage and the Gaeltacht

MAP 6:
MWEELREA/SHEEFFRY/ERRIFF SAC
CONSERVATION OBJECTIVES
INDICATIVE LAKE HABITATS

Map to be read in conjunction with the NPWS Conservation Objectives Document.


SITE CODE:
SAC 001932; version 3.01
CO. MAYO

The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
Ordnance Survey of Ireland Licence No EN 0059216. © Ordnance Survey of Ireland Government of Ireland

Níl sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbhreithnithe a déanamh ar theorainneacha na gceantar comharthaithe. Suirbhéarachta Ordonáis na hÉireann Ceadúnas







Map Version 1
Date: Aug 2017


Legend

 Mweelrea/Sheeffry/Erriff Complex SAC 001932

 OSi Discovery Series County Boundary

Cover of 4010 Northern Atlantic wet heaths with *Erica tetralix*

-  0%
-  0.1% - 20%
-  20.1% - 40%
-  40.1% - 60%
-  60.1% - 80%
-  80.1% - 100%




An Roinn
Cultúir, Oidhreacht agus Gaeltachta
Department of
Culture, Heritage and the Gaeltacht

**MAP 7:
MWEELREA/SHEEFFRY/ERRIFF SAC
CONSERVATION OBJECTIVES
WET HEATH**

Map to be read in conjunction with the NPWS Conservation Objectives Document.


**SITE CODE:
SAC 001932; version 3.01
CO. MAYO**



The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
Ordnance Survey of Ireland Licence No EN 0059216. © Ordnance Survey of Ireland Government of Ireland
Níl sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbhreithnithe a déanamh ar theorainneacha na gceantar comharthaithe. Suirbhéarachta Ordonáis na hÉireann Ceadúnas



**Map Version 1
Date: Aug 2017**




Legend

- Mweelrea/Sheeffry/Erriff Complex SAC 001932
- OSi Discovery Series County Boundary

Cover of 4030 European dry heaths

- 0%
- 0.1% - 20%
- 20.1% - 40%
- 40.1% - 60%
- 60.1% - 80%
- 80.1% - 100%




Legend

- Mweelrea/Sheeffry/Erriff Complex SAC 001932
- OSi Discovery Series County Boundary

Cover of 4060 Alpine and Boreal Heath

- 0%
- 0.1% - 20%
- 20.1% - 40%
- 40.1% - 60%
- 60.1% - 80%
- 80.1% - 100%



An Roinn
Cultúir, Oidhreacht agus Gaeltachta
Department of
Culture, Heritage and the Gaeltacht

MAP 10:
MWEELREA/SHEEFFRY/ERRIFF SAC
CONSERVATION OBJECTIVES
HYDROPHILOUS TALL HERB GRASSLAND

Map to be read in conjunction with the NPWS Conservation Objectives Document.

SITE CODE:
SAC 001932; version 3.01
CO. MAYO




The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
Ordnance Survey of Ireland Licence No EN 0059216. © Ordnance Survey of Ireland Government of Ireland

Nil sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbheithníthe a déanamh ar theorainneacha na gceantar comharthaithe. Suirbhéarachta Ordonáis na hÉireann Ceadúnas



Map Version 1
Date: Aug 2017




Legend

- Mweelrea/Sheeffry/Erriff Complex SAC 001932
- OSi Discovery Series County Boundary

Cover of 7130 Blanket Bog (active)*

- 0%
- 0.1% - 20%
- 20.1% - 40%
- 40.1% - 60%
- 60.1% - 80%
- 80.1% - 100%




Legend

- Mweelrea/Sheeffry/Erriff Complex SAC 001932
- OSi Discovery Series County Boundary

Coverage of 7140 Transition mires and quaking bogs

- 0%
- 0.1% - 20%
- 20.1% - 40%
- 40.1% - 60%
- 60.1% - 80%
- 80.1% - 100%



Legend

- Mweelrea/Sheeffry/Erriff Complex SAC 001932
- OSi Discovery Series County Boundary


Cover of 7150 Depressions on peat substrates of the *Rhynchosporion*


- 0%
- 0.1% - 20%
- 20.1% - 40%
- 40.1% - 60%
- 60.1% - 80%
- 80.1% - 100%



Legend

- ★ 7220 Petrifying springs with tufa formation (*Cratoneurion*)
- ▭ Mweelrea/Sheeffry/Erriff Complex SAC 001932
- ▭ OSi Discovery Series County Boundary






Legend

- Mweelrea/Sheeffry/Erriff Complex SAC 001932
- OSi Discovery Series County Boundary

Cover of 7230 Alkaline fens







- 0%
- 0.1% - 20%
- 20.1% - 40%
- 40.1% - 60%
- 60.1% - 80%
- 80.1% - 100%


Legend

 Mweelrea/Sheeffry/Erriff Complex SAC 001932


 OSi Discovery Series County Boundary

Cover of 8110 Siliceous scree of the montane to snow levels (*Androsacetalia alpinae* and *Galeopsietalia ladani*)

-  0%
-  0.1% - 20%
-  20.1% - 40%
-  40.1% - 60%
-  60.1% - 80%
-  80.1% - 100%










Legend

 Mweelrea/Sheeffry/Erriff Complex SAC 001932

 OSi Discovery Series County Boundary

Cover of 8210 Calcareous rocky slopes with chasmophytic vegetation

-  0%
-  0.1% - 20%
-  20.1% - 40%
-  40.1% - 60%
-  60.1% - 80%
-  80.1% - 100%



An Roinn
Cultúir, Oidhreacht agus Gaeltachta
Department of
Culture, Heritage and the Gaeltacht

**MAP 17:
MWEELREA/SHEEFFRY/ERRIFF SAC
CONSERVATION OBJECTIVES
CALCAREOUS ROCKY SLOPES**

Map to be read in conjunction with the NPWS Conservation Objectives Document.

**SITE CODE:
SAC 001932; version 3.01
CO. MAYO**




The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
Ordnance Survey of Ireland Licence No EN 0059216. © Ordnance Survey of Ireland Government of Ireland

Níl sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbheithníthe a déanamh ar theorainneacha na gceantar comharthaithe. Suirbhéarachta Ordonáis na hÉireann Ceadúnas









**Map Version 1
Date: Aug 2017**



Legend

 Mweelrea/Sheeffry/Erriff Complex SAC 001932

 OSi Discovery Series County Boundary




Cover of 8220 Siliceous rocky slopes with chasmophytic vegetation


-  0%
-  0.1% - 20%
-  20.1% - 40%
-  40.1% - 60%
-  60.1% - 80%
-  80.1% - 100%



Legend

- Mweelrea/Sheeffry/Erriff Complex SAC 001932
- OSi Discovery Series County Boundary
- 1013 Geyer's Whorl Snail *Vertigo geyeri*
- 1014 Narrow-mouthed Whorl Snail *Vertigo angustior*








Legend




- 1029 Freshwater Pearl Mussel *Margaritifera margaritifera* - Distribution Target
- 1029 Freshwater Pearl Mussel *Margaritifera margaritifera* - Suitable Habitat Target
- Fin Lough
- Mweelrea/Sheeffry/Erriff Complex SAC 001932
- 1029 Freshwater Pearl Mussel - *Margaritifera margaritifera* Catchment
- OSi Discovery Series County Boundary




Legend

-  Mweelrea/Sheeffry/Erriff Complex SAC 001932
-  1355 Otter - *Lutra lutra* Commuting 250m Buffer



Legend


-  Mweelrea/Sheeffry/Erriff Complex SAC 001932
-  1395 Petalwort *Petalophyllum ralfsii*
-  OSi Discovery Series County Boundary





Legend

- 1833 Slender Naiad - *Najas flexilis*
- Mweelrea/Sheeffry/Erriff Complex SAC 001932
- OSi Discovery Series County Boundary





Conservation objectives for Croaghaun/Slievemore SAC [001955]

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Objective: To maintain or restore the favourable conservation condition of the Annex I habitat(s) and/or the Annex II species for which the SAC has been selected:

Code	Description
4010	Northern Atlantic wet heaths with <i>Erica tetralix</i>
4030	European dry heaths
4060	Alpine and Boreal heaths
7130	Blanket bogs (* if active bog)
8110	Siliceous scree of the montane to snow levels (<i>Androsacetalia alpinae</i> and <i>Galeopsietalia ladani</i>)
8220	Siliceous rocky slopes with chasmophytic vegetation

* denotes a priority habitat



Citation: NPWS (2020) Conservation objectives for Croaghaun/Slievemore SAC [001955]. Generic Version 7.0. Department of Culture, Heritage and the Gaeltacht.

National Parks and Wildlife Service

Conservation Objectives Series

Bellacragher Saltmarsh SAC 002005



An Roinn Ealaíon, Oidhreachta,
Gnóthaí Réigiúnacha, Tuaithe agus Gaeltachta

Department of Arts, Heritage,
Regional, Rural and Gaeltacht Affairs



**National Parks and Wildlife Service,
Department of Arts, Heritage, Regional, Rural and Gaeltacht Affairs,
7 Ely Place, Dublin 2, Ireland.
Web: www.npws.ie
E-mail: nature.conservation@ahg.gov.ie**

Citation:

**NPWS (2016) Conservation Objectives: Bellacragher Saltmarsh SAC 002005.
Version 1. National Parks and Wildlife Service, Department of Arts, Heritage,
Regional, Rural and Gaeltacht Affairs.**

**Series Editor: Rebecca Jeffrey
ISSN 2009-4086**

Introduction

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

** indicates a priority habitat under the Habitats Directive*

002005	Bellacragher Saltmarsh SAC
1330	Atlantic salt meadows (<i>Glauco-Puccinellietalia maritimae</i>)
1410	Mediterranean salt meadows (<i>Juncetalia maritimi</i>)

Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

NPWS Documents

Year :	2007
Title :	Saltmarsh Monitoring Project 2006
Author :	McCorry, M.
Series :	Unpublished report to NPWS
<hr/>	
Year :	2009
Title :	Saltmarsh monitoring project 2007-2008
Author :	McCorry, M.; Ryle, T.
Series :	Unpublished report to NPWS
<hr/>	
Year :	2016
Title :	Bellacragher Saltmarsh SAC (site code: 2005) Conservation objectives supporting document-coastal habitats V1
Author :	NPWS
Series :	Conservation objectives supporting document
<hr/>	

Spatial data sources

Year : Revision 2010

Title : Saltmarsh Monitoring Project 2007-2008. Version 1

GIS Operations : QIs selected; clipped to SAC boundary. Expert opinion used as necessary to resolve any issues arising

Used For : 1330, 1410 (map 2)

Conservation Objectives for : Bellacragher Saltmarsh SAC [002005]

1330 Atlantic salt meadows (*Glauco-Puccinellietalia maritimae*)

To restore the favourable conservation condition of Atlantic salt meadows (*Glauco-Puccinellietalia maritimae*) in Bellacragher Saltmarsh SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes, including erosion and succession. For the sub-site mapped: Bellacragher Bay - 0.80ha. See map 2	Based on data from the Saltmarsh Monitoring Project (SMP) (McCorry, 2007; McCorry and Ryle, 2009). The sub-site Bellacragher Bay (site ID: SMP0021) that supports Atlantic salt meadows (ASM) was mapped to give a total estimated area of 0.80ha within Bellacragher Saltmarsh SAC. NB further unsurveyed areas may be present within the SAC. See the Bellacragher Saltmarsh SAC conservation objectives supporting document for coastal habitats for further details
Habitat distribution	Occurrence	No decline or change in habitat distribution, subject to natural processes. See map 2 for known distribution	Based on data from McCorry (2007) and McCorry and Ryle (2009). NB further unsurveyed areas may be present within the SAC. See the coastal habitats supporting document for further details
Physical structure: sediment supply	Presence/absence of physical barriers	Maintain natural circulation of sediments and organic matter, without any physical obstructions	Based on data from McCorry (2007) and McCorry and Ryle (2009). The saltmarsh occurs mostly on a peat substrate, though some patches occur on the stony shoreline. Within the SAC it mainly occurs as a thin band of vegetation generally between 3-6m wide on a thin band of mud/peat generally overlaying rocky/shingle deposits. See the coastal habitats supporting document for further details
Physical structure: creeks and pans	Occurrence	Maintain creek and pan structure, subject to natural processes, including erosion and succession	Based on data from McCorry (2007) and McCorry and Ryle (2009). The ASM structure is poorly developed, which is typical of these fringe marshes and no creek or pans are present. See the coastal habitats supporting document for further details
Physical structure: flooding regime	Hectares flooded; frequency	Maintain natural tidal regime	Based on data from McCorry (2007) and McCorry and Ryle (2009). See the coastal habitats supporting document for further details
Vegetation structure: zonation	Occurrence	Maintain the range of coastal habitats including transitional zones, subject to natural processes including erosion and succession	Based on data from McCorry (2007) and McCorry and Ryle (2009). The saltmarsh vegetation that occurs on peat generally transitions to blanket bog, wet heath, wet grassland, dry grassland, bracken (<i>Pteridium aquilinum</i>) scrub and mosaics of these habitats. The seaward edge of the saltmarsh usually borders intertidal mud or rocky shoreline. See the coastal habitats supporting document for further details
Vegetation structure: vegetation height	Centimetres	Maintain structural variation within sward	Based on data from McCorry (2007) and McCorry and Ryle (2009). The saltmarsh is mainly grazed by sheep and some areas are noticeably close-cropped. See the coastal habitats supporting document for further details
Vegetation structure: vegetation cover	Percentage cover at a representative number of monitoring stops	Maintain more than 90% of the area outside of creeks vegetated	Based on data from McCorry (2007) and McCorry and Ryle (2009). There are signs of erosion around the edge of the shoreline, some of which is caused by overgrazing by sheep with bare ground (<10%) significant in places. See the coastal habitats supporting document for further details


Vegetation composition: typical species and sub-communities	Percentage cover at a representative number of monitoring stops	Maintain range of sub-communities with typical species listed in McCorry and Ryle (2009)	Based on data from McCorry (2007) and McCorry and Ryle (2009). The species typical of ASM are present including saltmarsh rush (<i>Juncus gerardii</i>), common saltmarsh-grass (<i>Puccinellia maritima</i>), red fescue (<i>Festuca rubra</i>), buck's-horn plantain (<i>Plantago coronopus</i>), sea plantain (<i>P. maritima</i>) and thrift (<i>Armeria maritima</i>). A feature of the marsh is the presence of 'turf fucoids', a term given to various species of brown algae which occur in miniature forms on saltmarshes, especially in western Ireland. See the coastal habitats supporting document for further details
Vegetation composition: negative indicator species - <i>Spartina anglica</i>	Hectares	There is no record of common cordgrass (<i>Spartina anglica</i>) in the SAC and its establishment should be prevented	Based on data from McCorry (2007) and McCorry and Ryle (2009). Common cordgrass (<i>Spartina anglica</i>) was not recorded in Bellacragher Saltmarsh SAC. See the coastal habitats supporting document for further details

Conservation Objectives for : Bellacragher Saltmarsh SAC [002005]


1410 Mediterranean salt meadows (*Juncetalia maritimi*)


To maintain the favourable conservation condition of Mediterranean salt meadows (*Juncetalia maritimi*) in Bellacragher Saltmarsh SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes, including erosion and succession. For the sub-site mapped: Bellacragher Bay - 0.88ha. See map 2	Based on data from the Saltmarsh Monitoring Project (SMP) (McCorry, 2007; McCorry and Ryle, 2009). The sub-site Bellacragher Bay (site ID: SMP0021) that supports Mediterranean salt meadows (MSM) was mapped to give a total estimated area of 0.88ha within Bellacragher Saltmarsh SAC. NB further unsurveyed areas may be present within the SAC. See the Bellacragher Saltmarsh SAC conservation objective supporting document for coastal habitats for further details
Habitat distribution	Occurrence	No decline or change in habitat distribution, subject to natural processes. See map 2 for known distribution	Based on data from McCorry (2007) and McCorry and Ryle (2009). NB further unsurveyed areas may be present within the SAC. See the coastal habitats supporting document for further details
Physical structure: sediment supply	Presence/absence of physical barriers	Maintain natural circulation of sediments and organic matter, without any physical obstructions	Based on data from McCorry (2007) and McCorry and Ryle (2009). See the coastal habitats supporting document for further details
Physical structure: creeks and pans	Occurrence	Maintain creek and pan structure, subject to natural processes, including erosion and succession	Based on data from McCorry (2007) and McCorry and Ryle (2009). See the coastal habitats supporting document for further details
Physical structure: flooding regime	Hectares flooded; frequency	Maintain natural tidal regime	Based on data from McCorry (2007) and McCorry and Ryle (2009). MSM is found high up in the saltmarsh but requires occasional tidal inundation. See the coastal habitats supporting document for further details
Vegetation structure: zonation	Occurrence	Maintain the range of coastal habitats including transitional zones, subject to natural processes including erosion and succession	Based on data from McCorry (2007) and McCorry and Ryle (2009). Patches of sea rush (<i>Juncus maritimus</i>) occur in rocky and muddy areas in narrow bands at the foot of peat cliff faces and may form mosaics with ASM in places. There are also natural transitions from MSM to wet and dry grassland, blanket bog and wet heath. See the coastal habitats supporting document for further details
Vegetation structure: vegetation height	Centimetres	Maintain structural variation in the sward	Based on data from McCorry (2007) and McCorry and Ryle (2009). MSM generally is not significantly overgrazed in the SAC. See the coastal habitats supporting document for further details
Vegetation structure: vegetation cover	Percentage cover at a representative number of monitoring stops	Maintain more than 90% of the area outside of creeks vegetated	Based on data from McCorry (2007) and McCorry and Ryle (2009). Sheep grazing is causing some localised damage to MSM in the SAC although the area affected is generally quite small (<5%). See the coastal habitats supporting document for further details
Vegetation composition: typical species and sub-communities	Percentage cover at a representative number of monitoring stops	Maintain range of sub-communities with typical species listed in McCorry and Ryle (2009)	Based on data from McCorry (2007) and McCorry and Ryle (2009). MSM habitat is dominated by dense sea rush (<i>Juncus maritimus</i>) which forms some of the largest areas of saltmarsh vegetation. See the coastal habitats supporting document for further details
Vegetation composition: negative indicator species - <i>Spartina anglica</i>	Hectares	There is no record of common cordgrass (<i>Spartina anglica</i>) in the SAC and its establishment should be prevented	Based on data from McCorry (2007) and McCorry and Ryle (2009). Common cordgrass (<i>Spartina anglica</i>) was not recorded in Bellacragher Saltmarsh SAC. See the coastal habitats supporting document for further details



Legend

 Bellacragher Saltmarsh SAC 002005



SMP Site Code: SMP0021

Legend

- Bellacragher Saltmarsh SAC 002005
- Saltmarsh Monitoring Project Survey

Annex I Habitats

- 1330 Atlantic salt meadows (*Glauco-Puccinellietalia maritimae*)
- 1330 / 1410 Atlantic salt meadows (*Glauco-Puccinellietalia maritimae*) / Mediterranean salt meadows (*Juncetalia maritimi*)
- 1410 Mediterranean salt meadows (*Juncetalia maritimi*)
- OSi Discovery Series County Boundary

National Parks and Wildlife Service

Conservation Objectives Series

Ox Mountains Bogs SAC 002006



An Roinn Ealaíon, Oidhreachta,
Gnóthaí Réigiúnacha, Tuaithe agus Gaeltachta

Department of Arts, Heritage,
Regional, Rural and Gaeltacht Affairs



**National Parks and Wildlife Service,
Department of Arts, Heritage, Regional, Rural and Gaeltacht Affairs,
7 Ely Place, Dublin 2, Ireland.
Web: www.npws.ie
E-mail: nature.conservation@ahg.gov.ie**

Citation:

NPWS (2016) Conservation Objectives: Ox Mountains Bogs SAC 002006. Version 1. National Parks and Wildlife Service, Department of Arts, Heritage, Regional, Rural and Gaeltacht Affairs.

**Series Editor: Rebecca Jeffrey
ISSN 2009-4086**

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

* indicates a priority habitat under the Habitats Directive

002006	Ox Mountains Bogs SAC
1013	Geyer's Whorl Snail <i>Vertigo geyeri</i>
1528	Marsh Saxifrage <i>Saxifraga hirculus</i>
3110	Oligotrophic waters containing very few minerals of sandy plains (<i>Littorelletalia uniflorae</i>)
3160	Natural dystrophic lakes and ponds
4010	Northern Atlantic wet heaths with <i>Cladonia</i> <i>Cladonia</i>
4030	European dry heaths
7130	Blanket bogs (* if active bog)
7140	Transition mires and quaking bogs
7150	Depressions on peat substrates of the Rhynchosporion

Please note that this SAC adjoins River Moy SAC (002298). See map 2. The conservation objectives for this site should be used in conjunction with those for the adjacent site as appropriate.

Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

NPWS Documents

Year :	1989
Title :	A survey to locate blanket bogs of scientific interest in County Kerry and County Sligo
Author :	Douglas, C.; Garvey, L.; Kelly, L.; O'Sullivan, A.
Series :	Unpublished report to NPWS
Year :	2011
Title :	Monitoring and condition assessment of populations of <i>Vertigo geyeri</i> , <i>Vertigo angustior</i> and <i>Vertigo moulinsiana</i> in Ireland
Author :	Moorkens, E.; Killeen, I.
Series :	Irish Wildlife Manual No. 55
Year :	2013
Title :	A survey of the benthic macrophytes of three hard-water lakes: Lough Bunny, Lough Carra and Lough Owel
Author :	Roden, C.; Murphy, P.
Series :	Irish Wildlife Manual No. 70
Year :	2013
Title :	The status of EU protected habitats and species in Ireland. Volume 2. Habitats assessments
Author :	NPWS
Series :	Conservation assessments
Year :	2013
Title :	The status of EU protected habitats and species in Ireland. Volume 3. Species assessments
Author :	NPWS
Series :	Conservation assessments
Year :	2013
Title :	National survey of upland habitats (phase 3, 2012-2013) Site report no. 10: Ox Mountains Bogs cSAC (002006), Cos. Mayo and Sligo
Author :	Perrin, P.M.; Roche, J.R.; Barron, S.J.; Daly, O.H.; Hodd, R.L.; Muldoon, C.S.; Leydon, K.L.
Series :	Unpublished report to NPWS
Year :	2014
Title :	Guidelines for a national survey and conservation assessment of upland vegetation and habitats in Ireland, Version 2.0
Author :	Perrin, P.M.; Barron, S.J.; Roche, J.R.; O'Hanrahan, B.
Series :	Irish Wildlife Manual No. 79
Year :	2015
Title :	Monitoring recommendations for Marsh Saxifrage (<i>Saxifraga hirculus</i> L.) in the Republic of Ireland
Author :	Muldoon, C.S.; Waldren, S.; Lynn, D.
Series :	Irish Wildlife Manual No. 88
Year :	2015
Title :	Habitats Directive Annex I lake habitats: a working interpretation for the purposes of site-specific conservation objectives and Article 17 reporting
Author :	O Connor, Á.
Series :	Unpublished document by NPWS
Year :	2016
Title :	Ox Mountains Bogs SAC (site code: 2006) Conservation objectives supporting document-upland habitats V1
Author :	NPWS
Series :	Conservation objectives supporting document

Other References

Year :	1982
Title :	Eutrophication of waters. Monitoring assessment and control
Author :	OECD
Series :	OECD, Paris
<hr/>	
Year :	2000
Title :	Colour in Irish lakes
Author :	Free, G.; Allott, N.; Mills, P.; Kennelly, C.; Day, S.
Series :	Verhandlungen Internationale Vereinigung für theoretische und angewandte Limnologie, 27: 2620-2623
<hr/>	
Year :	2002
Title :	Deterioration of Atlantic soft water macrophyte communities by acidification, eutrophication and alkanisation
Author :	Arts, G.H.P.
Series :	Aquatic Botany, 73: 373-393
<hr/>	
Year :	2006
Title :	A reference-based typology and ecological assessment system for Irish lakes. Preliminary investigations. Final report. Project 2000-FS-1-M1 Ecological assessment of lakes pilot study to establish monitoring methodologies EU (WFD)
Author :	Free, G.; Little, R.; Tierney, D.; Donnelly, K.; Coroni, R.
Series :	EPA, Wexford
<hr/>	
Year :	2008
Title :	Water Quality in Ireland 2004-2006
Author :	Clabby, K.J.; Bradley, C.; Craig, M.; Daly, D.; Lucey, J.; McGarrigle, M.; O'Boyle, S.; Tierney, D.; Bowman, J.
Series :	EPA, Wexford
<hr/>	
Year :	2009
Title :	The identification, characterization and conservation value of isoetid lakes in Ireland
Author :	Free, G.; Bowman, J.; McGarrigle, M.; Little, R.; Coroni, R.; Donnelly, K.; Tierney, D.; Trodd, W.
Series :	Aquatic Conservation: Marine and Freshwater Ecosystems, 19 (3): 264-273
<hr/>	
Year :	2010
Title :	Water quality in Ireland 2007-2009
Author :	McGarrigle, M.; Lucey, J.; Ó Cinnéide, M.
Series :	EPA, Wexford
<hr/>	
Year :	2011
Title :	Conservation biology of <i>Saxifraga hirculus</i> L. in Ireland
Author :	Muldoon, C.S.
Series :	Unpublished Ph.D. Thesis, Trinity College Dublin
<hr/>	
Year :	2012
Title :	The impact of conifer plantation forestry on the ecology of peatland lakes
Author :	Drinan, T.J.
Series :	Unpublished PhD thesis, University College Cork
<hr/>	
Year :	2013
Title :	Interpretation manual of European Union habitats- Eur 28
Author :	European Commission- DG Environment
Series :	European Commission

Year : 2014
Title : New vice-county record for Marsh Saxifrage (*Saxifraga hirculus*) in Sligo (H28)
Author : Muldoon, C.; Hodd, R.; Lockhart, N.; Douglas, C.; Roche, J.
Series : Irish Naturalists' Journal, 33 (2): 130-131

Year : in prep.
Title : Monitoring of hard-water lakes in Ireland using charophytes and other macrophytes
Author : Roden, C.; Murphy, P.
Series : Unpublished report to NPWS

Spatial data sources

Year :	2008
Title :	OSi 1:5000 IG vector dataset
GIS Operations :	WaterPolygons feature class clipped to the SAC boundary. Expert opinion used to identify Annex I habitat and to resolve any issues arising
Used For :	3110, 3160 (map 3)
<hr/>	
Year :	2013
Title :	National Survey of Upland Habitats
GIS Operations :	Habitat dataset for site clipped to SAC boundary. Relevant QI selected and exported to new dataset. Expert opinion used as necessary to resolve any issues arising
Used For :	4010, 4030, 7130, 7140, 7150 (maps 4-8)
<hr/>	
Year :	2016
Title :	NPWS rare and threatened species database
GIS Operations :	Dataset created from spatial references in database records. Expert opinion used as necessary to resolve any issues arising
Used For :	1013 (map 9)
<hr/>	

3110 Oligotrophic waters containing very few minerals of sandy plains (Littorelletalia uniflorae)

To maintain the favourable conservation condition of Oligotrophic waters containing very few minerals of sandy plains (Littorelletalia uniflorae) in Ox Mountains Bogs SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Lake habitat 3110 occurs in Lough Easky. It may also be present in other lakes in the SAC, where it is likely to co-occur with habitat 3160, however the exact distribution of habitat 3110 in the SAC is unknown. In line with Article 17 reporting (NPWS, 2013), all lakes larger than 1ha have been mapped as 'potential 3110' (see map 3). Two measures of extent should be used: 1. the area of the lake itself and; 2. the extent of the vegetation communities/zones that typify the habitat. Further information relating to all attributes is provided in the lake habitats supporting document for the purposes of site-specific conservation objectives and Article 17 reporting (O Connor, 2015)
Habitat distribution	Occurrence	No decline, subject to natural processes	As noted above, the exact distribution of habitat 3110 in the SAC is not known. In map 3, all lakes larger than 1ha (based on 1:5,000 data) have been mapped as potential 3110
Typical species	Occurrence	Typical species present, in good condition, and demonstrating typical abundances and distribution	For lists of typical plant species, see Article 17 habitat assessment for 3110 (NPWS, 2013) and the lake habitats supporting document (O Connor, 2015)
Vegetation composition: characteristic zonation	Occurrence	All characteristic zones should be present, correctly distributed and in good condition	The characteristic zonation of lake habitat 3140 has been described (Roden and Murphy, 2013; in prep.), however, significant further work is necessary to describe the characteristic zonation and other spatial patterns in the remaining four Annex I lake habitats
Vegetation distribution: maximum depth	Metres	Maintain maximum depth of vegetation, subject to natural processes	The maximum depth of vegetation is likely to be specific to the lake shoreline in question. An indicative target has not yet been set for this lake habitat type. Indicative targets will be developed for the other lake habitats with time
Hydrological regime: water level fluctuations	Metres	Maintain appropriate natural hydrological regime necessary to support the habitat	Fluctuations in lake water level are typical in Ireland, but can be amplified by activities such as abstraction and drainage. Increased water level fluctuations can increase wave action, up-root vegetation, increase turbidity, alter the substratum and lead to release of nutrients from the sediment. The hydrological regime of the lakes must be maintained so that the area, distribution and depth of the lake habitat and its constituent/characteristic vegetation zones and communities are not reduced
Lake substratum quality	Various	Maintain appropriate substratum type, extent and chemistry to support the vegetation	Research is required to further characterise the substratum types (particle size and origin) and substratum quality (notably pH, calcium, iron and nutrient concentrations) favoured by each of the five Annex I lake habitats in Ireland. It is likely that the lake habitat 3110 is associated with a range of nutrient-poor substrates, from stones, cobble and gravel, through sands, silt, clay and peat. Substratum particle size is likely to vary with depth and along the shoreline within a single lake

Water quality: transparency	Metres	Maintain appropriate Secchi transparency. There should be no decline in Secchi depth/transparency	Transparency relates to light penetration and, hence, to the depth of colonisation of vegetation. It can be affected by phytoplankton blooms, water colour and turbidity. A specific target has yet to be established for this Annex I lake habitat. Habitat 3110 is associated with very clear water. The OECD fixed boundary system set transparency targets for oligotrophic lakes of $\geq 6\text{m}$ annual mean Secchi disk depth, and $\geq 3\text{m}$ annual minimum Secchi disk depth. Free et al. (2009) found high isoetid abundance in lakes with Secchi depths of more than 3m
Water quality: nutrients	$\mu\text{g/l P}$; mg/l N	Maintain the concentration of nutrients in the water column to sufficiently low levels to support the habitat and its typical species	As a nutrient-poor habitat, oligotrophic and Water Framework Directive (WFD) 'high' status targets apply. Where a lake has nutrient concentrations that are lower than these targets, there should be no decline within class, i.e. no upward trend in nutrient concentrations. For lake habitat 3110, annual average TP concentration should be $\leq 10\mu\text{g/l TP}$, average annual total ammonia concentration should be $\leq 0.040\text{mg/l N}$ and annual 95th percentile for total ammonia should be $\leq 0.090\text{mg/l N}$. See also The European Communities Environmental Objectives (Surface Waters) Regulations 2009
Water quality: phytoplankton biomass	$\mu\text{g/l Chlorophyll } a$	Maintain appropriate water quality to support the habitat, including high chlorophyll <i>a</i> status	Oligotrophic and WFD 'high' status targets apply to the lake habitat 3110. Where a lake has a chlorophyll <i>a</i> concentration that is lower than this target, there should be no decline within class, i.e. no upward trend in phytoplankton biomass. The average growing season (March-October) chlorophyll <i>a</i> concentration must be $< 5.8\mu\text{g/l}$. The annual average chlorophyll <i>a</i> concentration should be $< 2.5\mu\text{g/l}$ and the annual peak chlorophyll <i>a</i> concentration should be $\leq 8.0\mu\text{g/l}$. See also The European Communities Environmental Objectives (Surface Waters) Regulations 2009
Water quality: phytoplankton composition	EPA phytoplankton composition metric	Maintain appropriate water quality to support the habitat, including high phytoplankton composition status	The EPA has developed a phytoplankton composition metric for nutrient enrichment of Irish lakes. As for other water quality indicators, habitat 3110 requires WFD high status
Water quality: attached algal biomass	Algal cover and EPA phytobenthos metric	Maintain trace/ absent attached algal biomass ($< 5\%$ cover) and high phytobenthos status	Nutrient enrichment can favour epiphytic and epipelagic algae that can out-compete the submerged vegetation. The cover abundance of attached algae in lake habitat 3110 should, therefore, be trace/ absent ($< 5\%$ cover). EPA phytobenthos can be used as an indicator of changes in attached algal biomass. As for other water quality indicators, habitat 3110 requires high phytobenthos status
Water quality: macrophyte status	EPA macrophyte metric (The Free Index)	Maintain high macrophyte status	Nutrient enrichment can favour more competitive submerged macrophyte species that out-compete the typical and characteristic species for the lake habitat. The EPA monitors macrophyte status for WFD purposes using the 'Free Index'. The target for lake habitat 3110 is high status or an Ecological Quality Ratio (EQR) for lake macrophytes of ≥ 0.90 , as defined in Schedule Five of the European Communities Environmental Objectives (Surface Waters) Regulations 2009

Acidification status	pH units; mg/l	Maintain appropriate water and sediment pH, alkalinity and cation concentrations to support the habitat, subject to natural processes	Acidification can impact on species abundance and composition in soft water lake habitats. In Europe, acidification of isoetid lakes can lead to loss of isoetids and dominance by submerged <i>Sphagnum</i> mosses and <i>Juncus bulbosus</i> (Arts, 2002). The specific requirements of lake habitat 3110, in terms of water and sediment pH, alkalinity and cation concentration, have not been determined. For lake habitat 3110, and adopting a precautionary approach based on Arts (2002), minimum pH should not be <5.5 pH units. Maximum pH should be <9.0 pH units, in line with the surface water standards established for soft waters (where water hardness is ≤100mg/l calcium carbonate). See Schedule Five of the European Communities Environmental Objectives (Surface Waters) Regulations 2009
Water colour	mg/l PtCo	Maintain appropriate water colour to support the habitat	Increased water colour and turbidity decrease light penetration and can reduce the area of available habitat for lake macrophytes, particularly at the lower euphotic depths. The primary source of increased water colour in Ireland is disturbance to peatland. No habitat-specific or national standards for water colour currently exist. Studies have shown median colour concentrations in Irish lakes of 38mg/l PtCo (Free et al., 2000) and 33mg/l PtCo (Free et al., 2006). It is likely that the water colour in all Irish lake habitats would naturally be <50mg/l PtCo. Water colour can be very low (<20mg/l PtCo or even <10mg/l PtCo) in lake habitat 3110, where the peatland in the lake's catchment is intact
Dissolved organic carbon (DOC)	mg/l	Maintain appropriate organic carbon levels to support the habitat	Dissolved (and particulate) organic carbon (OC) in the water column is linked to water colour and acidification (organic acids). Increasing DOC in water has been documented across the Northern Hemisphere, including afforested peatland catchments in Ireland. Damage and degradation of peatland, leading to decomposition of peat is likely to be the predominant source of OC in Ireland. OC in water promotes decomposition by fungi and bacteria that, in turn, releases dissolved nutrients. The increased biomass of decomposers can also impact directly on the characteristic lake communities through shading, competition, etc.
Turbidity	Nephelometric turbidity units/ mg/l SS/ other appropriate units	Maintain appropriate turbidity to support the habitat	Turbidity can significantly affect the quantity and quality of light reaching rooted and attached vegetation and can, therefore, impact on lake habitats. The settlement of higher loads of inorganic or organic material on lake vegetation communities may also have impacts on sensitive, delicate species. Turbidity can increase as a result of re-suspension of material within the lake, higher loads entering the lake, or eutrophication. Turbidity measurement and interpretation is challenging. As a result, it is likely to be difficult to set habitat-specific targets for turbidity in lakes
Fringing habitat: area	Hectares	Maintain the area and condition of fringing habitats necessary to support the natural structure and functioning of habitat 3110	Most lake shorelines have fringing habitats of reedswamp, other swamp, fen, marsh or wet-woodland that intergrade with and support the structure and functions of the lake habitat. In this SAC, blanket bog and heath communities are likely to dominate shorelines. Poor fen and flush may also occur. Equally, fringing habitats are dependent on the lake, particularly its water levels, and support wetland communities and species of conservation concern. Many of the fringing wetland habitats support higher invertebrate and plant species richness than the lake habitats themselves

3160 Natural dystrophic lakes and ponds

To maintain the favourable conservation condition of Natural dystrophic lakes and ponds in Ox Mountains Bogs SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	This SAC has extensive blanket bog pool systems. Douglas et al. (1989) said the interconnecting pool system in Letterunshin, NW of L. Easky was the largest recorded in the blanket bog survey and of very high scientific and conservation value. Not all of the pools are mapped in the 1:5,000 OSI data. The 201 pools less than 1ha in area have been mapped as potential 3160 (see map 3). As all lakes in the SAC are surrounded by blanket bog and wet heath, 3160 likely also occurs in the larger lakes. The habitat is considered to be of high conservation value in the site. Two measures of extent should be used: 1. the area of the lake itself and; 2. the extent of the vegetation communities/zones that typify the habitat. Further information relating to all attributes is provided in the lake habitats supporting document for the purposes of site-specific conservation objectives and Article 17 reporting (O Connor, 2015)
Habitat distribution	Occurrence	No decline, subject to natural processes	As noted above, the habitat is widespread and of high conservation value in the SAC (see map 3)
Typical species	Occurrence	Typical species present, in good condition, and demonstrating typical abundances and distribution	For lists of typical plant and invertebrate species, see Article 17 habitat assessment for 3160 (NPWS, 2013) and the lake habitats supporting document for the purposes of site-specific conservation objectives and Article 17 reporting (O Connor, 2015)
Vegetation composition: characteristic zonation	Occurrence	All characteristic zones should be present, correctly distributed and in good condition	The characteristic zonation of lake habitat 3140 has been described (Roden and Murphy, 2013; in prep.), however, significant further work is necessary to describe the characteristic zonation and other spatial patterns in the other four Annex I lake habitats. Spatial patterns are likely to be relatively simple in 3160 lakes and ponds, with limited zonation
Vegetation distribution: maximum depth	Metres	Maintain maximum depth of vegetation, subject to natural processes	The maximum depth of vegetation is likely to be specific to the lake shoreline in question. An indicative target has not yet been set for this lake habitat type. Lakes in the SAC typically have very clear water and, therefore, maximum depth is expected to be large
Hydrological regime: water level fluctuations	Metres	Maintain appropriate natural hydrological regime necessary to support the habitat	Fluctuations in lake water level are typical in Ireland, but can be amplified by activities such as abstraction and drainage. Increased water level fluctuations can increase wave action, up-root vegetation, increase turbidity, alter the substratum and lead to release of nutrients from the sediment. The hydrological regime of the lakes and pools must be maintained so that the area, distribution and depth of the lake habitat and its constituent/characteristic vegetation zones and communities are not reduced. Owing to their size and the sensitivity of peatland, 3160 lakes and pools can easily be damaged or destroyed by drainage
Lake substratum quality	Various	Maintain appropriate substratum type, extent and chemistry to support the vegetation	Research is required to further characterise the substratum types (particle size and origin) and substratum quality (notably pH, calcium, iron and nutrient concentrations) favoured by each of the five Annex I lake habitats in Ireland. It is likely that habitat 3160 is associated with nutrient-poor peat and silt substrates

Water quality: transparency	Metres	Maintain appropriate Secchi transparency. There should be no decline in Secchi depth/transparency	Transparency relates to light penetration and, hence, to the depth of colonisation of vegetation. It can be affected by phytoplankton blooms, water colour and turbidity. A specific target has yet to be established for this Annex I lake habitat. Habitat 3160 is associated with very clear water. The OECD fixed boundary system set transparency targets for ultra-oligotrophic lakes of $\geq 12\text{m}$ annual mean Secchi disk depth, and $\geq 6\text{m}$ annual minimum Secchi disk depth
Water quality: nutrients	$\mu\text{g/l P}$; mg/l N	Maintain the concentration of nutrients in the water column to sufficiently low levels to support the habitat and its typical species	As a nutrient-poor habitat, oligotrophic and Water Framework Directive (WFD) 'high' status targets apply. Where a lake has nutrient concentrations that are lower than these targets, there should be no decline within class, i.e. no upward trend in nutrient concentrations. For 3160 lakes and pools, annual average TP concentration should be $\leq 5\mu\text{g/l TP}$, average annual total ammonia concentration should be $\leq 0.040\text{mg/l N}$ and annual 95th percentile for total ammonia should be $\leq 0.090\text{mg/l N}$. See also The European Communities Environmental Objectives (Surface Waters) Regulations 2009
Water quality: phytoplankton biomass	$\mu\text{g/l Chlorophyll } a$	Maintain appropriate water quality to support the habitat, including high chlorophyll <i>a</i> status	Oligotrophic and WFD 'high' status targets apply to lake habitat 3160. The average growing season (March-October) chlorophyll <i>a</i> concentration must be $< 5.8\mu\text{g/l}$ (The European Communities Environmental Objectives (Surface Waters) Regulations 2009). Where a lake has a chlorophyll <i>a</i> concentration that is lower than this target, there should be no decline within class, i.e. no upward trend in phytoplankton biomass. The OECD targets may be more appropriate for habitat 3160: annual average chlorophyll <i>a</i> concentration $< 1\mu\text{g/l}$ and annual peak chlorophyll <i>a</i> concentration $\leq 2.5\mu\text{g/l}$. See also The European Communities Environmental Objectives (Surface Waters) Regulations 2009
Water quality: phytoplankton composition	EPA phytoplankton composition metric	Maintain appropriate water quality to support the habitat, including high phytoplankton composition status	The EPA has developed a phytoplankton composition metric for nutrient enrichment of Irish lakes. As for other water quality indicators, habitat 3160 requires WFD high status
Water quality: attached algal biomass	Algal cover and EPA phytobenthos metric	Maintain trace/ absent attached algal biomass ($< 5\%$ cover) and high phytobenthos status	Nutrient enrichment can favour epiphytic and epipelagic algae that can out-compete the submerged vegetation. The cover abundance of attached algae in 3160 lakes and pools should, therefore, be trace/ absent ($< 5\%$ cover). EPA phytobenthos can be used as an indicator of changes in attached algal biomass. As for other water quality indicators, habitat 3160 requires high phytobenthos status
Water quality: macrophyte status	EPA macrophyte metric (The Free Index)	Maintain high macrophyte status	Nutrient enrichment can favour more competitive submerged macrophyte species that out-compete the typical and characteristic species for the lake habitat. The EPA monitors macrophyte status for WFD purposes using the 'Free Index'. The target for 3160 lakes and pools is high status or an Ecological Quality Ratio (EQR) for lake macrophytes of ≥ 0.90 , as defined in Schedule Five of the European Communities Environmental Objectives (Surface Waters) Regulations 2009

Acidification status	pH units; mg/l	Maintain appropriate water and sediment pH, alkalinity and cation concentrations to support the habitat, subject to natural processes	Acidification can impact on species abundance and composition in soft water lake habitats. Although EC (2013) describes habitat 3160 as having pH 3-6, Drinan (2012) found mean pHs of 5.16 and 5.62 in upland and lowland 3160 lakes, respectively. The target for lake habitat 3160 is pH >4.5 and <9.0, in line with the surface water standards for soft waters (where water hardness is ≤100mg/l calcium carbonate). See Schedule Five of the European Communities Environmental Objectives (Surface Waters) Regulations 2009. The specific requirements of habitat 3160, in terms of water and sediment pH, alkalinity and cation concentration, have not been determined
Water colour	mg/l PtCo	Maintain appropriate water colour to support the habitat	Increased water colour and turbidity decrease light penetration and can reduce the area of available habitat for lake macrophytes, particularly at the lower euphotic depths. The primary source of increased water colour in Ireland is disturbance to peatland. No habitat-specific or national standards for water colour currently exist. Studies have shown median colour concentrations in Irish lakes of 38mg/l PtCo (Free et al., 2000) and 33mg/l PtCo (Free et al., 2006). It is likely that the water colour in all Irish lake habitats would naturally be <50mg/l PtCo. Water colour can be very low (<20mg/l PtCo or even <10mg/l PtCo) in 3160 lakes and pools where the peatland in the catchment is intact
Dissolved organic carbon (DOC)	mg/l	Maintain appropriate organic carbon levels to support the habitat	Dissolved (and particulate) organic carbon (OC) in the water column is linked to water colour and acidification (organic acids). Increasing DOC in water has been documented across the Northern Hemisphere, including afforested peatland catchments in Ireland. Damage and degradation of peatland, leading to decomposition of peat is likely to be the predominant source of OC in Ireland. OC in water promotes decomposition by fungi and bacteria that, in turn, releases dissolved nutrients. The increased biomass of decomposers can also impact directly on the characteristic lake communities through shading, competition, etc.
Turbidity	Nephelometric turbidity units/ mg/l SS/ other appropriate units	Maintain appropriate turbidity to support the habitat	Turbidity can significantly affect the quantity and quality of light reaching rooted and attached vegetation and can, therefore, impact on lake habitats. The settlement of higher loads of inorganic or organic material on lake vegetation communities may also have impacts on sensitive, delicate species. Turbidity can increase as a result of re-suspension of material within the lake, higher loads entering the lake, or eutrophication. Turbidity measurement and interpretation is challenging. As a result, it is likely to be difficult to set habitat-specific targets for turbidity in lakes
Fringing habitat: area	Hectares	Maintain the area and condition of fringing habitats necessary to support the natural structure and functioning of habitat 3160	Most 3160 lake and pool shorelines intergrade with blanket bog, flush, poor-fen or heath habitats and these support the structure and functions of the lake habitat. Equally, fringing habitats are dependent on the lake, particularly its water levels, and can support wetland communities and species of conservation concern

4010 Northern Atlantic wet heaths with *Erica tetralix*

To restore the favourable conservation condition of Northern Atlantic wet heaths with *Erica tetralix* in Ox Mountains Bogs SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area increasing, subject to natural processes	Ox Mountains Bogs SAC was surveyed as part of the National Survey of Upland Habitats (NSUH; see Perrin et al., 2013; 2014). The total current area of wet heath stated by Perrin et al. (2013) is 1083.2ha, covering 10.2% of the SAC. Perrin et al. (2013) report obvious losses of habitat since 1995 of less than 0.01ha through landslides. A summary of the mapping methodology and a brief discussion of restoration potential are presented in the uplands supporting document
Habitat distribution	Occurrence	No decline from current distribution, subject to natural processes. See map 4	Wet heath was recorded by Perrin et al. (2013) throughout the SAC mainly on the lower slopes. Extensive patches occur on the slopes above Cloonacool, and Carrownedeen in the east. It also occurs through Fiddenderry and on the slopes above Easkey Lough. A summary of the mapping methodology is presented in the uplands supporting document
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil nutrient status within natural range	See the uplands supporting document for further details
Community diversity	Abundance of variety of vegetation communities	Maintain variety of vegetation communities, subject to natural processes	Perrin et al. (2013) recorded six different wet heath communities within this SAC. Data on the abundance of these communities is reproduced in the uplands supporting document. Further information on these communities is presented in Perrin et al. (2014)
Vegetation composition: cross-leaved heath	Occurrence within 20m of a representative number of 2m x 2m monitoring stops	Cross-leaved heath (<i>Erica tetralix</i>) present near each monitoring stop	Based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation composition: positive indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of positive indicator species at least 50%	Based on Perrin et al. (2014). The list of positive indicator species for this habitat is presented in Perrin et al. (2014). Further details can be found in the uplands supporting document
Vegetation composition: lichens and bryophytes	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of <i>Cladonia</i> and <i>Sphagnum</i> species, <i>Racomitrium lanuginosum</i> and pleurocarpous mosses at least 10%	Based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation composition: ericoid species and crowberry	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of ericoid species and crowberry (<i>Empetrum nigrum</i>) at least 15%	Based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation composition: dwarf shrub species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of dwarf shrubs less than 75%	Based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation composition: negative indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of negative indicator species less than 1%	Based on Perrin et al. (2014). The list of negative indicator species is given in Perrin et al. (2014). See the uplands supporting document for further details
Vegetation composition: non-native species	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of non-native species less than 1%	Based on Perrin et al. (2014). See the uplands supporting document for further details. <i>Campylopus introflexus</i> was recorded within this habitat by Perrin et al. (2013) at two monitoring stops with extensive carpets at one of these. Scattered non-native conifers were also recorded

Vegetation composition: native trees and shrubs	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of scattered native trees and shrubs less than 20%	Based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation composition: bracken	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of bracken (<i>Pteridium aquilinum</i>) less than 10%	Based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation composition: soft rush	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of soft rush (<i>Juncus effusus</i>) less than 10%	Based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation structure: <i>Sphagnum</i> condition	Condition at a representative number of 2m x 2m monitoring stops	Less than 10% of the <i>Sphagnum</i> cover is crushed, broken and/or pulled up	Based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation structure: signs of browsing	Percentage of shoots browsed at a representative number of 2m x 2m monitoring stops	Less than 33% collectively of the last complete growing season's shoots of ericoids, crowberry (<i>Empetrum nigrum</i>) and bog-myrtle (<i>Myrica gale</i>) showing signs of browsing	Based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation structure: burning	Occurrence in local vicinity of a representative number of 2m x 2m monitoring stops	No signs of burning in sensitive areas, into the moss, liverwort or lichen layer or exposure of peat surface due to burning	Based on Perrin et al. (2014). The list of sensitive areas is presented in Perrin et al. (2014). See the uplands supporting document for further details
Physical structure: disturbed bare ground	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of disturbed bare ground less than 10%	Based on Perrin et al. (2014). See the uplands supporting document for further details
Physical structure: drainage	Percentage cover in local vicinity of a representative number of monitoring stops	Area showing signs of drainage from heavy trampling, tracking or ditches less than 10%	Based on Perrin et al. (2014). See the uplands supporting document for further details
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat	Perrin et al. (2013) compiled and mapped existing rare and notable plant records for the SAC and added any new records collected during the NSUH survey. No relevant species were recorded in this habitat, however, new records should be considered within this attribute. See the uplands supporting document for further details

Conservation Objectives for : Ox Mountains Bogs SAC [002006]

4030 European dry heaths

To maintain the favourable conservation condition of European dry heaths in Ox Mountains Bogs SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Ox Mountains Bogs SAC was surveyed as part of the National Survey of Upland Habitats (NSUH; see Perrin et al., 2013; 2014). The total current area of dry heath stated by Perrin et al. (2013) is 332.9ha, covering 3.1% of the SAC. It occurs at low frequency throughout the SAC, but is locally abundant on the rocky slopes above Easky Lough and above Cloonacool. Perrin et al. (2013) report no significant losses of area since 1995. A summary of the mapping methodology is presented in the uplands supporting document
Habitat distribution	Occurrence	No decline from current distribution, subject to natural processes. See map 5	Dry heath was recorded by Perrin et al. (2013) throughout the SAC, but was most abundant on the eastern slopes above Easky Lough. A summary of the mapping methodology is presented in the uplands supporting document
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil nutrient status within natural range	See the uplands supporting document for further details
Community diversity	Abundance of variety of vegetation communities	Maintain variety of vegetation communities, subject to natural processes	Perrin et al. (2013) recorded three different dry heath communities within this SAC. Data on the abundance of these communities is reproduced in the uplands supporting document. Further information on these communities is presented in Perrin et al. (2014)
Vegetation composition: lichens and bryophytes	Number of species at a representative number of 2m x 2m monitoring stops	Number of bryophyte or non-crustose lichen species present at each monitoring stop is at least three, excluding <i>Campylopus</i> and <i>Polytrichum</i> mosses	Based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation composition: number of positive indicator species	Number of species at a representative number of 2m x 2m monitoring stops	Number of positive indicator species present at each monitoring stop is at least two	Based on Perrin et al. (2014). The list of positive indicator species for this habitat, which is composed of dwarf shrubs, is presented in Perrin et al. (2014). See the uplands supporting document for further details
Vegetation composition: cover of positive indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of positive indicator species at least 50% for siliceous dry heath and 50-75% for calcareous dry heath	Based on Perrin et al. (2014). The list of positive indicator species for this habitat, which is composed of dwarf shrubs, is presented in Perrin et al. (2014). See the uplands supporting document for further details
Vegetation composition: dwarf shrub composition	Percentage cover at a representative number of 2m x 2m monitoring stops	Proportion of dwarf shrub cover composed collectively of bog-myrtle (<i>Myrica gale</i>), creeping willow (<i>Salix repens</i>) and western gorse (<i>Ulex gallii</i>) is less than 50%	Based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation composition: negative indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of negative indicator species less than 1%	Based on Perrin et al. (2014). The list of negative indicator species is given in Perrin et al. (2014). See the uplands supporting document for further details
Vegetation composition: non-native species	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of non-native species less than 1%	Based on Perrin et al. (2014). See the uplands supporting document for further details. Scattered non-native conifers were observed within the habitat but this was limited to a few individuals

Vegetation composition: native trees and shrubs	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of scattered native trees and shrubs less than 20%	Based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation composition: bracken	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of bracken (<i>Pteridium aquilinum</i>) less than 10%	Based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation composition: soft rush	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of soft rush (<i>Juncus effusus</i>) less than 10%	Based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation structure: senescent ling	Percentage cover at a representative number of 2m x 2m monitoring stops	Senescent proportion of ling (<i>Calluna vulgaris</i>) cover less than 50%	Based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation structure: signs of browsing	Percentage of shoots browsed at a representative number of 2m x 2m monitoring stops	Less than 33% collectively of the last complete growing season's shoots of ericoids showing signs of browsing	Based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation structure: burning	Occurrence in local vicinity of a representative number of monitoring stops	No signs of burning in sensitive areas	Based on Perrin et al. (2014). The list of sensitive areas is presented in Perrin et al. (2014). See the uplands supporting document for further details
Vegetation structure: growth phases of ling	Percentage cover in local vicinity of a representative number of monitoring stops	Outside sensitive areas, all growth phases of ling (<i>Calluna vulgaris</i>) should occur throughout, with at least 10% of cover in the mature phase	Based on Perrin et al. (2014). See the uplands supporting document for further details
Physical structure: disturbed bare ground	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of disturbed bare ground less than 10%	Based on Perrin et al. (2014). See the uplands supporting document for further details
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat	Perrin et al. (2013) compiled and mapped existing rare and notable plant records for the SAC and added any new records collected during the NSUH survey. No relevant species were recorded in this habitat, however, new records should be considered within this attribute. See the uplands supporting document for further details

Conservation Objectives for : Ox Mountains Bogs SAC [002006]

7130 Blanket bogs (* if active bog)

To restore the favourable conservation condition of Blanket bogs in Ox Mountains Bogs SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area increasing, subject to natural processes	Ox Mountains Bogs SAC was surveyed as part of the National Survey of Upland Habitats (NSUH; see Perrin et al., 2013; 2014). Perrin et al. (2013) state that the current total area of blanket bog is 7249.6ha (68.5% of the SAC). This comprises 7097.3ha of active blanket bog area and 152.3ha of inactive blanket bog. Perrin et al. (2014) also report obvious losses of habitat since 1995 of approximately 5.8ha. However, this is almost certainly an under-estimate, as chronic losses due to erosion since 1995 cannot be quantified (106.6ha were mapped as eroding blanket bog by Perrin et al. (2013)). It should be noted that further restoration of blanket bog would be required in order to fulfil the targets for peat formation and hydrology presented below. A summary of the mapping methodology and a brief discussion of restoration potential are presented in the uplands supporting document
Habitat distribution	Occurrence	No decline from current distribution, subject to natural processes. See map 6	Blanket bog was recorded by Perrin et al. (2013) across the SAC and was by far the most dominant habitat type. A summary of the mapping methodology is presented in the uplands supporting document
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil nutrient status within natural range	See the uplands supporting document for further details
Ecosystem function: peat formation	Active blanket bog as a proportion of the total area of Annex I blanket bog	At least 99% of the total Annex I blanket bog area is active	From the habitat areas given by Perrin et al. (2013) above, 97.9% of the Annex I blanket bog habitat is currently actively peat-forming. See the uplands supporting document for further details
Ecosystem function: hydrology	Flow direction, water levels, occurrence of drains and erosion gullies	Natural hydrology unaffected by drains and erosion	Further details and a brief discussion of restoration potential is presented in the uplands supporting document
Community diversity	Abundance of variety of vegetation communities	Maintain variety of vegetation communities, subject to natural processes	Perrin et al. (2013) recorded six different active blanket bog communities within this SAC. Data on the abundance of these communities is reproduced in the uplands supporting document. Further information on these communities is presented in Perrin et al. (2014)
Vegetation composition: positive indicator species	Number of species at a representative number of 2m x 2m monitoring stops	Number of positive indicator species at each monitoring stop is at least seven	Based on Perrin et al. (2014). The list of positive indicator species for this habitat is presented in Perrin et al. (2014). See the uplands supporting document for further details
Vegetation composition: lichens and bryophytes	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of bryophytes or lichens, excluding <i>Sphagnum fallax</i> , at least 10%	Based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation composition: potential dominant species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of each of the potential dominant species less than 75%	Based on Perrin et al. (2014). See the uplands supporting document for further details, including the list of potentially dominant species
Vegetation composition: negative indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of negative indicator species less than 1%	Based on Perrin et al. (2014). The list of negative indicator species is given in Perrin et al. (2014). See the uplands supporting document for further details

Vegetation composition: non-native species	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of non-native species less than 1%	Based on Perrin et al. (2014). See the uplands supporting document for further details. <i>Campylopus introflexus</i> was recorded within this habitat by Perrin et al. (2013) with extensive carpets recorded
Vegetation composition: native trees and scrub	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of scattered native trees and shrubs less than 10%	Based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation structure: <i>Sphagnum</i> condition	Condition at a representative number of 2m x 2m monitoring stops	Less than 10% of the <i>Sphagnum</i> cover is crushed, broken and/or pulled up	Based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation structure: signs of browsing	Percentage of shoots browsed at a representative number of 2m x 2m monitoring stops	Last complete growing season's shoots of ericoids, crowberry (<i>Empetrum nigrum</i>) and bog-myrtle (<i>Myrica gale</i>) showing signs of browsing collectively less than 33%	Based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation structure: burning	Occurrence in local vicinity of a representative number of monitoring stops	No signs of burning in sensitive areas, into the moss, liverwort or lichen layer or exposure of peat surface due to burning	Based on Perrin et al. (2014). The list of sensitive areas is presented in Perrin et al. (2014). See the uplands supporting document for further details
Physical structure: disturbed bare ground	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of disturbed bare ground less than 10%	Based on Perrin et al. (2014). See the uplands supporting document for further details
Physical structure: drainage	Occurrence in local vicinity of a representative number of monitoring stops	Area showing signs of drainage from heavy trampling, tracking or ditches less than 10%	Based on Perrin et al. (2014). See the uplands supporting document for further details
Physical structure: erosion	Occurrence in local vicinity of a representative number of monitoring stops	Less than 5% of the greater bog mosaic comprises erosion gullies and eroded areas	Based on Perrin et al. (2014). See the uplands supporting document for further details
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat	Perrin et al. (2013) compiled and mapped existing rare and notable plant records for the SAC and added any new records collected during the NSUH survey. No relevant species were recorded in this habitat, however, new records should be considered within this attribute. See the uplands supporting document for further details

Conservation Objectives for : Ox Mountains Bogs SAC [002006]

7140 Transition mires and quaking bogs

To maintain the favourable conservation condition of Transition mires and quaking bogs in Ox Mountains Bogs SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Ox Mountains Bogs SAC was surveyed as part of the National Survey of Upland Habitats (NSUH; see Perrin et al., 2013; 2014). The total current area of transition mires and quaking bogs in the SAC stated by Perrin et al. (2013) is 36.6ha. This covers 0.3% of the SAC. Perrin et al. (2013) report no significant losses of area since 1995. A summary of the mapping methodology is presented in the uplands supporting document
Habitat distribution	Occurrence	No decline from current distribution, subject to natural processes. See map 7	Transition mire was recorded by Perrin et al. (2013) scattered throughout the SAC. It is frequent in the small valleys of the upland plateau, along the eastern fringes of the SAC, through Letterunshion Bog and the southern section of Fiddenderry. A summary of the mapping methodology is presented in the uplands supporting document
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil nutrient status within natural range	See the uplands supporting document for further details
Community diversity	Abundance of variety of vegetation communities	Maintain variety of vegetation communities, subject to natural processes	Perrin et al. (2013) recorded three different transition mire communities within this SAC. Data on the abundance of these communities is reproduced in the uplands supporting document. Further information on these vegetation communities is presented in Perrin et al. (2014)
Vegetation composition: number of positive indicator species	Number at a representative number of 2m x 2m monitoring stops	Number of positive indicator species at least three for in-filling pools and flushes and at least six for fens	Based on Perrin et al. (2014), where the list of positive indicator species for this habitat is also presented. See the uplands supporting document for further details
Vegetation composition: number of core positive indicator species	Number of species at a representative number of 2m x 2m monitoring stops	At least one core positive indicator species present	Based on Perrin et al. (2014), where the list of positive indicator species for this habitat is also presented. See the uplands supporting document for further details
Vegetation composition: cover of positive indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of positive indicator species is at least 25%	Based on Perrin et al. (2014), where the list of positive indicator species for this habitat is also presented. See the uplands supporting document for further details
Vegetation composition: negative indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of negative indicator species less than 1%	Based on Perrin et al. (2014), where the list of negative indicator species for this habitat is also presented. See the uplands supporting document for further details
Vegetation composition: non-native species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of non-native species less than 1%	Based on Perrin et al. (2014). See the uplands supporting document for further details. No non-native species were recorded within this habitat by Perrin et al. (2013)
Vegetation structure: height	Percentage at a representative number of 2m x 2m monitoring stops	Proportion of live leaves and/or flowering shoots of vascular plants that are more than 15cm above the ground surface should be at least 50%	Based on Perrin et al. (2014). This attribute is only applicable to fen and flush examples, not to in-filling pool examples. See the uplands supporting document for further details
Physical structure: disturbed bare ground	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of disturbed bare ground less than 10%	Based on Perrin et al. (2014). See the uplands supporting document for further details

Physical structure: drainage	Percentage area in local vicinity of a representative number of monitoring stops	Area showing signs of drainage from heavy trampling, tracking or ditches less than 10%	Based on Perrin et al. (2014). See the uplands supporting document for further details
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat	Perrin et al. (2013) compiled and mapped existing rare and notable plant records for the SAC and added any new records collected during the NSUH survey. No relevant species were recorded in this habitat, however, new records should be considered within this attribute. See the uplands supporting document for further details

Conservation Objectives for : Ox Mountains Bogs SAC [002006]

7150 Depressions on peat substrates of the Rhynchosporion

To maintain the favourable conservation condition of Depressions on peat substrates of the Rhynchosporion in Ox Mountains Bogs SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Ox Mountains Bogs SAC was surveyed as part of the National Survey of Upland Habitats (NSUH; see Perrin et al., 2013; 2014). The total current area of Depressions on peat surfaces of the Rhynchosporion in the SAC stated by Perrin et al. (2013) is 49.6ha. This covers 0.5% of the SAC. Perrin et al. (2013) report no significant losses of area since 1995. A summary of the mapping methodology is presented in the uplands supporting document
Habitat distribution	Occurrence	No decline from current distribution, subject to natural processes. See map 8	Rhynchosporion depressions were recorded by Perrin et al. (2013) scattered through the western portion of the SAC, particularly through Letterunshion Bog and at Tawnamore. A summary of the mapping methodology is presented in the uplands supporting document
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil nutrient status within natural range	See the uplands supporting document for further details
Vegetation composition: positive indicator species	Number of species at a representative number of 2m x 2m monitoring stops	Number of positive indicator species at each monitoring stop is at least five	Based on Perrin et al. (2014), where the list of positive indicator species for this habitat is also presented. Further details can be found in the uplands supporting document
Vegetation composition: <i>Rhynchospora</i> spp.	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of white beaked sedge (<i>Rhynchospora alba</i>) and brown beaked sedge (<i>R. fusca</i>) at least 10%	Based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation composition: potential dominant species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of each of the potential dominant species less than 35%	Based on Perrin et al. (2014). See the uplands supporting document for further details, including the list of potentially dominant species
Vegetation composition: negative indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of negative indicator species less than 1%	Based on Perrin et al. (2014), where the list of negative indicator species is also presented. See the uplands supporting document for further details
Vegetation composition: non-native species	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of non-native species less than 1%	Based on Perrin et al. (2014). See the uplands supporting document for further details. <i>Campylopus introflexus</i> was recorded within this habitat by Perrin et al. (2013) but did not form extensive carpets
Vegetation composition: native trees and scrub	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of scattered native trees and shrubs less than 10%	Based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation structure: <i>Sphagnum</i> condition	Condition at a representative number of 2m x 2m monitoring stops	Less than 10% of the <i>Sphagnum</i> cover is crushed, broken and/or pulled up	Based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation structure: signs of browsing	Percentage of shoots browsed at a representative number of 2m x 2m monitoring stops	Last complete growing season's shoots of ericoids, crowberry (<i>Empetrum nigrum</i>) and bog-myrtle (<i>Myrica gale</i>) showing signs of browsing collectively less than 33%	Based on Perrin et al. (2014). See the uplands supporting document for further details

Vegetation structure: burning	Occurrence in local vicinity of a representative number of monitoring stops	No signs of burning in sensitive areas, into the moss, liverwort or lichen layer or exposure of peat surface due to burning	Based on Perrin et al. (2014), where the list of sensitive areas is also presented. See the uplands supporting document for further details
Physical structure: disturbed bare ground	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of disturbed bare ground less than 10%	Based on Perrin et al. (2014). See the uplands supporting document for further details
Physical structure: drainage	Percentage area in local vicinity of a representative number of monitoring stops	Area showing signs of drainage from heavy trampling, tracking or ditches less than 10%	Based on Perrin et al. (2014). See the uplands supporting document for further details
Physical structure: erosion	Occurrence in local vicinity of a representative number of monitoring stops	Less than 5% of the greater bog mosaic comprises erosion gullies and eroded areas	Based on Perrin et al. (2014). See the uplands supporting document for further details
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat	Perrin et al. (2013) compiled and mapped existing rare and notable plant records for the SAC and added any new records collected during the NSUH survey. No relevant species were recorded in this habitat, however, new records should be considered within this attribute. See the uplands supporting document for further details

Conservation Objectives for : Ox Mountains Bogs SAC [002006]

1013 Geyer's Whorl Snail *Vertigo geyeri*


To maintain the favourable conservation condition of Geyer's Whorl Snail in Ox Mountains Bogs SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Distribution: occupied sites	Number	No decline. There is one known site for this species in this SAC within the 1km square G4429. See map 9	From Moorkens and Killeen (2011) (site code VgCAM21)
Presence in suitable habitat	Occurrence	Snails (living or recently dead adults and/or juveniles) are present in at least 60% of samples defined as suitable habitat	Based on Moorkens and Killeen (2011)
Species abundance	Number of individuals per sample	No decline in adult abundance in appropriate number of samples	Based on Moorkens and Killeen (2011). There should be at least five adults detected in 40% of samples
Habitat area	Hectares	Stable or increasing, subject to natural processes. Suitable habitat is defined as areas of flushed fen with small sedges and saturated mosses	Based on Moorkens and Killeen (2011). Optimal habitat is defined as flushed fen with sedge/moss lawns and mounds 5-20cm tall, containing a high diversity of plant species such as small-fruited yellow-sedge (<i>Carex viridula</i>), grass-of-Parnassus (<i>Parnassia palustris</i>), marsh horsetail (<i>Equisetum palustre</i>), jointed rush (<i>Juncus articulatus</i>) and the mosses <i>Scorpidium revolvens</i> and <i>Campyllum stellatum</i> , with scattered tussocks of black bog-rush (<i>Schoenus nigricans</i>) no greater than 80cm tall
Habitat quality: optimal habitat	Hectares	At least 0.3ha of optimal habitat present	Based on Moorkens and Killeen (2011). There should be at least 0.3ha of optimal habitat present at the site to maintain the species. See description of optimal habitat above
Habitat quality: soil wetness	Water table level	Water table should be between 0-5cm of the soil surface, but not above ground level at time of sampling	Based on Moorkens and Killeen (2011)


1528 Marsh Saxifrage *Saxifraga hirculus*


To maintain the favourable conservation condition of *Saxifraga hirculus* in Ox Mountains Bogs SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Distribution	Number and geographical spread	No loss in geographical spread and number of populations, subject to natural processes	Three populations of <i>Saxifraga hirculus</i> , in three flushes (A-C) lying in close proximity to each other, were discovered in the Ox Mountains Bogs SAC in June 2012. See Muldoon et al. (2014) for further details
Population size: number of rosettes	Number	Maintain the size of each population, subject to natural processes. The target numbers of rosettes are: >40,000 rosettes in Flush A; >4,800 rosettes in Flush B; >480 rosettes in Flush C	The number of rosettes was estimated to be: c.50,000 in Flush A; c.6,000 in Flush B and c.600 in Flush C. The target figures are a 20% reduction of the recorded number to allow for a margin of error and variability over monitoring seasons
Population size: area of occupancy	Hectares	Maintain the extent of each population, subject to natural processes. The target areas are: > 0.0234 ha (> 234 m ²) in Flush A, > 0.0053 ha (> 52.5 m ²) in Flush B and > 0.0016 ha (> 16.2 m ²) in Flush C	The area of cover of <i>Saxifraga hirculus</i> was estimated as 260 m ² in Flush A, 58.3 m ² in Flush B and 18 m ² in Flush C. The target area figures are a 10% reduction of the recorded areas to allow for a margin of error
Hydrological conditions: water levels	Occurrence of high or fluctuating water levels	Maintain appropriate natural hydrological regime necessary to support the habitat for the species	In Ireland, <i>Saxifraga hirculus</i> is now restricted to mineral flushes in blanket bog where rising groundwater forms small streams and seepage areas suitable for the species. Based on Muldoon (2011) and Muldoon et al. (2015)
Vegetation structure: sward height	Centimetres	Maintain a mean vegetation height of less than 15cm	See Muldoon (2011) and Muldoon et al. (2015) for further details
Vegetation composition: associated species	Species composition and abundance	Maintain appropriate associated species and vegetation communities to support the populations of <i>Saxifraga hirculus</i>	Presence of knotted pearlwort (<i>Sagina nodosa</i>), a positive indicator species and low cover of purple moor-grass (<i>Molinia caerulea</i>) and Yorkshire-fog (<i>Holcus lanatus</i>), both negative indicator species, should be maintained. See Muldoon (2011) and Muldoon et al. (2015) for further details
Vegetation structure: grazing levels	Evidence of grazing	Maintain grazing at light to moderate levels to ensure an open vegetation structure and to allow flowering to occur	See Muldoon (2011) and Muldoon et al. (2015) for further details




Legend

 Ox Mountains Bogs SAC 002006



Legend

- Ox Mountains Bogs SAC 002006
- River Moy SAC 002298

 An Roinn Ealaíon, Oidhreachta, Gnóthaí Réigiúnacha, Tuisithe agus Gaeltachta
 Department of Arts, Heritage, Regional, Rural and Gaeltacht Affairs

MAP 2:
OX MOUNTAINS BOGS SAC
CONSERVATION OBJECTIVES
ADJACENT DESIGNATIONS

Map to be read in conjunction with the NPWS Conservation Objectives Document.


SITE CODE:
SAC 002006; version 3.01,
SAC 002298; version 3.01.
Counties Sligo / Mayo / Roscommon

0 4.5 9 13.5 km

The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision. Ordnance Survey of Ireland Licence No EN 0059216. © Ordnance Survey of Ireland Government of Ireland.

Nil sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbheithnithe a déanamh ar theorainneacha na gceantar comharthaithe. Suirbhéarachta Ordoináis na hÉireann Ceadúnas Uimh EN 0059216. © Suirbhéarachta Ordoináis na hÉireann Rialtas na hÉireann.


Map Version 1
Date: July 2016




Legend

- Ox Mountains Bogs SAC 002006
- OSi Discovery Series County Boundaries

Lake Habitats

- 3110 Oligotrophic waters containing very few minerals of sandy plains (*Littorelletalia unifloraræ*)
- Potential 3110 / Potential 3160, Potential Oligotrophic waters containing very few minerals of sandy plains (*Littorelletalia unifloraræ*) / Potential Natural dystrophic lakes and ponds
- Potential 3160 Natural dystrophic lakes and ponds




Legend

- Ox Mountains Bogs SAC 002006
- OSi Discovery Series County Boundaries

Cover of 4010 Northern Atlantic wet heaths with *Erica tetralix*

- 0%
- 0.1% - 20%
- 20.1% - 40%
- 40.1% - 60%
- 60.1% - 80%
- 80.1% - 100%



Legend

- Ox Mountains Bogs SAC 002006
- OSi Discovery Series County Boundaries


Cover of 4030 European dry heaths

- 0%
- 0.1% - 20%
- 20.1% - 40%
- 40.1% - 60%
- 60.1% - 80%
- 80.1% - 100%


**MAP 5:
OX MOUNTAINS BOGS SAC
CONSERVATION OBJECTIVES
DRY HEATHS**

Map to be read in conjunction with the NPWS Conservation Objectives Document.

**SITE CODE:
SAC 002006; version 3.01. Counties Sligo / Mayo**



The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
Ordnance Survey of Ireland Licence No EN 0059216. © Ordnance Survey of Ireland Government of Ireland.
Níl sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbheithnithe a déanamh ar theorainneacha na gceantar comharthaithe. Suirbhéarachta Ordonáis na hÉireann Ceadúnas Uimh EN 0059216. © Suirbhéarachta Ordonáis na hÉireann Rialtas na hÉireann.




Legend

- Ox Mountains Bogs SAC 002006
- OSi Discovery Series County Boundaries

Cover of 7130 Blanket bogs (* if active bog)

- 0%
- 0.1% - 20%
- 20.1% - 40%
- 40.1% - 60%
- 60.1% - 80%
- 80.1% - 100%




Legend

- Ox Mountains Bogs SAC 002006
- OSi Discovery Series County Boundaries

Cover of 7140 Transition mires and quaking bogs

- 0%
- 0.1% - 20%
- 20.1% - 40%
- 40.1% - 60%
- 60.1% - 80%
- 80.1% - 100%




Legend

- Ox Mountains Bogs SAC 002006
- OSi Discovery Series County Boundaries

Cover of 7150 Depressions on peat substrates of the Rhynchosporion

- 0%
- 0.1% - 20%
- 20.1% - 40%
- 40.1% - 60%
- 60.1% - 80%
- 80.1% - 100%



National Parks and Wildlife Service

Conservation Objectives Series

Maumturk Mountains SAC 002008



An Roinn Ealaíon, Oidhreachta,
Gnóthaí Réigiúnacha, Tuaithe agus Gaeltachta

Department of Arts, Heritage,
Regional, Rural and Gaeltacht Affairs



**National Parks and Wildlife Service,
Department of Arts, Heritage, Regional, Rural and Gaeltacht Affairs,
7 Ely Place, Dublin 2, Ireland.
Web: www.npws.ie
E-mail: nature.conservation@ahg.gov.ie**

Citation:

**NPWS (2017) Conservation Objectives: Maumturk Mountains SAC 002008.
Version 1. National Parks and Wildlife Service, Department of Arts, Heritage,
Regional, Rural and Gaeltacht Affairs.**

**Series Editor: Rebecca Jeffrey
ISSN 2009-4086**

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

* indicates a priority habitat under the Habitats Directive

002008	Maumturk Mountains SAC
1106	Salmon <i>Salmo salar</i>
1833	Slender Naiad <i>Najas flexilis</i>
3110	Oligotrophic waters containing very few minerals of sandy plains (<i>Littorelletalia uniflorae</i>)
4010	Northern Atlantic wet heaths with <i>Cladonia</i> <i>Cladonia</i>
4060	Alpine and Boreal heaths
7130	Blanket bogs (* if active bog)
7150	Depressions on peat substrates of the Rhynchosporion
8220	Siliceous rocky slopes with chasmophytic vegetation

Please note that this SAC is adjacent to Lough Corrib SAC (000297), The Twelve Bens/Garraun Complex SAC (002031), Connemara Bog Complex SAC (002034), Lough Corrib SPA (004042) and Connemara Bog Complex SPA (004181). See map 2. The conservation objectives for this site should be used in conjunction with those for the adjacent sites as appropriate.

Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

NPWS Documents

Year :	1989
Title :	A survey to locate blanket bogs in Co. Galway. Part 2
Author :	Douglas, C.; Garvey, L.; Kelly, L.; O'Sullivan, A.
Series :	Unpublished report to the Wildlife Service
Year :	2012
Title :	Ireland Red List No. 8: Bryophytes
Author :	Lockhart, N.; Hodgetts, N.; Holyoak, D.
Series :	Ireland Red List series, NPWS
Year :	2013
Title :	The status of EU protected habitats and species in Ireland. Volume 2. Habitats assessments
Author :	NPWS
Series :	Conservation assessments
Year :	2014
Title :	Guidelines for a national survey and conservation assessment of upland vegetation and habitats in Ireland, Version 2.0
Author :	Perrin, P.M.; Barron, S.J.; Roche, J.R.; O'Hanrahan, B.
Series :	Irish Wildlife Manual No. 79
Year :	2015
Title :	Habitats Directive Annex I lake habitats: a working interpretation for the purposes of site-specific conservation objectives and Article 17 reporting
Author :	O Connor, Á.
Series :	Unpublished document by NPWS
Year :	2016
Title :	Ireland Red List No. 10: Vascular Plants
Author :	Wyse Jackson, M.; FitzPatrick, Ú.; Cole, E.; Jebb, M.; McFerran, D.; Sheehy Skeffington, M.; Wright, M.
Series :	Ireland Red Lists series, NPWS
Year :	2017
Title :	Maumturk Mountains SAC (site code: 2008) Conservation objectives supporting document-blanket bogs and associated habitats V1
Author :	NPWS
Series :	Conservation objectives supporting document
Year :	2017
Title :	Maumturk Mountains SAC (site code: 2008) Conservation objectives supporting document- <i>Najas flexilis</i> V1
Author :	NPWS
Series :	Conservation objectives supporting document

Other References

Year :	1982
Title :	Eutrophication of waters. Monitoring assessment and control
Author :	OECD
Series :	OECD, Paris

Year :	1983
Title :	Flora of Connemara and the Burren
Author :	Webb, D.A.; Scannell, M.J.P.
Series :	Royal Dublin Society, Dublin and Cambridge University Press, Cambridge
Year :	1986
Title :	A survey of the flora of some mountain ranges in the west of Ireland
Author :	Roden, C.M.
Series :	The Irish Naturalists' Journal, 22: 52-59
Year :	2000
Title :	Colour in Irish lakes
Author :	Free, G.; Allott, N.; Mills, P.; Kennelly, C.; Day, S.
Series :	Verhandlungen Internationale Vereinigung für theoretische und angewandte Limnologie, 27: 2620-2623
Year :	2002
Title :	Deterioration of Atlantic soft water macrophyte communities by acidification, eutrophication and algalisation
Author :	Arts, G.H.P.
Series :	Aquatic Botany, 73: 373-393
Year :	2006
Title :	A reference-based typology and ecological assessment system for Irish lakes. Preliminary investigations. Final report. Project 2000-FS-1-M1 Ecological assessment of lakes pilot study to establish monitoring methodologies EU (WFD)
Author :	Free, G.; Little, R.; Tierney, D.; Donnelly, K.; Coroni, R.
Series :	EPA, Wexford
Year :	2008
Title :	Water Quality in Ireland 2004-2006
Author :	Clabby, K.J.; Bradley, C.; Craig, M.; Daly, D.; Lucey, J.; McGarrigle, M.; O'Boyle, S.; Tierney, D.; Bowman, J.
Series :	EPA, Wexford
Year :	2009
Title :	The identification, characterization and conservation value of isoetid lakes in Ireland
Author :	Free, G.; Bowman, J.; McGarrigle, M.; Little, R.; Coroni, R.; Donnelly, K.; Tierney, D.; Trodd, W.
Series :	Aquatic Conservation: Marine and Freshwater Ecosystems, 19(3): 264–273
Year :	2010
Title :	Water quality in Ireland 2007-2009
Author :	McGarrigle, M.; Lucey, J.; Ó Cinnéide, M.
Series :	EPA, Wexford
Year :	2013
Title :	A report on the aquatic macrophytes in 12 lakes along the Maam Cross to Ballynahinch Road
Author :	Roden, C.
Series :	Unpublished report to RPS Group
Year :	2015
Title :	Water quality in Ireland 2010-2012
Author :	Bradley, C.; Byrne, C.; Craig, M.; Free, G.; Gallagher, T.; Kennedy, B.; Little, R.; Lucey, J.; Mannix, A.; McCreesh, P.; McDermott, G.; McGarrigle, M.; Ní Longphuirt, S.; O'Boyle, S.; Plant, C.; Tierney, D.; Trodd, W.; Webster, P.; Wilkes, R.; Wynne, C.
Series :	EPA, Wexford

Year : 2016
Title : The Status of Irish Salmon Stocks in 2015 with Precautionary Catch Advice for 2016
Author : SSCS (Standing Scientific Committee on Salmon)
Series : Independent Scientific Report to Inland Fisheries Ireland

Spatial data sources

Year : 2008
Title : OSi 1:5000 IG vector dataset
GIS Operations : WaterPolygons feature class clipped to the SAC boundary. Expert opinion used to identify Annex I habitat and to resolve any issues arising
Used For : 3110 (map 3)

Year : 2013
Title : Najas flexilis data
GIS Operations : Lake habitat for species clipped to SAC boundary
Used For : 1833 (map 4)

Conservation Objectives for : Maumturk Mountains SAC [002008]

3110 Oligotrophic waters containing very few minerals of sandy plains (Littorelletalia uniflorae)

To maintain the favourable conservation condition of Oligotrophic waters containing very few minerals of sandy plains (Littorelletalia uniflorae) in Maumturk Mountains SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Lake habitat 3110 is considered likely to occur in Loughs Shindilla, Loughanillaun, Maumwee and others in Maumturk Mountains SAC. In line with Article 17 reporting (NPWS, 2013), all lakes larger than 1ha were mapped as potential 3110 (see map 3). Lake habitat 3130 may also occur in lakes on marble formations, such as Loughs Lehanagh and Derryneen, both of which have slender naiad (<i>Najas flexilis</i>), a characteristic species of lake habitat 3130. Two measures of extent should be used: 1. the area of the lake itself and; 2. the extent of the vegetation communities/zones that typify the habitat. Further information relating to all attributes is provided in the lake habitats supporting document for the purposes of site-specific conservation objectives and Article 17 reporting (O Connor, 2015)
Habitat distribution	Occurrence	No decline, subject to natural processes	As noted above, habitat 3110 is likely to be widespread in lakes in the SAC, and all lakes larger than 1ha have been mapped as potential 3110 (see map 3)
Typical species	Occurrence	Typical species present, in good condition, and demonstrating typical abundances and distribution	For lists of typical plant species, see the Article 17 habitat assessment for 3110 (NPWS, 2013) and the lake habitats supporting document (O Connor, 2015). Environmental Protection Agency (EPA) biologists have recorded waterwort (<i>Elatine hexandra</i>), pipewort (<i>Eriocaulon aquaticum</i>), quillwort (<i>Isoetes lacustris</i>), bulbous rush (<i>Juncus bulbosus</i>), shoreweed (<i>Littorella uniflora</i>), water lobelia (<i>Lobelia dortmanna</i>), alternate water-milfoil (<i>Myriophyllum alterniflorum</i>), broad-leaved pondweed (<i>Potamogeton natans</i>) and bladderwort (<i>Utricularia</i> sp.) in Loughs Loughanillaun, Maumwee and Shindilla. See the Maumturk Mountains SAC conservation objectives supporting document for slender naiad (<i>Najas flexilis</i>) (1833) for information on Lehanagh and Derryneen Loughs. See also Webb and Scannell (1983)
Vegetation composition: characteristic zonation	Occurrence	All characteristic zones should be present, correctly distributed and in good condition	Further work is necessary to describe the characteristic zonation and other spatial patterns in lake habitat 3110 (see O Connor, 2015)
Vegetation distribution: maximum depth	Metres	Maintain maximum depth of vegetation, subject to natural processes	The maximum depth of vegetation is likely to be specific to the lake shoreline in question. Further work is necessary to develop indicative targets for lake habitat 3110. See the slender naiad (<i>Najas flexilis</i>) supporting document for information on Lehanagh and Derryneen Loughs. Loughanillaun ('Anillaun'), Maumwee and Shindilla Loughs are Water Framework Directive (WFD) monitoring lakes and information on vegetation depth may be available from the EPA
Hydrological regime: water level fluctuations	Metres	Maintain appropriate natural hydrological regime necessary to support the habitat	Fluctuations in lake water level are typical in Ireland, but can be amplified by activities such as abstraction and drainage. Increased water level fluctuations can increase wave action, up-root vegetation, increase turbidity, alter the substratum and lead to release of nutrients from the sediment. The hydrological regime of the lakes must be maintained so that the area, distribution and depth of the lake habitat and its constituent/characteristic vegetation zones and communities are not reduced

Lake substratum quality	Various	Maintain appropriate substratum type, extent and chemistry to support the vegetation	Research is required to further characterise the substratum types (particle size and origin) and substratum quality (notably pH, calcium, iron and nutrient concentrations) favoured by each of the five Annex I lake habitats in Ireland. It is likely that lake habitat 3110 is associated with a range of nutrient-poor substrates, from stones, cobble and gravel, through sands, silt, clay and peat. Substratum particle size is likely to vary with depth and along the shoreline within a single lake
Water quality: transparency	Metres	Maintain appropriate Secchi transparency. There should be no decline in Secchi depth/transparency	Transparency relates to light penetration and, hence, to the depth of colonisation of vegetation. It can be affected by phytoplankton blooms, water colour and turbidity. Specific targets have yet to be established for lake habitat 3110 (O Connor, 2015). Habitat 3110 is associated with very clear water. The OECD fixed boundary system set transparency targets for oligotrophic lakes of $\geq 6\text{m}$ annual mean Secchi disk depth, and $\geq 3\text{m}$ annual minimum Secchi disk depth. Free et al. (2009) found high isoetid abundance in lakes with Secchi depths of more than 3m. Maumwee Lough had transparency of 7.1m in 2007 (McGarrigle et al., 2010)
Water quality: nutrients	$\mu\text{g/l P}$; mg/l N	Maintain/restore the concentration of nutrients in the water column to sufficiently low levels to support the habitat and its typical species	As a nutrient-poor habitat, oligotrophic and WFD 'high' status targets apply. Where a lake has nutrient concentrations that are lower than these targets, there should be no decline within class, i.e. no upward trend in nutrient concentrations. For lake habitat 3110, annual average total phosphorus (TP) concentration should be $\leq 10\mu\text{g/l TP}$, average annual total ammonia concentration should be $\leq 0.040\text{mg/l N}$ and annual 95th percentile for total ammonia should be $\leq 0.090\text{mg/l N}$. See also The European Communities Environmental Objectives (Surface Waters) Regulations 2009. Loughs Maumwee and Loughanillaun were at high nutrient status in the 2007-09 and 2010-12 reporting periods, however, Shindilla Lough exceeded the TP target in 2004-06 and 2007-09 (Clabby et al., 2008; McGarrigle et al., 2010; Bradley et al., 2015)
Water quality: phytoplankton biomass	$\mu\text{g/l Chlorophyll } a$	Maintain/restore appropriate water quality to support the habitat, including high chlorophyll <i>a</i> status	Oligotrophic and WFD 'high' status targets apply to lake habitat 3110. Where a lake has a chlorophyll <i>a</i> concentration that is lower than this target, there should be no decline within class, i.e. no upward trend in phytoplankton biomass. The average growing season (March-October) chlorophyll <i>a</i> concentration must be $< 5.8\mu\text{g/l}$. The annual average chlorophyll <i>a</i> concentration should be $< 2.5\mu\text{g/l}$ and the annual peak chlorophyll <i>a</i> concentration should be $\leq 8.0\mu\text{g/l}$. See also The European Communities Environmental Objectives (Surface Waters) Regulations 2009. Loughanillaun, Maumwee and Shindilla Loughs were oligotrophic in 2004-06 (Clabby et al., 2008) and had high chlorophyll <i>a</i> status in 2007-09 (McGarrigle et al., 2010), however, Loughanillaun declined to good chlorophyll <i>a</i> status in 2010-12 (Bradley et al., 2015)
Water quality: phytoplankton composition	EPA phytoplankton composition metric	Maintain/restore appropriate water quality to support the habitat, including high phytoplankton composition status	The EPA has developed a phytoplankton composition metric for nutrient enrichment of Irish lakes. As for other water quality indicators, lake habitat 3110 requires WFD high status. Phytoplankton composition was high at Maumwee Lough and good at Shindilla Lough in 2010-12 (Bradley et al., 2015)

Water quality: attached algal biomass	Algal cover and EPA phytobenthos metric	Maintain trace/absent attached algal biomass (<5% cover) and high phytobenthos status	Nutrient enrichment can favour epiphytic and epipelic algae that can out-compete the submerged vegetation. The cover abundance of attached algae in lake habitat 3110 should, therefore, be trace/absent (<5% cover). EPA phytobenthos can be used as an indicator of changes in attached algal biomass. As for other water quality indicators, lake habitat 3110 requires high phytobenthos status. Phytobenthos status was high at Maumwee and Shindilla Loughs in 2010-12 (Bradley et al., 2015)
Water quality: macrophyte status	EPA macrophyte metric (The Free Index)	Maintain high macrophyte status	Nutrient enrichment can favour more competitive submerged macrophyte species that out-compete the typical and characteristic species for the lake habitat. The EPA monitors macrophyte status for WFD purposes using the 'Free Index'. The target for lake habitat 3110 is high status or an Ecological Quality Ratio (EQR) for lake macrophytes of ≥ 0.90 , as defined in Schedule Five of the European Communities Environmental Objectives (Surface Waters) Regulations 2009. All three monitored lakes had high macrophyte status in 2007-09 and 2010-12 (McGarrigle et al., 2010; Bradley et al., 2015)
Acidification status	pH units; mg/l	Maintain appropriate water and sediment pH, alkalinity and cation concentrations to support the habitat, subject to natural processes	Acidification can impact on species abundance and composition in soft water lake habitats. In Europe, acidification of isoetid lakes can lead to loss of isoetids and dominance by submerged <i>Sphagnum</i> mosses and <i>Juncus bulbosus</i> (Arts, 2002). The specific requirements of lake habitat 3110, in terms of water and sediment pH, alkalinity and cation concentration, have not been determined. For lake habitat 3110, and adopting a precautionary approach based on Arts (2002), minimum pH should not be <5.5 pH units. Maximum pH should be <9.0 pH units, in line with the surface water standards established for soft waters (where water hardness is ≤ 100 mg/l calcium carbonate). See Schedule Five of the European Communities Environmental Objectives (Surface Waters) Regulations 2009. All three monitored lakes passed in 2007-09 and 2010-12 (McGarrigle et al., 2010; Bradley et al., 2015). Maumwee Lough is an acid sensitive water monitoring site
Water colour	mg/l PtCo	Maintain/restore appropriate water colour to support the habitat	Increased water colour and turbidity decrease light penetration and can reduce the area of available habitat for lake macrophytes, particularly at the lower euphotic depths. The primary source of increased water colour in Ireland is disturbance to peatland. No habitat-specific or national standards for water colour currently exist. Studies have shown median colour concentrations in Irish lakes of 38mg/l PtCo (Free et al., 2000) and 33mg/l PtCo (Free et al., 2006). It is likely that the water colour in all Irish lake habitats would naturally be <50mg/l PtCo. Water colour can be very low (<20mg/l PtCo or even <10mg/l PtCo) in lake habitat 3110, where the peatland in the lake's catchment is intact. Free et al. (2006) reported colour of 20mg/l PtCo in Loughanillaun, 26mg/l PtCo in Maumwee and 27mg/l PtCo in Shindilla Lough
Dissolved organic carbon (DOC)	mg/l	Maintain/restore appropriate organic carbon levels to support the habitat	Dissolved (and particulate) organic carbon (OC) in the water column is linked to water colour and acidification (organic acids). Increasing DOC in water has been documented across the Northern Hemisphere, including afforested peatland catchments in Ireland. Damage and degradation of peatland, leading to decomposition of peat is likely to be the predominant source of OC in Ireland. OC in water promotes decomposition by fungi and bacteria that, in turn, releases dissolved nutrients. The increased biomass of decomposers can also impact directly on the characteristic lake communities through shading, competition, etc.

Turbidity	Nephelometric turbidity units/ mg/l SS/ other appropriate units	Maintain appropriate turbidity to support the habitat	Turbidity can significantly affect the quantity and quality of light reaching rooted and attached vegetation and can, therefore, impact on lake habitats. The settlement of higher loads of inorganic or organic material on lake vegetation communities may also have impacts on sensitive, delicate species. Turbidity can increase as a result of re-suspension of material within the lake, higher loads entering the lake, or eutrophication. Turbidity measurement and interpretation is challenging. As a result, it is likely to be difficult to set habitat-specific targets for turbidity in lakes
Fringing habitat: area and condition	Hectares	Maintain the area and condition of fringing habitats necessary to support the natural structure and functioning of habitat 3110	Most lake shorelines have fringing habitats of reedswamp, other swamp, fen, marsh or wet woodland that intergrade with and support the structure and functions of the lake habitat. In Maumturk Mountains SAC, active blanket bog and heath communities are likely to dominate lake shorelines. Transition mire, fen, flush and grassland may also occur. Fringing habitats are dependent on the lake, particularly its water levels, and support wetland communities and species of conservation concern. Many of the fringing wetland habitats support higher invertebrate and plant species richness than the lake habitats themselves

4010 Northern Atlantic wet heaths with *Erica tetralix*

To restore the favourable conservation condition of Northern Atlantic wet heaths with *Erica tetralix* in Maumturk Mountains SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Northern Atlantic wet heaths with <i>Erica tetralix</i> has not been mapped in detail for Maumturk Mountains SAC, but from current available data the total area of the qualifying habitat is estimated to be approximately 4,460ha, covering 33% of the SAC (NPWS internal files). Further details on this and the following attributes can be found in the Maumturk Mountains SAC conservation objectives supporting document for blanket bogs and associated habitats
Habitat distribution	Occurrence	No decline, subject to natural processes	The habitat occurs on the lower slopes of the mountains and in other areas with gentle/moderately sloping ground in the SAC. It occurs in association with other habitats including blanket bog, dry heath and upland grassland (NPWS internal files). Further information can be found within NPWS internal files and the blanket bogs and associated habitats supporting document
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil nutrient status within natural range	See the blanket bogs and associated habitats supporting document for further details
Community diversity	Abundance of variety of vegetation communities	Maintain variety of vegetation communities, subject to natural processes	The diversity of wet heath communities within this SAC is unknown. Further information on vegetation communities associated with this habitat is presented in Perrin et al. (2014)
Vegetation composition: cross-leaved heath	Occurrence within 20m of a representative number of monitoring stops	Cross-leaved heath (<i>Erica tetralix</i>) present within a 20m radius of each monitoring stop	Attribute and target based on Perrin et al. (2014)
Vegetation composition: positive indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of positive indicator species at least 50%	Attribute and target based on Perrin et al. (2014), where the list of positive indicator species for this habitat is also presented
Vegetation composition: lichens and bryophytes	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of <i>Cladonia</i> and <i>Sphagnum</i> species, <i>Racomitrium lanuginosum</i> and pleurocarpous mosses at least 10%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: ericoid species and crowberry	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of ericoid species and crowberry (<i>Empetrum nigrum</i>) at least 15%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: dwarf shrub species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of dwarf shrubs less than 75%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: negative indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of negative indicator species less than 1%	Attribute and target based on Perrin et al. (2014), where the list of negative indicator species for this habitat is also presented
Vegetation composition: non-native species	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of non-native species less than 1%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: native trees and shrubs	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of scattered native trees and shrubs less than 20%	Attribute and target based on Perrin et al. (2014)

Vegetation composition: bracken	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of bracken (<i>Pteridium aquilinum</i>) less than 10%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: soft rush	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of soft rush (<i>Juncus effusus</i>) less than 10%	Attribute and target based on Perrin et al. (2014)
Vegetation structure: <i>Sphagnum</i> condition	Condition at a representative number of 2m x 2m monitoring stops	Less than 10% of the <i>Sphagnum</i> cover is crushed, broken and/or pulled up	Attribute and target based on Perrin et al. (2014)
Vegetation structure: signs of browsing	Percentage of shoots browsed at a representative number of 2m x 2m monitoring stops	Less than 33% collectively of the last complete growing season's shoots of ericoids, crowberry (<i>Empetrum nigrum</i>) and bog-myrtle (<i>Myrica gale</i>) showing signs of browsing	Attribute and target based on Perrin et al. (2014)
Vegetation structure: burning	Occurrence in local vicinity of a representative number of monitoring stops	No signs of burning in sensitive areas, into the moss, liverwort or lichen layer or exposure of peat surface due to burning	Attribute and target based on Perrin et al. (2014), where the list of sensitive areas for this habitat is also presented
Physical structure: disturbed bare ground	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of disturbed bare ground less than 10%	Attribute and target based on Perrin et al. (2014)
Physical structure: drainage	Percentage area in local vicinity of a representative number of monitoring stops	Area showing signs of drainage from heavy trampling, tracking or ditches less than 10%	Attribute and target based on Perrin et al. (2014)
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat	This includes species listed in the Flora (Protection) Order, 2015 (FPO) and/or the red data lists (Lockhart et al., 2012; Wyse Jackson et al., 2016). There are recent records for the FPO listed and Vulnerable marsh clubmoss (<i>Lycopodiella inundata</i>) (Wyse Jackson et al., 2016) from wet heath habitat along the shores of Maumwee Lough at Lackavrea (O. Daly, pers. comm.)

Conservation Objectives for : Maumturk Mountains SAC [002008]

4060 Alpine and Boreal heaths

To restore the favourable conservation condition of Alpine and Boreal heaths in Maumturk Mountains SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Alpine and Boreal heaths habitat has not been mapped in detail for Maumturk Mountains SAC but from current available data the total area of the qualifying habitat is estimated to be approximately 1,301ha, covering 10% of the SAC (NPWS internal files). Further details on this and the following attributes can be found in the Maumturk Mountains SAC conservation objectives supporting document for blanket bogs and associated habitats
Habitat distribution	Occurrence	No decline, subject to natural processes	The habitat occurs on summits and ridges within the SAC. Examples are present at Knocknahillion and on a hill west of Corcogemore (NPWS internal files). Further information can be found within NPWS internal files and the blanket bogs and associated habitats supporting document
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil nutrient status within natural range	See the blanket bogs and associated habitats supporting document for further details
Community diversity	Abundance of variety of vegetation communities	Maintain variety of vegetation communities, subject to natural processes	The diversity of Alpine and Boreal heath communities within this SAC is unknown. Further information on vegetation communities associated with this habitat is presented in Perrin et al. (2014)
Vegetation composition: lichens and bryophytes	Number of species at a representative number of 2m x 2m monitoring stops	Number of bryophyte or non-crustose lichen species present at each monitoring stop is at least three	Attribute and target based on Perrin et al. (2014)
Vegetation composition: positive indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of positive indicator species at least 66%	Attribute and target based on Perrin et al. (2014), where the list of positive indicator species for this habitat is also presented
Vegetation composition: dwarf shrub species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of dwarf shrub species at least 10%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: negative indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of negative indicator species less than 10%	Attribute and target based on Perrin et al. (2014), where the list of negative indicator species for this habitat is also presented
Vegetation composition: non-native species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of non-native species less than 1%	Attribute and target based on Perrin et al. (2014)
Vegetation structure: signs of grazing	Percentage of leaves grazed at a representative number of 2m x 2m monitoring stops	Less than 10% collectively of the live leaves of specific graminoids showing signs of grazing	Attribute and target based on Perrin et al. (2014). See the blanket bogs and associated habitats supporting document for the list of specific graminoids
Vegetation structure: signs of browsing	Percentage of shoots browsed at a representative number of 2m x 2m monitoring stops	Less than 33% collectively of the last complete growing season's shoots of ericoids and crowberry (<i>Empetrum nigrum</i>) showing signs of browsing	Attribute and target based on Perrin et al. (2014)
Vegetation structure: burning	Occurrence in local vicinity of a representative number of monitoring stops	No signs of burning within the habitat	Attribute and target based on Perrin et al. (2014)

Physical structure: disturbed bare ground	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of disturbed bare ground less than 10%	Attribute and target based on Perrin et al. (2014)
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat and no decline in status of hepatic mats associated with this habitat	This includes species listed in the Flora (Protection) Order, 2015 and/or the red data lists (Lockhart et al., 2012; Wyse Jackson et al., 2016). The Near Threatened alpine clubmoss (<i>Diphasiastrum alpinum</i>) (Wyse Jackson et al., 2016) was recorded from this habitat. Further information can be found in Roden (1986)

Conservation Objectives for : Maumturk Mountains SAC [002008]

7130 Blanket bogs (* if active bog)

To restore the favourable conservation condition of Blanket bogs (* if active bog) in Maumturk Mountains SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Blanket bog has not been mapped in detail for Maumturk Mountains SAC, but from current available data the total area of the qualifying habitat is estimated to be approximately 1,926ha, covering 14% of the SAC (NPWS internal files). Further details on this and the following attributes can be found in the Maumturk Mountains SAC conservation objectives supporting document for blanket bogs and associated habitats
Habitat distribution	Occurrence	No decline, subject to natural processes	The habitat typically occurs in areas of lower ground along the western, eastern and south-eastern sections of the SAC. An excellent example of this habitat type is found at Caher (Douglas et al., 1989; NPWS internal files). Further information can be found within Douglas et al. (1989), NPWS internal files and the blanket bogs and associated habitats supporting document
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil nutrient status within natural range	See the blanket bogs and associated habitats supporting document for further details
Ecosystem function: peat formation	Active blanket bog as a proportion of the total area of Annex I blanket bog habitat	At least 99% of the total Annex I blanket bog area is active	See the blanket bogs and associated habitats supporting document for further details
Ecosystem function: hydrology	Flow direction, water levels, occurrence of drains and erosion gullies	Natural hydrology unaffected by drains and erosion	Further details and a brief discussion of restoration potential is presented in the blanket bogs and associated habitats supporting document
Community diversity	Abundance of variety of vegetation communities	Maintain variety of vegetation communities, subject to natural processes	A variety of blanket bog vegetation communities have been recorded in this SAC (Douglas et al., 1989; NPWS internal files), three of which correspond to communities recorded in the National Survey of Upland Habitats and listed in the provisional list of vegetation communities described in Perrin et al. (2014). Further information on vegetation communities associated with this habitat is presented in Perrin et al. (2014)
Vegetation composition: positive indicator species	Number of species at a representative number of 2m x 2m monitoring stops	Number of positive indicator species present at each monitoring stop is at least seven	Attribute and target based on Perrin et al. (2014), where the list of positive indicator species for this habitat is also presented
Vegetation composition: lichens and bryophytes	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of bryophytes or lichens, excluding <i>Sphagnum fallax</i> , at least 10%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: potential dominant species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of each of the potential dominant species less than 75%	Attribute and target based on Perrin et al. (2014). See the blanket bogs and associated habitats supporting document for the list of potential dominant species
Vegetation composition: negative indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of negative indicator species less than 1%	Attribute and target based on Perrin et al. (2014), where the list of negative indicator species for this habitat is also presented
Vegetation composition: non-native species	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of non-native species less than 1%	Attribute and target based on Perrin et al. (2014)

Vegetation composition: native trees and shrubs	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of scattered native trees and shrubs less than 10%	Attribute and target based on Perrin et al. (2014)
Vegetation structure: <i>Sphagnum</i> condition	Condition at a representative number of 2m x 2m monitoring stops	Less than 10% of the <i>Sphagnum</i> cover is crushed, broken and/or pulled up	Attribute and target based on Perrin et al. (2014)
Vegetation structure: signs of browsing	Percentage of shoots browsed at a representative number of 2m x 2m monitoring stops	Last complete growing season's shoots of ericoids, crowberry (<i>Empetrum nigrum</i>) and bog-myrtle (<i>Myrica gale</i>) showing signs of browsing collectively less than 33%	Attribute and target based on Perrin et al. (2014)
Vegetation structure: burning	Occurrence in local vicinity of a representative number of monitoring stops	No signs of burning in sensitive areas, into the moss, liverwort or lichen layer or exposure of peat surface due to burning	Attribute and target based on Perrin et al. (2014), where the list of sensitive areas for this habitat is also presented
Physical structure: disturbed bare ground	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of disturbed bare ground less than 10%	Attribute and target based on Perrin et al. (2014)
Physical structure: drainage	Percentage area in local vicinity of a representative number of monitoring stops	Area showing signs of drainage from heavy trampling, tracking or ditches less than 10%	Attribute and target based on Perrin et al. (2014)
Physical structure: erosion	Percentage area in local vicinity of a representative number of monitoring stops	Less than 5% of the greater bog mosaic comprises erosion gullies and eroded areas	Attribute and target based on Perrin et al. (2014)
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat and no decline in status of hepatic mats associated with this habitat	This includes species listed in the Flora (Protection) Order, 2015 (FPO) and/or the red data lists (Lockhart et al., 2012; Wyse Jackson et al., 2016). The Near Threatened brown beak-sedge (<i>Rhynchospora fusca</i>) (Wyse Jackson et al., 2016) has been recorded from blanket bog in the SAC (NPWS internal files). The FPO listed and Vulnerable marsh clubmoss (<i>Lycopodiella inundata</i>) and the FPO listed and Near Threatened slender cottongrass (<i>Eriophorum gracile</i>) (Wyse Jackson et al., 2016) have been recorded within the SAC (NPWS internal files), but these species cannot be assigned specifically to blanket bog

Conservation Objectives for : Maumturk Mountains SAC [002008]

7150 Depressions on peat substrates of the Rhynchosporion

To restore the favourable conservation condition of Depressions on peat substrates of the Rhynchosporion in Maumturk Mountains SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Depressions on peat substrates of the Rhynchosporion has not been mapped in detail for Maumturk Mountains SAC and thus the total area of the qualifying habitat in the SAC is unknown. Further details on this and the following attributes can be found in the Maumturk Mountains SAC conservation objectives supporting document for blanket bogs and associated habitats
Habitat distribution	Occurrence	No decline, subject to natural processes	The habitat is documented to occur amongst wet areas in lowland blanket bog in the SAC. The best examples of this habitat occur at Caher and to the south-east of Maam bridge (Douglas et al., 1989; NPWS internal files). Further information can be found within Douglas et al. (1989), NPWS internal files and the blanket bogs and associated habitats supporting document
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil nutrient status within natural range	See the blanket bogs and associated habitats supporting document for further details
Vegetation composition: positive indicator species	Number of species at a representative number of 2m x 2m monitoring stops	Number of positive indicator species at each monitoring stop is at least five	Attribute and target based on Perrin et al. (2014), where the list of positive indicator species for this habitat is also presented
Vegetation composition: <i>Rhynchospora</i> spp.	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of white beaked sedge (<i>Rhynchospora alba</i>) and brown beaked sedge (<i>R. fusca</i>) at least 10%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: potential dominant species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of each of the potential dominant species individually less than 35%	Attribute and target based on Perrin et al. (2014). See the blanket bogs and associated habitats supporting document for the list of potential dominant species
Vegetation composition: negative indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of negative indicator species less than 1%	Attribute and target based on Perrin et al. (2014), where the list of negative indicator species for this habitat is also presented
Vegetation composition: non-native species	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of non-native species less than 1%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: native trees and shrubs	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of scattered native trees and shrubs less than 10%	Attribute and target based on Perrin et al. (2014)
Vegetation structure: <i>Sphagnum</i> condition	Condition at a representative number of 2m x 2m monitoring stops	Less than 10% of the <i>Sphagnum</i> cover is crushed, broken and/or pulled up	Attribute and target based on Perrin et al. (2014)
Vegetation structure: signs of browsing	Percentage of shoots browsed at a representative number of 2m x 2m monitoring stops	Last complete growing season's shoots of ericoids, crowberry (<i>Empetrum nigrum</i>) and bog-myrtle (<i>Myrica gale</i>) showing signs of browsing collectively less than 33%	Attribute and target based on Perrin et al. (2014)

Vegetation structure: burning	Occurrence in local vicinity of a representative number of monitoring stops	No signs of burning in sensitive areas, into the moss, liverwort or lichen layer or exposure of peat surface due to burning	Attribute and target based on Perrin et al. (2014), where the list of sensitive areas for this habitat is also presented
Physical structure: disturbed bare ground	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of disturbed bare ground less than 10%	Attribute and target based on Perrin et al. (2014)
Physical structure: drainage	Percentage area in local vicinity of a representative number of monitoring stops	Area showing signs of drainage from heavy trampling, tracking or ditches less than 10%	Attribute and target based on Perrin et al. (2014)
Physical structure: erosion	Percentage area in local vicinity of a representative number of monitoring stops	Less than 5% of the greater bog mosaic comprises erosion gullies and eroded areas	Attribute and target based on Perrin et al. (2014)
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat	This includes species listed in the Flora (Protection) Order, 2015 (FPO) and/or the red data lists (Lockhart et al., 2012; Wyse Jackson et al., 2016). The Near Threatened brown beak-sedge (<i>Rhynchospora fusca</i>) and the FPO listed and Vulnerable marsh clubmoss (<i>Lycopodiella inundata</i>) (Wyse Jackson et al., 2016) have been recorded from the SAC (NPWS internal files), but these species cannot be assigned specifically to this habitat

Conservation Objectives for : Maumturk Mountains SAC [002008]

8220 Siliceous rocky slopes with chasmophytic vegetation

To restore the favourable conservation condition of Siliceous rocky slopes with chasmophytic vegetation in Maumturk Mountains SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Siliceous rocky slopes with chasmophytic vegetation has not been mapped in detail for Maumturk Mountains SAC, but from current available data the total area of the qualifying habitat is estimated to be approximately 202ha, covering c.1% of the SAC (NPWS internal files). Further details on this and the following attributes can be found in the Maumturk Mountains SAC conservation objectives supporting document for blanket bogs and associated habitats
Habitat distribution	Occurrence	No decline, subject to natural processes	The habitat is present amongst the steep slopes and corries of the Maumturk Mountains, Teernakill South, Lackavrea and Knocknagur (NPWS internal files). Further information can be found within NPWS internal files and the blanket bogs and associated habitats supporting document
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil nutrient status within natural range	See the blanket bogs and associated habitats supporting document for further details
Vegetation composition: positive indicator species	Number of species in local vicinity of a representative number of monitoring stops	At least one positive indicator species present in vicinity of each monitoring stop	Attribute and target based on Perrin et al. (2014). The list of positive indicator species for this habitat is also presented in Perrin et al. (2014) and is the same as for 8110 Siliceous screes
Vegetation composition: non-native species	Percentage cover in local vicinity of a representative number of monitoring stops	Proportion of vegetation composed of non-native species less than 1%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: bracken, native trees and shrubs	Percentage cover in local vicinity of a representative number of monitoring stops	Total cover of bracken (<i>Pteridium aquilinum</i>), native trees and shrubs less than 25%	Attribute and target based on Perrin et al. (2014)
Vegetation structure: grazing and browsing	Percentage of leaves/shoots grazed/browsed in local vicinity of a representative number of monitoring stops	Live leaves of forbs and shoots of dwarf shrubs showing signs of grazing or browsing collectively less than 50%	Attribute and target based on Perrin et al. (2014)
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat and no decline in status of hepatic mats associated with this habitat	This includes species listed in the Flora (Protection) Order, 2015 and/or the red data lists (Lockhart et al., 2012; Wyse Jackson et al., 2016). The Near Threatened beech fern (<i>Phegopteris connectilis</i>) (Wyse Jackson et al., 2016) has been recorded within the SAC (NPWS internal files), but this species cannot be assigned specifically to this habitat

Conservation Objectives for : Maumturk Mountains SAC [002008]

1106 Salmon *Salmo salar*

To maintain the favourable conservation condition of Atlantic Salmon in Maumturk Mountains SAC, which is defined by the following list of attributes and targets:


Attribute	Measure	Target	Notes
Distribution: extent of anadromy	Percentage of river accessible	100% of river channels down to second order accessible from estuary	Artificial barriers block salmons' upstream migration, thereby limiting species to lower stretches and restricting access to spawning areas
Adult spawning fish	Number	Conservation limit (CL) for each system consistently exceeded	A conservation limit (CL) is defined by the North Atlantic Salmon Conservation Organisation (NASCO) as "the spawning stock level that produces long-term average maximum sustainable yield as derived from the adult to adult stock and recruitment relationship". The target is based on the Standing Scientific Committee on Salmon (SSCS) annual model output of CL attainment levels. See SSCS (2016). Attainment of CL estimates are derived from direct counts of adults (rod catch, fish counter) or indirectly by fry abundance counts. Rivers in the Corrib catchment are currently achieving CL
Salmon fry abundance	Number of fry/5 minutes electrofishing	Maintain or exceed 0+ fry mean catchment-wide abundance threshold value. Currently set at 17 salmon fry/5 minutes sampling	The target is the threshold value for rivers currently exceeding their conservation limit (CL)
Out-migrating smolt abundance	Number	No significant decline	Smolt abundance can be negatively affected by a number of impacts such as estuarine pollution, predation and sea lice (<i>Lepeophtheirus salmonis</i>)
Number and distribution of redds	Number and occurrence	No decline in number and distribution of spawning redds due to anthropogenic causes	Salmon spawn in clean gravels
Water quality	EPA Q value	At least Q4 at all sites sampled by EPA	Q values based on triennial water quality surveys carried out by the Environmental Protection Agency (EPA)


Conservation Objectives for : Maumturk Mountains SAC [002008]


1833 Slender Naiad *Najas flexilis*

To maintain the favourable conservation condition of Slender Naiad in Maumturk Mountains SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Population extent	Hectares; distribution	No change to the spatial extent of <i>Najas flexilis</i> within Loughs Lehanagh and Derryneen, subject to natural processes. See map 4 for known locations	The selection of Maumturk Mountains SAC for <i>Najas flexilis</i> (slender naiad) was based on its presence in Lehanagh Lough. In 2013, Dr Cilian Roden discovered a small population in Derryneen Lough in the SAC (Roden, 2013). It is possible that the species occurs in other lakes within the SAC. See the Maumturk Mountains SAC conservation objectives supporting document for <i>Najas flexilis</i> for further details
Population depth	Metres	No change to the depth range of <i>Najas flexilis</i> within each lake, subject to natural processes	See the <i>Najas flexilis</i> supporting document for further details
Population viability	Plant traits	No decline in plant fitness, subject to natural processes	See the <i>Najas flexilis</i> supporting document for further details
Population abundance	Square metres	No change to the cover abundance of <i>Najas flexilis</i> , subject to natural processes	See the <i>Najas flexilis</i> supporting document for further details
Species distribution	Occurrence	No decline, subject to natural processes	See the <i>Najas flexilis</i> supporting document for further details
Habitat extent	Hectares	No decline, subject to natural processes	See the <i>Najas flexilis</i> supporting document for further details
Hydrological regime: water level fluctuations	Metres	Maintain appropriate natural hydrological regime necessary to support the habitat for the species	See the <i>Najas flexilis</i> supporting document for further details
Lake substratum quality	Various	Maintain appropriate substratum type, extent and chemistry to support the populations of the species	See the <i>Najas flexilis</i> supporting document for further details
Water quality	Various	Maintain appropriate water quality to support the populations of the species	See the <i>Najas flexilis</i> supporting document for further details
Acidification status	pH units; mg/l	Maintain appropriate water and sediment pH, alkalinity and cation concentrations to support the populations of <i>Najas flexilis</i> , subject to natural processes	See the <i>Najas flexilis</i> supporting document for further details
Water colour	mg/l PtCo	Maintain appropriate water colour to support the populations of <i>Najas flexilis</i>	See the <i>Najas flexilis</i> supporting document for further details
Associated species	Species composition and abundance	Maintain appropriate associated species and vegetation communities to support the populations of <i>Najas flexilis</i>	See the <i>Najas flexilis</i> supporting document for further details
Fringing habitat: area and condition	Hectares	Maintain the area and condition of fringing habitats necessary to support the populations of <i>Najas flexilis</i>	See the <i>Najas flexilis</i> supporting document for further details







Legend
 Maumturk Mountains SAC 002008



Legend

- Maumturk Mountains SAC 002008
- Connemara Bog Complex SAC 002034
- The Twelve Bens/Garraun Complex SAC 002031
- Lough Corrib SAC 000297
- Lough Corrib SPA 004042
- Connemara Bog Complex SPA 004181





Legend

- Maumturk Mountains SAC 002008
- OSi Discovery Series County Boundary

Indicative Lake Habitats

- Potential 3110 Potential oligotrophic waters containing very few minerals of sandy plains: *Littorelletalia uniflorae*
- Potential 3110 / Potential 3130* Potential oligotrophic waters containing very few minerals of sandy plains: *Littorelletalia uniflorae* / Potential oligotrophic to mesotrophic standing waters with vegetation of the *Littorelletalia uniflorae* and/or of the *Isoëto-Nanojuncetea*
*Habitat 3130 not a qualifying interest





Legend

- 1833 Slender Naiad *Najas flexilis*
- Maumturk Mountains SAC 002008
- OSi Discovery Series County Boundary

National Parks and Wildlife Service

Conservation Objectives Series

The Twelve Bens/Garraun Complex SAC 002031



An Roinn Ealaíon, Oidhreachta,
Gnóthaí Réigiúnacha, Tuaithe agus Gaeltachta

Department of Arts, Heritage,
Regional, Rural and Gaeltacht Affairs



**National Parks and Wildlife Service,
Department of Arts, Heritage, Regional, Rural and Gaeltacht Affairs,
7 Ely Place, Dublin 2, Ireland.
Web: www.npws.ie
E-mail: nature.conservation@ahg.gov.ie**

Citation:

**NPWS (2017) Conservation Objectives: The Twelve Bens/Garraun Complex SAC
002031. Version 1. National Parks and Wildlife Service, Department of Arts,
Heritage, Regional, Rural and Gaeltacht Affairs.**

**Series Editor: Rebecca Jeffrey
ISSN 2009-4086**

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

* indicates a priority habitat under the Habitats Directive

002031	The Twelve Bens/Garraun Complex SAC
1029	Freshwater Pearl Mussel <i>Margaritifera margaritifera</i>
1106	Salmon <i>Salmo salar</i>
1355	Otter <i>Lutra lutra</i>
1833	Slender Naiad <i>Najas flexilis</i>
3110	Oligotrophic waters containing very few minerals of sandy plains (Littorelletalia uniflorae)
3130	Oligotrophic to mesotrophic standing waters with vegetation of the Littorelletea uniflorae and/or Isoeto-Nanojuncetea
4060	Alpine and Boreal heaths
7130	Blanket bogs (* if active bog)
7150	Depressions on peat substrates of the Rhynchosporion
8110	Siliceous scree of the montane to snow levels (Androsacetalia alpinae and Galeopsietalia ladani)
8210	Calcareous rocky slopes with chasmophytic vegetation
8220	Siliceous rocky slopes with chasmophytic vegetation
91A0	Old sessile oak woods with <i>Q. robur</i> and <i>Q. petraea</i> in the British Isles

Please note that this SAC is adjacent to Maumturk Mountains SAC (002008), Connemara Bog Complex SAC (002034) and West Connacht Coast SAC (002998). See map 2. The conservation objectives for this site should be used in conjunction with those for the adjacent sites as appropriate.

Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

NPWS Documents

Year :	1984
Title :	The vegetation of Irish lakes
Author :	Heuff, H.
Series :	Unpublished report to NPWS
Year :	1987
Title :	Survey to locate lowland blanket bogs of scientific interest in Connemara, Co. Galway
Author :	Douglas, C.; Grogan, H.
Series :	Unpublished report to the Wildlife Service
Year :	1989
Title :	A survey to locate blanket bogs in Co. Galway. Part 2
Author :	Douglas, C.; Garvey, L.; Kelly, L.; O'Sullivan, A.
Series :	Unpublished report to the Wildlife Service
Year :	2002
Title :	<i>Najas flexilis</i> in Donegal
Author :	Roden, C.M.
Series :	Unpublished report to NPWS
Year :	2004
Title :	The distribution of <i>Najas flexilis</i> in Ireland 2002-2004
Author :	Roden, C.M.
Series :	Unpublished report to NPWS
Year :	2006
Title :	Otter survey of Ireland 2004/2005
Author :	Bailey, M.; Rochford, J.
Series :	Irish Wildlife Manual No. 23
Year :	2007
Title :	Supporting documentation for the Habitats Directive Conservation Status Assessment - backing documents. Article 17 forms and supporting maps
Author :	NPWS
Series :	Unpublished report to NPWS
Year :	2008
Title :	Rapid assessment of rivers with prior records of <i>Margaritifera margaritifera</i>
Author :	Moorkens, E.; Killeen, I.
Series :	Unpublished report to NPWS
Year :	2008
Title :	National survey of native woodlands 2003-2008
Author :	Perrin, P.M.; Martin, J.; Barron, S.; O'Neill, F.H.; McNutt, K.E.; Delaney, A.
Series :	Unpublished report to NPWS
Year :	2008
Title :	Survey of Dawros river
Author :	Moorkens, E.
Series :	unpublished report to NPWS

Year :	2009
Title :	NS II freshwater pearl mussel sub-basin management plans: fisheries survey. Stage 1 report
Author :	Paul Johnston Associates
Series :	Unpublished report to NPWS
Year :	2009
Title :	NS II Freshwater Pearl Mussel Sub-basin Management Plans: Report on Biological Monitoring of Surface Water Quality in Dawros Catchment, Co. Galway
Author :	Williams, L.
Series :	Unpublished report to NPWS
Year :	2010
Title :	A provisional inventory of ancient and long-established woodland in Ireland
Author :	Perrin, P.M.; Daly, O.H.
Series :	Irish Wildlife Manual No. 46
Year :	2010
Title :	Second draft Dawros freshwater pearl mussel sub-basin management plan (2009-2015). March 2010
Author :	NPWS
Series :	Unpublished document to the Department of Environment, Heritage and Local Government
Year :	2012
Title :	Ireland Red List No. 8: Bryophytes
Author :	Lockhart, N.; Hodgetts, N.; Holyoak, D.
Series :	Ireland Red List series, NPWS
Year :	2012
Title :	Catchment Assessment Report Manual
Author :	Monaghan, F.
Series :	Unpublished report to NPWS
Year :	2013
Title :	National otter survey of Ireland 2010/12
Author :	Reid, N.; Hayden, B.; Lundy, M.G.; Pietravalle, S.; McDonald, R.A.; Montgomery, W.I.
Series :	Irish Wildlife Manual No. 76
Year :	2013
Title :	The status of EU protected habitats and species in Ireland. Volume 2. Habitats assessments
Author :	NPWS
Series :	Conservation assessments
Year :	2013
Title :	The status of EU protected habitats and species in Ireland. Volume 3. Species assessments
Author :	NPWS
Series :	Conservation assessments
Year :	2014
Title :	Guidelines for a national survey and conservation assessment of upland vegetation and habitats in Ireland, Version 2.0
Author :	Perrin, P.M.; Barron, S.J.; Roche, J.R.; O'Hanrahan, B.
Series :	Irish Wildlife Manual No. 79
Year :	2014
Title :	Targeted survey of <i>Najas flexilis</i>
Author :	Roden, C.; Murphy, P.
Series :	Unpublished report to NPWS

Year :	2015
Title :	Habitats Directive Annex I lake habitats: a working interpretation for the purposes of site-specific conservation objectives and Article 17 reporting
Author :	O Connor, Á.
Series :	Unpublished document by NPWS
Year :	2016
Title :	Ireland Red List No. 10: Vascular Plants
Author :	Wyse Jackson, M.; FitzPatrick, Ú.; Cole, E.; Jebb, M.; McFerran, D.; Sheehy Skeffington, M.; Wright, M.
Series :	Ireland Red Lists series, NPWS
Year :	2016
Title :	Survey and Condition Assessment of the population of the freshwater pearl mussel <i>Margaritifera margaritifera</i> in the Dawros River, County Galway
Author :	Moorkens, E.
Series :	Unpublished report to NPWS
Year :	2017
Title :	The Twelve Bens/Garraun Complex SAC (site code: 2031) Conservation objectives supporting document- blanket bogs and associated habitats V1
Author :	NPWS
Series :	Conservation objectives supporting document
Year :	2017
Title :	The Twelve Bens/Garraun Complex SAC (site code: 2031) Conservation objectives supporting document- <i>Najas flexilis</i> V1
Author :	NPWS
Series :	Conservation objectives supporting document

Other References

Year :	1970
Title :	The lichens and lichen parasites of Derryclare Wood, Connemara
Author :	Folan, A.C.M.; Mitchell, M.E.
Series :	Proceedings of the Royal Irish Academy. Section B: Biological, Geological, and Chemical Science, 70: 163-170
Year :	1982
Title :	Otter survey of Ireland
Author :	Chapman, P.J.; Chapman, L.L.
Series :	Unpublished report to Vincent Wildlife Trust
Year :	1982
Title :	Eutrophication of waters. Monitoring assessment and control
Author :	OECD
Series :	OECD, Paris
Year :	1987
Title :	An account of the flora and vegetation of Derryclare Wood, Connemara (Co. Galway), Western Ireland
Author :	Ferguson, D.K.; Westhoff, V.
Series :	Proceedings of the Koninklijke Nederlandse Akademie van Wetenschappen, Series C 90(2): 139-172
Year :	1988
Title :	The reproductive biology of freshwater mussels in Ireland, with observations on their distribution and demography
Author :	Ross, E.D.
Series :	Unpublished Ph.D. Thesis, National University of Ireland, Galway

Year :	1991
Title :	The spatial organization of otters (<i>Lutra lutra</i>) in Shetland
Author :	Kruuk, H.; Moorhouse, A.
Series :	Journal of Zoology, 224: 41-57
Year :	1999
Title :	Diet of otters (<i>Lutra lutra</i>) on Inishmore, Aran Islands, west coast of Ireland
Author :	Kingston, S.; O'Connell, M.; Fairley, J.S.
Series :	Biology and Environment: Proceedings of the Royal Irish Academy, 99B: 173-182
Year :	1999
Title :	Derryclare Woods: a multidisciplinary study of its structure and age
Author :	Fahy, O.; Fuller, M.; Gabbett, M.; Gormally, M.; Sheehy Skeffington, M.
Series :	Report to the Heritage Council, Kilkenny
Year :	2000
Title :	Colour in Irish lakes
Author :	Free, G.; Allott, N.; Mills, P.; Kennelly, C.; Day, S.
Series :	Verhandlungen Internationale Vereinigung für theoretische und angewandte Limnologie, 27: 2620-2623
Year :	2001
Title :	Aquatic plants in Britain and Ireland
Author :	Preston, C.D.; Croft, J.M.
Series :	Harley Books, Colchester
Year :	2002
Title :	Reversing the habitat fragmentation of British woodlands
Author :	Peterken, G.
Series :	WWF-UK, London
Year :	2002
Title :	Deterioration of Atlantic soft water macrophyte communities by acidification, eutrophication and alkanisation
Author :	Arts, G.H.P.
Series :	Aquatic Botany, 73: 373-393
Year :	2003
Title :	Survey of the Pearl Mussel Population - 2003. Dawros River Co. Galway
Author :	Aster Environmental Consultants
Series :	Report to The Heritage Council, Kilkenny
Year :	2004
Title :	The ecology of <i>Najas flexilis</i>
Author :	Wingfield, R.A.; Murphy, K.J.; Hollingsworth, P.; Gaywood, M.J.
Series :	Scottish Natural Heritage Commissioned Report No. 017 (ROAME No. F98PA02)
Year :	2006
Title :	Otters - ecology, behaviour and conservation
Author :	Kruuk, H.
Series :	Oxford University Press
Year :	2006
Title :	The status of host fish populations and fish species richness in European freshwater pearl mussel (<i>Margaritifera margaritifera</i>) streams
Author :	Geist, J.; Porkka, M.; Kuehn, R.
Series :	Aquatic Conservation: Marine and Freshwater Ecosystems, 16: 251-266

Year :	2006
Title :	A reference-based typology and ecological assessment system for Irish lakes. Preliminary investigations. Final report. Project 2000-FS-1-M1 Ecological assessment of lakes pilot study to establish monitoring methodologies EU (WFD)
Author :	Free, G.; Little, R.; Tierney, D.; Donnelly, K.; Coroni, R.
Series :	EPA, Wexford
Year :	2009
Title :	The identification, characterization and conservation value of isoetid lakes in Ireland
Author :	Free, G.; Bowman, J.; McGarrigle, M.; Little, R.; Coroni, R.; Donnelly, K.; Tierney, D.; Trodd, W.
Series :	Aquatic Conservation: Marine and Freshwater Ecosystems, 19(3): 264–273
Year :	2010
Title :	Otter tracking study of Roaringwater Bay
Author :	De Jongh, A.; O'Neill, L.
Series :	Unpublished draft report to NPWS
Year :	2010
Title :	Addressing the conservation and rehabilitation of <i>Margaritifera margaritifera</i> populations in the Republic of Ireland within the framework of the habitats and species directive
Author :	Moorkens, E.
Series :	Journal of Conchology, 40: 339
Year :	2010
Title :	Water quality in Ireland 2007-2009
Author :	McGarrigle, M.; Lucey, J.; Ó Cinnéide, M.
Series :	EPA, Wexford
Year :	2014
Title :	Assessing near-bed velocity in a recruiting population of the endangered freshwater pearl mussel (<i>Margaritifera margaritifera</i>) in Ireland
Author :	Moorkens, E.; Killeen, I.
Series :	Aquatic Conservation: Marine and Freshwater Ecosystems, 24(6): 853-862
Year :	2014
Title :	The impact of rural land management changes on soil hydraulic properties and runoff processes: results from experimental plots in upland UK
Author :	Marshall, M.R.; Ballard, C.E.; Frogbrook, Z.L.; Solloway, I.; McIntyre, N.; Reynolds, B.; Wheeler, H.S.
Series :	Hydrological Processes, 28: 2617–2629
Year :	2015
Title :	Water quality in Ireland 2010-2012
Author :	Bradley, C.; Byrne, C.; Craig, M.; Free, G.; Gallagher, T.; Kennedy, B.; Little, R.; Lucey, J.; Mannix, A.; McCreesh, P.; McDermott, G.; McGarrigle, M.; Ní Longphuirt, S.; O'Boyle, S.; Plant, C.; Tierney, D.; Trodd, W.; Webster, P.; Wilkes, R.; Wynne, C.
Series :	EPA, Wexford
Year :	2016
Title :	The Status of Irish Salmon Stocks in 2015 with Precautionary Catch Advice for 2016
Author :	SSCS (Standing Scientific Committee on Salmon)
Series :	Independent Scientific Report to Inland Fisheries Ireland

Spatial data sources

Year :	2008
Title :	OSi 1:5000 IG vector dataset
GIS Operations :	WaterPolygons feature class clipped to the SAC boundary. Expert opinion used to identify Annex I habitat and to resolve any issues arising
Used For :	3110, 3130 (map 3)
Year :	Revision 2010
Title :	National Survey of Native Woodlands 2003-2008. Version 1
GIS Operations :	QIs selected; clipped to SAC boundary. Expert opinion used as necessary to resolve any issues arising
Used For :	91A0 (map 4)
Year :	2017
Title :	NPWS rare and threatened species database
GIS Operations :	Dataset created from spatial references in database records. Expert opinion used as necessary to resolve any issues arising
Used For :	1029 (map 5)
Year :	Revision 2012
Title :	Margaritifera Sensitive Areas data
GIS Operations :	Relevant catchment boundaries identified. Expert opinion used as necessary to resolve any issues arising
Used For :	1029 (map 5)
Year :	2010
Title :	OSi 1:5000 IG vector dataset
GIS Operations :	Creation of 80m buffer on the aquatic side of lake data; creation of 10m buffer on the terrestrial side of lake data. These datasets combined with the derived OSi Discovery Series river and canal datasets. Overlapping regions investigated and resolved; resulting dataset clipped to SAC boundary. Expert opinion used as necessary to resolve any issues arising. Creation of 250m buffer on aquatic side of the lake boundary to highlight potential commuting points
Used For :	1355 (map 6)
Year :	2005
Title :	OSi Discovery series vector data
GIS Operations :	Creation of a 10m buffer on the terrestrial side of river banks data; creation of 20m buffer applied to canal centreline data. Creation of a 20m buffer applied to river and stream centreline data; These datasets combined with the derived OSi 1:5000 vector lake buffer data. Overlapping regions investigated and resolved; resulting dataset clipped to SAC boundary. Expert opinion used as necessary to resolve any issues arising
Used For :	1355 (no map)
Year :	2013
Title :	Najas flexilis data
GIS Operations :	Lake habitat for species clipped to SAC boundary
Used For :	1833 (map 7)

3110 Oligotrophic waters containing very few minerals of sandy plains (Littorelletalia uniflorae)

To maintain the favourable conservation condition of Oligotrophic waters containing very few minerals of sandy plains (Littorelletalia uniflorae) in The Twelve Bens/Garraun Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Lake habitat 3110 is considered likely to occur in most lakes in The Twelve Bens/Garraun Complex SAC, notably in Loughs Inagh, Derryclare, Fee, Nahillion and Muck. In line with Article 17 reporting (NPWS, 2013), all lakes larger than 1ha have been mapped as potential 3110 (see map 3). It is likely, however, that the habitat also occurs in many of the smaller lakes and ponds in the SAC. In lakes/ponds in blanket bog, lake habitat 3160 may co-occur. Lake habitat 3110 is likely to co-occur with lake habitat 3130 in Pollacappul and Kylemore Loughs. Two measures of extent should be used: 1. the area of the lake itself and; 2. the extent of the vegetation communities/zones that typify the habitat. Further information relating to all attributes is provided in the lake habitats supporting document for the purposes of site-specific conservation objectives and Article 17 reporting (O Connor, 2015)
Habitat distribution	Occurrence	No decline, subject to natural processes	As noted above, all lakes larger than 1ha have been mapped as potential 3110, but the habitat is likely to be even more widespread in the SAC (see map 3)
Typical species	Occurrence	Typical species present, in good condition, and demonstrating typical abundances and distribution	For lists of typical plant species, see the Article 17 habitat assessment for 3110 (NPWS, 2013) and the lake habitats supporting document (O Connor, 2015). See Heuff (1984) and The Twelve Bens/Garraun Complex SAC conservation objectives supporting document for <i>Najas flexilis</i> for information on Pollacappul and Kylemore Loughs. Derryclare, Kylemore and Pollacappul Loughs are Water Framework Directive (WFD) monitoring lakes and regular macrophyte surveys are conducted by the Environmental Protection Agency (EPA)
Vegetation composition: characteristic zonation	Occurrence	All characteristic zones should be present, correctly distributed and in good condition	Further work is necessary to describe the characteristic zonation and other spatial patterns in lake habitat 3110 (see O Connor, 2015)
Vegetation distribution: maximum depth	Metres	Maintain maximum depth of vegetation, subject to natural processes	The maximum depth of vegetation is likely to be specific to the lake shoreline in question. Further work is necessary to develop indicative targets for lake habitat 3110. Maximum depth should be large in lakes in The Twelve Bens/Garraun Complex SAC within undisturbed peatland and uplands; however, pressures such as overgrazing, forestry and peat-cutting may have reduced vegetation depth in some lakes
Hydrological regime: water level fluctuations	Metres	Maintain appropriate natural hydrological regime necessary to support the habitat	Fluctuations in lake water level are typical in Ireland, but can be amplified by activities such as abstraction, drainage and overgrazing. Increased water level fluctuations can increase wave action, up-root vegetation, increase turbidity, alter the substratum and lead to release of nutrients from the sediment. The hydrological regime of the lakes must be maintained so that the area, distribution and depth of the lake habitat and its constituent/characteristic vegetation zones and communities are not reduced. It is possible that the hydrological regimes of some of the lakes have been altered by historic overgrazing (faster run-off, higher flood peaks, lower base flows, etc.; see Marshall et al., 2014)

Lake substratum quality	Various	Maintain appropriate substratum type, extent and chemistry to support the vegetation	Research is required to further characterise the substratum types (particle size and origin) and substratum quality (notably pH, calcium, iron and nutrient concentrations) favoured by each of the five Annex I lake habitats in Ireland. It is likely that lake habitat 3110 is associated with a range of nutrient-poor substrates, from stones, cobble and gravel, through sands, silt, clay and peat. Substratum particle size is likely to vary with depth and along the shoreline within a single lake
Water quality: transparency	Metres	Maintain appropriate Secchi transparency. There should be no decline in Secchi depth/transparency	Transparency relates to light penetration and, hence, to the depth of colonisation of vegetation. It can be affected by phytoplankton blooms, water colour and turbidity. Specific targets have yet to be established for lake habitat 3110 (O Connor, 2015). Habitat 3110 is associated with very clear water. The OECD fixed boundary system set transparency targets for oligotrophic lakes of $\geq 6\text{m}$ annual mean Secchi disk depth, and $\geq 3\text{m}$ annual minimum Secchi disk depth. Free et al. (2009) found high isoetid abundance in lakes with Secchi depths of more than 3m. Heuff (1984) recorded transparency greater than lake depth (1.75m) in Pollacappul and 2.2m in Kylemore. Roden (2004) stated transparency in Pollacappul was excellent. Free et al. (2006) give Secchi depth of 4.1m in Kylemore
Water quality: nutrients	$\mu\text{g/l P}$; mg/l N	Maintain the concentration of nutrients in the water column at sufficiently low levels to support the habitat and its typical species	As a nutrient-poor habitat, oligotrophic and WFD 'high' status targets apply. Where a lake has nutrient concentrations that are lower than these targets, there should be no decline within class, i.e. no upward trend in nutrient concentrations. For lake habitat 3110, annual average total phosphorus (TP) concentration should be $\leq 10\mu\text{g/l TP}$, average annual total ammonia concentration should be $\leq 0.040\text{mg/l N}$ and annual 95th percentile for total ammonia should be $\leq 0.090\text{mg/l N}$. See also The European Communities Environmental Objectives (Surface Waters) Regulations 2009. Derryclare, Kylemore and Pollacappul all had high nutrient status in 2007-09 and 2010-12 (McGarrigle et al., 2010; Bradley et al., 2015)
Water quality: phytoplankton biomass	$\mu\text{g/l Chlorophyll } a$	Maintain appropriate water quality to support the habitat, including high chlorophyll <i>a</i> status	Oligotrophic and WFD 'high' status targets apply to lake habitat 3110. Where a lake has a chlorophyll <i>a</i> concentration that is lower than this target, there should be no decline within class, i.e. no upward trend in phytoplankton biomass. The average growing season (March-October) chlorophyll <i>a</i> concentration must be $< 5.8\mu\text{g/l}$. The annual average chlorophyll <i>a</i> concentration should be $< 2.5\mu\text{g/l}$ and the annual peak chlorophyll <i>a</i> concentration should be $\leq 8.0\mu\text{g/l}$. See also The European Communities Environmental Objectives (Surface Waters) Regulations 2009. Derryclare, Kylemore and Pollacappul all had high chlorophyll <i>a</i> status in 2007-09 and 2010-12 (McGarrigle et al., 2010; Bradley et al., 2015)
Water quality: phytoplankton composition	EPA phytoplankton composition metric	Maintain appropriate water quality to support the habitat, including high phytoplankton composition status	The EPA has developed a phytoplankton composition metric for nutrient enrichment of Irish lakes. As for other water quality indicators, lake habitat 3110 requires WFD high status. Kylemore had high phytoplankton composition status in 2010-12 (Bradley et al., 2015)
Water quality: attached algal biomass	Algal cover and EPA phytobenthos metric	Maintain trace/absent attached algal biomass ($< 5\%$ cover) and high phytobenthos status	Nutrient enrichment can favour epiphytic and epipelic algae that can out-compete the submerged vegetation. The cover abundance of attached algae in lake habitat 3110 should, therefore, be trace/absent ($< 5\%$ cover). EPA phytobenthos can be used as an indicator of changes in attached algal biomass. As for other water quality indicators, habitat 3110 requires high phytobenthos status. Kylemore had high phytobenthos status in 2010-12 (Bradley et al., 2015)

Water quality: macrophyte status	EPA macrophyte metric (The Free Index)	Maintain/restore high macrophyte status	Nutrient enrichment can favour more competitive submerged macrophyte species that out-compete the typical and characteristic species for the lake habitat. The EPA monitors macrophyte status for WFD purposes using the 'Free Index'. The target for lake habitat 3110 is high status or an Ecological Quality Ratio (EQR) for lake macrophytes of ≥ 0.90 , as defined in Schedule Five of the European Communities Environmental Objectives (Surface Waters) Regulations 2009. Kylemore Lough failed the target in 2007-2009 and 2010-2012, having good macrophyte status (McGarrigle et al., 2010; Bradley et al., 2015). Derryclare and Pollacappul had high macrophyte status in both periods
Acidification status	pH units; mg/l	Maintain appropriate water and sediment pH, alkalinity and cation concentrations to support the habitat, subject to natural processes	Acidification can impact on species abundance and composition in soft water lake habitats. In Europe, acidification of isoetid lakes can lead to loss of isoetids and dominance by submerged <i>Sphagnum</i> mosses and <i>Juncus bulbosus</i> (Arts, 2002). The specific requirements of lake habitat 3110, in terms of water and sediment pH, alkalinity and cation concentration, have not been determined. For lake habitat 3110, and adopting a precautionary approach based on Arts (2002), minimum pH should not be < 5.5 pH units. Maximum pH should be < 9.0 pH units, in line with the surface water standards established for soft waters (where water hardness is ≤ 100 mg/l calcium carbonate). See Schedule Five of the European Communities Environmental Objectives (Surface Waters) Regulations 2009. See McGarrigle et al. (2010) and Bradley et al. (2015) for acidification status for the WFD monitoring lakes in the SAC in the 2007-09 and 2010-12 periods
Water colour	mg/l PtCo	Maintain appropriate water colour to support the habitat	Increased water colour and turbidity decrease light penetration and can reduce the area of available habitat for lake macrophytes, particularly at the lower euphotic depths. The primary source of increased water colour in Ireland is disturbance to peatland. No habitat-specific or national standards for water colour currently exist. Studies have shown median colour concentrations in Irish lakes of 38mg/l PtCo (Free et al., 2000) and 33mg/l PtCo (Free et al., 2006). It is likely that the water colour in all Irish lake habitats would naturally be < 50 mg/l PtCo. Water colour can be very low (< 20 mg/l PtCo or even < 10 mg/l PtCo) in lake habitat 3110, where the peatland in the lake's catchment is intact. Free et al. (2006) reported colour of 27mg/l PtCo in Derryclare, 25mg/l PtCo in Fee, 21mg/l PtCo in Inagh and 23mg/l PtCo in Kylemore. Overgrazing and other peatland degradation may have increased colour in some lakes in the SAC
Dissolved organic carbon (DOC)	mg/l	Maintain appropriate organic carbon levels to support the habitat	Dissolved (and particulate) organic carbon (OC) in the water column is linked to water colour and acidification (organic acids). Increasing DOC in water has been documented across the Northern Hemisphere, including afforested peatland catchments in Ireland. Damage and degradation of peatland, leading to decomposition of peat is likely to be the predominant source of OC in Ireland. OC in water promotes decomposition by fungi and bacteria that, in turn, releases dissolved nutrients. The increased biomass of decomposers can also impact directly on the characteristic lake communities through shading, competition, etc. Overgrazing and other peatland degradation is also likely to have increased DOC in some lakes in The Twelve Bens/Garraun Complex SAC

Turbidity	Nephelometric turbidity units/ mg/l SS/ other appropriate units	Maintain appropriate turbidity to support the habitat	Turbidity can significantly affect the quantity and quality of light reaching rooted and attached vegetation and can, therefore, impact on lake habitats. The settlement of higher loads of inorganic or organic material on lake vegetation communities may also have impacts on sensitive, delicate species. Turbidity can increase as a result of re-suspension of material within the lake, higher loads entering the lake, or eutrophication. Turbidity measurement and interpretation is challenging. As a result, it is likely to be difficult to set habitat-specific targets for turbidity in lakes. Increased loads of fine organic and inorganic particles from overgrazing may have increased turbidity in lakes in the SAC
Fringing habitat: area and condition	Hectares	Maintain the area and condition of fringing habitats necessary to support the natural structure and functioning of habitat 3110	Most lake shorelines have fringing habitats of reedswamp, other swamp, fen, marsh or wet woodland that intergrade with and support the structure and functions of the lake habitat. In The Twelve Bens/Garraun Complex SAC, lake shorelines are likely to have acid grassland, swamp, heath, blanket bog and rock communities. Fringing habitats are dependent on the lake, particularly its water levels, and support wetland communities and species of conservation concern. Many of the fringing wetland habitats support higher invertebrate and plant species richness than the lake habitats themselves

3130 Oligotrophic to mesotrophic standing waters with vegetation of the Littorelletea uniflorae and/or Isoeto-Nanojuncetea

To maintain the favourable conservation condition of Oligotrophic to mesotrophic standing waters with vegetation of the Littorelletea uniflorae and/or of the Isoeto-Nanojuncetea in The Twelve Bens/Garraun Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Lake habitat 3130 is considered likely to occur in Pollacappul Lough, and possibly also Kylemore Lough in The Twelve Bens/Garraun Complex SAC (see map 3). In both lakes, habitat 3130 is likely to co-occur with lake habitat 3110. Two measures of extent should be used: 1. the area of the lake itself and; 2. the extent of the vegetation communities/zones that typify the habitat. Further information relating to all attributes is provided in the lake habitats supporting document for the purposes of site-specific conservation objectives and Article 17 reporting (O Connor, 2015) and The Twelve Bens/Garraun Complex SAC conservation objectives supporting document for <i>Najas flexilis</i> (1833)
Habitat distribution	Occurrence	No decline, subject to natural processes	The characteristics and distribution of lake habitat 3130 in Ireland are not yet fully understood. The Annex II macrophyte <i>Najas flexilis</i> (slender naiad) is considered to be characteristic of the habitat and occurs in Pollacappul and Kylemore Loughs. As noted above, Pollacappul and Kylemore have been mapped as potential 3130, but the habitat may be more widespread in the SAC (see map 3). See O Connor (2015) and The Twelve Bens/Garraun Complex SAC conservation objectives supporting document for <i>Najas flexilis</i> for further information
Typical species	Occurrence	Typical species present, in good condition, and demonstrating typical abundances and distribution	For lists of typical plant species, see the Article 17 habitat assessment for 3130 (NPWS, 2013), O Connor (2015) and The Twelve Bens/Garraun Complex SAC <i>Najas flexilis</i> conservation objectives supporting document. Kylemore and Pollacappul Loughs are Water Framework Directive (WFD) monitoring lakes and regular macrophyte surveys are conducted by the Environmental Protection Agency (EPA)
Vegetation composition: characteristic zonation	Occurrence	All characteristic zones should be present, correctly distributed and in good condition	Further work is necessary to describe the characteristic zonation and other spatial patterns in lake habitat 3130 (see O Connor, 2015)
Vegetation distribution: maximum depth	Metres	Maintain maximum depth of vegetation, subject to natural processes	The maximum depth of vegetation is likely to be specific to the lake shoreline in question. Further work is necessary to develop indicative targets for lake habitat 3130. Pollacappul is a relatively shallow lake, Kylemore is deeper. Maximum depth should be large in lakes in the SAC within undisturbed peatland and uplands; however, pressures such as overgrazing, forestry and peat-cutting may have reduced vegetation depth in some lakes

Hydrological regime: water level fluctuations	Metres	Maintain appropriate natural hydrological regime necessary to support the habitat	Fluctuations in lake water level are typical in Ireland, but can be amplified by activities such as abstraction, drainage and overgrazing. Increased water level fluctuations can increase wave action, up-root vegetation, increase turbidity, alter the substratum and lead to release of nutrients from the sediment. The hydrological regime of the lakes must be maintained so that the area, distribution and depth of the lake habitat and its constituent/characteristic vegetation zones and communities are not reduced. It is possible that the hydrological regimes of some of the lakes have been altered by historic overgrazing (faster run-off, higher flood peaks, lower base flows, etc.; see Marshall et al., 2014)
Lake substratum quality	Various	Maintain appropriate substratum type, extent and chemistry to support the vegetation	Research is required to further characterise the substratum types (particle size and origin) and substratum quality (notably pH, calcium, iron and nutrient concentrations) favoured by each of the five Annex I lake habitats in Ireland. It is likely that lake habitat 3130 is associated with a range of substrate types that are more productive/base-rich relative to the substratum of lake habitat 3110. Substratum particle size is likely to vary with depth and along the shoreline within a single lake; however, it should be noted that <i>Najas flexilis</i> is typically found on soft substrata of mud, silt or fine sand (Preston and Croft, 2001; Roden, 2002, 2004). For further information see the lake habitats supporting document (O Connor, 2015) and the <i>Najas flexilis</i> supporting document
Water quality: transparency	Metres	Maintain appropriate Secchi transparency. There should be no decline in Secchi depth/transparency	Transparency relates to light penetration and, hence, to the depth of colonisation of vegetation. It can be affected by phytoplankton blooms, water colour and turbidity. Specific targets have yet to be established for lake habitat 3130 (O Connor, 2015). Habitat 3130 is associated with clear water, as evidenced by the growth of the character species <i>Najas flexilis</i> at depths of up to 10m. There is likely to be some variation in Secchi depth across lakes with habitat 3130, and site-specific conditions should also be considered. The OECD fixed boundary system set transparency targets for oligotrophic lakes of $\geq 6\text{m}$ annual mean Secchi disk depth, and $\geq 3\text{m}$ annual minimum Secchi disk depth. Heuff (1984) recorded transparency greater than lake depth (1.75m) in Pollacappul and 2.2m in Kylemore. Roden (2004) stated transparency in Pollacappul was excellent. Free et al. (2006) give Secchi depth of 4.1m in Kylemore
Water quality: nutrients	$\mu\text{g/l P}$; mg/l N	Maintain the concentration of nutrients in the water column at sufficiently low levels to support the habitat and its typical species	Lake habitat 3130 is associated with high water quality, with naturally low dissolved nutrients. It is naturally more productive than lake habitat 3110, probably reflecting higher concentrations of nutrients such as calcium, rather than P alone. Lake habitat 3130 may reach favourable condition slightly above the oligotrophic boundary for nutrients, but in the absence of habitat-specific targets, the targets are WFD 'High Status' or oligotrophic (OECD, 1982). The "good-moderate" boundary is too enriched to support the habitat. Annual average total phosphorus (TP) concentration should be $\leq 10\mu\text{g/l}$ TP, average annual total ammonia concentration should be $\leq 0.040\text{mg/l N}$ and annual 95th percentile for total ammonia should be $\leq 0.090\text{mg/l N}$. See also The European Communities Environmental Objectives (Surface Waters) Regulations 2009. Kylemore and Pollacappul had high nutrient status in 2007-09 and 2010-12 (McGarrigle et al., 2010; Bradley et al., 2015)

Water quality: phytoplankton biomass	µg/l Chlorophyll <i>a</i>	Maintain appropriate water quality to support the habitat, including high chlorophyll <i>a</i> status	Lake habitat 3130 is associated with high water quality, and naturally low algal growth. As for nutrients, the targets are WFD 'High Status' or oligotrophic (OECD, 1982). The "good-moderate" boundary is too enriched to support the habitat. The average growing season (March-October) chlorophyll <i>a</i> concentration must be <5.8µg/l. The annual average chlorophyll <i>a</i> concentration should be <2.5µg/l and the annual peak chlorophyll <i>a</i> concentration should be ≤8.0µg/l. Where a lake has a chlorophyll <i>a</i> concentration that is lower than this target, there should be no decline within class, i.e. no upward trend in phytoplankton biomass. See also The European Communities Environmental Objectives (Surface Waters) Regulations 2009. Kylemore and Pollacappul had high chlorophyll <i>a</i> status in 2007-09 and 2010-12 (McGarrigle et al., 2010; Bradley et al., 2015)
Water quality: phytoplankton composition	EPA phytoplankton composition metric	Maintain appropriate water quality to support the habitat, including high phytoplankton composition status	The EPA has developed a phytoplankton composition metric for nutrient enrichment of Irish lakes. As for other water quality indicators, lake habitat 3130 requires WFD high status. Kylemore had high phytoplankton composition status in 2010-12 (Bradley et al., 2015)
Water quality: attached algal biomass	Algal cover and EPA phytobenthos metric	Maintain trace/absent attached algal biomass (<5% cover) and high phytobenthos status	Nutrient enrichment can favour epiphytic and epipelic algae that can out-compete the submerged vegetation. The cover abundance of attached algae in lake habitat 3130 should, therefore, be trace/absent (<5% cover). EPA phytobenthos can be used as an indicator of changes in attached algal biomass. As for other water quality indicators, lake habitat 3130 requires high phytobenthos status. Kylemore had high phytobenthos status in 2010-12 (Bradley et al., 2015)
Water quality: macrophyte status	EPA macrophyte metric (The Free Index)	Maintain/restore high macrophyte status	Nutrient enrichment can favour more competitive submerged macrophyte species that out-compete the typical and characteristic species for the lake habitat. The EPA monitors macrophyte status for WFD purposes using the 'Free Index'. The target for lake habitat 3130 is high status or an Ecological Quality Ratio (EQR) for lake macrophytes of ≥0.90, as defined in Schedule Five of the European Communities Environmental Objectives (Surface Waters) Regulations 2009. Kylemore failed the target in 2007-2009 and 2010-2012, having good macrophyte status (McGarrigle et al., 2010; Bradley et al., 2015). Pollacappul had high macrophyte status in both periods
Acidification status	pH units; mg/l	Maintain appropriate water and sediment pH, alkalinity and cation concentrations to support the habitat, subject to natural processes	Acidification can impact on species abundance and composition in lake habitat 3130. Acidification reduces the abundance and reproductive capacity of <i>Najas flexilis</i> (Wingfield et al., 2004). The specific requirements of lake habitat 3130, in terms of water and sediment pH, alkalinity and cation concentration, have not been determined. In line with targets for <i>Najas flexilis</i> , median pH values should be greater than 7 pH units. Water and sediment alkalinity and concentrations of cations (notably calcium) should be appropriate to the habitat. The target for WFD Acidification/Alkalisiation status is high. Maximum pH should be <9.0 pH units, in line with the surface water standards. See Schedule Five of the European Communities Environmental Objectives (Surface Waters) Regulations 2009. See McGarrigle et al. (2010) and Bradley et al. (2015) for acidification status for the WFD monitoring lakes in the SAC in the 2007-09 and 2010-12 periods

Water colour	mg/l PtCo	Maintain appropriate water colour to support the habitat	Increased water colour and turbidity decrease light penetration and can reduce the area of available habitat for lake macrophytes, particularly at the lower euphotic depths. The primary source of increased water colour in Ireland is disturbance to peatland. No habitat-specific or national standards for water colour currently exist. Studies have shown median colour concentrations in Irish lakes of 38mg/l PtCo (Free et al., 2000) and 33mg/l PtCo (Free et al., 2006). It is likely that the water colour in all Irish lake habitats would naturally be <50mg/l PtCo. Water colour is generally <30mg/l PtCo or, more naturally, <20mg/l PtCo in lakes with habitat 3130, where the peatland in the lake's catchment is intact. Free et al. (2006) reported colour of 23mg/l PtCo in Kylemore. Overgrazing and other peatland degradation may have increased colour in some lakes in The Twelve Bens/Garraun Complex SAC
Dissolved organic carbon (DOC)	mg/l	Maintain appropriate organic carbon levels to support the habitat	Dissolved (and particulate) organic carbon (OC) in the water column is linked to water colour and acidification (organic acids). Increasing DOC in water has been documented across the Northern Hemisphere, including afforested peatland catchments in Ireland. Damage and degradation of peatland, leading to decomposition of peat is likely to be the predominant source of OC in Ireland. OC in water promotes decomposition by fungi and bacteria that, in turn, releases dissolved nutrients. The increased biomass of decomposers can also impact directly on the characteristic lake communities through shading, competition, etc. Overgrazing and other peatland degradation is also likely to have increased DOC in some lakes in the SAC
Turbidity	Nephelometric turbidity units/ mg/l SS/ other appropriate units	Maintain appropriate turbidity to support the habitat	Turbidity can significantly affect the quantity and quality of light reaching rooted and attached vegetation and can, therefore, impact on lake habitats. The settlement of higher loads of inorganic or organic material on lake vegetation communities may also have impacts on sensitive, delicate species. Turbidity can increase as a result of re-suspension of material within the lake, higher loads entering the lake, or eutrophication. Turbidity measurement and interpretation is challenging. As a result, it is likely to be difficult to set habitat-specific targets for turbidity in lakes. Increased loads of fine organic and inorganic particles from overgrazing may have increased turbidity in lakes in the SAC
Fringing habitat: area and condition	Hectares	Maintain the area and condition of fringing habitats necessary to support the natural structure and functioning of habitat 3130	Most lake shorelines have fringing habitats of reedswamp, other swamp, fen, marsh or wet woodland that intergrade with and support the structure and functions of the lake habitat. In The Twelve Bens/Garraun Complex SAC, lake shorelines are likely to have acid grassland, swamp, heath, blanket bog and rock communities. Fringing habitats are dependent on the lake, particularly its water levels, and support wetland communities and species of conservation concern. Many of the fringing wetland habitats support higher invertebrate and plant species richness than the lake habitats themselves

Conservation Objectives for : The Twelve Bens/Garraun Complex SAC [002031]

4060 Alpine and Boreal heaths

To restore the favourable conservation condition of Alpine and Boreal heaths in The Twelve Bens/Garraun Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Alpine and Boreal heaths habitat has not been mapped in detail for The Twelve Bens/Garraun Complex SAC, but from current available data the total area of the qualifying habitat is estimated to be approximately 225ha, covering c.1% of the SAC (NPWS internal files). Further details on this and the following attributes can be found in The Twelve Bens/Garraun Complex SAC conservation objectives supporting document for blanket bogs and associated habitats
Habitat distribution	Occurrence	No decline, subject to natural processes	The habitat occurs at high altitudes throughout the SAC, and is associated with exposed rock and scree. It is present on several hillsides and summits including Benglenisky, Benbaun, Muckanaght, Bencullagh, Maumonght, Benbrack, Knockbrack, Bengooria, Doughruagh and Altnagaighera (NPWS internal files). Further information can be found within NPWS internal files and the blanket bogs and associated habitats supporting document
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil nutrient status within natural range	See the blanket bogs and associated habitats supporting document for further details
Community diversity	Abundance of variety of vegetation communities	Maintain variety of vegetation communities, subject to natural processes	A variety of Alpine and Boreal heath vegetation communities have been recorded in this SAC (NPWS internal files; R. Hodd, pers. comm.), two of which correspond to communities recorded in the National Survey of Upland Habitats and listed in the provisional list of vegetation communities described in Perrin et al. (2014). Further information on vegetation communities associated with this habitat is presented in Perrin et al. (2014)
Vegetation composition: lichens and bryophytes	Number of species at a representative number of 2m x 2m monitoring stops	Number of bryophyte or non-crustose lichen species present at each monitoring stop is at least three	Attribute and target based on Perrin et al. (2014)
Vegetation composition: positive indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of positive indicator species at least 66%	Attribute and target based on Perrin et al. (2014), where the list of positive indicator species for this habitat is also presented
Vegetation composition: dwarf shrub species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of dwarf shrub species at least 10%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: negative indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of negative indicator species less than 10%	Attribute and target based on Perrin et al. (2014), where the list of negative indicator species for this habitat is also presented
Vegetation composition: non-native species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of non-native species less than 1%	Attribute and target based on Perrin et al. (2014)
Vegetation structure: signs of grazing	Percentage of leaves grazed at a representative number of 2m x 2m monitoring stops	Less than 10% collectively of the live leaves of specific graminoids showing signs of grazing	Attribute and target based on Perrin et al. (2014). See the blanket bogs and associated habitats supporting document for the list of specific graminoids

Vegetation structure: signs of browsing	Percentage of shoots browsed at a representative number of 2m x 2m monitoring stops	Less than 33% collectively of the last complete growing season's shoots of ericoids and crowberry (<i>Empetrum nigrum</i>) showing signs of browsing	Attribute and target based on Perrin et al. (2014)
Vegetation structure: burning	Occurrence in local vicinity of a representative number of monitoring stops	No signs of burning within the habitat	Attribute and target based on Perrin et al. (2014)
Physical structure: disturbed bare ground	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of disturbed bare ground less than 10%	Attribute and target based on Perrin et al. (2014)
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat and no decline in status of hepatic mats associated with this habitat	This includes species listed in the Flora (Protection Order, 2015 (FPO) and/or the red data lists (Lockhart et al., 2012; Wyse Jackson et al., 2016). The Near Threatened alpine clubmoss (<i>Diphasiastrum alpinum</i>) (Wyse Jackson et al., 2016) is present within the SAC (NPWS internal files). This species is known to be associated with this habitat type. The FPO listed and Vulnerable bryophytes <i>Adelanthus lindenbergianus</i> and <i>Bazzania pearsonii</i> , the FPO and Near Threatened <i>Mastigophora woodsii</i> and the FPO and Endangered <i>Plagiochila carringtonii</i> (Lockhart et al., 2012) are all associated with this habitat (NPWS internal files). Hepatic mats are present in this habitat in the SAC (R. Hodd, pers. comm.)

Conservation Objectives for : The Twelve Bens/Garraun Complex SAC [002031]

7130 Blanket bogs (* if active bog)

To restore the favourable conservation condition of Blanket bogs (* if active bog) in The Twelve Bens/Garraun Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Blanket bog has not been mapped in detail for The Twelve Bens/Garraun Complex SAC, but from current available data the total area of the qualifying habitat is estimated to be approximately 5,325ha, covering 33% of the SAC (NPWS internal files). Further details on this and the following attributes can be found in The Twelve Bens/Garraun Complex SAC conservation objectives supporting document for blanket bogs and associated habitats
Habitat distribution	Occurrence	No decline, subject to natural processes	The habitat is documented to occur throughout the SAC. It is best developed in low-lying basins and is present on the flanks of mountains up to an altitude of c.200m. Examples of this habitat can be found at Glenmore, Kylemore, Tooreenacoona, Owenglin, east of the Dawros River, on the western slopes of Bealnascarpa Mountain and on the flanks of Cregg Hill (NPWS internal files). Further information can be found in Douglas and Grogan (1987), Douglas et al. (1989), NPWS internal files and the blanket bogs and associated habitats supporting document
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil nutrient status within natural range	See the blanket bogs and associated habitats supporting document for further details
Ecosystem function: peat formation	Active blanket bog as a proportion of the total area of Annex I blanket bog habitat	At least 99% of the total Annex I blanket bog area is active	See the blanket bogs and associated habitats supporting document for further details
Ecosystem function: hydrology	Flow direction, water levels, occurrence of drains and erosion gullies	Natural hydrology unaffected by drains and erosion	Further details and a brief discussion of restoration potential is presented in the blanket bogs and associated habitats supporting document
Community diversity	Abundance of variety of vegetation communities	Maintain variety of vegetation communities, subject to natural processes	A variety of blanket bog vegetation communities have been recorded in this SAC (Douglas and Grogan, 1987; Douglas et al., 1989; NPWS internal files), six of which correspond to communities recorded in the National Survey of Upland Habitats and listed in the provisional list of vegetation communities described in Perrin et al. (2014). Further information on vegetation communities associated with this habitat is presented in Perrin et al. (2014)
Vegetation composition: positive indicator species	Number of species at a representative number of 2m x 2m monitoring stops	Number of positive indicator species present at each monitoring stop is at least seven	Attribute and target based on Perrin et al. (2014), where the list of positive indicator species for this habitat is also presented
Vegetation composition: lichens and bryophytes	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of bryophytes or lichens, excluding <i>Sphagnum fallax</i> , at least 10%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: potential dominant species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of each of the potential dominant species less than 75%	Attribute and target based on Perrin et al. (2014). See the blanket bogs and associated habitats supporting document for the list of potential dominant species
Vegetation composition: negative indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of negative indicator species less than 1%	Attribute and target based on Perrin et al. (2014), where the list of negative indicator species for this habitat is also presented

Vegetation composition: non-native species	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of non-native species less than 1%	Attribute and target based on Perrin et al. (2014). Rhododendron (<i>Rhododendron ponticum</i>) was recorded as encroaching on this habitat (Douglas et al., 1989; NPWS internal files)
Vegetation composition: native trees and shrubs	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of scattered native trees and shrubs less than 10%	Attribute and target based on Perrin et al. (2014)
Vegetation structure: <i>Sphagnum</i> condition	Condition at a representative number of 2m x 2m monitoring stops	Less than 10% of the <i>Sphagnum</i> cover is crushed, broken and/or pulled up	Attribute and target based on Perrin et al. (2014)
Vegetation structure: signs of browsing	Percentage of shoots browsed at a representative number of 2m x 2m monitoring stops	Last complete growing season's shoots of ericoids, crowberry (<i>Empetrum nigrum</i>) and bog-myrtle (<i>Myrica gale</i>) showing signs of browsing collectively less than 33%	Attribute and target based on Perrin et al. (2014)
Vegetation structure: burning	Occurrence in local vicinity of a representative number of monitoring stops	No signs of burning in sensitive areas, into the moss, liverwort or lichen layer or exposure of peat surface due to burning	Attribute and target based on Perrin et al. (2014), where the list of sensitive areas for this habitat is also presented
Physical structure: disturbed bare ground	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of disturbed bare ground less than 10%	Attribute and target based on Perrin et al. (2014)
Physical structure: drainage	Percentage area in local vicinity of a representative number of monitoring stops	Area showing signs of drainage from heavy trampling, tracking or ditches less than 10%	Attribute and target based on Perrin et al. (2014)
Physical structure: erosion	Percentage area in local vicinity of a representative number of monitoring stops	Less than 5% of the greater bog mosaic comprises erosion gullies and eroded areas	Attribute and target based on Perrin et al. (2014)
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat	This includes species listed in the Flora (Protection) Order, 2015 (FPO) and/or the red data lists (Lockhart et al., 2012; Wyse Jackson et al., 2016). The FPO listed and Vulnerable marsh clubmoss (<i>Lycopodiella inundata</i>) (Wyse Jackson et al., 2016) is present within the SAC (NPWS internal files), but this species cannot be specifically assigned to this habitat

7150 Depressions on peat substrates of the Rhynchosporion

To restore the favourable conservation condition of Depressions on peat substrates of the Rhynchosporion in The Twelve Bens/Garraun Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Depressions on peat substrates of the Rhynchosporion has not been mapped in detail for The Twelve Bens/Garraun Complex SAC and thus the total area of the qualifying habitat in the SAC is unknown. Further details on this and the following attributes can be found in The Twelve Bens/Garraun Complex SAC conservation objectives supporting document for blanket bogs and associated habitats
Habitat distribution	Occurrence	No decline, subject to natural processes	The habitat is documented to occur around pools, in wet hollows, quaking areas and in flushed areas within the SAC (NPWS internal files). Examples of this habitat can be found at Tooreenacoona, Glenmore, Kylemore and Owenglin. Further information can be found within Douglas and Grogan (1987), Douglas et al. (1989), NPWS internal files and the blanket bogs and associated habitats supporting document
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil nutrient status within natural range	See the blanket bogs and associated habitats supporting document for further details
Vegetation composition: positive indicator species	Number of species at a representative number of 2m x 2m monitoring stops	Number of positive indicator species at each monitoring stop is at least five	Attribute and target based on Perrin et al. (2014), where the list of positive indicator species for this habitat is also presented
Vegetation composition: <i>Rhynchospora</i> spp.	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of white beaked sedge (<i>Rhynchospora alba</i>) and brown beaked sedge (<i>R. fusca</i>) at least 10%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: potential dominant species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of each of the potential dominant species individually less than 35%	Attribute and target based on Perrin et al. (2014). See the blanket bogs and associated habitats supporting document for the list of potential dominant species
Vegetation composition: negative indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of negative indicator species less than 1%	Attribute and target based on Perrin et al. (2014), where the list of negative indicator species for this habitat is also presented
Vegetation composition: non-native species	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of non-native species less than 1%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: native trees and shrubs	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of scattered native trees and shrubs less than 10%	Attribute and target based on Perrin et al. (2014)
Vegetation structure: <i>Sphagnum</i> condition	Condition at a representative number of 2m x 2m monitoring stops	Less than 10% of the <i>Sphagnum</i> cover is crushed, broken and/or pulled up	Attribute and target based on Perrin et al. (2014)
Vegetation structure: signs of browsing	Percentage of shoots browsed at a representative number of 2m x 2m monitoring stops	Last complete growing season's shoots of ericoids, crowberry (<i>Empetrum nigrum</i>) and bog-myrtle (<i>Myrica gale</i>) showing signs of browsing collectively less than 33%	Attribute and target based on Perrin et al. (2014)

Vegetation structure: burning	Occurrence in local vicinity of a representative number of monitoring stops	No signs of burning in sensitive areas, into the moss, liverwort or lichen layer or exposure of peat surface due to burning	Attribute and target based on Perrin et al. (2014), where the list of sensitive areas for this habitat is also presented
Physical structure: disturbed bare ground	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of disturbed bare ground less than 10%	Attribute and target based on Perrin et al. (2014)
Physical structure: drainage	Percentage area in local vicinity of a representative number of monitoring stops	Area showing signs of drainage from heavy trampling, tracking or ditches less than 10%	Attribute and target based on Perrin et al. (2014)
Physical structure: erosion	Percentage area in local vicinity of a representative number of monitoring stops	Less than 5% of the greater bog mosaic comprises erosion gullies and eroded areas	Attribute and target based on Perrin et al. (2014)
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat	This includes species listed in the Flora (Protection) Order, 2015 (FPO) and/or the red data lists (Lockhart et al., 2012; Wyse Jackson et al., 2016). The Near Threatened brown beak-sedge (<i>Rhynchospora fusca</i>) (Wyse Jackson et al., 2016) can be specifically assigned to this habitat in the SAC (Douglas et al., 1989). The FPO and Vulnerable marsh clubmoss (<i>Lycopodiella inundata</i>) (Wyse Jackson et al., 2016) is present within the SAC (NPWS internal files), but this species cannot be specifically assigned to this habitat

Conservation Objectives for : The Twelve Bens/Garraun Complex SAC [002031]

8110 Siliceous scree of the montane to snow levels (*Androsacetalia alpinae* and *Galeopsietalia ladani*)

To restore the favourable conservation condition of Siliceous scree of the montane to snow levels (*Androsacetalia alpinae* and *Galeopsietalia ladani*) in The Twelve Bens/Garraun Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Siliceous scree of the montane to snow levels (<i>Androsacetalia alpinae</i> and <i>Galeopsietalia ladani</i>) has not been mapped in detail for The Twelve Bens/Garraun Complex SAC but from current available data the total area of the qualifying habitat is estimated to be approximately 162ha, covering 1% of the SAC (NPWS internal files). Further details on this and the following attributes can be found in The Twelve Bens/Garraun Complex SAC conservation objectives supporting document for blanket bogs and associated habitats
Habitat distribution	Occurrence	No decline, subject to natural processes	The habitat occurs on hillsides throughout the SAC. Example locations include Bencullagh, Muckanaght, Benbaun, Bengower, Derryclare and Doughruagh (NPWS internal files). Further information can be found within NPWS internal files and the blanket bogs and associated habitats supporting document
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil nutrient status within natural range	See the blanket bogs and associated habitats supporting document for further details
Vegetation composition: lichens and bryophytes	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of bryophytes and non-crustose lichen species at least 5%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: negative indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Proportion of vegetation composed of negative indicator species less than 1%	Attribute and target based on Perrin et al. (2014), where the list of negative indicator species for this habitat is also presented
Vegetation composition: non-native species	Percentage cover at a representative number of 2m x 2m monitoring stops	Proportion of vegetation composed of non-native species less than 1%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: positive indicator species	Number of species in local vicinity of a representative number of monitoring stops	At least one positive indicator species present in vicinity of each monitoring stop in block scree	Attribute and target based on Perrin et al. (2014). The list of positive indicator species for this habitat is also presented in Perrin et al. (2014) and is the same as for 8220 Siliceous rocky slopes
Vegetation composition: grass species and dwarf shrubs	Percentage cover in local vicinity of a representative number of monitoring stops	Total cover of grass species and dwarf shrubs less than 20%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: bracken, native trees and shrubs	Percentage cover in local vicinity of a representative number of monitoring stops	Total cover of bracken (<i>Pteridium aquilinum</i>), native trees and shrubs less than 25%	Attribute and target based on Perrin et al. (2014)
Vegetation structure: grazing and browsing	Percentage of leaves/shoots grazed/browsed at a representative number of 2m x 2m monitoring stops	Live leaves of forbs and shoots of dwarf shrubs showing signs of grazing or browsing collectively less than 50%	Attribute and target based on Perrin et al. (2014)
Physical structure: disturbance	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Ground disturbed by human and animal paths, scree running, vehicles less than 10%	Attribute and target based on Perrin et al. (2014)

Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat and no decline in status of hepatic mats associated with this habitat	This includes species listed in the Flora (Protection) Order, 2015 (FPO) and/or the red data lists (Lockhart et al., 2012; Wyse Jackson et al., 2016). The FPO listed and Vulnerable parsley fern (<i>Cryptogramma crispera</i>) (Wyse Jackson et al., 2016) and the Near Threatened beech fern (<i>Phegopteris connectilis</i>) (Wyse Jackson et al., 2016) have previously been recorded within the SAC (NPWS internal files), but these species cannot be assigned specifically to this habitat. Hepatic mats are present within this habitat in the SAC (R. Hodd, pers. comm.)
-------------------------------------	--------------------------------	--	--

Conservation Objectives for : The Twelve Bens/Garraun Complex SAC [002031]

8210 Calcareous rocky slopes with chasmophytic vegetation

To restore the favourable conservation condition of Calcareous rocky slopes with chasmophytic vegetation in The Twelve Bens/Garraun Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Calcareous rocky slopes with chasmophytic vegetation has not been mapped in detail for The Twelve Bens/Garraun Complex SAC, but from current available data the total area of the qualifying habitat is estimated to be approximately 105ha, covering c.1% of the SAC (NPWS internal files). Further details on this and the following attributes can be found in The Twelve Bens/Garraun Complex SAC conservation objectives supporting document for blanket bogs and associated habitats
Habitat distribution	Occurrence	No decline, subject to natural processes	The habitat occurs throughout the SAC. Locations where this habitat is present include the northern slopes of Muckanaght, west of Bengower, Benbaun, Barrlugwaum on the south-east slopes of Benchoona, Altnagaighera, Garraun, Doughruagh and Benlettery (NPWS internal files). Further information can be found within NPWS internal files and the blanket bogs and associated habitats supporting document
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil nutrient status within natural range	See the blanket bogs and associated habitats supporting document for further details
Vegetation composition: positive indicator fern and <i>Saxifraga</i> species	Number of species in local vicinity of a representative number of monitoring stops	Number of ferns and <i>Saxifraga</i> indicators at each monitoring stop is at least one	Attribute and target based on Perrin et al. (2014), where the list of positive indicator species for this habitat is also presented
Vegetation composition: positive indicator species	Number of species in local vicinity of a representative number of monitoring stops	Number of positive indicator species at each monitoring stop is at least three	Attribute and target based on Perrin et al. (2014), where the list of positive indicator species for this habitat is also presented
Vegetation composition: non-native species	Percentage cover in local vicinity of a representative number of monitoring stops	Proportion of vegetation composed of non-native species less than 1%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: bracken, native trees and shrubs	Percentage cover in local vicinity of a representative number of monitoring stops	Total cover of bracken (<i>Pteridium aquilinum</i>), native trees and shrubs less than 25%	Attribute and target based on Perrin et al. (2014)
Vegetation structure: grazing and browsing	Percentage of leaves/shoots grazed/browsed in local vicinity of a representative number of monitoring stops	Live leaves of forbs and shoots of dwarf shrubs showing signs of grazing or browsing collectively less than 50%	Attribute and target based on Perrin et al. (2014)
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat	This includes species listed in the Flora (Protection) Order, 2015 and/or the red data lists (Lockhart et al., 2012; Wyse Jackson et al., 2016). The Vulnerable alpine saw-wort (<i>Saussurea alpina</i>) and holly-fern (<i>Polystichum lonchitis</i>), and the Near Threatened beech fern (<i>Phegopteris connectilis</i>) and Irish saxifrage (<i>Saxifraga rosacea</i> subsp. <i>rosacea</i>) (Wyse Jackson et al., 2016) were recorded within this habitat in the SAC (NPWS internal files). The Near Threatened bryophytes <i>Mnium thomsonii</i> and <i>Eremonotus myriocarpus</i> (Lockhart et al., 2012) are present within the SAC (NPWS internal files) and are known to be associated with this habitat type

Conservation Objectives for : The Twelve Bens/Garraun Complex SAC [002031]

8220 Siliceous rocky slopes with chasmophytic vegetation

To restore the favourable conservation condition of Siliceous rocky slopes with chasmophytic vegetation in The Twelve Bens/Garraun Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Siliceous rocky slopes with chasmophytic vegetation has not been mapped in detail for The Twelve Bens/Garraun Complex SAC, but from current available data the total area of the qualifying habitat is estimated to be approximately 121ha, covering c.1% of the SAC (NPWS internal files). Further details on this and the following attributes can be found in The Twelve Bens/Garraun Complex SAC conservation objectives supporting document for blanket bogs and associated habitats
Habitat distribution	Occurrence	No decline, subject to natural processes	The habitat occurs throughout the SAC including the slopes of Bengower, Benbreen, Bencollaghduff, Bencorr, Bencorbeg, Derryclare, Bencullagh, Muckanaght, Benbaun, Doughruagh, Benchoona and Garraun (NPWS internal files). Further information can be found within NPWS internal files and the blanket bogs and associated habitats supporting document
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil nutrient status within natural range	See the blanket bogs and associated habitats supporting document for further details
Vegetation composition: positive indicator species	Number of species in local vicinity of a representative number of monitoring stops	At least one positive indicator species present in vicinity of each monitoring stop	Attribute and target based on Perrin et al. (2014). The list of positive indicator species for this habitat is also presented in Perrin et al. (2014) and is the same as for 8110 Siliceous screes
Vegetation composition: non-native species	Percentage cover in local vicinity of a representative number of monitoring stops	Proportion of vegetation composed of non-native species less than 1%	Attribute and target based on Perrin et al. (2014). New Zealand willowherb (<i>Epilobium brunnescens</i>) is present within this habitat in the SAC (NPWS internal files)
Vegetation composition: bracken, native trees and shrubs	Percentage cover in local vicinity of a representative number of monitoring stops	Total cover of bracken (<i>Pteridium aquilinum</i>), native trees and shrubs less than 25%	Attribute and target based on Perrin et al. (2014)
Vegetation structure: grazing and browsing	Percentage of leaves/shoots grazed/browsed in local vicinity of a representative number of monitoring stops	Live leaves of forbs and shoots of dwarf shrubs showing signs of grazing or browsing collectively less than 50%	Attribute and target based on Perrin et al. (2014)
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat and no decline in status of hepatic mats associated with this habitat	This includes species listed in the Flora (Protection) Order, 2015 (FPO) and/or the red data lists (Lockhart et al., 2012; Wyse Jackson et al., 2016). The Near Threatened beech fern (<i>Phegopteris connectilis</i>) (Wyse Jackson et al., 2016) and the Vulnerable bryophytes <i>Arctoa fulvella</i> , <i>Dicranodontium uncinatum</i> and <i>Lophozia opacifolia</i> (Lockhart et al., 2012) are present within the SAC (NPWS internal files). These species are known to be associated with this habitat type. The FPO listed and Vulnerable parsley fern (<i>Cryptogramma crispá</i>) (Wyse Jackson et al., 2016) has previously been recorded within the SAC (NPWS internal files), but it cannot be assigned specifically to this habitat. Hepatic mats are present in this habitat in the SAC (R. Hodd, pers. comm.)

Conservation Objectives for : The Twelve Bens/Garraun Complex SAC [002031]

91A0 Old sessile oak woods with *Ilex* and *Blechnum* in the British Isles

To maintain the favourable conservation condition of Old sessile oak woods with *Ilex* and *Blechnum* in the British Isles in The Twelve Bens/Garraun Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes; at least 9.5ha for the sub-site (Derryclare Wood, NSNW site code 1601) surveyed. See map 4 for surveyed area	The best examples of old sessile oak woods with <i>Ilex</i> and <i>Blechnum</i> in the British Isles in The Twelve Bens/Garraun Complex SAC occur at Kylemore and Derryclare Woods, but scattered fragments also occur near Salrock House at the head of Killary Bay Little, in steep ravines north of Ballinahinch Lake and on islands in Lough Inagh and Derryclare Lough (NPWS internal files). As part of the National Survey of Native Woodlands (NSNW), Derryclare Wood (NSNW site code 1601) was surveyed by Perrin et al. (2008). Derryclare Wood is also a statutory Nature Reserve (Statutory Instrument No. 177 of 1980) and is well-studied. Map 4 shows the woodland area surveyed by the NSNW, including the area classified as 91A0 (9.5ha). NB further unsurveyed areas are present within the SAC
Habitat distribution	Occurrence	No decline, subject to natural processes. The surveyed woodland location is shown on map 4	Distribution based on Perrin et al. (2008). It is important to note that there are additional areas of oak woodland, which were not mapped by the NSNW, present within the SAC
Woodland size	Hectares	Area stable or increasing. Where topographically possible, "large" woods at least 25ha in size and "small" woods at least 3ha in size	The target areas for individual woodlands aim to reduce habitat fragmentation and benefit those species requiring 'deep' woodland conditions (Peterken, 2002). In some cases, topographical constraints may restrict expansion
Woodland structure: cover and height	Percentage and metres	Diverse structure with a relatively closed canopy containing mature trees; subcanopy layer with semi-mature trees and shrubs; and well-developed herb layer	Described in Perrin et al. (2008) and NPWS internal files. See also Fahy et al. (1999)
Woodland structure: community diversity and extent	Hectares	Maintain diversity and extent of community types	Described in Perrin et al. (2008) and NPWS internal files. See also Ferguson and Westhoff (1987)
Woodland structure: natural regeneration	Seedling:sapling:pole ratio	Seedlings, saplings and pole age-classes occur in adequate proportions to ensure survival of woodland canopy	Oak (<i>Quercus petraea</i>) generally regenerates poorly. In suitable sites, ash (<i>Fraxinus excelsior</i>) can regenerate in large numbers although few seedlings reach pole size
Woodland structure: dead wood	m ³ per hectare; number per hectare	At least 30m ³ /ha of fallen timber greater than 10cm diameter; 30 snags/ha; both categories should include stems greater than 40cm diameter	Dead wood is a valuable resource and an integral part of a healthy, functioning woodland ecosystem
Woodland structure: veteran trees	Number per hectare	No decline	Mature and veteran trees are important habitats for bryophytes, lichens, saproxylic organisms and some bird species. Their retention is important to ensure continuity of habitats/niches and propagule sources

Woodland structure: indicators of local distinctiveness	Occurrence	No decline	Includes ancient or long-established woodlands (Perrin and Daly, 2010), archaeological and geological features as well as red data and other rare or localised species. Perrin and Daly (2010) identified Derryclare Wood (NSNW site code 1601) as possible ancient woodland. There is a well-developed fungus and lichen flora present (Folan and Mitchell, 1970). Many rare and protected bryophytes occur in the Kylemore woods, e.g. the Endangered liverwort <i>Cephalozia crassifolia</i> and the Near Threatened bryophytes <i>Lejeunea hibernica</i> , <i>L. eckloniana</i> , <i>Grimmia funalis</i> and <i>Telaranea europaea</i> (Lockhart et al., 2012). <i>Lejeunea hibernica</i> and <i>L. eckloniana</i> are listed on the Flora (Protection) Order, 2015
Vegetation composition: native tree cover	Percentage	No decline. Native tree cover not less than 95%	Species reported in Perrin et al. (2008) and NPWS internal files
Vegetation composition: typical species	Occurrence	A variety of typical native species present, depending on woodland type, including oak (<i>Quercus petraea</i>) and birch (<i>Betula pubescens</i>)	Species reported in Perrin et al. (2008) and NPWS internal files
Vegetation composition: negative indicator species	Occurrence	Negative indicator species, particularly non-native invasive species, absent or under control	The following are the most common non-native invasive species in this woodland type: beech (<i>Fagus sylvatica</i>), sycamore (<i>Acer pseudoplatanus</i>) and rhododendron (<i>Rhododendron ponticum</i>). The woods at Kylemore are infested with rhododendron. Beech is also present and substantial areas of the woods are underplanted with conifers (NPWS internal files)

Conservation Objectives for : The Twelve Bens/Garraun Complex SAC [002031]

1029 Freshwater Pearl Mussel *Margaritifera margaritifera*

To restore the favourable conservation condition of Freshwater Pearl Mussel in The Twelve Bens/Garraun Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Distribution	Kilometres	Maintain distribution at 6.43km. See map 5	The conservation objective applies to the Dawros freshwater pearl mussel (<i>Margaritifera margaritifera</i>) population in The Twelve Bens/Garraun Complex SAC, which is of international importance and one of eight Irish populations prioritised for conservation action (Moorkens, 2010; NPWS, 2010). It is listed on the European Communities Environmental Objectives (Freshwater Pearl Mussel) Regulations 2009 (Statutory Instrument No. 296 of 2009). The distribution and abundance of the species in the Dawros was mapped in 2003 and 2008 (Aster, 2003; Moorkens, 2008; Moorkens and Killeen, 2008). The species is known to extend from above Tullywee Bridge to the tidal limit downstream of Dawros Bridge; however, the stretch upstream of Tullywee Bridge has been little surveyed and requires further investigation. The target is for the species to be sufficiently widespread to maintain itself on a long-term basis as a viable component of the Dawros system. See NPWS (2010) for further information
Population size	Number of adult mussels	Restore Dawros population to at least 800,000 adult mussels	Aster (2003) estimated a population of between 10,750 and 19,322 per linear kilometre of the Dawros River. Moorkens (2008) estimated the total Dawros population at 800,000 to 1 million. Aster (2003) mapped the distribution and abundance from downstream of Tullywee Bridge to Dawros Bridge and categorised mussels as frequent or abundant throughout most of this length. The maximum density recorded in 2008 was 256 per square metre and an average abundance of 148 mussels per linear metre calculated from 9 samples (Moorkens, 2008; Moorkens and Killeen, 2008). NPWS (2013) in producing a national population estimate, assumed that all priority populations had declined at a rate of 1% per year from 2007 to 2012. The target is for the species to be sufficiently abundant to maintain itself on a long-term basis as a viable component of the Dawros system
Population structure: recruitment	Percentage per size class	Restore to at least 20% of population no more than 65mm in length; and at least 5% of population no more than 30mm in length	Mussels of no more than 65mm are considered 'young mussels' and may be found buried in the substratum and/or beneath adult mussels. Mussels of no more than 30mm are 'juvenile mussels' and are always buried in the substratum. See the European Communities Environmental Objectives (Freshwater Pearl Mussel) Regulations 2009. Size profile studies in the Dawros by Ross (1988) and Moorkens (2008) are summarised in NPWS (2010). In 2008, the smallest mussel was 22mm, and the Dawros failed both targets with 3.7% $\leq 65\text{mm}$ and 1.86% $\leq 30\text{mm}$ (Moorkens, 2008; Moorkens and Killeen, 2008). No juveniles were found in 2016, and 10% was $\leq 65\text{mm}$ (Moorkens, 2016). The Dawros population is unsustainable owing to lack of survival of juvenile mussels. The target is for sufficient juvenile recruitment to allow the species to maintain itself on a long-term basis as a viable component of the Dawros system

Population structure: adult mortality	Percentage	No more than 5% decline from previous number of live adults counted; dead shells less than 1% of the adult population and scattered in distribution	5% is considered the cut-off between the combined errors associated with natural fluctuations and sampling methods and evidence of true population decline. 1% of dead shells is considered to be indicative of natural losses. The Dawros passed both targets in 2008 (Moorkens, 2008; Moorkens and Killeen, 2008; NPWS, 2010). The occurrence of dead and moribund mussels in 2016 indicated an unnatural decline in adult numbers (Moorkens, 2016). The target is for sufficient survival of adults to allow the species to maintain itself on a long-term basis as a viable component of the Dawros system
Suitable habitat: extent	Kilometres	Restore suitable habitat in more than 4.8km in the Dawros system (see map 5) and any additional stretches necessary for salmonid spawning	The extent of the mussel habitat in the Dawros system downstream of Tullywee Bridge is well-documented and corresponds with its distribution (Aster, 2003; Moorkens, 2008; Moorkens and Killeen, 2008). Stretches upstream of Tullywee Bridge require further survey. Most of the mapped habitat in the Dawros system is occupied by adult mussels; however, it is unsuitable for juvenile recruitment (Moorkens, 2008, 2016; Moorkens and Killeen, 2008; NPWS, 2010). Hydromorphological changes, including bank erosion/slumping and siltation, are key impacts on the Dawros mussel habitat. The target is for sufficient habitat in favourable condition to allow the species to maintain itself on a long-term basis as a viable component of the Dawros system
Suitable habitat: condition	Kilometres	Restore condition of suitable habitat	The species' habitat is a combination of 1) the area of habitat adult and juvenile mussels can occupy; 2) the area of spawning and nursery habitats the host fish can occupy. Fish nursery habitat typically overlaps with mussel habitat. Fish spawning habitat is generally adjacent to mussel habitat, but may lie upstream of the generalised mussel distribution. Only those spawning areas that can regularly contribute juvenile fish to areas occupied by adult mussels should be considered. The availability of mussel and fish spawning/nursery habitats is determined by flow and substratum conditions. It is highly sensitive to hydromorphological changes, sedimentation and nutrient enrichment. Pressures throughout the catchment (map 5) contribute to such impacts. The habitat in the Dawros cannot support sufficient juvenile survival. The target is for sufficient habitat in favourable condition to allow the species to maintain itself on a long-term basis as a viable component of the Dawros system
Water quality: macroinvertebrate and phytobenthos (diatoms)	Ecological quality ratio (EQR)	Restore water quality - macroinvertebrates: EQR greater than 0.90 (Q4-5 or Q5); phytobenthos: EQR greater than 0.93	The EQR targets correspond to high ecological status for these two Water Framework Directive biological quality elements. They represent high water quality with very low nutrient concentrations (oligotrophic conditions). In 2009, the habitat in the Dawros system failed the macroinvertebrate target, but passed the diatom target (Williams, 2009; NPWS, 2010). See also The European Communities Environmental Objectives (Surface Waters) Regulations 2009. The target is for sufficient habitat in favourable condition to allow the species to maintain itself on a long-term basis as a viable component of the Dawros system

Substratum quality: filamentous algae (macroalgae); macrophytes (rooted higher plants)	Percentage	Restore substratum quality - filamentous algae: absent or trace (less than 5%); macrophytes: absent or trace (less than 5%)	The Dawros failed both targets in 2009 (NPWS, 2010). Up to 70% algal cover was recorded in mussel habitat, while high macrophyte cover (particularly <i>Myriophyllum</i>) was frequent (Williams, 2009; NPWS, 2010). In 2016, 91% of quadrats were in moderate or poor condition, mostly owing to algal cover (Moorkens, 2016). Bacterial and fungal growth and detritus accumulation also require further investigation in the Dawros, given the loading of organic fine sediment entering the river (Williams, 2009; Monaghan, 2012). Sufficient recruitment of juvenile mussels is being prevented by the poor condition of the river substratum. The target is for sufficient habitat in favourable condition to allow the species to maintain itself on a long-term basis as a viable component of the Dawros system
Substratum quality: sediment	Occurrence	Restore substratum quality - stable cobble and gravel substrate with very little fine material; no artificially elevated levels of fine sediment	The Dawros failed the target for the Sub-basin Management Plan, with fine sediment recorded throughout the mussel habitat in 2008 and 2009 (Moorkens, 2008; Moorkens and Killeen, 2008; Williams, 2009; NPWS, 2010). There was a clear relationship between heavy siltation and higher macrophyte cover abundance. In 2016, the Dawros passed the target and silt plumes were restricted to marginal areas, suggesting siltation is episodic and deposited fine sediments are periodically flushed-out by high flows (Moorkens, 2016). Sufficient survival of juvenile mussels is being prevented by the poor condition of the river substratum. The target is for sufficient habitat in favourable condition to allow the species to maintain itself on a long-term basis as a viable component of the Dawros system
Substratum quality: oxygen availability	Redox potential	Restore to no more than 20% decline from water column to 5cm depth in substrate	Differences in redox potential between the water column and the substrate correlate with differences in oxygen levels. Juvenile mussels require full oxygenation while buried in gravel. In suitable habitat, there should be very little loss of redox potential between the water column and underlying gravels. The Dawros failed the redox target in 2008, with an average loss of 35% redox potential at 5cm (Moorkens, 2008; Moorkens and Killeen, 2008; NPWS, 2010). In 2016, 92% of redox measurements in the Dawros passed the target and the overall average was 16% (Moorkens, 2016). The target is for sufficient habitat in favourable condition to allow the species to maintain itself on a long-term basis as a viable component of the Dawros system
Hydrological regime: flow variability	Metres per second	Restore appropriate hydrological regime	The availability of suitable freshwater pearl mussel habitat is largely determined by flow (catchment geology being the other important factor). In order to restore the habitat for the species, flow variability over the annual cycle must be such that: 1) high flows can wash fine sediments from the substratum; 2) high flows are not artificially increased so as to cause excessive scour of mussel habitat; 3) low flows do not exacerbate the deposition of fine sediment or growth of algae/macrophytes and 4) low flows do not cause stress to mussels in terms of exposure, water temperatures, food availability or aspects of the reproductive cycle; see Moorkens and Killeen (2014). Groundwater inflow to the substratum also contributes to water-cycling and favourable habitat condition. The target is for sufficient habitat in favourable condition to allow the species to maintain itself on a long-term basis as a viable component of the Dawros system

Host fish	Number	Maintain sufficient juvenile salmonids to host glochidial larvae	Salmonid fish are host to the larval stage of the freshwater pearl mussel and, thus, are essential to completion of the life cycle. 0+ and 1+ fish are typically used, both because of habitat overlaps and the development of immunity with age in fish. Fish presence is sufficient, as higher fish density and biomass is indicative of enriched conditions in mussel rivers. Geist et al. (2006) found that higher densities of host fish coincided with eutrophication, poor substrate quality for mussels and a lack of mussel recruitment, while significantly lower densities and biomass of host fish were associated with high numbers of juvenile mussels. Fish movements must be such that 0+ fish remain in the mussel habitat until their 1+ summer. No fish stocking should occur within the mussel habitat, nor any works that may change the salmonid balance or residency time. In 2009, glochidia were found on salmon, but not trout, in the Dawros (Johnston, 2009; NPWS, 2010)
Fringing habitat: area and condition	Hectares	Maintain the area and condition of fringing habitats necessary to support the population	Riparian habitats, including those along lake fringes, particularly natural/semi-natural woodlands and wetlands, even where they do not form part of a natural floodplain, are an integral part of the structure and functioning of river systems. Fringing habitats aid in the settlement of fine suspended material, protect banks from erosion and contribute to nutrient cycling and to the aquatic food web (e.g. allochthonous matter such as leaf fall), and provide habitat (refuge and resources) for certain life-stages of fish, birds and aquatic invertebrates. Shade may also be important in suppressing algal and macrophyte growth in enriched rivers and moderating temperatures. Equally, fringing habitats are dependent on rivers/lakes, particularly their water levels, and support wetland communities and species of conservation concern. The target is for sufficient habitat in favourable condition to allow the species to maintain itself on a long-term basis as a viable component of the Dawros system

Conservation Objectives for : The Twelve Bens/Garraun Complex SAC [002031]

1106 Salmon *Salmo salar*

To maintain the favourable conservation condition of Atlantic Salmon in The Twelve Bens/Garraun Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Distribution: extent of anadromy	Percentage of river accessible	100% of river channels down to second order accessible from estuary	Artificial barriers block salmons' upstream migration, thereby limiting species to lower stretches and restricting access to spawning areas
Adult spawning fish	Number	Conservation limit (CL) for each system consistently exceeded	A conservation limit (CL) is defined by the North Atlantic Salmon Conservation Organisation (NASCO) as "the spawning stock level that produces long-term average maximum sustainable yield as derived from the adult to adult stock and recruitment relationship". The target is based on the Standing Scientific Committee on Salmon (SSCS) annual model output of CL attainment levels. See SSCS (2016). Attainment of CL estimates are derived from direct counts of adults (rod catch, fish counter) or indirectly by fry abundance counts. The Owenglin River is currently achieving CL
Salmon fry abundance	Number of fry/5 minutes electrofishing	Maintain or exceed 0+ fry mean catchment-wide abundance threshold value. Currently set at 17 salmon fry/5 minutes sampling	The target is the threshold value for rivers currently exceeding their conservation limit (CL)
Out-migrating smolt abundance	Number	No significant decline	Smolt abundance can be negatively affected by a number of impacts such as estuarine pollution, predation and sea lice (<i>Lepeophtheirus salmonis</i>)
Number and distribution of redds	Number and occurrence	No decline in number and distribution of spawning redds due to anthropogenic causes	Salmon spawn in clean gravels
Water quality	EPA Q value	At least Q4 at all sites sampled by EPA	Q values based on triennial water quality surveys carried out by the Environmental Protection Agency (EPA)

Conservation Objectives for : The Twelve Bens/Garraun Complex SAC [002031]

1355 Otter *Lutra lutra*

To maintain the favourable conservation condition of Otter in The Twelve Bens/Garraun Complex SAC, which is defined by the following list of attributes and targets:


Attribute	Measure	Target	Notes
Distribution	Percentage positive survey sites	No significant decline	Measure based on standard otter survey technique. Favourable Conservation Status (FCS) target, based on 1980/81 survey findings, is 88% in SACs. Current range is estimated at 93.6% (Reid et al., 2013)
Extent of terrestrial habitat	Hectares	No significant decline. Area mapped and calculated as 854.66ha	No field survey. Areas mapped to include 10m terrestrial buffer along shoreline (above high water mark (HWM) and along river banks) identified as critical for otters (NPWS, 2007)
Extent of marine habitat	Hectares	No significant decline. Area mapped and calculated as 53.81ha	No field survey. Area mapped based on evidence that otters tend to forage within 80m of the shoreline (HWM) (Kruuk, 2006; NPWS, 2007)
Extent of freshwater (river) habitat	Kilometres	No significant decline. Length mapped and calculated as 382.7km	No field survey. River length calculated on the basis that otters will utilise freshwater habitats from estuary to headwaters (Chapman and Chapman, 1982)
Extent of freshwater (lake) habitat	Hectares	No significant decline. Area mapped and calculated as 540.7ha	No field survey. Area mapped based on evidence that otters tend to forage within 80m of the shoreline (NPWS, 2007)
Couching sites and holts	Number	No significant decline	Otters need lying up areas throughout their territory where they are secure from disturbance (Kruuk and Moorhouse, 1991; Kruuk, 2006)
Fish biomass available	Kilograms	No significant decline	Broad diet that varies locally and seasonally, but dominated by fish, in particular salmonids, eels and sticklebacks in freshwater (Bailey and Rochford, 2006; Reid et al., 2013) and wrasse and rockling in coastal waters (Kingston et al., 1999)
Barriers to connectivity	Number	No significant increase. For guidance, see map 6	Otters will regularly commute across stretches of open water up to 500m e.g. between the mainland and an island; between two islands; across an estuary (De Jongh and O'Neill, 2010). It is important that such commuting routes are not obstructed

Conservation Objectives for : The Twelve Bens/Garraun Complex SAC [002031]


1833 Slender Naiad *Najas flexilis*


To maintain the favourable conservation condition of Slender Naiad in The Twelve Bens/Garraun Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Population extent	Hectares; distribution	No change to the spatial extent of <i>Najas flexilis</i> within Pollacappul and Kylemore Loughs, subject to natural processes. See map 7 for known locations	The selection of The Twelve Bens/Garraun Complex SAC for <i>Najas flexilis</i> (slender naiad) was based on its presence in Loughs Pollacappul and Kylemore. Pollacappul is considered to support a large population of the species, while the species' status in Kylemore Lough (Roden and Murphy, 2014). See The Twelve Bens/Garraun Complex SAC conservation objectives supporting document for <i>Najas flexilis</i> for further details
Population depth	Metres	No change to the depth range of <i>Najas flexilis</i> within the lakes, subject to natural processes	See the <i>Najas flexilis</i> supporting document for further details
Population viability	Plant traits	No decline in plant fitness, subject to natural processes	See the <i>Najas flexilis</i> supporting document for further details
Population abundance	Square metres	No change to the cover abundance of <i>Najas flexilis</i> , subject to natural processes	See the <i>Najas flexilis</i> supporting document for further details
Species distribution	Occurrence	No decline, subject to natural processes	See the <i>Najas flexilis</i> supporting document for further details
Habitat extent	Hectares	No decline, subject to natural processes	See the <i>Najas flexilis</i> supporting document for further details
Hydrological regime: water level fluctuations	Metres	Maintain/restore appropriate natural hydrological regime necessary to support the habitat for the species	See the <i>Najas flexilis</i> supporting document for further details
Lake substratum quality	Various	Maintain appropriate substratum type, extent and chemistry to support the populations of the species	See the <i>Najas flexilis</i> supporting document for further details
Water quality	Various	Maintain/restore appropriate water quality to support the populations of the species	See the <i>Najas flexilis</i> supporting document for further details
Acidification status	pH units; mg/l	Maintain appropriate water and sediment pH, alkalinity and cation concentrations to support the populations of <i>Najas flexilis</i> , subject to natural processes	See the <i>Najas flexilis</i> supporting document for further details
Water colour	mg/l PtCo	Maintain appropriate water colour to support the populations of <i>Najas flexilis</i>	See the <i>Najas flexilis</i> supporting document for further details
Associated species	Species composition and abundance	Maintain appropriate associated species and vegetation communities to support the populations of <i>Najas flexilis</i>	See the <i>Najas flexilis</i> supporting document for further details
Fringing habitat: area and condition	Hectares	Maintain the area and condition of fringing habitats necessary to support the populations of <i>Najas flexilis</i>	See the <i>Najas flexilis</i> supporting document for further details



Legend

 The Twelve Bens/Garraun Complex SAC 002031




An Roinn Ealaíon, Cúlchreacht, Gnóthaí Réigiúnacha, Tuairhe agus Gaeltachta
 Department of Arts, Heritage, Regional, Rural and Gaeltacht Affairs

MAP 1:
THE TWELVE BENS / GARRAUN COMPLEX SAC
CONSERVATION OBJECTIVES
SAC DESIGNATION

Map to be read in conjunction with the NPWS Conservation Objectives Document.

SITE CODE:
SAC 002031; version 3.01. CO. GALWAY




The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
 Ordnance Survey of Ireland Licence No EN 0059216. © Ordnance Survey of Ireland Government of Ireland

Níl sna teorainneacha ar na léarscáileanna ach nod garshuimhach ginearálta. Féadfar athbheithníthe a déanamh ar theorainneacha na gceantar comharthaithe. Suirbhéarachta Ordonáis na hÉireann Ceadúnas

© Ordnance Survey Ireland




Map Version 1
Date: May 2017



Legend	
	The Twelve Bens / Garraun Complex SAC 002031
	Maumturk Mountains SAC 002008
	Connemara Bog Complex SAC 002034
	West Connacht Coast SAC 002998

© Ordnance Survey Ireland



MAP 2:
THE TWELVE BENS / GARRAUN COMPLEX SAC
CONSERVATION OBJECTIVES
ADJACENT / ADJOINING AND
OVERLAPPING DESIGNATIONS
 Map to be read in conjunction with the NPWS Conservation Objectives Document.

SITE CODE:
 SAC 002031; version 3.01. SAC 002998; version 3.03
 SAC 002008; version 3. SAC 002034 version 3.01.
 CO. GALWAY


0 2.5 5 7.5 10 12.5 km

The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
 Ordnance Survey of Ireland Licence No EN 0059216. © Ordnance Survey of Ireland Government of Ireland

Níl sna teorainneacha ar na léarscáilleana ach nod garshuíomhach ginearálta. Féadfar athbhreithnithe a déanamh ar theorainneacha na gceantar comharthaíthe. Suirbhéarachtá Ordónais na hÉireann Ceadúnas

N

Map Version 1
Date: May 2017




Legend

- The Twelve Bens / Garraun Complex SAC 002031
- OSi Discovery Series County Boundary


Indicative Lake Habitats

- Potential 3110 Potential oligotrophic waters containing very few minerals of sandy plains: *Littorelletalia uniflorae*
- Potential 3130 / Potential 3110 Potential oligotrophic to mesotrophic standing waters with vegetation of the *Littorelletalia uniflorae* and/or of the *Isoëto-Nanojuncetea* / Potential oligotrophic waters containing very few minerals of sandy plains: *Littorelletalia uniflorae*




Legend

Woodland Habitats Qualifying Interests


 91A0 Old sessile oak woods with *Ilex* and *Blechnum* in the British Isles

Woodland Habitats Non Qualifying Interests

 91E0 Alluvial forests with *Alnus glutinosa* and *Fraxinus excelsior* (*Alno-Padion*, *Alnion incanae*, *Salicion albae*)


 The Twelve Bens / Garraun Complex SAC 002031

 OSi Discovery Series County Boundary




Legend

- 1029 Freshwater Pearl Mussel *Margaritifera margaritifera* Distribution Target
- 1029 Freshwater Pearl Mussel *Margaritifera margaritifera* Suitable Habitat Target
- 1029 Freshwater Pearl Mussel *Margaritifera margaritifera* Catchment
- The Twelve Bens / Garraun Complex SAC 002031
- OSi Discovery Series County Boundary



Legend

- 1355 Otter *Lutra lutra* Commuting 250m
- The Twelve Bens / Garraun Complex SAC 002031
- OSi Discovery Series County Boundary




An Roinn Ealaíon, Oidhreachta,
Gnóthaí Réigiúnacha, Tuaithe agus Gaeltachta
Department of Arts, Heritage,
Regional, Rural and Gaeltacht Affairs

**MAP 6:
THE TWELVE BENS / GARRAUN COMPLEX SAC
CONSERVATION OBJECTIVES
OTTER COMMUTING**

Map to be read in conjunction with the NPWS Conservation Objectives Document.


**SITE CODE:
SAC 002031; version 3.01. CO. GALWAY**



The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
Ordnance Survey of Ireland Licence No EN 0059216. © Ordnance Survey of Ireland Government of Ireland
Níl sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbhreithnithe a déanamh ar theorainneacha na gceantar comharthaithe. Suirbhéarachta Ordonáis na hÉireann Ceadúnas



**Map Version 1
Date: May 2017**



Legend

- ★ 1833 Sender Naiad *Najas flexilis*
- The Twelve Bens / Garraun Complex SAC 002031
- OSi Discovery Series County Boundary




An Roinn Ealaíon, Oidhreacht,
Gnóthaí Réigiúnacha, Tuaithe agus Gaeltachta
Department of Arts, Heritage,
Regional, Rural and Gaeltacht Affairs

**MAP 7:
THE TWELVE BENS / GARRAUN COMPLEX SAC
CONSERVATION OBJECTIVES
SLENDER NAIAID**

Map to be read in conjunction with the NPWS Conservation Objectives Document.

**SITE CODE:
SAC 002031; version 3.01. CO. GALWAY**



0 1 2 3 4 5 km

The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
Ordnance Survey of Ireland Licence No EN 0059216. © Ordnance Survey of Ireland Government of Ireland

Níl sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbheithnithé a déanamh ar theorainneacha na gceantar comharthaithe. Suirbhéarachta Ordonáis na hÉireann Ceadúnas



**Map Version 1
Date: May 2017**

National Parks and Wildlife Service

Conservation Objectives Series

Connemara Bog Complex SAC 002034



*An Roinn
Ealaíon, Oidhreachta agus Gaeltachta*

*Department of
Arts, Heritage and the Gaeltacht*



**National Parks and Wildlife Service,
Department of Arts, Heritage and the Gaeltacht,**

7 Ely Place, Dublin 2, Ireland.

Web: www.npws.ie

E-mail: nature.conservation@ahg.gov.ie

Citation:

**NPWS (201) Conservation Objectives: Connemara Bog Complex SAC 002034.
Version 1. National Parks and Wildlife Service, Department of Arts, Heritage
and the Gaeltacht.**

Series Editor: Rebecca Jeffrey

ISSN 2009-4086

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

* indicates a priority habitat under the Habitats Directive

002034 Connemara Bog Complex SAC	
1065	Marsh Fritillary <i>Euphydryas aurinia</i>
1106	Salmon <i>Salmo salar</i>
1150	Coastal lagoonsE
1170	Reefs
1355	Otter <i>Lutra lutra</i>
1833	Slender Naiad <i>Najas flexilis</i>
3110	Oligotrophic waters containing very few minerals of sandy plains (Littorelletalia uniflorae)
3130	Oligotrophic to mesotrophic standing waters with vegetation of the Littorelletea uniflorae and/or Isoeto-Nanojuncetea
3160	Natural dystrophic lakes and ponds
3260	Water courses of plain to montane levels with the Ranunculion fluitantis and Callitriche-Batrachion vegetation
4010	Northern Atlantic wet heaths with <i>Calluna vulgaris</i>
4030	European dry heaths
6410	<i>Tilandsia</i> meadows on calcareous, peaty or clayey-silt-laden soils (Molinion caeruleae)
7130	Blanket bogs (* if active bog)
7140	Transition mires and quaking bogs
7150	Depressions on peat substrates of the Rhynchosporion
7230	Alkaline fens
91A0	Old sessile oak woods with <i>Quercus petraea</i> and <i>Q. robur</i> in the British Isles

Please note that this SAC overlaps with Connemara Bog Complex SPA (004181). It adjoins Lough Corrib SAC (000297), Maumturk Mountains SAC (002008), The Twelve Bens/Garraun Complex SAC (002031) and Kilkieran Bay and Islands SAC (002111). See map 2. The conservation objectives for this site should be used in conjunction with those for the overlapping and adjoining sites as appropriate.

Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

NPWS Documents

Year :	1984
Title :	The vegetation of Irish lakes
Author :	Heuff, H.
Series :	Unpublished report to NPWS
Year :	1987
Title :	Survey to locate lowland blanket bogs of scientific interest in Connemara, Co. Galway
Author :	Douglas, C.; Grogan, H.
Series :	Unpublished report to the Wildlife Service
Year :	1989
Title :	A survey to locate blanket bogs in Co. Galway. Part 2
Author :	Douglas, C.; Garvey, L.; Kelly, L.; O'Sullivan, A.
Series :	Unpublished report to the Wildlife Service
Year :	1998
Title :	Survey of Irish coastal lagoons, 1996 and 1998
Author :	Healy, B.; Roden, C.; Oliver, G.; Good, J.
Series :	Unpublished report to NPWS
Year :	2006
Title :	Otter survey of Ireland 2004/2005
Author :	Bailey, M.; Rochford, J.
Series :	Irish Wildlife Manual No. 23
Year :	2007
Title :	Supporting documentation for the Habitats Directive Conservation Status Assessment - backing documents. Article 17 forms and supporting maps
Author :	NPWS
Series :	Unpublished report to NPWS
Year :	2007
Title :	Inventory of Irish coastal lagoons (version 2)
Author :	Oliver, G.
Series :	Unpublished report to NPWS
Year :	2008
Title :	National survey of native woodlands 2003-2008
Author :	Perrin, P.M.; Martin, J.; Barron, S.; O'Neill, F.H.; McNutt, K.E.; Delaney, A.
Series :	Unpublished Report to NPWS
Year :	2009
Title :	NS II freshwater pearl mussel sub-basin management plans: Report on biological monitoring of surface water quality in the Owenriff catchment (Corrib sub-catchment), Co. Galway
Author :	Williams, L.
Series :	Unpublished report to NPWS
Year :	2010
Title :	A provisional inventory of ancient and long-established woodland in Ireland
Author :	Perrin, P.M.; Daly, O.H.
Series :	Irish Wildlife Manual No. 46

Year :	2013
Title :	National otter survey of Ireland 2010/12
Author :	Reid, N.; Hayden, B.; Lundy, M.G.; Pietravalle, S.; McDonald, R.A.; Montgomery, W.I.
Series :	Irish Wildlife Manual No. 76
Year :	2013
Title :	Irish semi-natural grasslands survey 2007-2012
Author :	O'Neill, F.H.; Martin, J.R.; Devaney, F.M.; Perrin, P.M.
Series :	Irish Wildlife Manual No. 78
Year :	2013
Title :	A survey of the benthic macrophytes of three hard-water lakes: Lough Bunny, Lough Carra and Lough Owel
Author :	Roden, C.; Murphy, P.
Series :	Irish Wildlife Manual No. 70
Year :	2013
Title :	The status of EU protected habitats and species in Ireland. Volume 2. Habitats assessments
Author :	NPWS
Series :	Conservation assessments
Year :	2014
Title :	Guidelines for a national survey and conservation assessment of upland vegetation and habitats in Ireland, Version 2.0
Author :	Perrin, P.M.; Barron, S.J.; Roche, J.R.; O'Hanrahan, B.
Series :	Irish Wildlife Manual No. 79
Year :	2015
Title :	Habitats Directive Annex I lake habitats: a working interpretation for the purposes of site-specific conservation objectives and Article 17 reporting
Author :	O Connor, Á.
Series :	Unpublished document by NPWS
Year :	2015
Title :	Connemara Bog Complex SAC (site code: 2034) Conservation objectives supporting document- coastal lagoons V1
Author :	NPWS
Series :	Conservation objectives supporting document
Year :	2015
Title :	Connemara Bog Complex SAC (site code: 2034) Conservation objectives supporting document- marine habitats V1
Author :	NPWS
Series :	Conservation objectives supporting document
Year :	2015
Title :	Connemara Bog Complex SAC (site code: 2034) Conservation objectives supporting document- <i>Najas flexilis</i> V1
Author :	NPWS
Series :	Conservation objectives supporting document

Other References

Year :	1982
Title :	Otter survey of Ireland
Author :	Chapman, P.J.; Chapman, L.L.
Series :	Unpublished report to Vincent Wildlife Trust

Year :	1982
Title :	Eutrophication of waters. Monitoring assessment and control
Author :	OECD
Series :	OECD, Paris
Year :	1991
Title :	The spatial organization of otters (<i>Lutra lutra</i>) in Shetland
Author :	Kruuk, H.; Moorhouse, A.
Series :	J. Zool, 224: 41-57
Year :	1991
Title :	Records of aquatic plants from Connemara and the Burren
Author :	Preston, C.D.; Stewart, N.F.; Webster, S.D.
Series :	Irish Naturalists' Journal 23(11): 464-467
Year :	1997
Title :	The BioMar biotope viewer: a guide to marine habitats, fauna and flora in Britain and Ireland
Author :	Picton, B.E.; Costello, M.J.
Series :	Environmental Science Unit, Trinity College Dublin
Year :	1999
Title :	Diet of otters (<i>Lutra lutra</i>) on Inishmore, Aran Islands, west coast of Ireland
Author :	Kingston, S.; O'Connell, M.; Fairley, J.S.
Series :	Biol & Environ Proc R Ir Acad B 99B:173-182
Year :	2000
Title :	Colour in Irish lakes
Author :	Free, G.; Allott, N.; Mills, P.; Kennelly, C.; Day, S.
Series :	Verhandlungen Internationale Vereinigung für theoretische und angewandte Limnologie. 27: 2620-2623
Year :	2001
Title :	Aquatic plants in Britain and Ireland
Author :	Preston, C.D.; Croft, J.M.
Series :	Harley Books, Colchester
Year :	2002
Title :	Reversing the habitat fragmentation of British woodlands
Author :	Peterken, G.
Series :	WWF-UK, London
Year :	2002
Title :	Deterioration of Atlantic soft water macrophyte communities by acidification, eutrophication and alkanisation
Author :	Arts, G.H.P.
Series :	Aquatic Botany, 73: 373-393
Year :	2003
Title :	Ecology of watercourses characterised by Ranunculion fluitantis and Callitriche-Batrachion Vegetation
Author :	Hatton-Ellis, T.W.; Grieve, N.
Series :	Conserving Natura 2000 Rivers Ecology Series No. 11. English Nature, Peterborough
Year :	2004
Title :	The ecology of <i>Najas flexilis</i>
Author :	Wingfield, R.A.; Murphy, K.J.; Hollingsworth, P.; Gaywood, M.J.
Series :	Scottish Natural Heritage Commissioned Report No. 017 (ROAME No. F98PA02)

Year :	2006
Title :	Otters - ecology, behaviour and conservation
Author :	Kruuk, H.
Series :	Oxford University Press
Year :	2006
Title :	A reference-based typology and ecological assessment system for Irish lakes. Preliminary investigations. Final report. Project 2000-FS-1-M1 Ecological assessment of lakes pilot study to establish monitoring methodologies EU (WFD)
Author :	Free, G.; Little, R.; Tierney, D.; Donnelly, K.; Coroni, R.
Series :	EPA, Wexford
Year :	2009
Title :	The identification, characterization and conservation value of isoetid lakes in Ireland
Author :	Free G.; Bowman, J.; McGarrigle, M.; Little, R.; Coroni, R.; Donnelly, K.; Tierney, D.; Trodd, W.
Series :	Aquatic Conservation: Marine and Freshwater Ecosystems 19 (3): 264–273
Year :	2010
Title :	Second draft Owenriff freshwater pearl mussel sub-basin management plan (2009-2015). March 2010
Author :	Department of the Environment, Heritage and Local Government
Series :	Unpublished report
Year :	2012
Title :	Subtidal reef survey of Salt Lake, Connemara Bog Complex
Author :	MERC
Series :	Unpublished report to the Marine Institute and NPWS
Year :	2012
Title :	The impact of conifer plantation forestry on the ecology of peatland lakes
Author :	Drinan, T.J.
Series :	Unpublished PhD thesis, University College Cork
Year :	2013
Title :	Monitoring and assessment of Irish lagoons for the purposes of the EU Water Framework Directive, 2009-2011. Parts 1 and 2
Author :	Roden, C.M; Oliver, G.A.
Series :	Unpublished report to the Environmental Protection Agency
Year :	2013
Title :	Management strategies for the protection of high status water bodies
Author :	Ní Chatháin, B.; Moorkens, E.; Irvine, K.
Series :	Strive Report Series No. 99. EPA, Wexford
Year :	2013
Title :	Interpretation manual of European Union habitats- Eur 28
Author :	European Commission- DG Environment
Series :	Reference document
Year :	in prep.
Title :	Monitoring of hard-water lakes in Ireland using charophytes and other macrophytes
Author :	Roden, C.; Murphy, P.
Series :	Unpublished report to NPWS

Spatial data sources

Year :	Revision 2011
Title :	Inventory of Irish Coastal Lagoons. Version 3
GIS Operations :	Clipped to SAC boundary
Used For :	1150 (map 3)
Year :	Interpolated 2015
Title :	1994, 1995 BioMar surveys; 2011 subtidal reef survey
GIS Operations :	Polygon feature classes from marine community types base data sub-divided based on interpolation of marine survey data. Expert opinion used as necessary to resolve any issues arising
Used For :	1170, marine community types (maps 4 and 5)
Year :	2005
Title :	OSi Discovery series vector data
GIS Operations :	High water mark (HWM) and low water mark (LWM) polyline feature classes converted into polygon feature classes and combined; EU Annex I Saltmarsh and Coastal data erased out if present
Used For :	Marine community types base data (map 5)
Year :	2008
Title :	OSi 1:5000 IG vector dataset
GIS Operations :	WaterPolygons feature class clipped to the SAC boundary. Expert opinion used to identify Annex I habitat and to resolve any issues arising
Used For :	3110, 3160 (map 6)
Year :	Revision 2012
Title :	Margaritifera Sensitive Areas
GIS Operations :	Relevant catchment boundaries identified. Expert opinion used as necessary to resolve any issues arising
Used For :	3260 (map 7)
Year :	Revision 2010
Title :	National Survey of Native Woodlands 2003-2008. Version 1
GIS Operations :	QIs selected; clipped to SAC boundary. Expert opinion used as necessary to resolve any issues arising
Used For :	91A0 (map 8)
Year :	2013
Title :	Najas flexilis data
GIS Operations :	Lake habitat for species clipped to SAC boundary
Used For :	1833 (map 9)

Conservation Objectives for : Connemara Bog Complex SAC [002034]

1150 Coastal lagoons

To maintain the favourable conservation condition of Coastal lagoons in Connemara Bog Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable, subject to slight natural variation. Favourable reference area 151.4ha. See map 3	Area calculated from spatial data derived from Oliver (2007) for IL061 (Loch Doire Bhanbh), IL062 (Loch an tSáile) and IL063 (Loch Conaortha). Area of Ardbear Salt Lake calculated using the same methodology. See lagoons supporting document for further details
Habitat distribution	Occurrence	No decline, subject to natural processes. See map 3	IL061 (Loch Doire Bhanbh), IL062 (Loch an tSáile) and IL063 (Loch Conaortha) listed in Oliver (2007) and Ardbear Salt Lake in Healy et al. (1998). See lagoons supporting document for further details
Salinity regime	Practical salinity units (psu)	Median annual salinity and temporal variation within natural ranges	Loch Doire Bhanbh and Ardbear Salt Lake are recorded as polyhaline lagoons, Loch an tSáile as oligohaline/mesohaline and Loch Conaortha as mesohaline. See lagoons supporting document for further details
Hydrological regime	Metres	Annual water level fluctuations and minima within natural ranges	Maximum depth of Loch Doire Bhanbh is recorded as 3m, Loch an tSáile as 13m, Loch Conaortha as 5m; the average depth of Ardbear Salt Lake is between 4-6m with a maximum depth of 27m. See lagoons supporting document for further details
Barrier: connectivity between lagoon and sea	Permeability	Appropriate hydrological connections between lagoons and sea, including where necessary, appropriate management	Loch Doire Bhanbh is described as a saltmarsh lagoon, Loch an tSáile and Loch Conaortha as rock/peat barrier lagoons and Ardbear Salt Lake as having a rock barrier. See lagoons supporting document for further details
Water quality: Chlorophyll <i>a</i>	µg/L	Annual median chlorophyll <i>a</i> within natural ranges and less than 5µg/L	Target based on Roden and Oliver (2013). See lagoons supporting document for further details
Water quality: Molybdate Reactive Phosphorus (MRP)	mg/L	Annual median MRP within natural ranges and less than 0.1mg/L	Target based on Roden and Oliver (2013). See lagoons supporting document for further details
Water quality: Dissolved Inorganic Nitrogen (DIN)	mg/L	Annual median DIN within natural ranges and less than 0.15mg/L	Target based on Roden and Oliver (2013). See lagoons supporting document for further details
Depth of macrophyte colonisation	Metres	Macrophyte colonisation to at least 4m depth	It is expected that macrophyte colonisation would extend down to the full depths of Loch Doire and Loch Conaortha. In Loch an tSáile and Ardbear Salt Lake macrophytes would be expected to extend to 4m depth; thereafter the dark water colour prohibits growth. See lagoons supporting document for further details
Typical plant species	Number and m ²	Maintain number and extent of listed lagoonal specialists, subject to natural variation	Species listed in Oliver (2007), Healy et al. (1997) and MERC (2012). See lagoons supporting document for further details
Typical animal species	Number	Maintain listed lagoon specialists, subject to natural variation	Species listed in Oliver (2007) and Healy et al. (1997). See lagoons supporting document for further details. The conservation objective and supporting document for reefs (1170) gives information on the <i>Serpula vermicularis</i> -dominated community complex
Negative indicator species	Number and percentage cover	Negative indicator species absent or under control	Low salinity, shallow water and elevated nutrient levels increase the threat of unnatural encroachment by reedbeds. See lagoons supporting document for further details

1170 Reefs

To maintain the favourable conservation condition of Reefs in Connemara Bog Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	The permanent habitat area is stable or increasing, subject to natural processes. See map 4	Habitat area estimated as 6ha from a 2011 subtidal reef survey (MERC, 2012) and an intertidal walkover in 2014
Distribution	Occurrence	The distribution of reefs is stable or increasing, subject to natural processes. See map 4	Based on information from a 2011 subtidal reef survey (MERC, 2012) and an intertidal walkover in 2014
Community extent	Hectares	Maintain the extent of the <i>Serpula vermicularis</i> -dominated community complex, subject to natural processes. See map 5	The likely extent of the <i>Serpula vermicularis</i> -dominated community complex was derived from dive surveys undertaken in 2011 (MERC, 2012). See marine supporting document for further details
Community structure	Biological composition	Conserve the high quality of the <i>Serpula vermicularis</i> -dominated community complex, subject to natural processes	Based on 2011 diver observation (MERC, 2012). See marine supporting document for further details
Community structure	Biological composition	Conserve the following community type in a natural condition: Intertidal reef community complex. See map 5	Based on information from 1994 and 1995 BioMar surveys (Picton and Costello, 1997) and an intertidal walkover in 2014. See marine supporting document for further details

Conservation Objectives for : Connemara Bog Complex SAC [002034]

3110 Oligotrophic waters containing very few minerals of sandy plains (Littorelletalia uniflorae)

To maintain the favourable conservation condition of Oligotrophic waters containing very few minerals of sandy plains (Littorelletalia uniflorae) in Connemara Bog Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	This SAC has a very large number of lakes and ponds (more than 1,100, based on 1:5,000 OSI data). Most are surrounded by peatland and are likely to contain lake habitats 3110 and/or 3160. Records for <i>Najas flexilis</i> indicates lake habitat 3130 also occurs in some lakes. The exact distribution of habitat 3110 in the site is unknown, but it is considered to be widespread and of high conservation value, particularly in Roundstone Blanket Bog (see indicative distribution in map 6). 3110 lakes in the SAC typically have shallow margins, often with rock outcrops along the shore, and a mixed rocky/peaty/sandy substratum. Two measures of extent should be used: 1. the area of the lake itself and; 2. the extent of the vegetation communities/zones that typify the habitat. Further information relating to all attributes is provided in the lake habitats supporting document for the purposes of site-specific conservation objectives and Article 17 reporting (O Connor, 2015)
Habitat distribution	Occurrence	No decline, subject to natural processes	As noted above, it is likely that the habitat is widespread and of high conservation value in the SAC (see map 6)
Typical species	Occurrence	Typical species present, in good condition, and demonstrating typical abundances and distribution	For lists of typical plant species, see Article 17 habitat assessment for 3110 (NPWS, 2013) and the lake habitats supporting document (O Connor, 2015). The only known Irish sites for <i>Luronium natans</i> are in the SAC, and include at least three lakes (Rich et al., 1995; Curtis et al., 2012). The status of <i>Luronium natans</i> in Ireland is currently uncertain, as it is unclear whether it is native to these sites, or was introduced. The protected <i>Pilularia globulifera</i> is known from a number of lakes in the SAC and is likely to be widespread
Vegetation composition: characteristic zonation	Occurrence	All characteristic zones should be present, correctly distributed and in good condition	The characteristic zonation of lake habitat 3140 has been described (Roden and Murphy, 2013; in prep.), however significant further work is necessary to describe the characteristic zonation and other spatial patterns in the remaining four Annex I lake habitats
Vegetation distribution: maximum depth	Metres	Maintain maximum depth of vegetation, subject to natural processes	The maximum depth of vegetation is likely to be specific to the lake shoreline in question. An indicative target of >6m has been developed for hard water lakes (3140) (see Roden and Murphy, 2013; in prep.). Indicative targets will be developed for the other lake habitats with time. Lakes in the SAC typically have very clear water and, therefore, maximum depth is expected to be large
Hydrological regime: water level fluctuations	Metres	Maintain appropriate natural hydrological regime necessary to support the habitat	Fluctuations in lake water level are typical in Ireland, but can be amplified by activities such as abstraction and drainage. Increased water level fluctuations can increase wave action, up-root vegetation, increase turbidity, alter the substratum and lead to release of nutrients from the sediment. The hydrological regime of the lakes must be maintained so that the area, distribution and depth of the lake habitat and its constituent/characteristic vegetation zones and communities are not reduced

Lake substratum quality	Various	Maintain appropriate substratum type, extent and chemistry to support the vegetation	Research is required to further characterise the substratum types (particle size and origin) and substratum quality (notably pH, calcium, iron and nutrient concentrations) favoured by each of the five Annex I lake habitats in Ireland. It is likely that lake habitat 3110 is associated with a range of nutrient-poor substrates, from stones, cobble and gravel, through sands, silt, clay and peat. Substratum particle size is likely to vary with depth and along the shoreline within a single lake
Water quality: transparency	Metres	Maintain appropriate Secchi transparency. There should be no decline in Secchi depth/transparency	Transparency relates to light penetration and, hence, to the depth of colonisation of vegetation. It can be affected by phytoplankton blooms, water colour and turbidity. A target has been set for hard water lakes (3140), however specific targets have yet to be established for the remaining lake habitats. Habitat 3110 is associated with very clear water. The OECD fixed boundary system set transparency targets for oligotrophic lakes of $\geq 6\text{m}$ annual mean Secchi disk depth, and $\geq 3\text{m}$ annual minimum Secchi disk depth. Free et al. (2009) found high isoetid abundance in lakes with Secchi depths of more than 3m
Water quality: nutrients	$\mu\text{g/l P}$; mg/l N	Maintain the concentration of nutrients in the water column to sufficiently low levels to support the habitat and its typical species	As a nutrient poor habitat, oligotrophic and Water Framework Directive (WFD) 'high' status targets apply. Where a lake has nutrient concentrations that are lower than these targets, there should be no decline within class, i.e. no upward trend in nutrient concentrations. For lake habitat 3110, annual average TP concentration should be $\leq 10\mu\text{g/l TP}$, average annual total ammonia concentration should be $\leq 0.040\text{mg/l N}$ and annual 95th percentile for total ammonia should be $\leq 0.090\text{mg/l N}$. See also The European Communities Environmental Objectives (Surface Waters) Regulations 2009
Water quality: phytoplankton biomass	$\mu\text{g/l Chlorophyll } a$	Maintain appropriate water quality to support the habitat, including high chlorophyll status	Oligotrophic and WFD 'high' status targets apply to the lake habitat 3110. Where a lake has a chlorophyll <i>a</i> concentration that is lower than this target, there should be no decline within class, i.e. no upward trend in phytoplankton biomass. The average growing season (March-October) chlorophyll <i>a</i> concentration must be $< 5.8\mu\text{g/l}$. The annual average chlorophyll <i>a</i> concentration should be $< 2.5\mu\text{g/l}$ and the annual peak chlorophyll <i>a</i> concentration should be $\leq 8.0\mu\text{g/l}$. See also The European Communities Environmental Objectives (Surface Waters) Regulations 2009
Water quality: phytoplankton composition	EPA phytoplankton composition metric	Maintain appropriate water quality to support the habitat, including high phytoplankton composition status	The EPA has developed a phytoplankton composition metric for nutrient enrichment of Irish lakes. As for other water quality indicators, habitat 3110 requires WFD high status
Water quality: attached algal biomass	Algal cover and EPA phytobenthos metric	Maintain trace/ absent attached algal biomass ($< 5\%$ cover) and high phytobenthos status	Nutrient enrichment can favour epiphytic and epipelic algae that can out-compete the submerged vegetation. The cover abundance of attached algae in the oligotrophic soft water habitat should, therefore, be trace/ absent ($< 5\%$ cover). EPA phytobenthos can be used as an indicator of changes in attached algal biomass. As for other water quality indicators, habitat 3110 requires high phytobenthos status
Water quality: macrophyte status	EPA macrophyte metric (The Free Index)	Maintain high macrophyte status	Nutrient enrichment can favour more competitive submerged macrophyte species that out-compete the typical and characteristic species for the lake habitat. The EPA monitors macrophyte status for Water Framework Directive purposes using the 'Free Index'. The target for lake habitat 3110 is high status or an Ecological Quality Ratio (EQR) for lake macrophytes of ≥ 0.90 , as defined in Schedule Five of the European Communities Environmental Objectives (Surface Waters) Regulations 2009

Acidification status	pH units, mg/l	Maintain appropriate water and sediment pH, alkalinity and cation concentrations to support the habitat, subject to natural processes	Acidification can impact on species abundance and composition in soft water lake habitats. In Europe, acidification of isoetid lakes can lead to loss of isoetids and dominance by submerged <i>Sphagnum</i> mosses and <i>Juncus bulbosus</i> (Arts, 2002). The specific requirements of lake habitat 3110, in terms of water and sediment pH, alkalinity and cation concentration, have not been determined. For this habitat, and adopting a precautionary approach based on Arts (2002), minimum pH should not be <5.5 pH units. Maximum pH should be <9.0 pH units, in line with the surface water standards established for soft waters (where water hardness is ≤100mg/l CaCO ₃). See Schedule Five of the European Communities Environmental Objectives (Surface Waters) Regulations 2009
Water colour	mg/l PtCo	Maintain appropriate water colour to support the habitat	Increased water colour and turbidity decrease light penetration and can reduce the area of available habitat for lake macrophytes, particularly at the lower euphotic depths. The primary source of increased water colour in Ireland is disturbance to peatland. No habitat-specific or national standards for water colour currently exist. Studies have shown median colour concentrations in Irish lakes of 38mg/l PtCo (Free et al., 2000) and 33mg/l PtCo (Free et al., 2006). It is likely that the water colour in all Irish lake habitats would naturally be <50mg/l PtCo. Water colour can be very low (<20mg/l PtCo or even <10mg/l PtCo) in lake habitat 3110, where the peatland in the lake's catchment is intact
Dissolved organic carbon (DOC)	mg/l	Maintain appropriate organic carbon levels to support the habitat	Dissolved (and particulate) organic carbon (OC) in the water column is linked to water colour and acidification (organic acids). Increasing DOC in water has been documented across the Northern Hemisphere, including afforested peatland catchments in Ireland. Damage and degradation of peatland, leading to decomposition of peat is likely to be the predominant source of OC in Ireland. OC in water promotes decomposition by fungi and bacteria that, in turn, releases dissolved nutrients. The increased biomass of decomposers can also impact directly on the characteristic lake communities through shading, competition, etc.
Turbidity	nephelometric turbidity units/ mg/l SS/ other appropriate units	Maintain appropriate turbidity to support the habitat	Turbidity can significantly affect the quantity and quality of light reaching rooted and attached vegetation and can, therefore, impact on lake habitats. The settlement of higher loads of inorganic or organic material on lake vegetation communities may also have impacts on sensitive, delicate species. Turbidity can increase as a result of re-suspension of material within the lake, higher loads entering the lake, or eutrophication. Turbidity measurement and interpretation is challenging. As a result, it is likely to be difficult to set habitat-specific targets for turbidity in lakes
Fringing habitat	Area	Maintain the area and condition of fringing habitats necessary to support the natural structure and functioning of habitat 3110	Most lake shorelines have fringing habitats of reedswamp, other swamp, fen, marsh or wet-woodland that intergrade with and support the structure and functions of the lake habitat. In this SAC, 3110 lakes are most likely to be fringed by poor fen, flush, blanket bog and heath communities. Equally, fringing habitats are dependent on the lake, particularly its water levels, and support wetland communities and species of conservation concern. Many of the fringing wetland habitats support higher invertebrate and plant species richness than the lake habitats themselves

3130 Oligotrophic to mesotrophic standing waters with vegetation of the Littorelletea uniflorae and/or Isoeto-Nanojuncetea

To maintain the favourable conservation condition of Oligotrophic to mesotrophic standing waters with vegetation of the Littorelletea uniflorae and/or Isoeto-Nanojuncetea in Connemara Bog Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	This SAC has a very large number of lakes and ponds (>1,100, based on 1:5,000 OSi data). Most are likely to contain lake habitats 3110 and/or 3160; however, records of <i>Najas flexilis</i> demonstrate that lake habitat 3130 also occurs. The indicative distribution of habitat 3130 in the SAC is based on records for this characteristic species (see maps 6 and 9). Potential 3130 lakes are also identified overlying lake marbles or the Metagabbro Suite. The habitat is likely to be more widespread, as a result of basic influences from such geologies or coastal sands. Two measures of extent should be used: 1. the area of the lake itself and; 2. the extent of the vegetation communities/zones that typify the habitat. Further information relating to all attributes is provided in the lake habitats supporting document for the purposes of site-specific conservation objectives and Article 17 reporting (O Connor, 2015) and the <i>Najas flexilis</i> supporting document
Habitat distribution	Occurrence	No decline, subject to natural processes	The characteristics and distribution of habitat 3130 in Ireland are not yet fully understood. The Annex II macrophyte slender naiad (<i>Najas flexilis</i>) is considered to be characteristic of the habitat and has been recorded from 11 lakes across the SAC
Typical species	Occurrence	Typical species present, in good condition, and demonstrating typical abundances and distribution	For lists of typical plant species, see Article 17 habitat assessment for lake habitat 3130 (NPWS, 2013b), O Connor (2015) and the <i>Najas flexilis</i> supporting document
Vegetation composition: characteristic zonation	Occurrence	All characteristic zones should be present, correctly distributed and in good condition	The characteristic zonation of lake habitat 3140 has been described (Roden and Murphy, 2013; in prep.), however significant further work is necessary to describe the characteristic zonation and other spatial patterns in the remaining four lake habitats
Vegetation distribution: maximum depth	Metres	Maintain maximum depth of vegetation, subject to natural processes	The maximum depth of vegetation is likely to be specific to the lake shoreline in question. An indicative target of >6m has been developed for hard water lakes (3140) (see Roden and Murphy, 2013; in prep.). Indicative targets will be developed for the other lake habitats with time
Hydrological regime: water level fluctuations	Metres	Maintain appropriate natural hydrological regime necessary to support the habitat	Fluctuations in lake water level are typical in Ireland, but can be amplified by activities such as abstraction and drainage. Increased water level fluctuations can increase wave action, up-root vegetation, increase turbidity, alter the substratum and lead to release of nutrients from the sediment. The hydrological regime of the lakes must be maintained so that the area, distribution and depth of the lake habitat and its constituent/characteristic vegetation zones and communities are not reduced

Lake substratum quality	Various	Maintain appropriate substratum type, extent and chemistry to support the vegetation	Research is required to further characterise the substratum types (particle size and origin) and substratum quality (notably pH, calcium, iron and nutrient concentrations) favoured by each of the five Annex I lake habitats in Ireland. It is likely that the soft water lake habitat with base-rich influences (3130) is associated with a range of substrate types that are more productive/base-rich relative to the substratum of 3110. Substratum particle size is likely to vary with depth and along the shoreline within a single lake, however it should be noted that <i>Najas flexilis</i> is typically found on soft substrata of mud, silt or fine sand (Preston and Croft, 2001, Roden, 2002; 2004)
Water quality: transparency	Metres	Maintain appropriate Secchi transparency. There should be no decline in Secchi depth/transparency	Transparency relates to light penetration and, hence, to the depth of colonisation of vegetation. It can be affected by phytoplankton blooms, water colour and turbidity. A target has been set for hard water lakes (3140), however specific targets have yet to be established for the remaining lake habitats. Habitat 3130 is associated with clear water, as evidenced by the growth of the character species <i>Najas flexilis</i> at depths of up to 10m. The OECD fixed boundary system set transparency targets for oligotrophic lakes of $\geq 6\text{m}$ annual mean Secchi disk depth, and $\geq 3\text{m}$ annual minimum Secchi disk depth. There is likely to be some variation across lakes with habitat 3130 in Secchi depth and site-specific conditions should also be considered
Water quality: nutrients	$\mu\text{g/l P}$; mg/l N	The concentration of nutrients in the water column should be sufficiently low to prevent changes in species composition or habitat condition	3130 is associated with high water quality, with naturally low dissolved nutrients. It is naturally more productive than 3110, probably reflecting higher concentrations of nutrients such as calcium, rather than P alone. 3130 may reach favourable condition slightly above the oligotrophic boundary for nutrients, but in the absence of habitat-specific targets, the targets are Water Framework Directive) WFD 'High Status' or oligotrophic (OECD, 1982). The "good-moderate" boundary is too enriched to support the habitat. Annual average TP concentration should be $\leq 10\mu\text{g/l TP}$, average annual total ammonia should be $\leq 0.040\text{mg/l N}$ and annual 95th percentile for total ammonia should be $\leq 0.090\text{mg/l N}$. Where nutrient concentrations are lower, there should be no upward trend in nutrient concentrations. See also The European Communities Environmental Objectives (Surface Waters) Regulations 2009
Water quality: phytoplankton biomass	$\mu\text{g/l Chlorophyll } a$	Maintain appropriate water quality to support the habitat, including high chlorophyll <i>a</i> status	3130 is associated with high water quality, and naturally low algal growth. As for nutrients, the targets are WFD 'High Status' or oligotrophic (OECD, 1982). The "good-moderate" boundary is too enriched to support the habitat. The average growing season (March-October) chlorophyll <i>a</i> concentration must be $< 5.8\mu\text{g/l}$. The annual average chlorophyll <i>a</i> should be $< 2.5\mu\text{g/l}$ and the annual peak chlorophyll <i>a</i> should be $< 8.0\mu\text{g/l}$. Where a lake has a chlorophyll <i>a</i> concentration that is lower than this target, there should be no decline within class, i.e. no upward trend in phytoplankton biomass. See also The European Communities Environmental Objectives (Surface Waters) Regulations 2009
Water quality: phytoplankton composition	EPA phytoplankton composition metric	Maintain appropriate water quality to support the habitat, including high phytoplankton composition status	The EPA has developed a phytoplankton composition metric for nutrient enrichment of Irish lakes. As for other water quality indicators, habitat 3130 is considered to require WFD high status

Water quality: attached algal biomass	Algal cover and EPA phytobenthos metric	Maintain trace/ absent attached algal biomass (< 5% cover) and high phytobenthos status	Nutrient enrichment can favour epiphytic and epipellic algae that can out-compete the submerged vegetation. The cover abundance of attached algae in lake habitat 3130 should, therefore, be trace/ absent (<5% cover). EPA phytobenthos can be used as an indicator of changes in attached algal biomass. As for other water quality indicators, habitat 3130 requires high phytobenthos status
Water quality: macrophyte status	EPA macrophyte metric (The Free Index)	Maintain high macrophyte status	Nutrient enrichment can favour more competitive submerged macrophyte species that out-compete the typical and characteristic species for the soft water lake habitat with base-rich influences (3130). The EPA monitors macrophyte status for Water Framework Directive purposes using the 'Free Index'. The target for habitat 3130 is high status or an Ecological Quality Ratio (EQR) for lake macrophytes of ≥ 0.90 , as defined in Schedule Five of the European Communities Environmental Objectives (Surface Waters) Regulations 2009
Acidification status	pH units; mg/l	Maintain appropriate water and sediment pH, alkalinity and cation concentrations to support the habitat, subject to natural processes	Acidification can impact on species abundance and composition in habitat 3130. Acidification reduces the abundance and reproductive capacity of <i>Najas flexilis</i> (Wingfield et al., 2004). The specific requirements of habitat 3130, in terms of water and sediment pH, alkalinity and cation concentration, have not been determined. In line with targets for <i>Najas flexilis</i> , median pH values should be greater than 7 pH units. Water and sediment alkalinity and concentrations of cations (notably calcium) should be appropriate to the habitat. The target for WFD Acidification/Alkalisiation status is high. Maximum pH should be <9.0 pH units, in line with the surface water standards. See The European Communities Environmental Objectives (Surface Waters) Regulations 2009
Water colour	mg/l PtCo	Maintain appropriate water colour to support the habitat	Increased water colour and turbidity decrease light penetration and can reduce the area of available habitat for lake macrophytes, particularly at the lower euphotic depths. The primary source of increased water colour in Ireland is disturbance to peatland. No habitat-specific or national standards for water colour currently exist. Studies have shown median colour concentrations in Irish lakes of 38mg/l PtCo (Free, et al., 2000) and 33mg/l PtCo (Free et al., 2006). It is likely that the water colour in all Irish lake habitats would naturally be <50mg/l PtCo. Water colour is generally <30mg/l PtCo or, more naturally, <20mg/l PtCo in lakes with habitat 3130, where the peatland in the lake's catchment is intact
Dissolved organic carbon (DOC)	mg/l	Maintain appropriate organic carbon levels to support the habitat	Dissolved (and particulate) organic carbon (OC) in the water column is linked to water colour and acidification (organic acids). Increasing DOC in water has been documented across the Northern Hemisphere, including afforested peatland catchments in Ireland. Damage and degradation of peatland, leading to decomposition of peat is likely to be the predominant source of OC in Ireland. OC in water promotes decomposition by fungi and bacteria that, in turn, releases dissolved nutrients. The increased biomass of decomposers can also impact directly on the characteristic lake communities through shading, competition, etc.

Turbidity	Nephelometric turbidity units/ mg/l SS/ other appropriate unit	Maintain appropriate turbidity to support the habitat	Turbidity can significantly affect the quantity and quality of light reaching rooted and attached vegetation and can, therefore, impact on lake habitats. The settlement of higher loads of inorganic or organic material on lake vegetation communities may also have impacts on sensitive, delicate species. Turbidity can increase as a result of re-suspension of material within the lake, higher loads entering the lake, or eutrophication. Turbidity measurement and interpretation is challenging. As a result, it is likely to be difficult to set habitat-specific targets for turbidity in lakes
Fringing habitat: area	Hectares	Maintain the area and condition of fringing habitats necessary to support the natural structure and functioning of habitat 3130	Most lake shorelines have fringing habitats of reedswamp, other swamp, fen, marsh or wet-woodland that intergrade with and support the structure and functions of the lake habitat. Equally, fringing habitats are dependent on the lake, particularly its water levels, and support wetland communities and species of conservation concern. Many of the fringing wetland habitats support higher invertebrate and plant species richness than the lake habitats themselves

3160 Natural dystrophic lakes and ponds

To maintain the favourable conservation condition of Natural dystrophic lakes and ponds in Connemara Bog Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	This SAC has a very large number of lakes and ponds (more than 1,100 from 1:5,000 OSI data). Most are surrounded by peatland and are likely to contain lake habitats 3110 and/or 3160. The selection of the SAC for habitat 3160 was based on its widespread distribution in areas of low-lying blanket bog; dystrophic ponds being found wherever the peat is deep and the water table high. The habitat is considered to be of high conservation value in the SAC. Two measures of extent should be used: 1. the area of the lake itself and; 2. the extent of the vegetation communities/zones that typify the habitat. For further information on the distribution, vegetation and morphology of the habitat in the site, see Douglas and Grogan (1987) and Douglas et al. (1989). Further information relating to all attributes is provided in the lake habitats supporting document for the purposes of site-specific conservation objectives and Article 17 reporting (O Connor, 2015)
Habitat distribution	Occurrence	No decline, subject to natural processes	As noted above, it is likely that the habitat is widespread and of high conservation value in the site (see map 6)
Typical species	Occurrence	Typical species present, in good condition, and demonstrating typical abundances and distribution	For lists of typical plant and invertebrate species, see Article 17 habitat assessment for 3160 (NPWS, 2013) and the lake habitats supporting document for the purposes of site-specific conservation objectives and Article 17 reporting (O Connor, 2015)
Vegetation composition: characteristic zonation	Occurrence	All characteristic zones should be present, correctly distributed and in good condition	The characteristic zonation of lake habitat 3140 has been described (Roden and Murphy, 2013; in prep.), however significant further work is necessary to describe the characteristic zonation and other spatial patterns in the other Annex I lake habitats. Spatial patterns are likely to be relatively simple in 3160 lakes and ponds, with limited zonation
Vegetation distribution: maximum depth	Metres	Maintain maximum depth of vegetation, subject to natural processes	The maximum depth of vegetation is likely to be specific to the lake shoreline in question. An indicative target of >6m has been developed for hard water lakes (3140) (see Roden and Murphy, 2013; in prep.). Indicative targets will be developed for the other lake habitats with time. Lakes in the SAC typically have very clear water and, therefore, maximum depth is expected to be large
Hydrological regime: water level fluctuations	Metres	Maintain appropriate natural hydrological regime necessary to support the habitat	Fluctuations in lake water level are typical in Ireland, but can be amplified by activities such as abstraction and drainage. Increased water level fluctuations can increase wave action, up-root vegetation, increase turbidity, alter the substratum and lead to release of nutrients from the sediment. The hydrological regime of the lakes must be maintained so that the area, distribution and depth of the lake habitat and its constituent/characteristic vegetation zones and communities are not reduced. Owing to their size and the sensitivity of peatland, 3160 lakes and ponds can easily be damaged or destroyed by drainage
Lake substratum quality	Various	Maintain appropriate substratum type, extent and chemistry to support the vegetation	Research is required to further characterise the substratum types (particle size and origin) and substratum quality (notably pH, calcium, iron and nutrient concentrations) favoured by each of the five Annex I lake habitats in Ireland. It is likely that habitat 3160 is associated with nutrient-poor peat and silt substrates

Water quality: transparency	Metres	Maintain appropriate Secchi transparency. There should be no decline in Secchi depth/transparency	Transparency relates to light penetration and, hence, to the depth of colonisation of vegetation. It can be affected by phytoplankton blooms, water colour and turbidity. A target has been set for hard water lakes (3140), however specific targets have yet to be established for the remaining lake habitats. Habitat 3160 is associated with very clear water. The OECD fixed boundary system set transparency targets for ultra-oligotrophic lakes of $\geq 12\text{m}$ annual mean Secchi disk depth, and $\geq 6\text{m}$ annual minimum Secchi disk depth
Water quality: nutrients	$\mu\text{g/l P}$ or mg/l N	Maintain the concentration of nutrients in the water column to sufficiently low levels to support the habitat and its typical species	As a nutrient poor habitat, oligotrophic and Water Framework Directive 'high' status targets apply. Where a lake has nutrient concentrations that are lower than these targets, there should be no decline within class, i.e. no upward trend in nutrient concentrations. For 3160 lakes and ponds, annual average TP concentration should be $\leq 5\mu\text{g/l TP}$, average annual total ammonia concentration should be $\leq 0.040\text{mg/l N}$ and annual 95th percentile for total ammonia should be $\leq 0.090\text{mg/l N}$. See also The European Communities Environmental Objectives (Surface Waters) Regulations 2009
Water quality: phytoplankton biomass	$\mu\text{g/l Chlorophyll } a$	Maintain appropriate water quality to support the habitat, including high chlorophyll <i>a</i> status	Oligotrophic and Water Framework Directive 'high' status targets apply to habitat 3160. The average growing season (March-October) chlorophyll <i>a</i> concentration must be $< 5.8\mu\text{g/l}$ (The European Communities Environmental Objectives (Surface Waters) Regulations 2009). Where a lake has a chlorophyll <i>a</i> concentration that is lower than this target, there should be no decline within class, i.e. no upward trend in phytoplankton biomass. The OECD targets may be more appropriate for habitat 3160: annual average chlorophyll <i>a</i> concentration $< 1\mu\text{g/l}$ and annual peak chlorophyll <i>a</i> concentration $\leq 2.5\mu\text{g/l}$. See also The European Communities Environmental Objectives (Surface Waters) Regulations 2009
Water quality: phytoplankton composition	EPA phytoplankton composition metric	Maintain appropriate water quality to support the habitat, including high phytoplankton composition status	The EPA has developed a phytoplankton composition metric for nutrient enrichment of Irish lakes. As for other water quality indicators, habitat 3160 requires WFD high status
Water quality: attached algal biomass	Algal cover and EPA phytobenthos metric	Maintain trace/ absent attached algal biomass ($< 5\%$ cover) and high phytobenthos status	Nutrient enrichment can favour epiphytic and epipelagic algae that can out-compete the submerged vegetation. The cover abundance of attached algae in 3160 lakes and ponds should, therefore, be trace/ absent ($< 5\%$ cover). EPA phytobenthos can be used as an indicator of changes in attached algal biomass. As for other water quality indicators, habitat 3160 requires high phytobenthos status
Water quality: macrophyte status	EPA macrophyte metric (The Free Index)	Maintain high macrophyte status	Nutrient enrichment can favour more competitive submerged macrophyte species that out-compete the typical and characteristic species for the lake habitat. The EPA monitors macrophyte status for Water Framework Directive purposes using the 'Free Index'. The target for 3160 lakes and ponds is high status or an Ecological Quality Ratio (EQR) for lake macrophytes of ≥ 0.90 , as defined in Schedule Five of the European Communities Environmental Objectives (Surface Waters) Regulations 2009

Acidification status	pH units; mg/l	Maintain appropriate water and sediment pH, alkalinity and cation concentrations to support the habitat, subject to natural processes	Acidification can impact on species abundance and composition in soft water lake habitats. Although EC (2013) describes habitat 3160 as having pH 3-6, Drinan (2012) found mean pHs of 5.16 and 5.62 in upland and lowland 3160 lakes, respectively. The target for habitat 3160 is pH >4.5 and <9.0, in line with the surface water standards for soft waters (where water hardness is ≤100mg/l CaCO ₃). See Schedule Five of the European Communities Environmental Objectives (Surface Waters) Regulations 2009. The specific requirements of lake habitat 3160, in terms of water and sediment pH, alkalinity and cation concentration, have not been determined
Water colour	mg/l PtCo	Maintain appropriate water colour to support the habitat	Increased water colour and turbidity decrease light penetration and can reduce the area of available habitat for lake macrophytes, particularly at the lower euphotic depths. The primary source of increased water colour in Ireland is disturbance to peatland. No habitat-specific or national standards for water colour currently exist. Studies have shown median colour concentrations in Irish lakes of 38mg/l PtCo (Free, et al., 2000) and 33mg/l PtCo (Free et al., 2006). It is likely that the water colour in all Irish lake habitats would naturally be <50mg/l PtCo. Water colour can be very low (<20mg/l PtCo or even <10mg/l PtCo) in 3160 lakes and ponds, where the peatland in the lake's catchment is intact
Dissolved organic carbon (DOC)	mg/l	Maintain appropriate organic carbon levels to support the habitat	Dissolved (and particulate) organic carbon (OC) in the water column is linked to water colour and acidification (organic acids). Increasing DOC in water has been documented across the Northern Hemisphere, including afforested peatland catchments in Ireland. Damage and degradation of peatland, leading to decomposition of peat is likely to be the predominant source of OC in Ireland. OC in water promotes decomposition by fungi and bacteria that, in turn, releases dissolved nutrients. The increased biomass of decomposers can also impact directly on the characteristic lake communities through shading, competition, etc.
Turbidity	Nephelometric turbidity units/ mg/l SS/ other appropriate units	Maintain appropriate turbidity to support the habitat	Turbidity can significantly affect the quantity and quality of light reaching rooted and attached vegetation and can, therefore, impact on lake habitats. The settlement of higher loads of inorganic or organic material on lake vegetation communities may also have impacts on sensitive, delicate species. Turbidity can increase as a result of re-suspension of material within the lake, higher loads entering the lake, or eutrophication. Turbidity measurement and interpretation is challenging. As a result, it is likely to be difficult to set habitat-specific targets for turbidity in lakes
Fringing habitat	Area	Maintain the area and condition of fringing habitats necessary to support the natural structure and functioning of habitat 3160	Most 3160 lake and pond shorelines intergrade with blanket bog, flush, poor fen or heath habitats and these support the structure and functions of the lake habitat. Equally, fringing habitats are dependent on the lake, particularly its water levels, and can support wetland communities and species of conservation concern

Conservation Objectives for : Connemara Bog Complex SAC [002034]

3260 Water courses of plain to montane levels with the *Ranunculion fluitantis* and *Callitricho-Batrachion* vegetation

To maintain the favourable conservation condition of Water courses of plain to montane levels with the *Ranunculion fluitantis* and *Callitricho-Batrachion* vegetation in Connemara Bog Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Kilometres	Area stable or increasing, subject to natural processes	Selection of this SAC for 3260 used a broad interpretation and little is known of the distribution of high conservation value sub-types. Rivers and streams are widespread and abundant in the SAC (see map 7). Note: rooted macrophytes should be absent or trace (<5% cover) in freshwater pearl mussel (<i>Margaritifera margaritifera</i>) habitat. The SAC overlaps with three <i>Margaritifera</i> sensitive areas: Owenriff (priority SAC catchment), Knock and Ballynahinch/Recess (see map 7). Mussels occur within the SAC in the Derrygauna River (tributary of the Owenriff), and the Owentooey and Recess Rivers. Within the Owenriff catchment, the freshwater pearl mussel (1029) conservation objective for Lough Corrib SAC (000297) takes precedence, because the mussel requires environmental conditions close to natural background levels
Habitat distribution	Occurrence	No decline, subject to natural processes	The description of 3260 is broad, from upland bryophyte/macroalgal dominated river stretches, to lowland depositing rivers with pondweeds and starworts (EC, 2013), and further study is needed of Irish sub-types and their conservation value. As noted above, little is known about the distribution of the habitat and its sub-types in the site. Heuff (1987) surveyed the Vougheen Stream, the outflow from Lough Cam (both in Roundstone Blanket Bog) and the Owenboliska. Rivers in the SAC are, naturally, very nutrient-poor, with five High Status sites on four rivers noted by Ni Chathain et al. (2013). Rivers and streams can be peaty and slow-flowing in flat blanket bog, and spatey, often with cascades, in more sloping areas. They are frequently base-poor, but <i>Margaritifera</i> rivers tend to have base-rich influences. See Williams (2009) and DEHLG (2010) for information on macrophytes in the Owenriff catchment
Hydrological regime: river flow	Metres per second	Maintain appropriate hydrological regimes	Any high conservation value sub-types in the site will be associated with natural hydrology. A natural flow regime is required for both plant communities and channel geomorphology to be in favourable condition, exhibiting typical dynamics for the river type (Hatton-Ellis and Grieve, 2003). For many of the sub-types of this habitat, high flows are required to maintain the substratum necessary for the characteristic species. Flow variation can be particularly important, with high and flood flows being critical to the hydromorphology. Peatlands can also have slow-flowing or ponded streams and rivers, with biotic communities likely to resemble those in associated lakes
Hydrological regime: groundwater discharge	Metres per second	Maintain appropriate hydrological regimes	The groundwater contribution to rivers in the SAC is likely to be small, owing to the geology and dominance of blanket peat soils. Even small groundwater contributions, however, can significantly alter the hydrochemistry, particularly where there is basic bedrock (e.g. lake marbles or the Metagabbro Suite) and/or subsoils

Substratum composition: particle size range	Millimetres	Maintain appropriate substratum particle size range, quantity and quality, subject to natural processes	Although many of the high conservation value sub-types are dominated by coarse substrata, for certain sub-types, notably tidal forms, fine substrata are required. Peat is a common substratum in gently sloping rivers within the site. The size and distribution of particles is largely determined by the river flow. The chemical composition (particularly minerals and nutrients) of the substratum is also important. The quality of finer sediment particles is a notable driver for rooted plant communities
Water quality	Various	Maintain appropriate water quality to support the natural structure and functioning of the habitat	The specific targets may vary among sub-types. The rivers within the SAC are considered to be naturally very nutrient poor and, therefore, to typically require Water Framework Directive high status, in terms of nutrient and oxygenation standards, and EQRs (Ecological Quality Ratios) for macroinvertebrates and phytobenthos
Vegetation composition: typical species	Occurrence	Typical species of the relevant habitat sub-type should be present and in good condition	The sub-types of this habitat are poorly understood and their typical species have not yet been fully defined. The typical species may include higher plants, bryophytes, macroalgae and microalgae, and invertebrates. The only known Irish sites for <i>Luronium natans</i> are in the SAC, and although generally considered to be a lake species, it has been found in two streams (Rich et al., 1995; Curtis et al., 2012). The status of <i>Luronium natans</i> in Ireland is currently uncertain, as it is unclear whether it is native to these sites, or was introduced (Preston and Croft, 2001). The uncommon pondweed hybrid, <i>Potamogeton x sparganiifolius</i> is known from the Owenmore (Ballynahinch) River (Preston et al., 1991)
Floodplain connectivity: area	Hectares	Area of active floodplain at, and upstream of the habitat, necessary to support all sub-types of the habitat, should be maintained	River connectivity with the floodplain is important for the functioning of this habitat. Channels with a naturally functioning floodplain are better able to maintain habitat and water quality (Hatton-Ellis and Grieve, 2003). Floodplain connectivity is particularly important in terms of sediment sorting and nutrient deposition. High conservation value rivers are intimately connected to floodplain habitats and function as important wildlife corridors, connecting otherwise isolated or fragmented habitats in the wider countryside (Hatton-Ellis and Grieve, 2003)
Riparian habitat: area	Hectares	Maintain the area and condition of fringing habitats necessary to support the habitat and its sub-types	Riparian habitats, including those along lake fringes, even where they do not form part of a natural floodplain, are an integral part of the structure and functioning of river systems. Fringing habitats can contribute to the aquatic food web (e.g. allochthonous matter such as leaf fall), provide habitat (refuge and resources) for certain life-stages of fish, birds and aquatic invertebrates, assist in the settlement of fine suspended material, protect banks from erosion and contribute to nutrient cycling. Shade may also be important in suppressing algal growth in enriched rivers and moderating temperatures. Equally, fringing habitats are dependent on rivers/lakes, particularly their water levels, and support wetland communities and species of conservation concern

Conservation Objectives for : Connemara Bog Complex SAC [002034]

4010 Northern Atlantic wet heaths with *Erica tetralix*

To restore the favourable conservation condition of Northern Atlantic wet heaths with *Erica tetralix* in Connemara Bog Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Stable or increasing, subject to natural processes	Northern Atlantic wet heaths with <i>Erica tetralix</i> has not been mapped in detail for this SAC and thus total area of the qualifying habitat is unknown. It occurs in mosaic/in association with with other habitats, including other heath habitats (4030), blanket bogs (7130), lakes (3110, 3160), rocky outcrops and grasslands. Further information can be found in Douglas and Grogan (1987); Douglas et al. (1989); NPWS internal files; commonage framework plans for part or all of the following agricultural units: GA2, GA4, GA5, GA6, GA7, GA8, GA9, GA11, GA15, GA16, GA18, GA21
Habitat distribution	Occurrence	No decline, subject to natural processes	See note on area above
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil nutrient status within natural range	Relevant nutrients and their natural ranges are yet to be defined. However, nitrogen deposition is noted as being relevant to this habitat in NPWS (2013)
Community diversity	Abundance of variety of vegetation communities	Maintain variety of vegetation communities, subject to natural processes	Further information on vegetation communities associated with this habitat is presented in Perrin et al. (2014)
Vegetation composition: cross-leaved heath	Occurrence within 20m of a representative number of 2m x 2m monitoring stops	Presence of cross-leaved heath (<i>Erica tetralix</i>) near each monitoring stop	Attribute and target based on Perrin et al. (2014)
Vegetation composition: positive indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of positive indicator species at least 50%	Attribute and target based on Perrin et al. (2014) where the list of positive indicator species for this habitat is also given
Vegetation composition: lichens and bryophytes	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of <i>Cladonia</i> and <i>Sphagnum</i> species, <i>Racomitrium lanuginosum</i> and pleurocarpous mosses at least 10%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: ericoid species and crowberry	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of ericoid species and crowberry (<i>Empetrum nigrum</i>) at least 15%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: dwarf shrub species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of dwarf shrubs less than 75%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: negative indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of negative indicator species less than 1%	Attribute and target based on Perrin et al. (2014) where the list of negative indicator species is also given
Vegetation composition: non-native species	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of non-native species less than 1%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: native trees and shrubs	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of scattered native trees and shrubs less than 20%	Attribute and target based on Perrin et al. (2014)

Vegetation composition: bracken	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of bracken (<i>Pteridium aquilinum</i>) less than 10%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: soft rush	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of soft rush (<i>Juncus effusus</i>) less than 10%	Attribute and target based on Perrin et al. (2014)
Vegetation structure: <i>Sphagnum</i> condition	Condition at a representative number of 2m x 2m monitoring stops	Less than 10% of the <i>Sphagnum</i> cover is crushed, broken and/or pulled up	Attribute and target based on Perrin et al. (2014)
Vegetation structure: signs of browsing	Percentage of shoots browsed at a representative number of 2m x 2m monitoring stops	Less than 33% collectively of the last complete growing season's shoots of ericoids, crowberry (<i>Empetrum nigrum</i>) and bog-myrtle (<i>Myrica gale</i>) showing signs of browsing	Attribute and target based on Perrin et al. (2014)
Vegetation structure: burning	Occurrence in local vicinity of a representative number of 2m x 2m monitoring stops	No signs of burning in sensitive areas, into the moss, liverwort or lichen layer or exposure of peat surface due to burning	Attribute and target based on Perrin et al. (2014) where the list of sensitive areas is also presented
Physical structure: disturbed bare ground	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of disturbed bare ground less than 10%	Attribute and target based on Perrin et al. (2014)
Physical structure: drainage	Percentage cover in local vicinity of a representative number of monitoring stops	Area showing signs of drainage from heavy trampling, tracking or ditches less than 10%	Attribute and target based on Perrin et al. (2014)
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat	This includes species listed in the Flora (Protection) Order 2015 and/or the red data book (Curtis and McGough, 1988)

Conservation Objectives for : Connemara Bog Complex SAC [002034]

4030 European dry heaths

To restore the favourable conservation condition of European dry heaths in Connemara Bog Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	European dry heaths has not been mapped in detail for this SAC and thus total area of the qualifying habitat is unknown. It occurs in mosaic/in association with other habitats, including other heath habitats (4010, 4060), rocky outcrops and grasslands. Further information can be found in Douglas and Grogan (1987); Douglas et al. (1989); NPWS internal files; commonage framework plans for part or all of the following agricultural units: GA2, GA4, GA5, GA6, GA7, GA8, GA9, GA11, GA15, GA16, GA18, GA21
Habitat distribution	Occurrence	No decline, subject to natural processes	See note on area above
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil nutrient status within natural range	Relevant nutrients and their natural ranges are yet to be defined. However, nitrogen deposition is noted as being relevant to this habitat in NPWS (2013)
Community diversity	Abundance of variety of vegetation communities	Maintain variety of vegetation communities, subject to natural processes	Further information on these communities is presented in Perrin et al. (2014)
Vegetation composition: lichens and bryophytes	Number of species at a representative number of 2m x 2m monitoring stops	Number of bryophyte or non-crustose lichen species present at each monitoring stop is least three, excluding <i>Campylopus</i> and <i>Polytrichum</i> mosses	Attribute and target based on Perrin et al. (2014)
Vegetation composition: number of positive indicator species	Number at a representative number of 2m x 2m monitoring stops	Number of positive indicator species present at each monitoring stop is at least two	Attribute and target based on Perrin et al. (2014) where the list of positive indicator species for this habitat, which is composed of dwarf shrubs, is also presented
Vegetation composition: cover of positive indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of positive indicator species at least 50% for siliceous dry heath and 50-75% for calcareous dry heath	Attribute and target based on Perrin et al. (2014) where the list of positive indicator species for this habitat, which is composed of dwarf shrubs, is also given
Vegetation structure: dwarf shrub composition	Percentage cover at a representative number of 2m x 2m monitoring stops	Proportion of dwarf shrub cover composed collectively of bog-myrtle (<i>Myrica gale</i>), creeping willow (<i>Salix repens</i>) and western gorse (<i>Ulex gallii</i>) is less than 50%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: negative indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of negative indicator species less than 1%	Attribute and target based on Perrin et al. (2014) where the list of negative indicator species is also given
Vegetation composition: non-native species	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of non-native species less than 1%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: native trees and shrubs	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of scattered native trees and shrubs less than 20%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: bracken	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of bracken (<i>Pteridium aquilinum</i>) less than 10%	Attribute and target based on Perrin et al. (2014)

Vegetation composition: soft rush	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of soft rush (<i>Juncus effusus</i>) less than 10%	Attribute and target based on Perrin et al. (2014)
Vegetation structure: senescent ling	Percentage cover at a representative number of 2m x 2m monitoring stops	Senescent proportion of ling (<i>Calluna vulgaris</i>) cover less than 50%	Attribute and target based on Perrin et al. (2014)
Vegetation structure: signs of browsing	Percentage of shoots browsed at a representative number of 2m x 2m monitoring stops	Less than 33% collectively of the last complete growing season's shoots of ericoids showing signs of browsing	Attribute and target based on Perrin et al. (2014)
Vegetation structure: burning	Occurrence in local vicinity of a representative number of monitoring stops	No signs of burning in sensitive areas	Attribute and target based on Perrin et al. (2014) where the list of sensitive areas is also presented
Vegetation structure: growth phases of ling	Percentage cover in local vicinity of a representative number of monitoring stops	Outside sensitive areas, all growth phases of ling should occur throughout, with at least 10% of cover in the mature phase	Attribute and target based on Perrin et al. (2014)
Vegetation structure: disturbed bare ground	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of disturbed bare ground less than 10%	Attribute and target based on Perrin et al. (2014)
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat and no decline in status of hepatic mats associated with this habitat	This includes species listed in the Flora (Protection) Order 2015 and/or the red data book (Curtis and McGough, 1988)

Conservation Objectives for : Connemara Bog Complex SAC [002034]

6410 *Molinia* meadows on calcareous, peaty or clayey-silt-laden soils (*Molinion caeruleae*)

To maintain the favourable conservation condition of *Molinia* meadows on calcareous, peaty or clayey-silt-laden soils (*Molinion caeruleae*) in Connemara Bog Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	This Annex I grassland habitat has not been mapped for this SAC and thus the total area of the qualifying habitat is unknown. It is likely to occur as rather small fragmented areas on wet acid soils (NPWS internal files)
Habitat distribution	Occurrence	No decline, subject to natural processes	See note for area above
Vegetation composition: typical species	Number at a representative number of monitoring stops	At least seven positive indicator species present, including one "high quality" species as listed in O'Neill et al. (2013)	List of positive indicator species, including high quality species, identified by O'Neill et al. (2013). Note that purple moor-grass (<i>Molinia caerulea</i>) is a positive indicator species, but not necessarily an essential component of the habitat
Vegetation composition: negative indicator species	Percentage at a representative number of monitoring stops	Negative indicator species collectively not more than 20% cover, with cover by an individual species not more than 10%	Attribute and target based on O'Neill et al. (2013)
Vegetation composition: non-native species	Percentage at a representative number of monitoring stops	Cover of non-native species not more than 1%	List of negative indicator species identified by O'Neill et al. (2013)
Vegetation composition: moss species	Percentage at a representative number of monitoring stops	Hair mosses (<i>Polytrichum</i> spp.) not more than 25% cover	Attribute and target based on O'Neill et al. (2013)
Vegetation structure: woody species and bracken	Percentage at a representative number of monitoring stops	Cover of woody species and bracken (<i>Pteridium aquilinum</i>) not more than 5% cover	Attribute and target based on O'Neill et al. (2013)
Vegetation structure: broadleaf herb: grass ratio	Percentage at a representative number of monitoring stops	Broadleaf herb component of vegetation between 40 and 90%	Attribute and target based on O'Neill et al. (2013)
Vegetation structure: sward height	Percentage at a representative number of monitoring stops	At least 30% of sward between 10 and 80cm tall	Attribute and target based on O'Neill et al. (2013)
Vegetation structure: litter	Percentage at a representative number of monitoring stops	Litter cover not more than 25%	Attribute and target based on O'Neill et al. (2013)
Physical structure: bare ground	Percentage	Not more than 10% bare ground	Attribute and target based on O'Neill et al. (2010)
Physical structure: bare soil	Percentage at a representative number of monitoring stops	Not more than 10% bare soil	Attribute and target based on O'Neill et al. (2013)
Physical structure: disturbance	Square metres	Area showing signs of serious grazing or other disturbance less than 20m ²	Attribute and target based on O'Neill et al. (2013)

Conservation Objectives for : Connemara Bog Complex SAC [002034]

7130 Blanket bogs (* if active bog)

To restore the favourable conservation condition of Blanket bogs in Connemara Bog Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Stable or increasing, subject to natural processes	Blanket bogs has not been mapped in detail for this SAC and thus total area of the qualifying habitat is unknown. It occurs in mosaic/in association with other habitats, including lakes (3110, 3160), heath habitats (4010, 4030), transition mires (7150) and rocky outcrops. Further information can be found in Douglas and Grogan (1987); Douglas et al. (1989); NPWS internal files; commonage framework plans for part or all of the following agricultural units: GA2, GA4, GA5, GA6, GA7, GA8, GA9, GA11, GA15, GA16, GA18, GA21
Habitat distribution	Occurrence	No decline, subject to natural processes	See note on area above
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil nutrient status within natural range	Relevant nutrients and their natural ranges are yet to be defined. However, nitrogen deposition is noted as being relevant to this habitat in NPWS (2013)
Ecosystem function: peat formation	Active blanket bog as a proportion of the total area of Annex I blanket bog habitat	Active bog at least 99% of the total Annex I blanket bog area	Blanket bogs are considered active when "still supporting a significant area of vegetation that is normally peat forming" (EC, 2013)
Ecosystem function: hydrology	Flow direction, water levels, occurrence of drains and erosion gullies	Natural hydrology unaffected by drains and erosion	Drains and erosion gullies can affect the natural hydrological processes of blanket bog
Community diversity	Abundance of variety of vegetation communities	Maintain variety of vegetation communities, subject to natural processes	Further information on vegetation communities associated with this habitat is presented in Perrin et al. (2014)
Vegetation composition: positive indicator species	Number of species at a representative number of 2m x 2m monitoring stops	Number of positive indicator species at each monitoring stop is at least seven	Attribute and target based on Perrin et al. (2014) where the list of positive indicator species for this habitat is also given
Vegetation composition: lichens and bryophytes	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of bryophytes or lichens, excluding <i>Sphagnum fallax</i> , at least 10%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: potential dominant species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of each of the potential dominant species less than 75%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: negative indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of negative indicator species less than 1%	Attribute and target based on Perrin et al. (2014) where the list of negative indicator species for this habitat is also given
Vegetation composition: non-native species	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of non-native species less than 1%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: native trees and scrub	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of scattered native trees and shrubs less than 10%	Attribute and target based on Perrin et al. (2014)
Vegetation structure: <i>Sphagnum</i> condition	Condition of <i>Sphagnum</i> at a representative number of 2m x 2m monitoring stops	Less than 10% of the <i>Sphagnum</i> cover is crushed, broken and/or pulled up	Attribute and target based on Perrin et al. (2014)

Vegetation structure: signs of browsing	Percentage of shoots browsed at a representative number of 2m x 2m monitoring stops	Last complete growing season's shoots of ericoids, crowberry (<i>Empetrum nigrum</i>) and bog-myrtle (<i>Myrica gale</i>) showing signs of browsing collectively less than 33%	Attribute and target based on Perrin et al. (2014)
Vegetation structure: burning	Occurrence in local vicinity of a representative number of monitoring stops	No signs of burning in sensitive areas, into the moss, liverwort or lichen layer or exposure of peat surface due to burning	Attribute and target based on Perrin et al. (2014) where the list of sensitive areas is also presented
Physical structure: disturbed bare ground	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of disturbed bare ground less than 10%	Attribute and target based on Perrin et al. (2014)
Physical structure: drainage	Percentage area in local vicinity of a representative number of monitoring stops	Area showing signs of drainage from heavy trampling, tracking or ditches less than 10%	Attribute and target based on Perrin et al. (2014)
Physical structure: erosion	Occurrence in local vicinity of a representative number of monitoring stops	Less than 5% of the greater bog mosaic comprises erosion gullies and eroded areas	Attribute and target based on Perrin et al. (2014). The greater bog mosaic incorporates the blanket bog itself and associated vegetation types as well as non-vegetation cover types that appear to have been derived from former blanket bog including gravel, rock and running water
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat	This includes species listed in the Flora (Protection) Order 2015 and/or the red data book (Curtis and McGough, 1988)

Conservation Objectives for : Connemara Bog Complex SAC [002034]

7140 Transition mires and quaking bogs

To restore the favourable conservation condition of Transition mires and quaking bogs in Connemara Bog Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Transition mires and quaking bogs has not been mapped in detail for this SAC and thus total area of the qualifying habitat is unknown. It occurs in mosaic/in association with other habitats including blanket bogs (7130) and heath habitats (4010) and fen (7230). Further information can be found in Douglas and Grogan (1987); Douglas et al. (1989); NPWS internal files; commonage framework plans for part or all of the following agricultural units: GA2, GA4, GA5, GA6, GA7, GA8, GA9, GA11, GA15, GA16, GA18, GA21
Habitat distribution	Occurrence	No decline from current distribution, subject to natural processes	See note on area above
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil nutrient status within natural range	Relevant nutrients and their natural ranges are yet to be defined. However, nitrogen deposition is noted as being relevant to this habitat in NPWS (2013)
Community diversity	Abundance of variety of vegetation communities	Maintain variety of vegetation communities, subject to natural processes	Further information on vegetation communities associated with this habitat is presented in Perrin et al. (2014)
Vegetation composition: number of positive indicator species	Number at a representative number of 2m x 2m monitoring stops	Number of positive indicator species at least three for in-filling pools and flushes and at least six for fens	Attribute and target based on Perrin et al. (2014) where the list of positive indicator species for this habitat is also given
Vegetation composition: number of core positive indicator species	Number of species at a representative number of 2m x 2m monitoring stops	At least one core positive indicator species present	Attribute and target based on Perrin et al. (2014) where the list of positive indicator species for this habitat is also given
Vegetation composition: cover of positive indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of positive indicator species is at least 25%	Attribute and target based on Perrin et al. (2014) where the list of positive indicator species for this habitat is also given
Vegetation composition: negative indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of negative indicator species less than 1%	Attribute and target based on Perrin et al. (2014) where the list of negative indicator species for this habitat is also given
Vegetation composition: non-native species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of non-native species less than 1%	Attribute and target based on Perrin et al. (2014)
Vegetation structure: height	Percentage at a representative number of 2m x 2m monitoring stops	Proportion of live leaves and/or flowering shoots of vascular plants that are more than 15cm above the ground surface should be at least 50%	Attribute and target based on Perrin et al. (2014). This attribute is only applicable to fen and flush examples of the habitat, not to infilling pool examples
Physical structure: disturbed bare ground	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of disturbed bare ground less than 10%	Attribute and target based on Perrin et al. (2014)
Physical structure: drainage	Percentage area in local vicinity of a representative number of monitoring stops	Area showing signs of drainage from heavy trampling, tracking or ditches less than 10%	Attribute and target based on Perrin et al. (2014)

Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat	This includes species listed in the Flora (Protection Order 2015 and/or the red data book (Curtis and McGough, 1988))
-------------------------------------	--------------------------------	--	---

Conservation Objectives for : Connemara Bog Complex SAC [002034]

7150 Depressions on peat substrates of the Rhynchosporion

To restore the favourable conservation condition of Depressions on peat substrates of the Rhynchosporion in Connemara Bog Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Depressions on peat substrates of the Rhynchosporion has not been mapped in detail for this SAC and thus total area of the qualifying habitat is unknown. It overlaps with blanket bogs (7130). Further information can be found in Douglas and Grogan (1987); Douglas et al. (1989); NPWS internal files
Habitat distribution	Occurrence	No decline, subject to natural processes	See note on area above
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil nutrient status within natural range	Relevant nutrients and their natural ranges are yet to be defined. However, nitrogen deposition is noted as being relevant to this habitat in NPWS (2013)
Vegetation composition: positive indicator species	Number of species at a representative number of 2m x 2m monitoring stops	Number of positive indicator species at each monitoring stop is at least five	Attribute and target based on Perrin et al. (2014) where the list of positive indicator species for this habitat is also given
Vegetation composition: <i>Rhynchospora</i> spp.	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of white beaked sedge (<i>Rhynchospora alba</i>) and brown beaked sedge (<i>R. fusca</i>) at least 10%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: potential dominant species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of each of the potential dominant species less than 35%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: negative indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of negative indicator species less than 1%	Attribute and target based on Perrin et al. (2014) where the list of negative indicator species for this habitat is also given
Vegetation composition: non-native species	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of non-native species less than 1%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: native trees and scrub	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of scattered native trees and shrubs less than 10%	Attribute and target based on Perrin et al. (2014)
Vegetation structure: <i>Sphagnum</i> condition	Condition of <i>Sphagnum</i> at a representative number of 2m x 2m monitoring stops	Less than 10% of the <i>Sphagnum</i> cover is crushed, broken and/or pulled up	Attribute and target based on Perrin et al. (2014)
Vegetation structure: signs of browsing	Percentage of shoots browsed at a representative number of 2m x 2m monitoring stops	Last complete growing season's shoots of ericoids, crowberry (<i>Empetrum nigrum</i>) and bog-myrtle (<i>Myrica gale</i>) showing signs of browsing collectively less than 33%	Attribute and target based on Perrin et al. (2014)
Vegetation structure: burning	Occurrence in local vicinity of a representative number of monitoring stops	No signs of burning in sensitive areas, into the moss, liverwort or lichen layer or exposure of peat surface due to burning	Attribute and target based on Perrin et al. (2014) where the list of sensitive areas is also presented

Physical structure: disturbed bare ground	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of disturbed bare ground less than 10%	Attribute and target based on Perrin et al. (2014)
Physical structure: drainage	Percentage area in local vicinity of a representative number of monitoring stops	Area showing signs of drainage from heavy trampling, tracking or ditches less than 10%	Attribute and target based on Perrin et al. (2014)
Physical structure: erosion	Occurrence in local vicinity of a representative number of monitoring stops	Less than 5% of the greater bog mosaic comprises erosion gullies and eroded areas	Attribute and target based on Perrin et al. (2014). The greater bog mosaic incorporates the blanket bog itself, associated vegetation types as well as non-vegetation cover types that appear to have been derived from former blanket bog including gravel, rock and running water
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat	This includes species listed in the Flora (Protection) Order 2015 and/or the red data book (Curtis and McGough, 1988)

Conservation Objectives for : Connemara Bog Complex SAC [002034]

7230 Alkaline fens

To restore the favourable conservation condition of Alkaline fens in Connemara Bog Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Alkaline fens has not been mapped in detail for this SAC and thus total area of the qualifying habitat is unknown. It occurs in mosaic/in association with other habitats including blanket bogs (7130), transition mires (7140) and heath habitats (4010). Further information can be found in Douglas and Grogan (1987); Douglas et al. (1989); NPWS internal files; commonage framework plans for part or all of the following agricultural units: GA2, GA4, GA5, GA6, GA7, GA8, GA9, GA11, GA15, GA16, GA18, GA21
Habitat distribution	Occurrence	No decline from current distribution, subject to natural processes	See note on area above
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil nutrient status within natural range	Relevant nutrients and their natural ranges are yet to be defined. However, nitrogen deposition is noted as being relevant to this habitat in NPWS (2013)
Community diversity	Abundance of variety of vegetation communities	Maintain variety of vegetation communities, subject to natural processes	Further information on the communities associated with this habitat is presented in Perrin et al. (2014)
Vegetation composition: number of positive indicator species (brown mosses)	Number at a representative number of 2m x 2m monitoring stops	Number of brown moss species present at each monitoring stop is at least one	Attribute and target based on Perrin et al. (2014) where the list of positive indicator species for this habitat is also given
Vegetation composition: number of positive indicator species (vascular plants)	Number at a representative number of 2m x 2m monitoring stops	Number of positive vascular plant indicator species at least two for small-sedge flushes and at least three for black bog-rush (<i>Schoenus nigricans</i>) flush and bottle sedge (<i>Carex rostrata</i>) fen	Attribute and target based on Perrin et al. (2014) where the list of positive indicator species for this habitat is also given
Vegetation composition: cover of positive indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of brown moss species and positive vascular plant indicator species at least 20% for small-sedge flushes and at least 75% cover for black bog-rush (<i>Schoenus nigricans</i>) flush and bottle sedge (<i>Carex rostrata</i>) fen	Attribute and target based on Perrin et al. (2014) where the list of positive indicator species for this habitat is also given
Vegetation composition: negative indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of negative indicator species less than 1%	Attribute and target based on Perrin et al. (2014) where the list of negative indicator species for this habitat is also given
Vegetation composition: non-native species	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of non-native species less than 1%	Attribute and target based on Perrin et al. (2014)
Vegetation composition: native trees and scrub	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of scattered native trees and shrubs less than 10%	Attribute and target based on Perrin et al. (2014)

Vegetation composition: soft rush and common reed	Percentage cover in local vicinity of a representative number of monitoring stops	Total cover of soft rush (<i>Juncus effusus</i>) and common reed (<i>Phragmites australis</i>) less than 10%	Attribute and target based on Perrin et al. (2014)
Vegetation structure: height	Percentage at a representative number of 2m x 2m monitoring stops	Proportion of live leaves and/or flowering shoots of vascular plants that are more than 5cm above the ground surface should be at least 50%	Attribute and target based on Perrin et al. (2014)
Physical structure: disturbed bare ground	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of disturbed bare ground less than 10%	Attribute and target based on Perrin et al. (2014)
Physical structure: drainage	Percentage area in local vicinity of a representative number of monitoring stops	Area showing signs of drainage from heavy trampling, tracking or ditches less than 10%	Attribute and target based on Perrin et al. (2014)
Physical structure: tufa formations	Occurrence in local vicinity of a representative number of monitoring stops	Disturbed proportion of vegetation cover is less than 1%	Attribute and target based on Perrin et al. (2014)
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat	This includes species listed in the Flora (Protection) Order 2015 and/or the red data book (Curtis and McGough, 1988)

Conservation Objectives for : Connemara Bog Complex SAC [002034]

91A0 Old sessile oak woods with Ilex and Blechnum in the British Isles

To maintain the favourable conservation condition of Old sessile oak woods with *Ilex* and *Blechnum* in the British Isles in Connemara Bog Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes. See map 8 for surveyed areas	This SAC includes a number of areas of this Annex I woodland type, some of which have been recently surveyed and mapped by the National Survey of Native Woodlands (NSNW) (Perrin et al., 2008): sites no. 1600, 1602, 1930. NB there are other areas of this habitat within the SAC, such as on lake islands (NPWS internal files)
Habitat distribution	Occurrence	No decline, subject to natural processes. Surveyed locations are shown on map 8	See note on area above
Woodland size	Hectares	Area stable or increasing. Where topographically possible, "large" woods at least 25ha in size and "small" woods at least 3ha in size	The target areas for individual woodlands aim to reduce habitat fragmentation and benefit those species requiring 'deep' woodland conditions (Peterken, 2002). In some cases, topographical constraints may restrict expansion
Woodland structure: cover and height	Percentage and metres	Diverse structure with a relatively closed canopy containing mature trees; subcanopy layer with semi-mature trees and shrubs; and well-developed herb layer	Described in Perrin et al (2008)
Woodland structure: community diversity and extent	Hectares	Maintain diversity and extent of community types	Based on data from Perrin et al. (2008)
Woodland structure: natural regeneration	Seedling: sapling: pole ratio	Seedlings, saplings and pole age-classes occur in adequate proportions to ensure survival of woodland canopy	Oak (<i>Quercus petraea</i>) generally regenerates poorly. In suitable sites, ash (<i>Fraxinus excelsior</i>) can regenerate in large numbers although few seedlings reach pole size
Woodland structure: dead wood	m ³ per hectare; number per hectare	At least 30m ³ /ha of fallen timber greater than 10cm diameter; 30 snags/ha; both categories should include stems greater than 40cm diameter	Dead wood is a valuable resource and an integral part of a healthy, functioning woodland ecosystem
Woodland structure: veteran trees	Number per hectare	No decline	Mature and veteran trees are important habitats for bryophytes, lichens, saproxylic organisms and some bird species. Their retention is important to ensure continuity of habitats/niches and propagule sources
Woodland structure: indicators of local distinctiveness	Occurrence	No decline	Includes ancient or long-established woodlands (see Perin and Daly, 2010), archaeological and geological features as well as red-data and other rare or localised species. Shannawoneen (site 1600) and Ballynahinch (site 1602) woods are listed as possible ancient woodlands by Perrin and Daly (2010)
Vegetation composition: native tree cover	Percentage	No decline. Native tree cover not less than 95%	Species reported in Perrin et al. (2008)
Vegetation composition: typical species	Occurrence	A variety of typical native species present, depending on woodland type, including oak (<i>Quercus petraea</i>) and birch (<i>Betula pubescens</i>)	Species reported in Perrin et al. (2008)

Vegetation composition:
negative indicator species

Occurrence

Negative indicator species, particularly non-native invasive species, absent or under control

The following are the most common non-native invasive species in this woodland type: beech (*Fagus sylvatica*), sycamore (*Acer pseudoplatanus*) and rhododendron (*Rhododendron ponticum*). Rhododendron and beech were noted in woodlands surveyed in this SAC by Perrin et al. (2008)

Conservation Objectives for : Connemara Bog Complex SAC [002034]**1065 Marsh Fritillary *Euphydryas aurinia***

To maintain the favourable conservation condition of Marsh Fritillary in Connemara Bog Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Distribution: occupied 1km squares	Number	No decline, subject to natural processes	The presence of marsh fritillary has not been mapped in this SAC so current distribution is unknown
Proof of breeding: larval webs	Number at a representative number of sub-sites	Proof of breeding, confirmed by detection of webs	There is currently no survey data for sub-sites within the SAC
Potential habitat: area	Hectares	Area of potential habitat stable or increasing, subject to natural processes	Potential habitat for marsh fritillary is defined as areas of vegetation where devil's-bit scabious (<i>Succisa pratensis</i>) is present, with mean height less than 50cm and with less than 10% cover of scrub more than 1m tall. There is no figure available for the total area of suitable habitat in the SAC

Conservation Objectives for : Connemara Bog Complex SAC [002034]

1106 Salmon *Salmo salar*

To restore the favourable conservation condition of Atlantic Salmon in Connemara Bog Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Distribution: extent of anadromy	percentage of river accessible	100% of river channels down to second order accessible from estuary	Artificial barriers block salmon's upstream migration, thereby limiting species to lower stretches and restricting access to spawning areas. There are no barriers to migration on the Cashla system. A new proposed regional water supply scheme below Glenicmurrin Lake will have a fish pass incorporated. On the Owenboliska River, there is a fish pass in place at the water regulating sluice below Boliska Lake. There are no barriers to migration on the Owenmore system
Adult spawning fish	Number	Conservation Limit (CL) for each system consistently exceeded	A conservation limit is defined by the North Atlantic Salmon Conservation Organisation (NASCO) as "the spawning stock level that produces long-term average maximum sustainable yield as derived from the adult to adult stock and recruitment relationship". The target is based on the Standing Scientific Committee of the National Salmon Commission's annual model output of CL attainment levels. See SSC (2015). Stock estimates are either derived from direct counts of adults (rod catch, fish counter) or indirectly by fry abundance counts. The Owenmore and Cashla rivers are currently exceeding CL while the Owenboliska is below CL
Salmon fry abundance	Number of fry/5 minutes electrofishing	Maintain or exceed 0+ fry mean catchment-wide abundance threshold value. Currently set at 17 salmon fry/5 minutes sampling	Target is threshold value for rivers currently exceeding their conservation limit (CL)
Out-migrating smolt abundance	Number	No significant decline	Smolt abundance can be negatively affected by a number of impacts such as estuarine pollution, predation and sea lice (<i>Lepeophtheirus salmonis</i>). Marine salmon farming takes place in the estuary of the Owenmore River
Number and distribution of redds	Number and occurrence	No decline in number and distribution of spawning redds due to anthropogenic causes	Salmon spawn in clean gravels. The habitat for salmon is good in the Owenmore and Cashla catchments and improvements in spawning areas and input of gravel has been ongoing in both catchments. Funding has been approved for habitat improvement works in the upper reaches of the Owenboliska system
Water quality	EPA Q value	At least Q4 at all sites sampled by EPA	Q values based on triennial water quality surveys carried out by the Environmental Protection Agency (EPA)

Conservation Objectives for : Connemara Bog Complex SAC [002034]

1355 Otter *Lutra lutra*

To maintain the favourable conservation condition of Otter in Connemara Bog Complex SAC, which is defined by the following list of attributes and targets:



Attribute	Measure	Target	Notes
Distribution	Percentage positive survey sites	No significant decline	Measure based on standard otter survey technique. FCS target, based on 1980/81 survey findings, is 88% in SACs. Current range estimated at 93.6% (Reid et al., 2013)
Extent of terrestrial habitat	Hectares	No significant decline. Area mapped and calculated as 2194.8ha	No field survey. Areas mapped to include 10m terrestrial buffer along shoreline (above HWM and along river banks) identified as critical for otters (NPWS, 2007)
Extent of marine habitat	Hectares	No significant decline. Area mapped and calculated as 139.0ha	No field survey. Area mapped based on evidence that otters tend to forage within 80m of the shoreline (HWM) (NPWS, 2007; Kruuk, 2006)
Extent of freshwater (river) habitat	Kilometres	No significant decline. Length mapped and calculated as 564.0km	River length calculated on the basis that otters will utilise freshwater habitats from estuary to headwaters (Chapman and Chapman, 1982)
Extent of freshwater (lake/lagoon) habitat	Hectares	No significant decline. Area mapped and calculated as 3908.6ha	No field survey. Area mapped based on evidence that otters tend to forage within 80m of the shoreline (NPWS, 2007)
Couching sites and holts	Number	No significant decline	Otters need lying up areas throughout their territory where they are secure from disturbance (Kruuk, 2006; Kruuk and Moorhouse, 1991)
Fish biomass available	Kilograms	No significant decline	Broad diet that varies locally and seasonally, but dominated by fish, in particular salmonids, eels and sticklebacks in freshwater (Bailey and Rochford, 2006; Reid et al., 2013) and wrasse and rockling in coastal waters (Kingston et al., 1999)

Conservation Objectives for : Connemara Bog Complex SAC [002034]


1833 Slender Naiad *Najas flexilis*

To maintain the favourable conservation condition of Slender Naiad in Connemara Bog Complex SAC, which is defined by the following list of attributes and targets:








Attribute	Measure	Target	Notes
Population extent	Hectares; distribution	No change to the spatial extent of <i>Najas flexilis</i> within each lake, subject to natural processes. See map 9 for known locations	See <i>Najas flexilis</i> supporting document for further details
Population depth	Metres	No change to the depth range of <i>Najas flexilis</i> within each lake, subject to natural processes	See <i>Najas flexilis</i> supporting document for further details
Population viability	Plant traits	No decline in plant fitness, subject to natural processes	See <i>Najas flexilis</i> supporting document for further details
Population abundance	Square metres	No change to the cover abundance of <i>Najas flexilis</i> , subject to natural processes	See <i>Najas flexilis</i> supporting document for further details
Species distribution	Occurrence	No decline, subject to natural processes	See <i>Najas flexilis</i> supporting document for further details
Habitat extent	Hectares; distribution	No decline, subject to natural processes	See <i>Najas flexilis</i> supporting document for further details
Hydrological regime: water level fluctuations	Metres	Maintain appropriate natural hydrological regime necessary to support the habitat for the species	See <i>Najas flexilis</i> supporting document for further details
Lake substratum quality	Various	Maintain appropriate substratum type, extent and chemistry to support the populations of the species	See <i>Najas flexilis</i> supporting document for further details
Water quality	Various	Maintain appropriate water quality to support the populations of the species	See <i>Najas flexilis</i> supporting document for further details
Acidification status	pH units, mg/l	Maintain appropriate water and sediment pH, alkalinity and cation concentrations to support the populations of <i>Najas flexilis</i> , subject to natural processes	See <i>Najas flexilis</i> supporting document for further details
Water colour	mg/L PtCo	Maintain appropriate water colour to support the populations of <i>Najas flexilis</i>	See <i>Najas flexilis</i> supporting document for further details
Associated species	Species composition; abundance	Maintain appropriate associated species and vegetation communities to support the populations of <i>Najas flexilis</i>	See <i>Najas flexilis</i> supporting document for further details
Fringing habitat: area	Hectares	Maintain the area and condition of fringing habitats necessary to support the populations of <i>Najas flexilis</i>	See <i>Najas flexilis</i> supporting document for further details



Legend
 Connemara Bog Complex SAC 002034




Legend

-  Connemara Bog Complex SAC 002034
-  Lough Corrib SAC 000297
-  Maumturk Mountains SAC 002008
-  The Twelve Bens/Garraun Complex SAC 002031
-  Kilkieran Bay and Islands SAC 002111
-  Connemara Bog Complex SPA 004181
-  OSi Discovery Series County Boundary




Legend

- Connemara Bog Complex SAC 002034
- 1150 *Coastal lagoons
- OSi Discovery Series County Boundary



- Legend**
- Connemara Bog Complex SAC 002034
 - 1170 Reefs
 - OSi Discovery Series County Boundary



**MAP 4:
CONNEMARA BOG COMPLEX SAC
CONSERVATION OBJECTIVES
REEFS**

Map to be read in conjunction with the NPWS Conservation Objectives Document.


**SITE CODE:
SAC 002034; version 3.01. CO. GALWAY**

0 2.5 5 7.5 10 km

The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
Ordnance Survey of Ireland Licence No EN 0059214. © Ordnance Survey of Ireland Government of Ireland.

Níl sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbheithnithe a déanamh ar theorainneacha na gceantar comharthaite. Suirbhéarachta Ordonáis na hÉireann Ceadúnas Uimh EN 0059214. © Suirbhéarachta Ordonáis na hÉireann Rialtas na hÉireann.






Legend

- Connemara Bog Complex SAC 002034
- OSi Discovery Series County Boundary

Marine Community Types

- Anoxic mud community
- Intertidal reef community complex
- Ruppia maritima*-dominated community
- Serpula vermicularis*-dominated subtidal reef community complex





Legend


- Connemara Bog Complex SAC 002034
- OSi Discovery Series County Boundary

Lake Habitats

- 3110 Oligotrophic waters containing very few minerals of sandy plains (*Littorelletalia uniflorarae*)
- 3110 / Potential 3130, Oligotrophic waters containing very few minerals of sandy plains (*Littorelletalia uniflorarae*) / Potential Oligotrophic waters containing very few minerals of sandy plains (*Littorelletalia uniflorarae*)
- 3130 Oligotrophic to mesotrophic standing waters with vegetation of the *Littorelletalia uniflorarae* and/or *Isoetes-Nanojuncetia*
- 3160 Natural dystrophic lakes and ponds
- Potential 3110 Oligotrophic waters containing very few minerals of sandy plains (*Littorelletalia uniflorarae*)
- Potential 3110 / Potential 3130, Potential Oligotrophic waters containing very few minerals of sandy plains (*Littorelletalia uniflorarae*) / Potential Oligotrophic to mesotrophic standing waters with vegetation of the *Littorelletalia uniflorarae* and/or *Isoetes-Nanojuncetia*
- Potential 3110 / Potential 3160, Potential Oligotrophic waters containing very few minerals of sandy plains (*Littorelletalia uniflorarae*) / Potential Natural dystrophic lakes and ponds
- Potential 3160 Natural dystrophic lakes and ponds
- Potential 3160 / Potential 3110, Potential Natural dystrophic lakes and ponds / Potential Oligotrophic waters containing very few minerals of sandy plains (*Littorelletalia uniflorarae*)







Legend

- Connemara Bog Complex SAC 002034
- 1833 Slender Naiad - *Najas flexilis*
- OSi Discovery Series County Boundary

National Parks and Wildlife Service

Conservation Objectives Series

Ballinafad SAC 002081



An Roinn Cultúir,
Oidhreacht agus Gaeltachta
Department of Culture,
Heritage and the Gaeltacht

**National Parks and Wildlife Service,
Department of Culture, Heritage and the Gaeltacht,
90 King Street North, Dublin 7, D07 N7CV, Ireland.**

**Web: www.npws.ie
E-mail: nature.conservation@chg.gov.ie**

Citation:

**NPWS (2018) Conservation Objectives: Ballinacorney SAC 002081. Version 1.
National Parks and Wildlife Service, Department of Culture, Heritage and the
Gaeltacht.**

**Series Editor: Rebecca Jeffrey
ISSN 2009-4086**

Introduction

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

** indicates a priority habitat under the Habitats Directive*

002081 Ballinafad SAC

1303 Lesser Horseshoe Bat *Rhinolophus hipposideros*

Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

NPWS Documents

Year :	2006
Title :	Bat mitigation guidelines for Ireland
Author :	Kelleher, C.; Marnell, F.
Series :	Irish Wildlife Manual No. 25
<hr/>	
Year :	2018
Title :	Conservation objectives supporting document – lesser horseshoe bat (<i>Rhinolophus hipposideros</i>)
Author :	NPWS
Series :	Conservation objectives supporting document
<hr/>	

Other References

Year :	2008
Title :	The lesser horseshoe bat conservation handbook
Author :	Schofield, H.W.
Series :	The Vincent Wildlife Trust
<hr/>	

Spatial data sources

Year :	2018
Title :	NPWS lesser horseshoe bat database
GIS Operations :	Roost identified, clipped to SAC boundary. Expert opinion used as necessary to resolve any issues arising
Used For :	1303 (map 2)
<hr/>	
Year :	2007
Title :	Forest Inventory and Planning System (FIPS)
GIS Operations :	Dataset clipped to 2.5km buffer centred on roost location
Used For :	1303 (map 2)
<hr/>	

1303 Lesser Horseshoe Bat *Rhinolophus hipposideros*


To maintain the favourable conservation condition of Lesser Horseshoe Bat in Ballinafad SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Population per roost	Number	Minimum number of 62 bats for the summer roost (roost id. 683 in NPWS database). See map 2	A figure of 100 bats for summer roosts and 50 bats for winter roosts was set as a minimum qualifying standard (MQS) when SACs were being selected for lesser horseshoe bat (<i>Rhinolophus hipposideros</i>). Where possible, NPWS conduct annual counts at each qualifying roost. For most summer roosts, qualified means from the 2012-2016 data have been calculated whereby the year with the highest maximum count and the year with the lowest maximum count were removed and the mean of the remaining years was calculated. This mean is usually set as the target figure for the roost. However, very limited counts have been taken at the summer roost (roost id. 683) in Ballinafad SAC. Also, this roost is situated at the northern extreme of the species' distribution in Ireland, and indeed Europe; thus, in this case, a figure below the MQS is justified and the most recent available count from 2014 is set as the target, i.e. 62 bats. See NPWS (2018) for further information on all attributes and targets
Summer roosts	Condition	No decline	Ballinafad SAC has been selected for lesser horseshoe bat because of the presence of one internationally important summer roost (roost id. 683 in NPWS database). Damage or disturbance to the roost or to the habitat immediately surrounding it will lead to a decline in its condition (Kelleher and Marnell, 2006). See the conservation objectives supporting document for lesser horseshoe bat (NPWS, 2018) for further information on all attributes and targets
Extent of potential foraging habitat	Hectares	No significant decline within 2.5km of qualifying roost	Lesser horseshoe bats normally forage in woodlands/scrub within 2.5km of their roosts (Schofield, 2008). See map 2 which shows a 2.5km zone around the above roost and identifies potential foraging grounds
Linear features	Kilometres	No significant loss within 2.5km of qualifying roost. See map 2	This species follows commuting routes from its roost to its foraging grounds. Lesser horseshoe bats will not cross open ground. Consequently, linear features such as hedgerows, treelines and stone walls provide vital connectivity for this species within 2.5km around each roost (Schofield, 2008)
Light pollution	Lux	No significant increase in artificial light intensity adjacent to named roost or along commuting routes within 2.5km of the roost. See map 2	Lesser horseshoe bats are very sensitive to light pollution and will avoid brightly lit areas. Inappropriate lighting around roosts may cause abandonment; lighting along commuting routes may cause preferred foraging areas to be abandoned, thus increasing energetic costs for bats (Schofield, 2008)



Legend

 Ballinafad SAC 002081



Legend

- OS Discovery Series County Boundary
- Ballinacrad SAC 002081
- 1303 Lesser Horseshoe Bat *Rhinolophus hipposideros***
- Roost Location
- Roost ID 683 Foraging Range
- Potential Foraging Grounds

National Parks and Wildlife Service

Conservation Objectives Series

Corliskea/Trien/Cloonfelliv Bog SAC 002110



An Roinn
Ealaíon, Oidhreachta agus Gaeltachta

Department of
Arts, Heritage and the Gaeltacht



**National Parks and Wildlife Service,
Department of Arts, Heritage and the Gaeltacht,**

7 Ely Place, Dublin 2, Ireland.

Web: www.npws.ie

E-mail: nature.conservation@ahg.gov.ie

Citation:

**NPWS (201) Conservation Objectives: Corliskea/Trien/Cloonfelliv Bog SAC
002110. Version 1. National Parks and Wildlife Service, Department of Arts,
Heritage and the Gaeltacht.**

Series Editor: Rebecca Jeffrey

ISSN 2009-4086

Introduction

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

** indicates a priority habitat under the Habitats Directive*

002110	Corliskea/Trien/Cloonfelliv Bog SAC
7110	Active raised bogsE
7120	Degraded raised bogs still capable of natural regeneration
7150	Depressions on peat substrates of the Rhynchosporion
91D0	Bog woodland

Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

NPWS Documents

Year :	2014
Title :	Raised Bog Monitoring and Assessment Survey 2013
Author :	Fernandez, F.; Connolly K.; Crowley W.; Denyer J.; Duff K.; Smith G.
Series :	Irish Wildlife Manual No. 81
<hr/>	
Year :	2014
Title :	National raised bog SAC management plan
Author :	Department of Arts, Heritage and the Gaeltacht
Series :	Draft for consultation. 15 January 2014
<hr/>	
Year :	2014
Title :	Cloonfelliv Bog (SAC 002110), Co. Galway/Roscommon, Site Report
Author :	Fernandez, F.; Connolly, K.; Crowley, W.; Denyer J.; Duff K.; Smith G.
Series :	Raised bog monitoring and assessment survey 2013
<hr/>	
Year :	2014
Title :	Corliskea Bog (SAC 002110), Co. Galway/Roscommon, Site Report
Author :	Fernandez, F.; Connolly, K.; Crowley, W.; Denyer J.; Duff K.; Smith G.
Series :	Raised bog monitoring and assessment survey 2013
<hr/>	
Year :	2014
Title :	Trien Bog (SAC 002110), Co. Roscommon, Site Report
Author :	Fernandez, F.; Connolly, K.; Crowley, W.; Denyer J.; Duff K.; Smith G.
Series :	Raised bog monitoring and assessment survey 2013
<hr/>	
Year :	2016
Title :	Corliskea/Trien/Cloonfelliv Bog SAC (site code: 2110) Conservation objectives supporting document- raised bog habitats V1
Author :	NPWS
Series :	Conservation objectives supporting document

Other References

Year :	2011
Title :	Review and revision of empirical critical loads and dose-response relationships. Proceedings of an expert workshop, Noordwijkerhout, 23-25 June 2010
Author :	Bobbink, R.; Hettelingh, J.P.
Series :	RIVM report 680359002, Coordination Centre for Effects, National Institute for Public Health and the Environment (RIVM)
<hr/>	
Year :	2014
Title :	Nitrogen deposition and exceedance of critical loads for nutrient nitrogen in Irish grasslands
Author :	Henry, J.; Aherne, J.
Series :	Science of the Total Environment 470–471: 216–223

Spatial data sources

Year :	2014
Title :	Scientific Basis for Raised Bog Conservation in Ireland
GIS Operations :	RBSB13_SACs_ARB_DRB dataset, RBSB13_SACs_2012_HB dataset, RBSB13_SACs_DrainagePatterns_5k dataset and RBSB13_SAC_LIDAR_DTMs dataset clipped to SAC boundary. Expert opinion used as necessary to resolve any issues arising
Used For :	potential 7110; digital elevation model; drainage patterns (maps 2 and 4)
<hr/>	
Year :	2013
Title :	Raised Bog Monitoring and Assessment Survey 2013
GIS Operations :	RBMA13_ecotope_map dataset clipped to SAC boundary. Appropriate ecotopes selected and exported to new dataset. Expert opinion used as necessary to resolve any issues arising
Used For :	7110 ecotopes (map 3)
<hr/>	

Conservation Objectives for : Corliskea/Trien/Cloonfelliv Bog SAC [002110]

7110 Active raised bogs

To restore the favourable conservation condition of Active raised bogs in Corliskea/Trien/Cloonfelliv Bog SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Restore area of active raised bog to 107.5ha, subject to natural processes	Active Raised Bog (ARB) habitat was mapped at 69.2ha by Fernandez et al. (2014). Area of Degraded Raised Bog (DRB) on the High Bog (HB) has been modelled as 54.8ha. See map 2. However, it is estimated that only 31.8ha is potentially restorable to ARB by drain blocking. The total potential ARB on the HB is therefore estimated to be 101.0ha. Eco-hydrological assessments of the cutover estimates that an additional 6.5ha of bog forming habitats could be restored. The long term target for ARB is therefore 107.5ha. See raised bog supporting document for further details on this and following attributes
Habitat distribution	Occurrence	Restore the distribution and variability of active raised bog across the SAC. See map 3 for distribution in 2013	ARB occurs at numerous locations on the high bog throughout Corliskea/Trien/Cloonfelliv Bog SAC. DRB occurs on the high bog surrounding ARB areas, which will require restoration measures. There is also potential for ARB restoration on cutover areas of the bog (see area target above)
High bog area	Hectares	No decline in extent of high bog necessary to support the development and maintenance of active raised bog. See map 2	The area of high bog within Corliskea/Trien/Cloonfelliv Bog SAC in 2012 (latest figure available) was 452.4ha (DAHG, 2014)
Hydrological regime: water levels	Centimetres	Restore appropriate water levels throughout the site	For ARB, mean water level needs to be near or above the surface of the bog lawns for most of the year. Seasonal fluctuations should not exceed 20cm, and should only be 10cm below the surface, except for very short periods of time. Open water is often characteristic of soak systems
Hydrological regime: flow patterns	Flow direction; slope	Restore, where possible, appropriate high bog topography, flow directions and slopes. See map 4 for current situation	ARB depends on mean water levels being near or above the surface of bog lawns for most of the year. Long and gentle slopes are the most favourable to achieve these conditions. Changes to flow directions due to subsidence of bogs can radically change water regimes and cause drying out of high quality ARB areas
Transitional areas between high bog and adjacent mineral soils (including cutover areas)	Hectares; distribution	Restore adequate transitional areas to support/protect active raised bog and the services it provides	ARB is threatened due to effects of past drainage and peat-cutting around the margins of Corliskea/Trien/Cloonfelliv Bog SAC. Natural marginal habitats no longer exist around these bogs. Eco-hydrological assessments have evaluated the potential for ARB restoration on cutover areas (see note for habitat area attribute above)
Vegetation quality: central ecotope, active flush, soaks, bog woodland	Hectares	Restore 53.8ha of central ecotope/active flush/soaks/bog woodland as appropriate	At least 50% of ARB habitat should be high quality (i.e. central ecotope, active flush, soaks, bog woodland). Target area of active raised bog for the site has been set at 107.5ha (see area target above)
Vegetation quality: microtopographical features	Hectares	Restore adequate cover of high quality microtopographical features	High quality microtopography (hummocks, hollows and pools) is well developed on Corliskea/Trien/Cloonfelliv Bog
Vegetation quality: bog moss (<i>Sphagnum</i>) species	Percentage cover	Restore adequate cover of bog moss (<i>Sphagnum</i>) species to ensure peat-forming capacity	<i>Sphagnum</i> cover varies naturally across Ireland with relatively high cover in the east to lower cover in the west. Hummock forming species such as <i>Sphagnum austini</i> are particularly good peat formers. <i>Sphagnum</i> cover and distribution also varies naturally across a site

Typical ARB species: flora	Occurrence	Restore, where appropriate, typical active raised bog flora	Typical flora species include widespread species, as well as those with more restricted distributions but typical of the habitat's subtypes or geographical range
Typical ARB species: fauna	Occurrence	Restore, where appropriate, typical active raised bog fauna	Typical fauna species include widespread species, as well as those with more restricted distributions but typical of the habitat's subtypes or geographical range
Elements of local distinctiveness	Occurrence	Maintain features of local distinctiveness, subject to natural processes	Corliskea/Trien/Cloonfelliv Bog SAC is noted for extensive ARB areas with central, sub-central ecotopes, active flush and bog woodland, together with a lake and swallow hole system
Negative physical indicators	Percentage cover	Negative physical features absent or insignificant	Negative physical indicators include: bare peat, algae dominated pools and hollows, marginal cracks, tear patterns, subsidence features such as dry mineral mounds /ridges emerging or expanding and evidence of burning
Vegetation composition: native negative indicator species	Percentage cover	Native negative indicator species at insignificant levels	Disturbance indicators include species indicative of conditions drying out such as abundant bog asphodel (<i>Narthecium ossifragum</i>), deergrass (<i>Trichophorum germanicum</i>) and harestail cotton-grass (<i>Eriophorum vaginatum</i>) forming tussocks; abundant magellanic bog-moss (<i>Sphagnum magellanicum</i>) in pools previously dominated by <i>Sphagnum</i> species typical of very wet conditions (e.g. feathery bog-moss (<i>S. cuspidatum</i>)); and indicators of frequent burning events such as abundant <i>Cladonia floerkeana</i> and high cover of carnation sedge (<i>Carex panicea</i>) (particularly in true midlands raised bogs)
Vegetation composition: non-native invasive species	Percentage cover	Non-native invasive species at insignificant levels and not more than 1% cover	Most common non-native invasive species include lodgepole pine (<i>Pinus contorta</i>), rhododendron (<i>Rhododendron ponticum</i>), and pitcherplant (<i>Sarracenia purpurea</i>)
Air quality: nitrogen deposition	kg N/ha/year	Air quality surrounding bog close to natural reference conditions. The total N deposition should not exceed 5kg N/ha/yr	Change in air quality can result from fertiliser drift; adjacent quarry activities; or other atmospheric inputs. The critical load range for ombrotrophic bogs has been set as between 5 and 10kg N/ha/yr (Bobbink and Hettelingh, 2011). The latest N deposition figures for the area around Corliskea/Trien/Cloonfelliv Bog SAC suggest that the current level is approximately 10.6kg N/ha/yr (Henry and Aherne, 2014)
Water quality	Hydrochemical measures	Water quality on the high bog and in transitional areas close to natural reference conditions	Water chemistry within raised bogs is influenced by atmospheric inputs (rainwater). However, within soak systems, water chemistry is influenced by other inputs such as focused flow or interaction with underlying substrates. Water chemistry in areas surrounding the high bog varies due to influences of different water types (bog water, regional groundwater and run-off from surrounding mineral lands)

Conservation Objectives for : Corliskea/Trien/Cloonfelloiv Bog SAC [002110]

7120 Degraded raised bogs still capable of natural regeneration

The long-term aim for Degraded raised bogs still capable of natural regeneration is that its peat-forming capability is re-established; therefore, the conservation objective for this habitat is inherently linked to that of Active raised bogs (7110) and a separate conservation objective has not been set in Corliskea/Trien/Cloonfelloiv Bog SAC

Attribute	Measure	Target	Notes
------------------	----------------	---------------	--------------

Conservation Objectives for : Corliskea/Trien/Cloonfelloiv Bog SAC [002110]

7150 Depressions on peat substrates of the Rhynchosporion

Depressions on peat substrates of the Rhynchosporion is an integral part of good quality Active raised bogs (7110) and thus a separate conservation objective has not been set for the habitat in Corliskea/Trien/Cloonfelloiv Bog SAC


Attribute	Measure	Target	Notes
-----------	---------	--------	-------

Conservation Objectives for : Corliskea/Trien/Cloonfelliv Bog SAC [002110]


91D0 Bog woodland


To maintain the favourable conservation condition of Bog woodland in Corliskea/Trien/Cloonfelliv Bog SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes. At least 0.29ha. See map 4	Bog woodland occurs on Corliskea Bog (0.25ha) and Trien Bog (0.04ha) and is regarded as a component of the Active Raised Bog (ARB) habitat (7110). Thus, the conservation objective and supporting document for ARB (7110) are also relevant to this habitat and common attributes have not been repeated here. The latest survey for bog woodland in Corliskea/Trien/Cloonfelliv Bog SAC is reported in Fernandez et al. (2014)
Habitat distribution	Occurrence	No decline, subject to natural processes. See map 4	Bog woodland occurs in active flushes on both Corliskea and Trien Bogs
Vegetation composition: positive indicator species	Number in a representative number of monitoring stops	Birch (<i>Betula pubescens</i>), bog moss (<i>Sphagnum</i> species) and at least five other species present	Bog woodland is typically species-poor but with a characteristic and distinctive flora. Positive indicator species are listed in bog woodland monitoring survey (Cross and Lynn, 2013)
Vegetation composition: negative indicator species	Percentage cover at a representative number of monitoring stops	Both native and non-native invasive species absent or under control. Total cover should be less than 10%	Negative indicator species include bracken (<i>Pteridium aquilinum</i>) and bramble (<i>Rubus fruticosus</i>), which can become invasive if the site begins drying out
Woodland structure: cover and height of birch	Percentage cover and metres at a representative number of monitoring stops	A minimum 30% cover of birch (<i>Betula pubescens</i>) with a median canopy height of 4m	Attribute and target based on Cross and Lynn (2013)
Woodland structure: dwarf shrub cover	Percentage cover at a representative number of monitoring stops	Dwarf shrub cover not more than 50%	Attribute and target based on Cross and Lynn (2013)
Woodland structure: ling cover	Percentage cover at a representative number of monitoring stops	Ling (<i>Calluna vulgaris</i>) cover not more than 40%	Attribute and target based on Cross and Lynn (2013)
Woodland structure: bryophyte cover	Percentage cover at a representative number of monitoring stops	Bryophyte cover at least 50%, with bog moss (<i>Sphagnum</i> spp.) cover at least 25%	Attribute and target based on Cross and Lynn (2013)
Woodland structure: tree size classes	Occurrence	Each size class present	Size classes are defined in Cross and Lynn (2013). The presence of all size classes suggests that a woodland has good structural variety with trees of varying ages
Woodland structure: senescent and dead wood	Occurrence	Senescent or dead wood present	Mature and veteran trees and dead wood are important for bryophytes, lichens, saproxylic organisms and some bird species. Their retention within a woodland is important to ensure continuity of habitats/niches and propagule sources over time. However, as birch (<i>Betula pubescens</i>) trees seldom exceed 30cm in diameter in this habitat and dead wood rots quickly and is engulfed by bog mosses (<i>Sphagnum</i> spp.), volume of dead wood may not be as high in bog woodland as in other woodland types



Legend

 Corliskea/Trien/Cloonfelliv Bog SAC 002110

 *An Roinn Ealaíon, Oidhreacht agus Gaeltachta*
Department of Arts, Heritage and the Gaeltacht

MAP 1:
CORLISKEA/TRIEN/CLOONFELLIV BOG SAC
CONSERVATION OBJECTIVES
SAC DESIGNATION


Map to be read in conjunction with the NPWS Conservation Objectives Document.

SITE CODE:
SAC 002110; version 3.02. Co. Galway / Roscommon


0 200 400 600 800 1,000 m

The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
Ordnance Survey of Ireland Licence No EN 0059216. © Ordnance Survey of Ireland Government of Ireland.

Níl sna teorainneacha ar na léarscáileanna ach nod garshuimhach ginearálta. Féadfar athbheithníthe a déanamh ar theorainneacha na gceantar comharthaite. Suirbhéarachta Ordonáis na hÉireann Ceadúnas Uimh EN 0059216. © Suirbhéarachta Ordonáis na hÉireann Rialtas na hÉireann.




Map Version 1
Date: Feb 2016



Legend

- Corliskea/Trien/Cloonfelliv Bog SAC 002110
- High Bog Boundary
- Potential 7110 *Active Raised Bogs
- OSi Discovery Series County Boundary




Legend

- Corliskea/Trien/Cloonfelliv Bog SAC 002110
- OSi Discovery Series County Boundary
- High Bog Boundary


Active Raised Bogs Ecotopes

- Bog Woodland
- Central ecotope
- Soaks / active flush
- Sub-central ecotope



Legend

- Corliskea/Trien/Cloonfellov Bog SAC 002110
- OSi Discovery Series County Boundary
- High Bog Boundary
- Drainage Patterns
- Contours





Conservation objectives for Tully Lough SAC [002130]

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Objective: To maintain or restore the favourable conservation condition of the Annex I habitat(s) and/or the Annex II species for which the SAC has been selected:

Code Description

3130 Oligotrophic to mesotrophic standing waters with vegetation of the Littorelletea uniflorae and/or Isoeto-Nanojuncetea

* denotes a priority habitat

Code Common Name Scientific Name

1833 Slender Naiad *Najas flexilis*



Citation: NPWS (2020) Conservation objectives for Tully Lough SAC [002130]. Generic Version 7.0.
Department of Culture, Heritage and the Gaeltacht.

Conservation objectives for Newport River SAC [002144]

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Objective: To maintain or restore the favourable conservation condition of the Annex I habitat(s) and/or the Annex II species for which the SAC has been selected:

ADD HABITATS

Code	Common Name	Scientific Name
1029	Freshwater Pearl Mussel	<i>Margaritifera margaritifera</i>
1106	Salmon	<i>Salmo salar</i>

Citation: NPWS (2018) Conservation objectives for Newport River SAC [002144]. Generic Version 6.0.
Department of Culture, Heritage and the Gaeltacht.



Conservation objectives for Lough Dahybaun SAC [002177]

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Objective: To maintain or restore the favourable conservation condition of the Annex I habitat(s) and/or the Annex II species for which the SAC has been selected:

ADD HABITATS

Code	Common Name	Scientific Name
1833	Slender Naiad	<i>Najas flexilis</i>

Citation: NPWS (2020) Conservation objectives for Lough Dahybaun SAC [002177]. Generic Version 7.0. Department of Culture, Heritage and the Gaeltacht.

For more information please go to: www.npws.ie/protected-sites/conservation-management-planning



National Parks and Wildlife Service

Conservation Objectives Series

Towerhill House SAC 002179



An Roinn Cultúir,
Oidhreacht agus Gaeltachta
Department of Culture,
Heritage and the Gaeltacht

**National Parks and Wildlife Service,
Department of Culture, Heritage and the Gaeltacht,
90 King Street North, Dublin 7, D07 N7CV, Ireland.
Web: www.npws.ie
E-mail: nature.conservation@chg.gov.ie**

Citation:

**NPWS (2018) Conservation Objectives: Towerhill House SAC 002179. Version 1.
National Parks and Wildlife Service, Department of Culture, Heritage and the
Gaeltacht.**

**Series Editor: Rebecca Jeffrey
ISSN 2009-4086**

Introduction

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

** indicates a priority habitat under the Habitats Directive*

002179 Towerhill House SAC

1303 Lesser Horseshoe Bat *Rhinolophus hipposideros*

Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

NPWS Documents

Year :	2006
Title :	Bat mitigation guidelines for Ireland
Author :	Kelleher, C.; Marnell, F.
Series :	Irish Wildlife Manual No. 25
<hr/>	
Year :	2018
Title :	Conservation objectives supporting document – lesser horseshoe bat (<i>Rhinolophus hipposideros</i>)
Author :	NPWS
Series :	Conservation objectives supporting document

Other References

Year :	2007
Title :	Protecting and managing underground sites for bats
Author :	Mitchell-Jones, A.J.; Bihari, Z.; Masing, M.; Rodrigues, L.
Series :	EUROBATS Publication Series No. 2
<hr/>	
Year :	2008
Title :	The lesser horseshoe bat conservation handbook
Author :	Schofield, H.W.
Series :	The Vincent Wildlife Trust
<hr/>	
Year :	2009
Title :	Importance of night roosts for bat conservation: roosting behaviour of the lesser horseshoe bat <i>Rhinolophus hipposideros</i>
Author :	Knight, T.; Jones, G.
Series :	Endangered Species Research, 8: 79-86

Spatial data sources

Year :	2018
Title :	NPWS lesser horseshoe bat database
GIS Operations :	Roosts identified, clipped to SAC boundary. Expert opinion used as necessary to resolve any issues arising
Used For :	1303 (map 2)
<hr/>	
Year :	2007
Title :	Forest Inventory and Planning System (FIPS)
GIS Operations :	Dataset clipped to 2.5km buffer centred on roost locations
Used For :	1303 (map 2)
<hr/>	


Conservation Objectives for : Towerhill House SAC [002179]

1303 Lesser Horseshoe Bat *Rhinolophus hipposideros*

To maintain the favourable conservation condition of Lesser Horseshoe Bat in Towerhill House SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Population per roost	Number	Minimum number of 373 bats for the summer roost (roost id. 668 in NPWS database); minimum number of 50 bats for the winter roost (roost id. 682). See map 2	A figure of 100 bats for summer roosts and 50 bats for winter roosts was set as a minimum qualifying standard (MQS) when SACs were being selected for lesser horseshoe bat (<i>Rhinolophus hipposideros</i>). NPWS conduct annual counts at each qualifying roost. The qualified mean from the 2012-2016 summer data has been calculated whereby the year with the highest maximum count and the year with the lowest maximum count were removed and the mean of the remaining years was calculated. This mean is set as the target figure for the summer roost (roost id. 668 in NPWS database) in Towerhill House SAC. There is no recent data available for the winter roost in the SAC (roost id. 682); therefore, the MQS is set as the target figure for the winter roost in the SAC. See the conservation objectives supporting document for lesser horseshoe bat (NPWS, 2018) for further information on all attributes and targets
Winter roosts	Condition	No decline	Towerhill House SAC has been selected for lesser horseshoe bat because of the presence of one internationally important winter roost (roost id. 682 in NPWS database). Damage or disturbance to the roost or to the habitat immediately surrounding it will lead to a decline in its condition (Mitchell-Jones et al., 2007)
Summer roosts	Condition	No decline	Towerhill House SAC has been selected for lesser horseshoe bat because of the presence of one internationally important summer roost (roost id. 668 in NPWS database). Damage or disturbance to the roost or to the habitat immediately surrounding it will lead to a decline in its condition (Kelleher and Marnell, 2006)
Auxiliary roosts	Number and condition	No decline	Lesser horseshoe bat populations will use a variety of roosts during the year besides the main summer maternity and winter hibernation roosts. Such additional roosts within the SAC may be important as night roosts, satellite roosts, etc. Night roosts are also considered an integral part of core foraging areas and require protection (Knight and Jones, 2009). In addition, in response to weather conditions for example, bats may use different seasonal roosts from year to year; this is particularly noticeable in winter. A database of all known lesser horseshoe bat roosts is available on the National Biodiversity Data Centre website. NB further unrecorded roosts may also be present within this SAC
Extent of potential foraging habitat	Hectares	No significant decline within 2.5km of qualifying roosts	Lesser horseshoe bats normally forage in woodlands/scrub within 2.5km of their roosts (Schofield, 2008). See map 2 which shows a 2.5km zone around the above roosts and identifies potential foraging grounds
Linear features	Kilometres	No significant loss within 2.5km of qualifying roosts. See map 2	This species follows commuting routes from its roost to its foraging grounds. Lesser horseshoe bats will not cross open ground. Consequently, linear features such as hedgerows, treelines and stone walls provide vital connectivity for this species within 2.5km around each roost (Schofield, 2008)

Light pollution	Lux	No significant increase in artificial light intensity adjacent to named roosts or along commuting routes within 2.5km of those roosts. See map 2	Lesser horseshoe bats are very sensitive to light pollution and will avoid brightly lit areas. Inappropriate lighting around roosts may cause abandonment; lighting along commuting routes may cause preferred foraging areas to be abandoned, thus increasing energetic costs for bats (Schofield, 2008)
-----------------	-----	--	---



Legend

 Towerhill House SAC002179

 An Roinn Cultúir,
Oidhreacht agus Gaeltachta
Department of Culture,
Heritage and the Gaeltacht

**MAP 1:
TOWERHILL HOUSE SAC
CONSERVATION OBJECTIVES
SAC DES GNATI ON**

Map to be read in conjunction with the NPWS Conservation Objectives Document.


**SITE CODE
SAC 002179; version 3.0. CO MAYO**

0 80 160 240 320 Meters

The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
Ordnance Survey of Ireland Licence No EN 0059216. © Ordnance Survey of Ireland and Government of Ireland

Ní snáiteoraí nneacha ar nal éarscáileanna ach nod garshú omhach g near áta. Féad ar at hbréithithe a déana mh ar theorá nneacha na gceantar comhatháthe. Súrbhéaracht a Ordonás na hÉireann Ceadúnas U mh EN 0059216. © Súrbhéaracht a Ordonás na hÉireann Rátas na hÉireann


**Map Version 1
Date: June 2018**



Legend

- Towerhill House SAC 002179
- OS Discovery Series County Boundary
- 1303 Lesser Horseshoe Bat *Rhinolophus hipposideros***
- Roost Location
- Roost ID 682 Foraging Range
- Roost ID 688 Foraging Range
- Potential Foraging Grounds

National Parks and Wildlife Service

Conservation Objectives Series

Clare Island Cliffs SAC 002243



An Roinn Ealaíon, Oidhreachta,
Gnóthaí Réigiúnacha, Tuaithe agus Gaeltachta

Department of Arts, Heritage,
Regional, Rural and Gaeltacht Affairs



**National Parks and Wildlife Service,
Department of Arts, Heritage, Regional, Rural and Gaeltacht Affairs,
7 Ely Place, Dublin 2, Ireland.
Web: www.npws.ie
E-mail: nature.conservation@ahg.gov.ie**

Citation:

**NPWS (2016) Conservation Objectives: Clare Island Cliffs SAC 002243. Version 1.
National Parks and Wildlife Service, Department of Arts, Heritage, Regional,
Rural and Gaeltacht Affairs.**

**Series Editor: Rebecca Jeffrey
ISSN 2009-4086**

Introduction

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

** indicates a priority habitat under the Habitats Directive*

002243	Clare Island Cliffs SAC
1230	Vegetated sea cliffs of the Atlantic and Baltic coasts
8210	Calcareous rocky slopes with chasmophytic vegetation
8220	Siliceous rocky slopes with chasmophytic vegetation

Please note that this SAC overlaps with Clare Island SPA (004136) and is adjacent to West Connaght Coast SAC (002998). See map 2. The conservation objectives for this site should be used in conjunction with those for the overlapping and adjacent sites as appropriate.

Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

NPWS Documents

Year :	2011
Title :	National survey and assessment of the conservation status of Irish sea cliffs
Author :	Barron, S.J.; Delaney, A.; Perrin, P.M.; Martin, J.; O'Neill, F.
Series :	Irish Wildlife Manual No. 53
Year :	2012
Title :	Ireland Red List No. 8: Bryophytes
Author :	Lockhart, N.; Hodgetts, N.; Holyoak, D.
Series :	Ireland Red List series, NPWS
Year :	2013
Title :	The status of EU protected habitats and species in Ireland. Volume 2. Habitats assessments
Author :	NPWS
Series :	Conservation assessments
Year :	2014
Title :	Guidelines for a national survey and conservation assessment of upland vegetation and habitats in Ireland, Version 2.0
Author :	Perrin, P.M.; Barron, S.J.; Roche, J.R.; O'Hanrahan, B.
Series :	Irish Wildlife Manual No. 79
Year :	2016
Title :	Clare Island Cliffs SAC (site code: 2243) Conservation objectives supporting document-coastal habitats V1
Author :	NPWS
Series :	Conservation objectives supporting document
Year :	2016
Title :	Ireland Red List No. 10: Vascular Plants
Author :	Wyse Jackson, M.; FitzPatrick, Ú.; Cole, E.; Jebb, M.; McFerran, D.; Sheehy Skeffington, M.; Wright, M.
Series :	Ireland Red Lists series, NPWS

Other References

Year :	1986
Title :	A resurvey of Clare Island flora
Author :	Doyle, G.J.; Foss, P.J.
Series :	Irish Naturalists' Journal, 22: 85-89
Year :	2005
Title :	National inventory of sea cliffs and coastal heaths
Author :	Browne, A.
Series :	Unpublished Report to NPWS

Spatial data sources

Year : 2011
Title : National survey and assessment of the conservation status of Irish sea cliffs
GIS Operations : Clipped to SAC boundary
Used For : 1230 (map 3)

1230 Vegetated sea cliffs of the Atlantic and Baltic coasts

To maintain the favourable conservation condition of Vegetated sea cliffs of the Atlantic and Baltic coasts in Clare Island Cliffs SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat length	Kilometres	Area stable, subject to natural processes, including erosion. For the sub-site mapped (Capnagower), total length of cliff sections: 14.69km. See map 3	Based on data from the Irish Sea Cliff Survey (ISCS) (Barron et al., 2011). Cliffs are linear features and are therefore measured in kilometres. The sub-site Capnagower (ISCS site ID: 08017) was identified using a combination of aerial photos and the DCENR helicopter viewer. The length of cliff was measured (in sections) to give a total estimated area of 14.69km within the SAC. The length of cliff is likely to be underestimated. See the Clare Island Cliffs SAC conservation objectives supporting document for coastal habitats for further details
Habitat distribution	Occurrence	No decline, subject to natural processes. See map 3	Sea cliffs are known to occur along the coastline from Kinnacorra in the east, along the north coast and around the south-western corner of the island as far east as Tonabrickill. Both hard and soft cliffs have been noted in this SAC, with hard cliffs being by far the most dominant type (Browne, 2005; Barron et al., 2011). See the coastal habitats supporting document for further details
Physical structure: functionality and hydrological regime	Occurrence of artificial barriers	No alteration to natural functioning of geomorphological and hydrological processes, including groundwater quality, due to artificial structures	Based on data from Barron et al. (2011). Maintaining natural geomorphological processes, including natural erosion, is important for the health of vegetated sea cliffs. Hydrological processes maintain flushes, and in some cases tufa formations, that can be associated with sea cliffs. Hydrological features such as gullies, streams and cascades were identified by the ISCS as occurring within Clare Island Cliffs SAC. See the coastal habitats supporting document for further details
Vegetation structure: zonation	Occurrence	Maintain the range of sea cliff habitat zonation including transitional zones, subject to natural processes including erosion and succession	Based on data from Barron et al. (2011). The cliffs in the south-west and eastern sectors are low-lying with a sward of maritime grassland vegetation. On the upper cliffs, there is a concentration of alpine vegetation, which is species-rich and includes a number of rarities, in association with a species-rich bryophyte flora. See the coastal habitats supporting document for further details
Vegetation structure: vegetation height	Centimetres	Maintain structural variation within sward	Based on data from Barron et al. (2011). The cliffs in the south-west and east of the SAC have low-growing vegetation along the cliff top. See the coastal habitats supporting document for further details
Vegetation composition: typical species and sub-communities	Percentage cover at a representative number of monitoring stops	Maintain range of sub-communities with typical species listed in the Irish Sea Cliff Survey (Barron et al., 2011)	The cliffs are well-vegetated, with a maritime sward of grasses and herbs. A plantain (<i>Plantago</i> spp.) dominated sward of low-growing vegetation occurs along the tops of some of the cliffs. On the upper cliffs, there is a concentration of species-rich alpine vegetation, which occurs in association with a species-rich bryophyte flora. See the coastal habitats supporting document for further details
Vegetation composition: negative indicator species	Percentage	Negative indicator species (including non-native species) to represent less than 5% cover	Based on data from Barron et al. (2011). See the coastal habitats supporting document for further details
Vegetation composition: bracken and woody species	Percentage	Cover of bracken (<i>Pteridium aquilinum</i>) on grassland and/or heath less than 10%. Cover of woody species on grassland and/or heath less than 20%	Based on data from Barron et al. (2011). See the coastal habitats supporting document for further details

Conservation Objectives for : Clare Island Cliffs SAC [002243]

8210 Calcareous rocky slopes with chasmophytic vegetation

To maintain the favourable conservation condition of Calcareous rocky slopes with chasmophytic vegetation in Clare Island Cliffs SAC, which is defined by the following list of attributes and targets:


Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Calcareous rocky slopes with chasmophytic vegetation has not been mapped in detail for Clare Island Cliffs SAC and thus the total area of the qualifying habitat is unknown. It occurs as a mosaic with siliceous rocky slopes with chasmophytic vegetation (8220) and is generally found on the sheer cliffs on the seaward face of Croaghmore and to the north and north-west of the island on some of the higher sea cliffs, above c.250m (NPWS internal files)
Habitat distribution	Occurrence	No decline, subject to natural processes	See note on area above
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil nutrient status within natural range	Relevant nutrients and their natural ranges are yet to be defined. However, nitrogen deposition is noted as being relevant to this habitat (NPWS, 2013)
Vegetation composition: positive indicator fern and <i>Saxifraga</i> species	Number of species in local vicinity of a representative number of monitoring stops	Number of ferns and <i>Saxifraga</i> indicators at each monitoring stop is at least one	Attribute and target based on Perrin et al. (2014), where the list of positive indicator species for this habitat is also presented
Vegetation composition: positive indicator species	Number of species in local vicinity of a representative number of monitoring stops	Number of positive indicator species at each monitoring stop is at least three	Attribute and target based on Perrin et al. (2014), where the list of positive indicator species for this habitat is also presented
Vegetation composition: non-native species	Percentage cover in local vicinity of a representative number of monitoring stops	Proportion of vegetation composed of non-native species less than 1%	Attribute and target based on Perrin et al. (2014). Non-native species can be invasive and have deleterious effects on native vegetation. A low target is set as non-native species can spread rapidly and are most easily dealt with when still at lower abundances
Vegetation composition: bracken, native trees and shrubs	Percentage cover in local vicinity of a representative number of monitoring stops	Total cover of bracken (<i>Pteridium aquilinum</i>), native trees and shrubs less than 25%	Attribute and target based on Perrin et al. (2014). High cover of bracken, native trees and shrubs would indicate that the rocky slopes are becoming more vegetated which would impact on the niches of the chasmophytic vegetation
Vegetation structure: grazing and browsing	Percentage of leaves/shoots grazed/browsed in local vicinity of a representative number of monitoring stops	Live leaves of forbs and shoots of dwarf shrubs showing signs of grazing or browsing collectively less than 50%	Attribute and target based on Perrin et al. (2014)
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat	This includes species listed in the Flora (Protection) Order, 2015 and/or the red data lists (Lockhart et al., 2016; Wyse Jackson et al., 2016). The red-listed (Wyse Jackson et al., 2016) species holly-fern (<i>Polystichum lonchitis</i>) and alpine saw-wort (<i>Saussurea alpina</i>) have been recorded from the rocky cliffs (NPWS internal files; Doyle and Foss, 1986)

Conservation Objectives for : Clare Island Cliffs SAC [002243]

8220 Siliceous rocky slopes with chasmophytic vegetation


To maintain the favourable conservation condition of Siliceous rocky slopes with chasmophytic vegetation in Clare Island Cliffs SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Siliceous rocky slopes with chasmophytic vegetation has not been mapped in detail for Clare Island Cliffs SAC and thus the total area of the qualifying habitat is unknown. It occurs as a mosaic with calcareous rocky slopes with chasmophytic vegetation (8210) and is generally found on the sheer cliffs on the seaward face of Croaghmore and to the north and north-west of the island on some of the higher sea cliffs, above c.250m (NPWS internal files)
Habitat distribution	Occurrence	No decline, subject to natural processes	See note on area above
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil nutrient status within natural range	Relevant nutrients and their natural ranges are yet to be defined. However, nitrogen deposition is noted as being relevant to this habitat (NPWS, 2013)
Vegetation composition: positive indicator species	Number of species in local vicinity of a representative number of monitoring stops	At least one positive indicator species present in vicinity of each monitoring stop	Attribute and target based on Perrin et al. (2014), where the list of positive indicator species for this habitat is also presented
Vegetation composition: non-native species	Percentage cover in local vicinity of a representative number of monitoring stops	Proportion of vegetation composed of non-native species less than 1%	Attribute and target based on Perrin et al. (2014). Non-native species can be invasive and have deleterious effects on native vegetation. A low target is set as non-native species can spread rapidly and are most easily dealt with when still at lower abundances
Vegetation composition: bracken, native trees and shrubs	Percentage cover in local vicinity of a representative number of monitoring stops	Total cover of bracken (<i>Pteridium aquilinum</i>), native trees and shrubs less than 25%	Attribute and target based on Perrin et al. (2014). High cover of bracken, native trees and shrubs would indicate that the rocky slopes are becoming more vegetated which would impact on the niches of the chasmophytic vegetation
Vegetation structure: grazing and browsing	Percentage of leaves/shoots grazed/browsed in local vicinity of a representative number of monitoring stops	Live leaves of forbs and shoots of dwarf shrubs showing signs of grazing or browsing collectively less than 50%	Attribute and target based on Perrin et al. (2014)
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat	This includes species listed in the Flora (Protection) Order, 2015 and/or the red data lists (Lockhart et al., 2016; Wyse Jackson et al., 2016). The red-listed (Wyse Jackson et al., 2016) species holly-fern (<i>Polystichum lonchitis</i>) and alpine saw-wort (<i>Saussurea alpina</i>) have been recorded from the rocky cliffs (NPWS internal files; Doyle and Foss, 1986)



Legend

 Clare Island Cliffs SAC 002243




An Roinn Ealaíon, Oidhreachta,
Gnóthaí Féilgíonacha, Tuaithe agus Gaeltachta
Department of Arts, Heritage,
Regional, Rural and Gaeltacht Affairs

**MAP 1:
CLARE ISLAND SAC
CONSERVATION OBJECTIVES
SAC DESIGNATION**

Map to be read in conjunction with the NPWS Conservation Objectives Document.

**SITE CODE: SAC 002243; version 3.02
CO. MAYO**




The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
Ordnance Survey of Ireland Licence No EN 0059216. © Ordnance Survey of Ireland Government of Ireland

Níl sna teorainneacha ar na léarscáileanna ach nod garshuimhach ginearálta. Féadfar athbheithníthe a déanamh ar theorainneacha na gceantar comharthaithe. Suirbhéarachta Ordonáis na hÉireann Ceadúnas




**Map Version 1
Date: Oct 2016**

© Ordnance Survey Ireland



Legend

- Clare Island Cliffs SAC 002243
- West Connacht Coast SAC 002998
- Clare Island SPA 004136
- OSi Discovery Series County Boundary



Legend

- 1230 Vegetated sea cliffs of the Atlantic and Baltic coasts
- Clare Island Cliffs SAC 002243
- OSi Discovery Series County Boundary

An Roinn Ealaíon, Oidhreachta,
Gnóthaí Réigiúnacha, Tuaithe agus Gaeltachta
Department of Arts, Heritage,
Regional, Rural and Gaeltacht Affairs

**MAP 3:
CLARE ISLAND SAC
CONSERVATION OBJECTIVES
VEGETATED SEA CLIFFS**

Map to be read in conjunction with the NPWS Conservation Objectives Document.

**SITE CODE:
SAC 002243; version 3.02,
CO. MAYO**

0 0.3 0.6 0.9 1.2 1.5 km

The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
Ordnance Survey of Ireland Licence No EN 0059216. © Ordnance Survey of Ireland Government of Ireland

Níl sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbhreithnithe a déanamh ar theorainneacha na gceantar comharthaithe. Suirbhéarachta Ordonáis na hÉireann Ceadúnas

N

**Map Version 1
Date: Oct 2016**

National Parks and Wildlife Service

Conservation Objectives Series

Achill Head SAC 002268



An Roinn
Ealaíon, Oidhreachta agus Gaeltachta

Department of
Arts, Heritage and the Gaeltacht



**National Parks and Wildlife Service,
Department of Arts, Heritage and the Gaeltacht,
7 Ely Place, Dublin 2, Ireland.
Web: www.npws.ie
E-mail: nature.conservation@ahg.gov.ie**

Citation:

**NPWS (2013) Conservation Objectives: Achill Head SAC 002268. Version 1.
National Parks and Wildlife Service, Department of Arts, Heritage and the
Gaeltacht.**

**Series Editor: Rebecca Jeffrey
ISSN 2009-4086**

Introduction

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

** indicates a priority habitat under the Habitats Directive*

002268	Achill Head SAC
1140	Mudflats and sandflats not covered by seawater at low tide
1160	Large shallow inlets and bays
1170	Reefs

Please note that this SAC adjoins Keel Machair/Menaun Cliffs SAC (001513) and Croaghaun/Slievemore SAC (001955). See map 2. The conservation objectives for this site should be used in conjunction with those for adjacent sites as appropriate.

Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

NPWS Documents

Year : 2013
Title : Achill Head SAC (site code 2268) Conservation objectives supporting document- marine habitats V1
Author : NPWS
Series : Conservation objectives supporting document

Other References

Year : 1997
Title : The BioMar biotope viewer: a guide to marine habitats, fauna and flora in Britain and Ireland
Author : Picton, B.E.; Costello, M.J.
Series : Environmental Science Unit, Trinity College Dublin

Year : 2012
Title : Intertidal benthic survey of Achill Head SAC
Author : MERC
Series : Unpublished report to the Marine Institute and NPWS

Year : 2012
Title : Subtidal sediment and subtidal and intertidal reef survey of Achill Head SAC
Author : MERC
Series : Unpublished report to the Marine Institute and NPWS

Spatial data sources

Year :	Interpolated 2013
Title :	1995 BioMar Survey; 2011 intertidal and subtidal surveys
GIS Operations :	Polygon feature classes from marine community types base data sub-divided based on interpolation of marine survey data. Expert opinion used as necessary to resolve any issues arising
Used For :	1140, 1170, Marine community types (maps 3, 5 and 6)
<hr/>	
Year :	2005
Title :	OSi Discovery series vector data
GIS Operations :	High Water Mark (HWM) polyline feature class converted into polygon feature class; clipped to SAC boundary. EPA WFD transitional waterbody data erased from extent. Expert opinion used as necessary to resolve any issues arising
Used For :	1160 (map 4)
<hr/>	
Year :	2005
Title :	OSi Discovery series vector data
GIS Operations :	High water mark (HWM) and low water mark (LWM) polyline feature classes converted into polygon feature classes and combined; EU Annex I Saltmarsh and Coastal data erased out if present
Used For :	Marine community types base data (map 6)
<hr/>	

Conservation Objectives for : Achill Head SAC [002268]

1140 Mudflats and sandflats not covered by seawater at low tide

To maintain the favourable conservation condition of Mudflats and sandflats not covered by seawater at low tide in Achill Head SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	The permanent habitat area is stable or increasing, subject to natural processes. See map 3	Habitat area was estimated using OSi data as 16ha
Community distribution	Hectares	Conserve the following community type in a natural condition: Intertidal fine sand community. See map 6	Based on an intertidal survey undertaken in 2011 (MERC, 2012). See marine supporting document for further information

Conservation Objectives for : Achill Head SAC [002268]

1160 Large shallow inlets and bays


To maintain the favourable conservation condition of Large shallow inlets and bays in Achill Head SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	The permanent habitat area is stable or increasing, subject to natural processes. See map 4	Habitat area was estimated as 6888ha using OSI data and the Transitional Water Body area as defined under the Water Framework Directive
Community distribution	Hectares	Conserve the following community types in a natural condition: Intertidal fine sand community; Mobile subtidal sand with <i>Gastrosaccus spinifer</i> community; Subtidal sand with <i>Bathyporeia elegans</i> and polychaetes community complex; Intertidal reef community complex; <i>Laminaria</i> -dominated community complex; Subtidal reef community. See map 5	Based on the BioMar survey in 1995 (Picton and Costello, 1997) and intertidal and subtidal surveys undertaken in 2011 (MERC, 2012). See marine supporting document for further information


1170 Reefs


To maintain the favourable conservation condition of Reefs in Achill Head SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	The permanent habitat area is stable or increasing, subject to natural processes. See map 5	Habitat area estimated as 1919ha from the 1995 BioMar Survey (Picton and Costello, 1997) and a reef survey in 2011 (MERC, 2012)
Distribution	Occurrence	The distribution of reefs remains stable, subject to natural processes. See map 5 for mapped distribution	Based on information from the 1995 BioMar Survey (Picton and Costello, 1997) and a reef survey in 2011 (MERC, 2012). See marine supporting document for further details
Community structure	Biological composition	Conserve the following community types in a natural condition: Intertidal reef community complex; <i>Laminaria</i> -dominated community complex; Subtidal reef community. See map 6	Reef mapping based on information from the 1995 BioMar Survey (Picton and Costello, 1997) and a reef survey in 2011 (MERC, 2012). See marine supporting document for further details



Legend

 Achill Head SAC 002268

 *An Roinn Ealaíon, Oidhreachta agus Gaeltachta*
 Department of Arts, Heritage and the Gaeltacht

MAP 1:
ACHILL HEAD SAC
CONSERVATION OBJECTIVES
SAC DESIGNATION


Map to be read in conjunction with the NPWS Conservation Objectives Document.


SITE CODE:
SAC 002268; version 3. CO. MAYO

0 1 2 3 km

The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision. Reproduced from Ordnance Survey material by permission of the Government (Permit number EN 0059212).


Níl sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbhréithníthe a déanamh ar theorainneacha na goeantar comharthaíthe. Macasamhail d'ábhar na Suirbhéarachta Ordoanáis le chead ón Rialtas (Ceadúnas Uimh. EN 0059212)


Map Version 1
Date: Oct 2013




Legend

- Keel Machair/Menaun Cliffs SAC 001513
- Croaghaun/Slievemore SAC 001955
- Achill Head SAC 002268
- OSi Discovery Series County Boundaries



Legend


- Achill Head SAC 002268
- 1140 Mudflats and sandflats not covered by sea water at low tide
- OSi Discovery Series County Boundaries



Legend

- Achill Head SAC 002268
- 1160 Large shallow inlets and bays
- OSi Discovery Series County Boundaries






Legend

- Achill Head SAC 002268
- 1170 Reefs
- OSi Discovery Series County Boundaries





Legend

- Achill Head SAC 002268
- OSi Discovery Series County Boundaries

Marine Community Types

- Intertidal fine sand community
- Intertidal reef community complex
- Laminaria*-dominated community complex
- Mobile subtidal sand with *Gastrosaccus spinifer* community
- Subtidal reef community
- Subtidal sand with *Bathyporeia elegans* and polychaetes community complex



National Parks and Wildlife Service

Conservation Objectives Series

Williamstown Turloughs SAC 002296



An Roinn
Cultúir, Oidhreachta agus Gaeltachta
Department of
Culture, Heritage and the Gaeltacht



**National Parks and Wildlife Service,
Department of Culture, Heritage and the Gaeltacht,
7 Ely Place, Dublin 2, Ireland.
Web: www.npws.ie
E-mail: nature.conservation@ahg.gov.ie**

Citation:

**NPWS (2018) Conservation Objectives: Williamstown Turloughs SAC 002296.
Version 1. National Parks and Wildlife Service, Department of Culture, Heritage
and the Gaeltacht.**

**Series Editor: Rebecca Jeffrey
ISSN 2009-4086**

Introduction

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

** indicates a priority habitat under the Habitats Directive*

002296 Williamstown Turloughs SAC

3180 TurloughsE

Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

NPWS Documents

Year :	1992
Title :	Turloughs over 10ha - Vegetation survey and evaluation
Author :	Goodwillie, R.N.
Series :	Unpublished report to NPWS
<hr/>	
Year :	2017
Title :	Conservation objectives supporting document: Turloughs* and Rivers with muddy banks with Chenopodium rubri p.p. and Bidention p.p. vegetation
Author :	O Connor, Á.
Series :	Conservation objectives supporting document
<hr/>	

Other References

Year :	1997
Title :	Williamstown Environmental Study
Author :	ECS (Environmental Consultancy Services)
Series :	Unpublished report to the Office of Public Works
<hr/>	

Spatial data sources

Year : 2017

Title : Internal NPWS data

GIS Operations : Paper map scanned and georectified. Turlough as outlined on map digitised and clipped to SAC boundary. Expert opinion used as necessary to resolve any issues arising

Used For : 3180 (map 2)

Conservation Objectives for : Williamstown Turloughs SAC [002296]



3180 Turloughs

To restore the favourable conservation condition of Turloughs* in Williamstown Turloughs SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable at c.110ha or increasing, subject to natural processes. See map 2	Williamstown Turloughs SAC is a complex of three turlough basins: Curragh, Polleagh (including Polleagh West) and Gortduff (see map 2). The total area of turlough habitat in the SAC is estimated to be 109.5ha (based on NPWS internal files). In the 1990s, drainage damaged the SAC and the medium- to long-term impacts on turlough area have not been assessed. Restoration of turlough area may therefore be required. See O Connor (2017) for information on all attributes and targets
Habitat distribution	Occurrence	No decline, subject to natural processes	As noted above, Williamstown Turloughs SAC contains three turloughs: Curragh, Polleagh and Gortduff. See map 2
Hydrological regime: groundwater contribution; flood duration, frequency, area and depth; permanently flooded/wet areas	Various	Restore appropriate natural hydrological regime necessary to support the natural structure and functioning of the habitat	See O Connor (2017) for details on the sub-divided attributes and targets of hydrological regime. The hydrology of the Williamstown turloughs was investigated by ECS (1997). The turloughs are in hydraulic continuity, their zone of contribution is small and groundwater flow is to the west. Drainage works in 1995 and 1996 impacted the natural hydrological regime, including by reducing the flood extent, the contributing catchment by c.20% and, at Polleagh, water level fluctuations by c.37%. A pond was also infilled. Further drainage was observed in 2005 (NPWS internal files). All three turloughs have permanent water: an estimated 13ha at Polleagh Lough and 5.25ha at Curragh (ECS, 1997). Permanent water at Polleagh and Gortduff is likely dependent on a perched water table in summer. Polleagh Lough merges with Polleagh West at times of high flood, completely surrounding a hill of glacial drift. Sinkholes have been observed at Curragh and Gortduff
Soil type	Hectares	Maintain variety, area and extent of soil types necessary to support turlough vegetation and other biota	ECS (1997) noted sub-soils of peat and glacial deposits and soils of peat, degraded grey brown podzolics, brown earths, gleys and podzols in the Williamstown study area. The Williamstown turloughs are dominated by wet soils, including peat and marl. ECS (1997) recorded peat, marl, mud and a stony shoreline at Polleagh turlough, and peat, soft mud and well-drained mineral soils at Curragh turlough
Soil nutrient status: nitrogen and phosphorus	N and P concentration in soil	Maintain/restore nutrient status appropriate to soil types and vegetation communities	Direct fertilisation and run-off of organic and chemical fertilisers may have artificially enriched soils in the Williamstown turloughs
Physical structure: bare ground	Presence	Maintain/restore sufficient wet bare ground, as appropriate	Bare soil was noted on the edge of Polleagh Lough during surveys in 1996 (ECS, 1997). A wet annual community, likely to be Annex I habitat 3270 (Rivers with muddy banks with <i>Chenopodium rubri</i> p.p. and <i>Bidention</i> p.p. vegetation), was recorded to the north-west of Polleagh Lough (ECS, 1997). This community was associated with permanently damp, soft, fine mud
Chemical processes: calcium carbonate deposition and concentration	Calcium carbonate deposition rate/soil concentration	Maintain appropriate calcium carbonate deposition rate and concentration in soil	As noted above, the Williamstown turloughs are dominated by wet soils, including peat and marl. Marl deposition was recorded at Polleagh and there was some associated with stony bays at the lake margins at Curragh (ECS, 1997)


Water quality	Various	Maintain/restore appropriate water quality to support the natural structure and functioning of the habitat	Water quality is sub-divided into more detailed attributes (nutrients, colour, phytoplankton and epiphyton biomass) and targets in O Connor (2017). The abundance of peaty substratum, marl, common sedge (<i>Carex nigra</i>) communities and species such as bulbous rush (<i>Juncus bulbosus</i>) and shoreweed (<i>Littorella uniflora</i>) at Polleagh and Curragh indicate highly calcareous, oligotrophic conditions and a requirement for high status/oligotrophic water. Limited water sampling in 1995/6 indicated generally oligotrophic conditions at Curragh and Polleagh; however, elevated chlorophyll <i>a</i> concentrations were measured in both (Curragh range <1-13.25µg/l; Polleagh <1-12.1µg/l), indicating enrichment. Organic and chemical agricultural fertilisation, animal housing, farmyards and domestic dwellings are potential nutrient sources
Active peat formation	Flood duration	Maintain active peat formation	As noted above, the Williamstown turloughs are dominated by wet soils, including peat and marl. Extensive peat development was noted to the south of Polleagh and at the south-eastern side of the basin at Curragh (ECS, 1997)
Vegetation composition: area of vegetation communities	Hectares	Maintain/restore area of sensitive and high conservation value vegetation communities/units	As noted, the impacts of drainage on the vegetation communities of the turloughs have not been monitored. The vegetation of Polleagh and Curragh was surveyed in detail (ECS, 1997) and classified in accordance with Goodwillie (1992). 13 Goodwillie (1992) turlough communities were recorded at Polleagh (P) and 8 at Curragh (C), a total of 15: Wet <i>Carex nigra</i> (P,C), Dry <i>C. nigra</i> (P,C), Peaty <i>C. nigra</i> (P), Reedbed (P), Marl pond (P), Peaty pond (C), Wet annuals (P), Open water (P), <i>Polygonum amphibium</i> (P,C), <i>P. amphibium</i> (grassy) (P,C), Tall herb (P), Peat grassland (P), <i>Potentilla reptans</i> (species-rich) (P), Poor grassland (C) and <i>Lolium</i> grassland (P,C) (ECS, 1997). A <i>Juncus</i> grassland community was also found at both turloughs (ECS, 1997). <i>Polygonum amphibium</i> , <i>P. amphibium</i> (grassy), Wet <i>Carex nigra</i> , Dry <i>C. nigra</i> and <i>Lolium</i> grassland were noted at Gortduff (ECS, 1997)
Vegetation composition: vegetation zonation	Distribution	Maintain/restore vegetation zonation/mosaic characteristic of the site	As detailed above, the vegetation of the turloughs in the SAC was surveyed and classified in accordance with Goodwillie (1992) (ECS, 1997). Good vegetation zonation was associated with the steep-sided Gortduff. Drainage works are likely to have impacted vegetation zonation through both direct vegetation removal/reclamation and in-filling, and hydrological change
Vegetation structure: sward height	Centimetres	Maintain/restore sward heights appropriate to the vegetation unit, and a variety of sward heights across the turloughs	It was noted that Curragh was more intensively grazed than Polleagh in 1996 (ECS, 1997)
Typical species	Presence	Maintain/restore typical species within and across the turloughs	Typical species is sub-divided into more detailed attributes (terrestrial, wetland and aquatic plants, invertebrates and birds) and targets in O Connor (2017). As noted above, the impacts of drainage on the species of the Williamstown turloughs have not been monitored. 112 vascular plant species have been recorded in the Williamstown turloughs, including northern yellow-cross (<i>Rorippa islandica</i>) and red goosefoot (<i>Chenopodium rubrum</i>) at Polleagh (ECS, 1997). The wet annuals community recorded at Polleagh requires further study to determine if it is Annex I habitat 3270 (Rivers with muddy banks with <i>Chenopodium rubri</i> p.p. and <i>Bidention</i> p.p. vegetation). Wintering waterfowl at Polleagh and Curragh include wigeon, mallard, teal, tufted duck, shoveler, whooper swan, lapwing, golden plover, curlew and dunlin (ECS, 1997). Breeding waterfowl included snipe, lapwing, redshank and ringed plover. Curragh has fish, including pike

Fringing habitats: area	Hectares	Maintain/restore marginal fringing habitats that support turlough vegetation, invertebrate, mammal and/or bird populations	NPWS site files show much of the land surrounding the Williamstown turloughs was 'improved' grassland in 2001 (NPWS internal files). Some lowland dry grassland occurred at all three turloughs, and also lowland wet grassland around Polleagh and Curragh. Cutover bog is widespread in the catchment
Vegetation structure: turlough woodland	Species diversity and woodland structure	Maintain appropriate turlough woodland diversity and structure	No turlough woodland or scrub has been recorded in Williamstown Turloughs SAC



Legend

 Williamstown Turloughs SAC 002296




An Roinn
Cultúir, Oidhreacht agus Gaeltachta
Department of
Culture, Heritage and the Gaeltacht

**MAP 1:
WILLIAMSTOWN TURLOUGHs SAC
CONSERVATION OBJECTIVES
SAC DESIGNATION**

Map to be read in conjunction with the NPWS Conservation Objectives Document.

**SITE CODE:
SAC 002296; version 3.02. CO. GALWAY**

0 140 280 420 560 Meters




The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
Ordnance Survey of Ireland Licence No EN 0059216. © Ordnance Survey of Ireland Government of Ireland

Níl sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbhreithnithe a déanamh ar theorainneacha na gceantar comharthaithe. Suirbhéarachta Ordonáis na hÉireann Ceadúnas Uimh EN 0059216. © Suirbhéarachta Ordonáis na hÉireann Rialtas na hÉireann



**Map Version 1
Date: Dec 2017**



Legend

- 3180 Turlough
- Williamstown Turloughs SAC 002296
- OSi Discovery Series County Boundary

National Parks and Wildlife Service

Conservation Objectives Series

River Moy SAC 002298



An Roinn Ealaíon, Oidhreachta,
Gnóthaí Réigiúnacha, Tuaithe agus Gaeltachta

Department of Arts, Heritage,
Regional, Rural and Gaeltacht Affairs



**National Parks and Wildlife Service,
Department of Arts, Heritage, Regional, Rural and Gaeltacht Affairs,
7 Ely Place, Dublin 2, Ireland.
Web: www.npws.ie
E-mail: nature.conservation@ahg.gov.ie**

Citation:

**NPWS (2016) Conservation Objectives: River Moy SAC 002298. Version 1.
National Parks and Wildlife Service, Department of Arts, Heritage, Regional,
Rural and Gaeltacht Affairs.**

**Series Editor: Rebecca Jeffrey
ISSN 2009-4086**

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

* indicates a priority habitat under the Habitats Directive

002298	River Moy SAC
--------	---------------

1092	White-clawed Crayfish <i>Austropotamobius pallipes</i>
1095	Sea Lamprey <i>Petromyzon marinus</i>
1096	Brook Lamprey <i>Lampetra planeri</i>
1106	Salmon <i>Salmo salar</i>
1355	Otter <i>Lutra lutra</i>
7110	Active raised bogs*
7120	Degraded raised bogs still capable of natural regeneration
7150	Depressions on peat substrates of the Rhynchosporion
7230	Alkaline fens
91A0	Old sessile oak woods with <i>Ilex</i> and <i>Blechnum</i> in the British Isles
91E0	Alluvial forests with <i>Alnus glutinosa</i> and <i>Fraxinus excelsior</i> (Alno-Padion, Alnion incanae, Salicion albae)*

Please note that this SAC overlaps with Killala Bay/Moy Estuary SPA (004036) and Lough Conn and Lough Cullin SPA (004228). It is adjacent to Killala Bay/Moy Estuary SAC (000458), Lough Hoe Bog SAC (000633), Bellacorick Bog Complex SAC (001922) and Ox Mountains Bogs SAC (002006). See map 2. The conservation objectives for this site should be used in conjunction with those for overlapping and adjacent sites as appropriate.

Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

NPWS Documents

Year :	1998
Title :	Conservation management of the white-clawed crayfish, (<i>Austropotamobius pallipes</i>)
Author :	Reynolds, J.D.
Series :	Irish Wildlife Manual No. 1
Year :	2004
Title :	The status and distribution of lamprey and shad in the Slaney and Munster Blackwater SACs
Author :	King, J.J.; Linnane, S.M.
Series :	Irish Wildlife Manuals No. 14
Year :	2004
Title :	A survey of juvenile lamprey populations in the Moy catchment
Author :	O'Connor, W.
Series :	Irish Wildlife Manuals No. 15
Year :	2006
Title :	Otter survey of Ireland 2004/2005
Author :	Bailey, M.; Rochford, J.
Series :	Irish Wildlife Manual No. 23
Year :	2006
Title :	Assessment of impacts of turf cutting on designated raised bogs
Author :	Fernandez Valverde, F.; MacGowan, F.; Farrell, M.; Crowley, W.; Croal, Y.; Fanning, M.; McKee, A-M.
Series :	Unpublished report to NPWS
Year :	2007
Title :	Supporting documentation for the Habitats Directive Conservation Status Assessment - backing documents. Article 17 forms and supporting maps
Author :	NPWS
Series :	Unpublished report to NPWS
Year :	2008
Title :	National survey of native woodlands 2003-2008
Author :	Perrin, P.M.; Martin, J.; Barron, S.; O'Neill, F.H.; McNutt, K.E.; Delaney, A.
Series :	Unpublished Report to NPWS
Year :	2010
Title :	A provisional inventory of ancient and long-established woodland in Ireland
Author :	Perrin, P.M.; Daly, O.H.
Series :	Irish Wildlife Manual No. 46
Year :	2010
Title :	A technical manual for monitoring white-clawed crayfish (<i>Austropotamobius pallipes</i>) in Irish lakes
Author :	Reynolds, J., O'Connor, W., O'Keeffe, C.; Lynn, D.
Series :	Irish Wildlife Manual No.45
Year :	2012
Title :	Killala Bay/Moy Estuary SAC (00458) Coastal Supporting doc V1
Author :	NPWS
Series :	Conservation objectives supporting document

Year :	2012
Title :	Killala Bay/Moy Estuary SAC (000458) Marine supporting doc v.1
Author :	NPWS
Series :	Conservation objectives supporting document
Year :	2013
Title :	National otter survey of Ireland 2010/12
Author :	Reid, N.; Hayden, B.; Lundy, M.G.; Pietravalle, S.; McDonald, R.A.; Montgomery, W.I.
Series :	Irish Wildlife Manual No. 76
Year :	2014
Title :	Guidelines for a national survey and conservation assessment of upland vegetation and habitats in Ireland, Version 2.0
Author :	Perrin, P.M.; Barron, S.J.; Roche, J.R.; O'Hanrahan, B.
Series :	Irish Wildlife Manual No. 79
Year :	2014
Title :	Raised Bog Monitoring and Assessment Survey 2013
Author :	Fernandez, F.; Connolly K.; Crowley W.; Denyer J.; Duff K.; Smith G.
Series :	Irish Wildlife Manual No. 81
Year :	2014
Title :	National raised bog SAC management plan
Author :	Department of Arts, Heritage and the Gaeltacht
Series :	Draft for consultation. 15 January 2014
Year :	2014
Title :	Derrynabrock Bog (SAC 002298), Co.Roscommon/Mayo, Site Report
Author :	Fernandez, F.; Connolly, K.; Crowley, W.; Denyer J.; Duff K.; Smith G.
Series :	Raised bog monitoring and assessment survey 2013
Year :	2014
Title :	Tawnaghbeg Bog (SAC 002298), Co. Mayo, Site Report
Author :	Fernandez, F.; Connolly, K.; Crowley, W.; Denyer J.; Duff K.; Smith G.
Series :	Raised bog monitoring and assessment survey 2013
Year :	2016
Title :	River Moy SAC (site code: 2298) Conservation objectives supporting document- raised bog habitats V1
Author :	NPWS
Series :	Conservation objectives supporting document

Other References

Year :	1982
Title :	Otter survey of Ireland
Author :	Chapman, P.J.; Chapman, L.L.
Series :	Unpublished report to Vincent Wildlife Trust
Year :	2002
Title :	Reversing the habitat fragmentation of British woodlands
Author :	Peterken, G.
Series :	WWF-UK, London

Year : 2003
Title : Monitoring the river, sea and brook lamprey, *Lampetra fluviatilis*, *L. planeri* and *Petromyzon marinus*
Author : Harvey, J.; Cowx, I.
Series : Conserving Natura 2000 Rivers Monitoring Series No. 5. English Nature, Peterborough

Year : 2003
Title : Identifying lamprey. A field key for sea, river and brook lamprey
Author : Gardiner, R.
Series : Conserving Natura 2000 rivers, Conservation techniques No. 4. English Nature, Peterborough

Year : 2007
Title : Evolutionary history of lamprey paired species *Lampetra fluviatilis* L. and *Lampetra planeri* Bloch as inferred from mitochondrial DNA variation
Author : Espanhol, R.; Almeida, P.R.; Alves, M.J.
Series : Molecular Ecology 16, 1909-1924

Year : 2010
Title : Otter tracking study of Roaringwater Bay
Author : De Jongh, A.; O'Neill, L.
Series : Unpublished draft report to NPWS

Year : 2015
Title : Behaviour of sea lamprey (*Petromyzon marinus* L.) at man-made obstacles during upriver spawning migration: use of telemetry to assess efficacy of weir modifications for improved passage
Author : Rooney, S.M.; Wightman, G.D.; O Conchuir, R.; King, J.J.
Series : Biology and Environment: Proc. R. Ir. Acad. 115 B, 1-12

Year : 2015
Title : River engineering works and lamprey ammocoetes; impacts, recovery, mitigation
Author : King, J.J.; Wightman, G.D.; Hanna, G.; Gilligan, N.
Series : Water and Environment Journal, 29, 482-488

Year : 2016
Title : The status of Irish salmon stocks in 2015 with precautionary catch advice for 2016
Author : Standing Scientific Committee on Salmon
Series : Independent scientific report to Inland Fisheries Ireland

Spatial data sources

Year :	2014
Title :	Scientific Basis for Raised Bog Conservation in Ireland
GIS Operations :	RBSB13_SACs_ARB_DRB dataset, RBSB13_SACs_2012_HB dataset, RBSB13_SACs_DrainagePatterns_5k dataset and RBSB13_SAC_LIDAR_DTMs dataset clipped to SAC boundary. Expert opinion used as necessary to resolve any issues arising
Used For :	Potential 7110; digital elevation model; drainage patterns (maps 3 and 5)
Year :	2013
Title :	Raised Bog Monitoring and Assessment Survey 2013
GIS Operations :	RBMA13_ecotope_map dataset clipped to SAC boundary. Appropriate ecotopes selected and exported to new dataset. Expert opinion used as necessary to resolve any issues arising
Used For :	7110 ecotopes (map 4)
Year :	Digitised 2003
Title :	Raised Bog Restoration Project 1999
GIS Operations :	Ecotope dataset clipped to SAC boundary. Appropriate ecotopes selected and exported to new dataset. Expert opinion used as necessary to resolve any issues arising
Used For :	7110 ecotopes (map 4)
Year :	Revision 2010
Title :	National Survey of Native Woodlands 2003-2008. Version 1
GIS Operations :	QIs selected; clipped to SAC boundary. Expert opinion used as necessary to resolve any issues arising
Used For :	91A0, 91E0 (map 6)
Year :	2005
Title :	OSi Discovery series vector data
GIS Operations :	Creation of a 10m buffer on the terrestrial side of river banks data; creation of 20m buffer applied to canal centreline data. Creation of a 20m buffer applied to river and stream centreline data; These datasets combined with the derived OSi 1:5000 vector lake buffer data. Overlapping regions investigated and resolved; resulting dataset clipped to SAC boundary. Expert opinion used as necessary to resolve any issues arising
Used For :	1355 (no map)
Year :	2010
Title :	OSi 1:5000 IG vector dataset
GIS Operations :	Creation of 80m buffer on the aquatic side of lake data; creation of 10m buffer on the terrestrial side of lake data. These datasets combined with the derived OSi Discovery Series river and canal datasets. Overlapping regions investigated and resolved; resulting dataset clipped to SAC boundary. Expert opinion used as necessary to resolve any issues arising. Creation of 250m buffer on aquatic side of the lake boundary to highlight potential commuting points
Used For :	1355 (map 8)
Year :	2016
Title :	NPWS rare and threatened species database
GIS Operations :	Dataset created from spatial references in database records. Expert opinion used as necessary to resolve any issues arising
Used For :	1092 (map 7)

Conservation Objectives for : River Moy SAC [002298]

7110 Active raised bogs

To restore the favourable conservation condition of Active raised bogs in River Moy SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Restore area of active raised bog to 132.4ha, subject to natural processes	There are five raised bogs listed for River Moy SAC. The total area of Active Raised Bog (ARB) habitat for these five bogs was mapped at 45.3ha. Area of Degraded Raised Bog (DRB) on the High Bog (HB) has been modelled as 152.4ha. See map 3. However, it is estimated that only 82.1ha is potentially restorable to ARB by drain blocking. The total potential ARB on the HB is therefore estimated to be 127.4ha. Eco-hydrological assessments of the cutover estimates that an additional 5.0ha of bog forming habitats could be restored. The long term target for ARB is therefore 132.4ha. See raised bog supporting document for further details on this and following attributes
Habitat distribution	Occurrence	Restore the distribution and variability of active raised bog across the SAC. See map 4 for most recently mapped distribution	ARB occurs on most of the bogs in the River Moy SAC. DRB occurs on all five bogs in the River Moy SAC. There is also potential for ARB restoration on cutover areas surrounding the bogs (see area target above)
High bog area	Hectares	No decline in extent of high bog necessary to support the development and maintenance of active raised bog. See map 3	The area of high bog within the five raised bogs listed for River Moy SAC in 2012 (latest figure available) was 498.4ha (DAHG 2014)
Hydrological regime: water levels	Centimetres	Restore appropriate water levels throughout the site	For ARB, mean water level needs to be near or above the surface of the bog lawns for most of the year. Seasonal fluctuations should not exceed 20cm, and should only be 10cm below the surface, except for very short periods of time. Open water is often characteristic of soak systems
Hydrological regime: flow patterns	Flow direction; slope	Restore, where possible, appropriate high bog topography, flow directions and slopes. See map 5 for current situation	ARB depends on mean water levels being near or above the surface of bog lawns for most of the year. Long and gentle slopes are the most favourable to achieve these conditions. Changes to flow directions due to subsidence of bogs can radically change water regimes and cause drying out of high quality ARB areas and soak systems
Transitional areas between high bog and adjacent mineral soils (including cutover areas)	Hectares; distribution	Restore adequate transitional areas to support/protect active raised bog and the services it provides	ARB is threatened due to effects of past drainage and peat-cutting around the margins of the bogs within the River Moy SAC. Natural marginal habitats no longer exist. Eco-hydrological assessments have evaluated the potential for ARB restoration on cutover areas (see note for habitat area attribute above)
Vegetation quality: central ecotope, active flush, soaks, bog woodland	Hectares	Restore 66.2ha of central ecotope/active flush/soaks/bog woodland as appropriate	At least 50% of ARB habitat should be high quality (i.e. central ecotope, active flush, soaks, bog woodland). Target area of active raised bog for the site has been set at 132.4ha (see area target above)
Vegetation quality: microtopographical features	Hectares	Restore adequate cover of high quality microtopographical features	High quality microtopography (hummocks, hollows and pools) is well developed in less disturbed parts of the bogs in River Moy SAC
Vegetation quality: bog moss (<i>Sphagnum</i>) species	Percentage cover	Restore adequate cover of bog moss (<i>Sphagnum</i>) species to ensure peat-forming capacity	<i>Sphagnum</i> cover varies naturally across Ireland with relatively high cover in the east to lower cover in the west. Hummock forming species such as <i>Sphagnum austini</i> are particularly good peat formers. <i>Sphagnum</i> cover and distribution also varies naturally across a site

Typical ARB species: flora	Occurrence	Restore, where appropriate, typical active raised bog flora	Typical flora species include widespread species, as well as those with more restricted distributions but typical of the habitat's subtypes or geographical range
Typical ARB species: fauna	Occurrence	Restore, where appropriate, typical active raised bog fauna	Typical fauna species include widespread species, as well as those with more restricted distributions but typical of the habitat's subtypes or geographical range
Elements of local distinctiveness	Occurrence	Maintain features of local distinctiveness, subject to natural processes	An important feature of interest in relation to the raised bogs in the River Moy SAC is the fact that they occur at the north-western edge of the geographic range of the habitat in Ireland
Negative physical indicators	Percentage cover	Negative physical features absent or insignificant	Negative physical indicators include: bare peat, algae dominated pools and hollows, marginal cracks, tear patterns, subsidence features such as dry mineral mounds/ridges emerging or expanding and evidence of burning
Vegetation composition: native negative indicator species	Percentage cover	Native negative indicator species at insignificant levels	Disturbance indicators include species indicative of conditions drying out such as abundant bog asphodel (<i>Narthecium ossifragum</i>), deergrass (<i>Trichophorum germanicum</i>) and harestail cotton-grass (<i>Eriophorum vaginatum</i>) forming tussocks; abundant magellanic bog-moss (<i>Sphagnum magellanicum</i>) in pools previously dominated by <i>Sphagnum</i> species typical of very wet conditions (e.g. feathery bog-moss (<i>S. cuspidatum</i>)); and indicators of frequent burning events such as abundant <i>Cladonia floerkeana</i> and high cover of carnation sedge (<i>Carex panicea</i>) (particularly in true midlands raised bogs)
Vegetation composition: non-native invasive species	Percentage cover	Non-native invasive species at insignificant levels and not more than 1% cover	Most common non-native invasive species include lodgepole pine (<i>Pinus contorta</i>), rhododendron (<i>Rhododendron ponticum</i>), and pitcherplant (<i>Sarracenia purpurea</i>)
Air quality: nitrogen deposition	kg N/ha/year	Air quality surrounding bog close to natural reference conditions. The total N deposition should not exceed 5kg N/ha/yr	Change in air quality can result from fertiliser drift; adjacent quarry activities; or other atmospheric inputs. The critical load range for ombrotrophic bogs has been set as between 5 and 10kg N/ha/yr (Bobbink and Hettelingh, 2011). The latest N deposition figures for the area around the bogs in River Moy SAC suggests that the current level is approximately 8.5kg N/ha/yr (Henry and Aherne, 2014)
Water quality	Hydrochemical measures	Water quality on the high bog and in transitional areas close to natural reference conditions	Water chemistry within raised bogs is influenced by atmospheric inputs (rainwater). However, within soak systems, water chemistry is influenced by other inputs such as focused flow or interaction with underlying substrates. Water chemistry in areas surrounding the high bog varies due to influences of different water types (bog water, regional groundwater and run-off from surrounding mineral lands)

Conservation Objectives for : River Moy SAC [002298]

7120 Degraded raised bogs still capable of natural regeneration

The long-term aim for Degraded raised bogs still capable of natural regeneration is that its peat-forming capability is re-established; therefore, the conservation objective for this habitat is inherently linked to that of Active raised bogs (7110) and a separate conservation objective has not been set in River Moy SAC

Attribute	Measure	Target	Notes
------------------	----------------	---------------	--------------

Conservation Objectives for : River Moy SAC [002298]

7150 Depressions on peat substrates of the Rhynchosporion

Depressions on peat substrates of the Rhynchosporion is an integral part of good quality Active raised bogs (7110) and thus a separate conservation objective has not been set for the habitat in River Moy SAC

Attribute	Measure	Target	Notes
------------------	----------------	---------------	--------------

7230 Alkaline fens

To maintain the favourable conservation condition of Alkaline fens in River Moy SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	The full extent of of this habitat within the SAC is unknown. An extensive area is known to occur as part of a wetland complex on the Glone River, north-west of Ballyhaunis but there are likely to be other areas present in the SAC
Habitat distribution	Occurrence	No decline, subject to natural processes	Full distribution of the habitat in this SAC is currently unknown- see note above
Hydrological regime	Metres	Appropriate natural hydrological regimes necessary to support the natural structure and functioning of the habitat	Maintenance of groundwater, surface water flows and water table levels within natural ranges is essential for this wetland habitat
Peat formation	Flood duration	Active peat formation, where appropriate	In order for peat to form, water levels need to be slightly below or above the soil surface for c.90% of the time (Jim Ryan, pers. comm.)
Water quality: nutrients	Water chemistry measures	Appropriate water quality to support the natural structure and functioning of the habitat	Fens receive natural levels of nutrients (e.g. iron, magnesium and calcium) from water sources. However, they are generally poor in nitrogen and phosphorus with the latter tending to be the limiting nutrient
Vegetation structure: typical species	Percentage	Maintain vegetation cover of typical species including brown mosses and vascular plants	Mosses listed for fen in this SAC include <i>Campylium stellatum</i> , <i>Aneura pinguis</i> and <i>Scorpidium scorpioides</i> while vascular plants include long-stalked yellow sedge (<i>Carex lepidocarpa</i>), black bog rush (<i>Schoenus nigricans</i>), blunt-flowered rush (<i>Juncus subnodulosus</i>), purple moor-grass (<i>Molinia caerulea</i>), grass of Parnassus (<i>Parnassia palustris</i>), butterwort (<i>Pinguicula vulgaris</i>), marsh helleborine (<i>Epipactis palustris</i>) and meadow thistle (<i>Cirsium dissectum</i>) (internal NPWS files)
Vegetation composition: trees and shrubs	Percentage	Cover of scattered native trees and shrubs less than 10%	Scrub and trees will tend to invade if fen conditions become drier. Attribute and target based on upland habitat conservation assessment criteria (Perrin et al., 2014)
Physical structure: disturbed bare ground	Percentage	Cover of disturbed bare ground less than 10%. Where tufa is present, disturbed bare ground less than 1%	While grazing may be appropriate in this habitat, excessive areas of disturbed bare ground may develop due to unsuitable grazing regimes. Attribute and target based on upland habitat conservation assessment criteria (Perrin et al., 2014)
Physical structure: drainage	Percentage	Areas showing signs of drainage as a result of drainage ditches or heavy trampling less than 10%	Attribute and target based on upland habitat conservation assessment criteria (Perrin et al., 2014)

91A0 Old sessile oak woods with Ilex and Blechnum in the British Isles

To maintain the favourable conservation condition of Old sessile oak woods with *Ilex* and *Blechnum* in the British Isles in River Moy SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Old sessile oakwoods are likely to occur as mosaics with other woodland types and the total extent within the SAC is unknown. Two sites (1763, 1800) in the SAC were surveyed as part of the the National Survey of Native Woodlands (NSNW) (Perrin et al., 2008). Site 1763 (Pontoon) is an extensive area of woodland and 106.3ha was mapped as this Annex I habitat type (or mosaics containing it). See map 6. NB further areas are likely to be present within the SAC
Habitat distribution	Occurrence	No decline. Woodlands surveyed as part of the NSNW are shown on map 6	The main location of this woodland type in the SAC is Pontoon Woods. See note on area above
Woodland size	Hectares	Area stable or increasing. Where topographically possible, "large"; woods at least 25ha in size and "small" woods at least 3ha in size	The sizes of at least some of the existing woodlands need to be increased in order to reduce habitat fragmentation and benefit those species requiring "deep" woodland conditions (Peterken, 2002). Topographical and land ownership constraints may restrict expansion
Woodland structure: cover and height	Percentage and metres	Diverse structure with a relatively closed canopy containing mature trees; subcanopy layer with semi-mature trees and shrubs; and well-developed herb layer	Described in Perrin et al (2008)
Woodland structure: community diversity and extent	Hectares	Maintain diversity and extent of community types	Described in Perrin et al. (2008)
Woodland structure: natural regeneration	Seedling: sapling: pole ratio	Seedlings, saplings and pole age-classes occur in adequate proportions to ensure survival of woodland canopy	Oak (<i>Quercus</i> spp.) regenerates poorly. In suitable sites ash (<i>Fraxinus excelsior</i>) can regenerate in large numbers although few seedlings reach pole size
Woodland structure: dead wood	m ³ per hectare; number per hectare	At least 30m ³ /ha of fallen timber greater than 10cm diameter; 30 snags/ha; both categories should include stems greater than 40cm diameter	Dead wood is a valuable resource and an integral part of a healthy, functioning woodland ecosystem
Woodland structure: veteran trees	Number per hectare	No decline	Mature and veteran trees are important habitats for bryophytes, lichens, saproxylic organisms and some bird species. Their retention is important to ensure continuity of habitats/niches and propagule sources
Woodland structure: indicators of local distinctiveness	Occurrence	No decline	Includes ancient or long-established woodlands, archaeological and geological features as well as red-data and other rare or localised species. Perrin and Daly (2010) list Pontoon Wood as possible ancient woodland
Vegetation composition: native tree cover	Percentage	No decline. Native tree cover not less than 95%	Species reported in Perrin et al. (2008)

Vegetation composition: typical species	Occurrence	A variety of typical native species present, depending on woodland type, including oak (<i>Quercus petraea</i>) and birch (<i>Betula pubescens</i>)	Species reported in Perrin et al. (2008)
Vegetation composition: negative indicator species	Occurrence	Negative indicator species, particularly non-native invasive species, absent or under control	The following are the most common invasive species in this woodland type: beech (<i>Fagus sylvatica</i>), sycamore (<i>Acer pseudoplatanus</i>), rhododendron (<i>Rhododendron ponticum</i>) and cherry laurel (<i>Prunus laurocerasus</i>)

Conservation Objectives for : River Moy SAC [002298]

91E0 Alluvial forests with *Alnus glutinosa* and *Fraxinus excelsior* (Alno-Padion, Alnion incanae, Salicion albae)

To maintain the favourable conservation condition of Alluvial forests with *Alnus glutinosa* and *Fraxinus excelsior* (Alno-Padion, Alnion incanae, Salicion albae) in River Moy SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Total extent of this habitat within the SAC is unknown and it may occur in mosaics with other woodland types. Two sites (1763, 1800) within the SAC were surveyed as part of the the National Survey of Native Woodlands (NSNW) (Perrin et al., 2008). Map 6 shows surveyed woodlands including areas classified as 91E0 (2.76ha). NB areas mapped as other wet woodland types may also correspond with this Annex I woodland type. There are also likely to be additional areas of this Annex I woodland type within the SAC
Habitat distribution	Occurrence	No decline. Woodlands surveyed as part of the NSNW are shown on map 6	The area of this habitat identified by the NSNW occurs at Prospect (site 1800) on the western shore of Lough Conn. See note on area above
Woodland size	Hectares	Area stable or increasing. Where topographically possible, "large" woods at least 25ha in size and "small" woods at least 3ha in size	The sizes of at least some of the existing woodlands need to be increased in order to reduce habitat fragmentation and benefit those species requiring 'deep' woodland conditions (Peterken, 2002). Topographical and land-ownership constraints may restrict expansion
Woodland structure: cover and height	Percentage and metres	Diverse structure with a relatively closed canopy containing mature trees; subcanopy layer with semi-mature trees and shrubs; and well-developed herb layer	Described in Perrin et al. (2008)
Woodland structure: community diversity and extent	Hectares	Maintain diversity and extent of community types	Described in Perrin et al. (2008)
Woodland structure: natural regeneration	Seedling: sapling: pole ratio	Seedlings, saplings and pole age-classes occur in adequate proportions to ensure survival of woodland canopy	Alder (<i>Alnus glutinosa</i>) and oak (<i>Quercus</i> spp.) regenerate poorly. Ash (<i>Fraxinus excelsior</i>) often regenerates in large numbers although few seedlings reach pole size
Hydrological regime: Flooding depth/height of water table	Metres	Appropriate hydrological regime necessary for maintenance of alluvial vegetation	Periodic flooding is essential to maintain alluvial woodlands along river floodplains and lakeshores
Woodland structure: dead wood	m ³ per hectare; number per hectare	At least 30m ³ /ha of fallen timber greater than 10cm diameter; 30 snags/ha; both categories should include stems greater than 40cm diameter (greater than 20cm diameter in the case of alder)	Dead wood is a valuable resource and an integral part of a healthy, functioning woodland ecosystem
Woodland structure: veteran trees	Number per hectare	No decline	Mature and veteran trees are important habitats for bryophytes, lichens, saproxylic organisms and some bird species. Their retention is important to ensure continuity of habitats/niches and propagule sources
Woodland structure: indicators of local distinctiveness	Occurrence	No decline	Includes ancient or long-established woodlands, archaeological and geological features as well as red-data and other rare or localised species

Vegetation composition: native tree cover	Percentage	No decline. Native tree cover not less than 95%	Species reported in Perrin et al. (2008)
Vegetation composition: typical species	Occurrence	A variety of typical native species present, depending on woodland type, including including alder (<i>Alnus glutinosa</i>), willows (<i>Salix</i> spp.), oak (<i>Quercus robur</i>) and ash (<i>Fraxinus excelsior</i>)	Species reported in Perrin et al. (2008)
Vegetation composition: negative indicator species	Occurrence	Negative indicator species, particularly non-native invasive species, absent or under control	The following are the most common invasive species in this woodland type: sycamore (<i>Acer pseudoplatanus</i>) and Himalayan balsam (<i>Impatiens glandulifera</i>). The NSNW notes rhododendron (<i>Rhododendron ponticum</i>) clearance in site 1800

Conservation Objectives for : River Moy SAC [002298]

1092 White-clawed Crayfish *Austropotamobius pallipes*

To maintain the favourable conservation condition of White-clawed Crayfish in River Moy SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Distribution	Occurrence	No reduction from baseline. See map 7	The general distribution of white-clawed crayfish in the SAC is that it is widespread in the upper tributaries of the River Moy and the rivers which feed Loughs Conn and Cullin. It is absent from the main River Moy. The named tributaries that it is recorded from are the following: Upstream of Lough Conn: River Deel and its tributaries of the Torean River, Rathnamagh River and Rappa Stream; Fiddaunglass; Addergoole River. Upstream of Lough Cullin: Tobergal River; Clydagh; tributaries of the Toormore and Manulla Rivers. Moy tributaries: Gweestion River; tributaries of the Pollagh, Gloré, Yellow and Geestaun Rivers; Killeen River; Spaddagh River; Sonnagh River; Owenaher River; Owengarve River
Population structure: recruitment	Occurrence of juveniles and females with eggs	Juveniles and/or females with eggs in all occupied tributaries	See Reynolds et al. (2010) for further details
Negative indicator species	Occurrence	No alien crayfish species	Alien crayfish species are identified as a major direct threat to this species and as a disease vector. See Reynolds (1998) for further details. Ireland is currently free of non-native invasive crayfish species
Disease	Occurrence	No instances of disease	Crayfish plague is identified as major threat and has occurred in Ireland even in the absence of alien vectors. See Reynolds (1998) for further details. Disease can in some circumstances be introduced through contaminated equipment and water in the absence of vector species
Water quality	EPA Q value	At least Q3-4 at all sites sampled by EPA	Target taken from Demers and Reynolds (2002). Q values based on triennial water quality surveys carried out by the EPA
Habitat quality: heterogeneity	Occurrence of positive habitat features	No decline in heterogeneity or habitat quality	Crayfish need high habitat heterogeneity. Larger crayfish must have stones to hide under, or an earthen bank in which to burrow. Hatchlings shelter in vegetation, gravel and among fine tree-roots. Smaller crayfish are typically found among weed and debris in shallow water. Larger juveniles in particular may also be found among cobbles and detritus such as leaf litter. These conditions must be available on the whole length of occupied habitat

Conservation Objectives for : River Moy SAC [002298]

1095 Sea Lamprey *Petromyzon marinus*

To maintain the favourable conservation condition of Sea Lamprey in River Moy SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Distribution: extent of anadromy	Percentage of river accessible	Greater than 75% of main stem length of rivers accessible from estuary	This SAC only covers the freshwater portion of the River Moy. The adjacent Killala Bay/Moy Estuary SAC (site code: 000485) encompasses the estuarine elements of sea lamprey habitat. Artificial barriers can block or cause difficulties to lampreys' upstream migration, thereby limiting species to lower stretches and restricting access to spawning areas (Rooney et al. 2015), however, there are no artificial barriers in the Moy catchment limiting lamprey access
Population structure of juveniles	Number of age/size groups	At least three age/size groups present	Attribute and target based on Harvey and Cowx (2003) and O'Connor (2007)
Juvenile density in fine sediment	Juveniles/m ²	Mean catchment juvenile density at least 1/m ²	Juveniles burrow in areas of fine sediment in still water. Attribute and target based on Harvey and Cowx (2003)
Extent and distribution of spawning habitat	m ² and occurrence	No decline in extent and distribution of spawning beds	Attribute and target based on spawning bed mapping by Inland Fisheries Ireland (IFI). Lampreys spawn in clean gravels
Availability of juvenile habitat	Number of positive sites in 3rd order channels (and greater), downstream of spawning areas	More than 50% of sample sites positive	Silting habitat is essential for larval lamprey and they can be severely impacted by sediment removal. Recovery can be rapid and newly-created habitat can be rapidly colonised (King et al., 2015). However, it is vital that such sedimenting habitats are retained. Occupancy in excess of 50% of sites would be 'reasonable' for the Irish catchments examined to date. (King and Linnane, 2004; King et al., unpublished data)

Conservation Objectives for : River Moy SAC [002298]

1096 Brook Lamprey *Lampetra planeri*

To maintain the favourable conservation condition of Brook Lamprey in River Moy SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Distribution	Percentage of river accessible	Access to all watercourses down to first order streams	Artificial barriers can block lampreys' migration both up- and downstream, thereby possibly limiting species to specific stretches, restricting access to spawning areas and creating genetically isolated populations (Espanhol et al., 2007). However, there are no artificial barriers in the Moy catchment limiting lamprey access
Population structure of juveniles	Number of age/size groups	At least three age/size groups of brook/river lamprey present	Attribute and target based on data from Harvey and Cowx (2003). It is impossible to distinguish between brook and river lamprey juveniles in the field (Gardiner, 2003), hence they are considered together in this target
Juvenile density in fine sediment	Juveniles/m ²	Mean catchment juvenile density of brook/river lamprey at least 2/m ²	Juveniles burrow in areas of fine sediment in still water. Attribute and target based on data from Harvey and Cowx (2003) who state 10/m ² in optimal conditions and more than 2/m ² on a catchment basis
Extent and distribution of spawning habitat	m ² and occurrence	No decline in extent and distribution of spawning beds	Attribute and target based on spawning bed mapping by Inland Fisheries Ireland (IFI). Lampreys spawn in clean gravels
Availability of juvenile habitat	Number of positive sites in 2nd order channels (and greater), downstream of spawning areas	More than 50% of sample sites positive	Silting habitat is essential for larval lamprey and they can be severely impacted by sediment removal. Recovery can be rapid and newly-created habitat can be rapidly colonised (King et al., 2015). However, it is vital that such sedimenting habitats are retained. Occupancy in excess of 50% of sites would be 'reasonable' for the Irish catchments examined to date. (King and Linnane, 2004; King et al., unpublished data)

Conservation Objectives for : River Moy SAC [002298]

1106 Salmon *Salmo salar*

To maintain the favourable conservation condition of Salmon in River Moy SAC, which is defined by the following list of attributes and targets:


Attribute	Measure	Target	Notes
Distribution: extent of anadromy	Percentage of river accessible	100% of river channels down to second order accessible from estuary	Artificial barriers block salmon's upstream migration, thereby limiting species to lower stretches and restricting access to spawning areas. There are no artificial barriers on the Moy catchment limiting salmon access
Adult spawning fish	Number	Conservation Limit (CL) for each system consistently exceeded	A conservation limit is defined by the North Atlantic Salmon Conservation Organisation (NASCO) as "the spawning stock level that produces long-term average maximum sustainable yield as derived from the adult to adult stock and recruitment relationship". The target is based on the Standing Scientific Committee of the National Salmon Commission's annual model output of CL attainment levels. See SSC (2016). Stock estimates are either derived from direct counts of adults (rod catch, fish counter) or indirectly by fry abundance counts. For the 2016 SSC advice, the Moy is currently exceeding its CL by 19,012 salmon
Salmon fry abundance	Number of fry/5 minutes electrofishing	Maintain or exceed 0+ fry mean catchment-wide abundance threshold value. Currently set at 17 salmon fry/5 minutes sampling	Target is threshold value for rivers currently exceeding their conservation limit (CL)
Out-migrating smolt abundance	Number	No significant decline	Smolt abundance can be negatively affected by a number of impacts such as estuarine pollution, predation and sea lice (<i>Lepeophtheirus salmonis</i>)
Number and distribution of redds	Number and occurrence	No decline in number and distribution of spawning redds due to anthropogenic causes	Salmon spawn in clean gravels. There are no artificial barriers preventing salmon from accessing suitable spawning habitat in this SAC
Water quality	EPA Q value	At least Q4 at all sites sampled by EPA	Q values based on triennial water quality surveys carried out by the Environmental Protection Agency (EPA)

Conservation Objectives for : River Moy SAC [002298]

1355 Otter *Lutra lutra*

To maintain the favourable conservation condition of Otter in River Moy SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Distribution	Percentage positive survey sites	No significant decline	Measure based on standard otter survey technique. FCS target, based on 1980/81 survey findings, is 88% in SACs. Current range is estimated at 93.6% (Reid et al., 2013)
Extent of terrestrial habitat	Hectares	No significant decline. Area mapped and calculated as 1068.8ha	No field survey. Areas mapped to include 10m terrestrial buffer along lake shorelines and along river banks identified as critical for otters (NPWS, 2007)
Extent of freshwater (river) habitat	Kilometres	No significant decline. Length mapped and calculated as 479.4km	No field survey. River length calculated on the basis that otters will utilise freshwater habitats from estuary to headwaters (Chapman and Chapman, 1982)
Extent of freshwater (lake) habitat	Hectares	No significant decline. Area mapped and calculated as 1248.2ha	No field survey. Area mapped based on evidence that otters tend to forage within 80m of the shoreline (NPWS, 2007)
Couching sites and holts	Number	No significant decline	Otters need lying up areas throughout their territory where they are secure from disturbance (Kruuk, 2006; Kruuk and Moorhouse, 1991)
Fish biomass available	Kilograms	No significant decline	Broad diet that varies locally and seasonally, but dominated by fish, in particular salmonids, eels and sticklebacks in freshwater (Bailey and Rochford, 2006; Reid et al., 2013)
Barriers to connectivity	Number	No significant increase. For guidance, see map 8	Otters will regularly commute across stretches of open water up to 500m e.g. between the mainland and an island; between two islands; across an estuary (De Jongh and O'Neill, 2010). It is important that such commuting routes are not obstructed



Legend

River Moy SAC 002298

An tAonán Ealaíon, Oidhreacht, Gnóthaí Réigiúnacha, Tuaithe agus Gaeltachta
 Department of Arts, Heritage, Regional, Rural and Gaeltacht Affairs

MAP 1:
RIVER MOY SAC
CONSERVATION OBJECTIVES
SAC DESIGNATION

Map to be read in conjunction with the NPWS Conservation Objectives Document.

SITE CODE:
SAC 002298; version 3.01. CO. MAYO,
CO. SLIGO & CO. ROSCOMMON


0 2 4 6 8 10 km

The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
 Ordnance Survey of Ireland Licence No EN 0059216. © Ordnance Survey of Ireland Government of Ireland

Níl sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbheithníthe a déanamh ar theorainneacha na gceantar comharthaíthe. Suirbhéarachta Ordonáis na hÉireann Ceadúnas


N

Map Version 1
Date: July 2016



Legend

- River Moy SAC 002298
- Bellacorick Bog Complex SAC 001922
- Killala Bay/Moy Estuary SAC 000458
- Lough Hoe Bog SAC 000633
- Ox Mountains Bogs SAC 002006
- Killala Bay/Moy Estuary SPA 004036
- Lough Conn and Lough Cullin SPA 004228

 An Roinn Ealaíon, Oidhreacht, Gnóthaí Réigiúnacha, Tuaithe agus Gaeltachta
Department of Arts, Heritage, Regional, Rural and Gaeltacht Affairs

**MAP 2:
RIVER MOY SAC
CONSERVATION OBJECTIVES
ADJACENT / ADJOINING & OVERLAPPING
DESIGNATIONS**

Map to be read in conjunction with the NPWS Conservation Objectives Document.


**SITE CODE:
SAC 002298; version 3.01. CO. MAYO,
CO. SLIGO & CO. ROSCOMMON**




0 2.5 5 7.5 10 12.5 km

The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
Ordnance Survey of Ireland Licence No EN 0059216. © Ordnance Survey of Ireland Government of Ireland


Nil sna teorainneacha ar na léarscálanna ach nod garshuíomhach ginearálta. Féadfar athbheirnithe a déanamh ar theorainneacha na gceantar comharthaithe. Suirbhéarachta Ordonáis na hÉireann Ceadúnas


**Map Version 1
Date: July 2016**











Cloongoonagh Bog








Legend


-  High Bog Boundary
 -  Drainage Patterns
 -  Contours
- Elevation (m)**
-  56.65 m
 -  39.86 m

Derrynabrock / Kilgarriff / Tawnabeg / Gowlaun Bogs





Legend



-  High Bog Boundary
 -  Drainage Patterns
 -  Contours
- Elevation (m)**
-  91.94 m
 -  71.07 m




Legend

-  River Moy SAC 002298








Legend

- River Moy SAC 002298
- 1092 White-clawed Crayfish




An Roinn Ealaíon, Oidhreachta,
Gnóthaí Réigiúnacha, Tuaithe agus Gaeltachta
Department of Arts, Heritage,
Regional, Rural and Gaeltacht Affairs

**MAP 7:
RIVER MOY SAC
CONSERVATION OBJECTIVES
WHITE-CLAWED CRAYFISH**

Map to be read in conjunction with the NPWS Conservation Objectives Document.

**SITE CODE:
SAC 002298; version 3.01. CO. MAYO,
CO. SLIGO & CO. ROSCOMMON**





The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
Ordnance Survey of Ireland Licence No EN 0059216. © Ordnance Survey of Ireland Government of Ireland


Níl sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbhreithnithe a déanamh ar theorainneacha na gceantar comharthaíthe. Suirbhéarachta Ordonáis na hÉireann Ceadúnas



Map Version 1
Date: July 2016

Legend

-  River Moy SAC 002298
-  1355 Otter - *Lutra lutra* commuting 250m buffer



National Parks and Wildlife Service

Conservation Objectives Series

Kildun Souterrain SAC 002320



An Roinn Cultúir,
Oidhreacht agus Gaeltachta
Department of Culture,
Heritage and the Gaeltacht

**National Parks and Wildlife Service,
Department of Culture, Heritage and the Gaeltacht,
90 King Street North, Dublin 7, D07 N7CV, Ireland.**

**Web: www.npws.ie
E-mail: nature.conservation@chg.gov.ie**

Citation:

**NPWS (2018) Conservation Objectives: Kildun Souterrain SAC 002320. Version
1. National Parks and Wildlife Service, Department of Culture, Heritage and the
Gaeltacht.**

**Series Editor: Rebecca Jeffrey
ISSN 2009-4086**

Introduction

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

** indicates a priority habitat under the Habitats Directive*

002320 Kildun Souterrain SAC

1303 Lesser Horseshoe Bat *Rhinolophus hipposideros*

Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

NPWS Documents

Year : 2018
Title : Conservation objectives supporting document – lesser horseshoe bat (*Rhinolophus hipposideros*)
Author : NPWS
Series : Conservation objectives supporting document

Other References

Year : 2007
Title : Protecting and managing underground sites for bats
Author : Mitchell-Jones, A.J.; Bihari, Z.; Masing, M.; Rodrigues, L.
Series : EUROBATS Publication Series No. 2

Year : 2008
Title : The lesser horseshoe bat conservation handbook
Author : Schofield, H.W.
Series : The Vincent Wildlife Trust

Year : 2009
Title : Importance of night roosts for bat conservation: roosting behaviour of the lesser horseshoe bat *Rhinolophus hipposideros*
Author : Knight, T.; Jones, G.
Series : Endangered Species Research, 8: 79-86

Spatial data sources

Year : 2018
Title : NPWS lesser horseshoe bat database
GIS Operations : Roost identified, clipped to SAC boundary. Expert opinion used as necessary to resolve any issues arising
Used For : 1303 (map 2)

Year : 2007
Title : Forest Inventory and Planning System (FIPS)
GIS Operations : Dataset clipped to 2.5km buffer centred on roost location
Used For : 1303 (map 2)

Conservation Objectives for : Kildun Souterrain SAC [002320]

1303 Lesser Horseshoe Bat *Rhinolophus hipposideros*

To restore the favourable conservation condition of Lesser Horseshoe Bat in Kildun Souterrain SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Population per roost	Number	Minimum number of 50 bats for the winter roost (roost id. 676 in NPWS database). See map 2	A figure of 100 bats for summer roosts and 50 bats for winter roosts was set as a minimum qualifying standard (MQS) when SACs were being selected for lesser horseshoe bat (<i>Rhinolophus hipposideros</i>). NPWS conduct annual counts at each qualifying roost. Qualified means from the 2013-2017 winter data have been calculated whereby the year with the highest maximum count and the year with the lowest maximum count were removed and the mean of the remaining years was calculated. This mean is usually set as the target figure for the roost. However, in the case of the winter roost (roost id. 676 in NPWS database) in Kildun Souterrain SAC, where a mean of 15 bats was recorded (2013-2017), the target is instead set at the MQS of 50 bats. See the conservation objectives supporting document for lesser horseshoe bat (NPWS, 2018) for further information on all attributes and targets
Winter roosts	Condition	No decline	Kildun Souterrain SAC has been selected for lesser horseshoe bat because of the presence of one internationally important winter roost (roost id. 676 in NPWS database). Damage or disturbance to the roost or to the habitat immediately surrounding it will lead to a decline in its condition (Mitchell-Jones et al., 2007)
Auxiliary roosts	Number and condition	No decline	Lesser horseshoe bat populations will use a variety of roosts during the year besides the main summer maternity and winter hibernation roosts. Such additional roosts within the SAC may be important as night roosts, satellite roosts, etc. Night roosts are also considered an integral part of core foraging areas and require protection (Knight and Jones, 2009). In addition, in response to weather conditions for example, bats may use different seasonal roosts from year to year; this is particularly noticeable in winter. A database of all known lesser horseshoe bat roosts is available on the National Biodiversity Data Centre website. NB further unrecorded roosts may also be present within this SAC
Extent of potential foraging habitat	Hectares	No significant decline within 2.5km of qualifying roost	Lesser horseshoe bats normally forage in woodlands/scrub within 2.5km of their roosts (Schofield, 2008). See map 2 which shows a 2.5km zone around the above roost and identifies potential foraging grounds
Linear features	Kilometres	No significant loss within 2.5km of qualifying roost. See map 2	This species follows commuting routes from its roost to its foraging grounds. Lesser horseshoe bats will not cross open ground. Consequently, linear features such as hedgerows, treelines and stone walls provide vital connectivity for this species within 2.5km around each roost (Schofield, 2008)
Light pollution	Lux	No significant increase in artificial light intensity adjacent to named roost or along commuting routes within 2.5km of the roost. See map 2	Lesser horseshoe bats are very sensitive to light pollution and will avoid brightly lit areas. Inappropriate lighting around roosts may cause abandonment; lighting along commuting routes may cause preferred foraging areas to be abandoned, thus increasing energetic costs for bats (Schofield, 2008)



Legend

 Kildun Souterrain SAC 002320



An Roinn Cultúir,
Oidhreacht agus Gaeltachta
Department of Culture,
Heritage and the Gaeltacht

**MAP 1:
KILDUN SOUTERRAIN SAC
CONSERVATION OBJECTIVES
SAC DESIGNATION**

Map to be read in conjunction with the NPWS Conservation Objectives Document.

**SITE CODE:
SAC 002320; version 3. CO. MAYO**


0 20 40 60 80 Meters

The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
Ordnance Survey of Ireland Licence No EN 0059216. © Ordnance Survey of Ireland Government of Ireland

Níl sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbhreithnithe a déanamh ar theorainneacha na gceantar comharthaite. Suirbhéarachta Ordonáis na hÉireann Ceadúnas Uimh EN 0059216. © Suirbhéarachta Ordonáis na hÉireann Rialtas na hÉireann



**Map Version 1
Date: June 2018**



Legend

- OSi Discovery Series County Boundary
- Kildun Souterrain SAC 002320
- 1303 Lesser Horseshoe Bat *Rhinolophus hipposideros***
- Roost Location
- Roost ID 676 Foraging Range
- Potential Foraging Grounds

National Parks and Wildlife Service

Conservation Objectives Series

Drumalough Bog SAC 002338



An Roinn Ealaíon, Oidhreachta,
Gnóthaí Réigiúnacha, Tuaithe agus Gaeltachta

Department of Arts, Heritage,
Regional, Rural and Gaeltacht Affairs



**National Parks and Wildlife Service,
Department of Arts, Heritage, Regional, Rural and Gaeltacht Affairs,
7 Ely Place, Dublin 2, Ireland.
Web: www.npws.ie
E-mail: nature.conservation@ahg.gov.ie**

Citation:

**NPWS (2016) Conservation Objectives: Drumalough Bog SAC 002338. Version 1.
National Parks and Wildlife Service, Department of Arts, Heritage, Regional,
Rural and Gaeltacht Affairs.**

**Series Editor: Rebecca Jeffrey
ISSN 2009-4086**

Introduction

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

** indicates a priority habitat under the Habitats Directive*

002338	Drumalough Bog SAC
7110	Active raised bogsE
7120	Degraded raised bogs still capable of natural regeneration
7150	Depressions on peat substrates of the Rhynchosporion

Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

NPWS Documents

Year :	2014
Title :	National raised bog SAC management plan
Author :	Department of Arts, Heritage and the Gaeltacht
Series :	Draft for consultation. 15 January 2014
<hr/>	
Year :	2016
Title :	Drumalough Bog SAC (site code: 2338) Conservation objectives supporting document- raised bog habitats V1
Author :	NPWS
Series :	Conservation objectives supporting document

Other References

Year :	2011
Title :	Review and revision of empirical critical loads and dose-response relationships. Proceedings of an expert workshop, Noordwijkerhout, 23-25 June 2010
Author :	Bobbink, R.; Hettelingh, J.P.
Series :	RIVM report 680359002, Coordination Centre for Effects, National Institute for Public Health and the Environment (RIVM)
<hr/>	
Year :	2014
Title :	Nitrogen deposition and exceedance of critical loads for nutrient nitrogen in Irish grasslands
Author :	Henry, J.; Aherne, J.
Series :	Science of the Total Environment 470–471: 216–223

Spatial data sources

Year :	2014
Title :	Scientific Basis for Raised Bog Conservation in Ireland
GIS Operations :	RBSB13_SACs_ARB_DRB dataset, RBSB13_SACs_2012_HB dataset, RBSB13_SACs_DrainagePatterns_5k dataset and RBSB13_SAC_LIDAR_DTMs dataset clipped to SAC boundary. Expert opinion used as necessary to resolve any issues arising
Used For :	Potential 7110; digital elevation model; drainage patterns (maps 2 and 4)
<hr/>	
Year :	Digitised 2003
Title :	Raised Bog Restoration Project 1999
GIS Operations :	Ecotope dataset clipped to SAC boundary. Appropriate ecotopes selected and exported to new dataset. Expert opinion used as necessary to resolve any issues arising
Used For :	7110 ecotopes (map 3)
<hr/>	

Conservation Objectives for : Drumalough Bog SAC [002338]

7110 Active raised bogs

To restore the favourable conservation condition of Active raised bogs in Drumalough Bog SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Restore area of active raised bog to 45.2ha, subject to natural processes	Active Raised Bog (ARB) habitat on Drumalough Bog West has been estimated as 5.1ha. The area of ARB on Drumalough Bog East is unknown. Area of Degraded Raised Bog (DRB) on the High Bog (HB) has been modelled as 46.6ha. See map 2. However, it is estimated that only 39.0ha of this area is potentially restorable to ARB by drain blocking. The total potential ARB on the HB is therefore estimated to be 44.1ha. Eco-hydrological assessments of the cutover estimates that an additional 1.1ha of bog forming habitats could be restored. The long term target for ARB is therefore 45.2ha. See raised bog supporting document for further details on this and following attributes
Habitat distribution	Occurrence	Restore the distribution and variability of active raised bog across the SAC. See map 3 for distribution mapped in 2003	ARB has been mapped on Drumalough Bog West. DRB occurs on both Drumalough Bog East and Drumalough Bog West, which will require restoration measures. There is also potential for ARB restoration on cutover areas of the bog (see area target above)
High bog area	Hectares	No decline in extent of high bog necessary to support the development and maintenance of active raised bog. See map 2	The area of high bog within Drumalough Bog SAC in 2012 (latest figure available) was 159.0ha (DAHG 2014)
Hydrological regime: water levels	Centimetres	Restore appropriate water levels throughout the site	For ARB, mean water level needs to be near or above the surface of the bog lawns for most of the year. Seasonal fluctuations should not exceed 20cm, and should only be 10cm below the surface, except for very short periods of time. Open water is often characteristic of soak systems
Hydrological regime: flow patterns	Flow direction; slope	Restore, where possible, appropriate high bog topography, flow directions and slopes. See map 4 for current situation	ARB depends on mean water levels being near or above the surface of bog lawns for most of the year. Long and gentle slopes are the most favourable to achieve these conditions. Changes to flow directions due to subsidence of bogs can radically change water regimes and cause drying out of high quality ARB areas and soak systems
Transitional areas between high bog and adjacent mineral soils (including cutover areas)	Hectares; distribution	Restore adequate transitional areas to support/protect active raised bog and the services it provides	ARB is threatened due to effects of past drainage and peat-cutting around the margins of Drumalough Bog. Eco-hydrological assessments have evaluated the potential for ARB restoration on cutover areas (see note for habitat area attribute above)
Vegetation quality: central ecotope, active flush, soaks, bog woodland	Hectares	Restore 22.6ha of central ecotope/active flush/soaks/bog woodland as appropriate	At least 50% of ARB habitat should be high quality (i.e. central ecotope, active flush, soaks, bog woodland). Target area of active raised bog for the SAC has been set at 45.2ha (see area target above)
Vegetation quality: microtopographical features	Hectares	Restore adequate cover of high quality microtopographical features	High quality microtopography (hummocks, hollows and pools) is well developed in the western part of Drumalough Bog
Vegetation quality: bog moss (Sphagnum) species	Percentage cover	Restore adequate cover of bog moss (<i>Sphagnum</i>) species to ensure peat-forming capacity	<i>Sphagnum</i> cover varies naturally across Ireland with relatively high cover in the east to lower cover in the west. Hummock forming species such as <i>Sphagnum austinii</i> are particularly good peat formers. <i>Sphagnum</i> cover and distribution also varies naturally across a site

Typical ARB species: flora	Occurrence	Restore, where appropriate, typical active raised bog flora	Typical flora species include widespread species, as well as those with more restricted distributions but typical of the habitat's subtypes or geographical range
Typical ARB species: fauna	Occurrence	Restore, where appropriate, typical active raised bog fauna	Typical fauna species include widespread species, as well as those with more restricted distributions but typical of the habitat's subtypes or geographical range
Elements of local distinctiveness	Occurrence	Maintain features of local distinctiveness, subject to natural processes	These include features of geological, topographical, archaeological and hydrological interest as well as noteworthy species of flora and fauna
Negative physical indicators	Percentage cover	Negative physical features absent or insignificant	Negative physical indicators include: bare peat, algae dominated pools and hollows, marginal cracks, tear patterns, subsidence features such as dry mineral mounds/ridges emerging or expanding and evidence of burning
Vegetation composition: native negative indicator species	Percentage cover	Native negative indicator species at insignificant levels	Disturbance indicators include species indicative of conditions drying out such as abundant bog asphodel (<i>Narthecium ossifragum</i>), deergrass (<i>Trichophorum germanicum</i>) and harestail cotton-grass (<i>Eriophorum vaginatum</i>) forming tussocks; abundant magellanic bog-moss (<i>Sphagnum magellanicum</i>) in pools previously dominated by <i>Sphagnum</i> species typical of very wet conditions (e.g. feathery bog-moss (<i>S. cuspidatum</i>)); and indicators of frequent burning events such as abundant <i>Cladonia floerkeana</i> and high cover of carnation sedge (<i>Carex panicea</i>) (particularly in true midlands raised bogs)
Vegetation composition: non-native invasive species	Percentage cover	Non-native invasive species at insignificant levels and not more than 1% cover	Most common non-native invasive species include lodgepole pine (<i>Pinus contorta</i>), rhododendron (<i>Rhododendron ponticum</i>) and pitcherplant (<i>Sarracenia purpurea</i>)
Air quality: nitrogen deposition	kg N/ha/year	Air quality surrounding bog close to natural reference conditions. The total N deposition should not exceed 5kg N/ha/yr	Change in air quality can result from fertiliser drift; adjacent quarry activities; or other atmospheric inputs. The critical load range for ombrotrophic bogs has been set as between 5 and 10kg N/ha/yr (Bobbink and Hettelingh, 2011). The latest N deposition figures for the area around Drumalough Bog suggests that the current level is approximately 9.9kg N/ha/yr (Henry and Aherne, 2014)
Water quality	Hydrochemical measures	Water quality on the high bog and in transitional areas close to natural reference conditions	Water chemistry within raised bogs is influenced by atmospheric inputs (rainwater). However, within soak systems, water chemistry is influenced by other inputs such as focused flow or interaction with underlying substrates. Water chemistry in areas surrounding the high bog varies due to influences of different water types (bog water, regional groundwater and run-off from surrounding mineral lands)

Conservation Objectives for : Drumalough Bog SAC [002338]

7120 Degraded raised bogs still capable of natural regeneration

The long-term aim for Degraded raised bogs still capable of natural regeneration is that its peat-forming capability is re-established; therefore, the conservation objective for this habitat is inherently linked to that of Active raised bogs (7110) and a separate conservation objective has not been set in Drumalough Bog SAC



Attribute	Measure	Target	Notes
-----------	---------	--------	-------

Conservation Objectives for : Drumalough Bog SAC [002338]


7150 Depressions on peat substrates of the Rhynchosporion



Depressions on peat substrates of the Rhynchosporion is an integral part of good quality Active raised bogs (7110) and thus a separate conservation objective has not been set for the habitat in Drumalough Bog SAC

Attribute	Measure	Target	Notes
-----------	---------	--------	-------








Legend

 Drumalough Bog SAC 002338








Legend

-  Drumalough Bog SAC 002338
-  High Bog Boundary
-  Potential 7110 *Active Raised Bogs



Legend



-  Drumalough Bog SAC 002338
-  High Bog Boundary
- Active Raised Bog Ecotopes**
-  Sub-central Ecotope



Legend

- Drumalough Bog SAC 002338
- High Bog Boundary
- Drainage Patterns
- Contours

Elevation



National Parks and Wildlife Service

Conservation Objectives Series

Tullaghanrock Bog SAC 002354



*An Roinn
Ealaíon, Oidhreachta agus Gaeltachta*

*Department of
Arts, Heritage and the Gaeltacht*



**National Parks and Wildlife Service,
Department of Arts, Heritage and the Gaeltacht,**

7 Ely Place, Dublin 2, Ireland.

Web: www.npws.ie

E-mail: nature.conservation@ahg.gov.ie

Citation:

**NPWS (201) Conservation Objectives: Tullaghanrock Bog SAC 002354. Version
1. National Parks and Wildlife Service, Department of Arts, Heritage and the
Gaeltacht.**

Series Editor: Rebecca Jeffrey

ISSN 2009-4086

Introduction

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

** indicates a priority habitat under the Habitats Directive*

002354	Tullaghanrock Bog SAC
7110	Active raised bogsE
7120	Degraded raised bogs still capable of natural regeneration
7150	Depressions on peat substrates of the Rhynchosporion

Please note that this SAC adjoins Callow Bog SAC (000595). See map 2. The conservation objectives for this site should be used in conjunction with those for the adjacent site as appropriate.

Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

NPWS Documents

Year :	2000
Title :	Raised bog restoration project. A continuation of the investigation into the conservation and restoration of selected raised bog sites in Ireland
Author :	Derwin, J.; Mac Gowan, F.
Series :	Unpublished report to Duchas, the Heritage Service
Year :	2014
Title :	National raised bog SAC management plan
Author :	Department of Arts, Heritage and the Gaeltacht
Series :	Draft for consultation. 15 January 2014
Year :	2015
Title :	Tullaghanrock SAC (site code: 2354) Conservation objectives supporting document- raised bog habitats V1
Author :	NPWS
Series :	Conservation objectives supporting document

Other References

Year :	2011
Title :	Review and revision of empirical critical loads and dose-response relationships. Proceedings of an expert workshop, Noordwijkerhout, 23-25 June 2010
Author :	Bobbink, R.; Hettelingh, J.P.
Series :	RIVM report 680359002, Coordination Centre for Effects, National Institute for Public Health and the Environment (RIVM)
Year :	2014
Title :	Nitrogen deposition and exceedance of critical loads for nutrient nitrogen in Irish grasslands
Author :	Henry, J.; Aherne, J.
Series :	Science of the Total Environment 470–471: 216–223

Spatial data sources

Year :	2014
Title :	Scientific Basis for Raised Bog Conservation in Ireland
GIS Operations :	RBSB13_SACs_ARB_DRB dataset, RBSB13_SACs_2012_HB dataset, RBSB13_SACs_DrainagePatterns_5k dataset and RBSB13_SAC_LIDAR_DTMs dataset clipped to SAC boundary. Expert opinion used as necessary to resolve any issues arising
Used For :	potential 7110; digital elevation model; drainage patterns (maps 3 and 5)
<hr/>	
Year :	Digitised 2006
Title :	Raised Bog Restoration Project 1999
GIS Operations :	Ecotope dataset clipped to SAC boundary. Appropriate ecotopes selected and exported to new dataset. Expert opinion used as necessary to resolve any issues arising
Used For :	7110 ecotopes (map 4)
<hr/>	

Conservation Objectives for : Tullaghanrock Bog SAC [002354]

7110 Active raised bogs

To restore the favourable conservation condition of Active raised bogs in Tullaghanrock Bog SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Restore area of active raised bog to 15.2ha, subject to natural processes	Active Raised Bog (ARB) habitat was mapped at 11.0ha (Derwin and MacGowan 2000). Area of Degraded Raised Bog (DRB) on the High Bog (HB) has been modelled as 3.9ha. See map 3. It is considered that all of this area is potentially restorable to ARB by drain blocking. The total potential ARB on the HB is therefore estimated to be 14.9ha. Eco-hydrological assessments of the cutover estimates that an additional 0.3ha of bog forming habitats could be restored. The long term target for ARB is therefore 15.2ha. See raised bog supporting document for further details on this and following attributes
Habitat distribution	Occurrence	Restore the distribution and variability of active raised bog across the SAC. See map 4 for distribution in 1999	ARB habitat at Tullaghanrock Bog is central and sub-central ecotopes only. There is also potential for ARB restoration on cutover areas of the bog (see area target above)
High bog area	Hectares	No decline in extent of high bog necessary to support the development and maintenance of active raised bog. See map 3	The area of high bog within Tullaghanrock Bog SAC in 2012 (latest figure available) was 62.8ha (DAHG 2014)
Hydrological regime: water levels	Centimetres	Restore appropriate water levels throughout the site	For ARB, mean water level needs to be near or above the surface of the bog lawns for most of the year. Seasonal fluctuations should not exceed 20cm, and should only be 10cm below the surface, except for very short periods of time
Hydrological regime: flow patterns	Flow direction; slope	Restore, where possible, appropriate high bog topography, flow directions and slopes. See map 5 for current situation	ARB depends on mean water levels being near or above the surface of bog lawns for most of the year. Long and gentle slopes are the most favourable to achieve these conditions. Changes to flow directions due to subsidence of bogs can radically change water regimes and cause drying out of high quality ARB
Transitional areas between high bog and adjacent mineral soils (including cutover areas)	Hectares; distribution	Restore adequate transitional areas to support / protect active raised bog and the services it provides	To the south and east there is a semi-natural margin between the high bog and the River Lung. Eco-hydrological assessments have evaluated the potential for ARB restoration on cutover areas (see note for habitat area attribute above)
Vegetation quality: central ecotope, active flush, soaks, bog woodland	Hectares	Restore 7.6ha of central ecotope/active flush/soaks/bog woodland as appropriate	At least 50% of ARB habitat should be high quality (i.e. central ecotope, active flush, soaks, bog woodland). Target area of active raised bog for the site has been set at 15.2ha (see area target above)
Vegetation quality: microtopographical features	Hectares	Restore adequate cover of high quality microtopographical features	Hummock and hollow microtopography with frequent pools is well developed on Tullaghanrock Bog
Vegetation quality: bog moss (<i>Sphagnum</i>) species	Percentage cover	Restore adequate cover of bog moss (<i>Sphagnum</i>) species to ensure peat-forming capacity	<i>Sphagnum</i> cover varies naturally across Ireland with relatively high cover in the east to lower cover in the west. Hummock forming species such as <i>Sphagnum austini</i> are particularly good peat formers. <i>Sphagnum</i> cover and distribution also varies naturally across a site
Typical ARB species: flora	Occurrence	Restore, where appropriate, typical active raised bog flora	Typical flora species include widespread species, as well as those with more restricted distributions but typical of the habitat's subtypes or geographical range

Typical ARB species: fauna	Occurrence	Restore, where appropriate, typical active raised bog fauna	Typical fauna species include widespread species, as well as those with more restricted distributions but typical of the habitat's subtypes or geographical range
Elements of local distinctiveness	Occurrence	Maintain features of local distinctiveness, subject to natural processes	These include features of geological, topographical, archaeological and hydrological interest as well as noteworthy species of flora and fauna
Negative physical indicators	Percentage cover	Negative physical features absent or insignificant	Negative physical indicators include: bare peat, algae dominated pools and hollows, marginal cracks, tear patterns, subsidence features such as dry mineral mounds /ridges emerging or expanding and evidence of burning
Vegetation composition: native negative indicator species	Percentage cover	Native negative indicator species at insignificant levels	Native negative indicator species that suggest drying out include abundant bog asphodel (<i>Narthecium ossifragum</i>), deergrass (<i>Trichophorum germanicum</i>) and haretail cotton-grass (<i>Eriophorum vaginatum</i>) forming tussocks; abundant magellanic bog-moss (<i>Sphagnum magellanicum</i>) in pools previously dominated by <i>Sphagnum</i> species typical of very wet conditions (e.g. feathery bog-moss (<i>S. cuspidatum</i>)). Indicators of frequent burning events include abundant <i>Cladonia floerkeana</i> and high cover of carnation sedge (<i>Carex panicea</i>) (particularly in true midlands raised bogs)
Vegetation composition: non-native invasive species	Percentage cover	Non-native invasive species at insignificant levels and not more than 1% cover	Most common non-native invasive species on raised bogs include lodgepole pine (<i>Pinus contorta</i>), rhododendron (<i>Rhododendron ponticum</i>), and pitcherplant (<i>Sarracenia purpurea</i>)
Air quality: nitrogen deposition	kg N/ha/year	Air quality surrounding bog close to natural reference conditions. The total N deposition should not exceed 5kg N/ha/yr	Change in air quality can result from fertiliser drift; adjacent quarry activities; or other atmospheric inputs. The critical load range for ombrotrophic bogs has been set as between 5 and 10kg N/ha/yr (Bobbink and Hettelingh, 2011). The latest N deposition figures for the area around Tullaghanrock Bog suggests that the current level is approximately 8.9kg N/ha/yr (Henry and Aherne, 2014)
Water quality	Hydrochemical measure	Water quality on the high bog and transitional areas close to natural reference conditions	Water chemistry within raised bogs is influenced by atmospheric inputs (rainwater). However, within soak systems, water chemistry is influenced by other inputs such as focused flow or interaction with underlying substrates. Water chemistry in marginal areas and lagg zone surrounding the high bog varies due to influences of different water types (bog water, regional groundwater, and runoff from surrounding mineral lands)

Conservation Objectives for : Tullaghanrock Bog SAC [002354]

7120 Degraded raised bogs still capable of natural regeneration

The long-term aim for Degraded raised bogs still capable of natural regeneration is that its peat-forming capability is re-established; therefore, the conservation objective for this habitat is inherently linked to that of Active raised bogs (7110) and a separate conservation objective has not been set in Tullaghanrock Bog SAC


Attribute	Measure	Target	Notes
-----------	---------	--------	-------


Conservation Objectives for : Tullaghanrock Bog SAC [002354]


7150 Depressions on peat substrates of the Rhynchosporion

Depressions on peat substrates of the Rhynchosporion is an integral part of good quality Active raised bogs (7110) and thus a separate conservation objective has not been set for the habitat in Tullaghanrock Bog SAC

Attribute	Measure	Target	Notes
-----------	---------	--------	-------



Legend
 Tullaghanrock Bog SAC 002354

 *An Roinn Ealaíon, Oidhreachta agus Gaeltachta*
 Department of Arts, Heritage and the Gaeltacht

MAP 1:
TULLAGHANROCK BOG SAC
CONSERVATION OBJECTIVES
SAC DESIGNATION

Map to be read in conjunction with the NPWS Conservation Objectives Document.


SITE CODE:
SAC 002354; version 3. Co. Roscommon

0 50 100 150 200 250 m




The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
 Ordnance Survey of Ireland Licence No EN 0059214. © Ordnance Survey of Ireland Government of Ireland.


Níl sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbheithníthe a déanamh ar theorainneacha na gceantar comharthaíthe. Suirbhéarachta Ordonáis na hÉireann Ceadúnas Uimh EN 0059214. © Suirbhéarachta Ordonáis na hÉireann Rialtas na hÉireann.


Map Version 1
Date: Dec 2015



Legend


-  Tullaghanrock Bog SAC 002354
-  Callow Bog SAC 000595
-  OSi Discovery Series County Boundary



**MAP 2:
TULLAGHANROCK BOG SAC
CONSERVATION OBJECTIVES
ADJOINING / OVERLAPPING
DESIGNATIONS**

Map to be read in conjunction with the NPWS Conservation Objectives Document.

**SITE CODE:
SAC 002354; version 3,
SAC 000595; version 3. Co. Roscommon**




The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
Ordnance Survey of Ireland Licence No EN 0059214. © Ordnance Survey of Ireland Government of Ireland.

Níl sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbheithníthe a déanamh ar theorainneacha na gceantar comharthaíthe. Suirbhéarachta Ordonáis na hÉireann Ceadúnas Uimh EN 0059214. © Suirbhéarachta Ordonáis na hÉireann Rialtas na hÉireann.




**Map Version 1
Date: Dec 2015**




Legend

- Tullaghanrock Bog SAC 002354
- High Bog Boundary
- Potential 7110 *Active Raised Bogs



Legend


- Tullaghanrock Bog SAC 002354
- Active Raised Bogs Ecotopes**
- Central ecotope
- Sub-central ecotope



Legend

- Tullaghanrock Bog SAC 002354
- High Bog Boundary
- Drainage Patterns
- Contours

Elevation



National Parks and Wildlife Service

Conservation Objectives Series

West Connacht Coast SAC 002998



An Roinn
Ealaíon, Oidhreachta agus Gaeltachta

Department of
Arts, Heritage and the Gaeltacht



**National Parks and Wildlife Service,
Department of Arts, Heritage and the Gaeltacht,
7 Ely Place, Dublin 2, Ireland.
Web: www.npws.ie
E-mail: nature.conservation@ahg.gov.ie**

Citation:

**NPWS (201) Conservation Objectives: West Connacht Coast SAC 002998.
Version 1. National Parks and Wildlife Service, Department of Arts, Heritage
and the Gaeltacht.**

**Series Editor: Rebecca Jeffrey
ISSN 2009-4086**

Introduction

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

* indicates a priority habitat under the Habitats Directive

002998	West Connacht Coast SAC
1349	Common Bottlenose Dolphin <i>Tursiops truncatus</i>

Please note that this SAC overlaps with the following SPAs: 004004, 004084, 004111, 004136, 004144, 004170 and adjoins the following SACs: 000278, 000328, 000330, 000470, 000472, 000484, 000495, 000507, 001228, 001309, 001311, 001501, 001529, 001932, 002031, 002074, 002243, 002265. See map 2. The conservation objectives for this site should be used in conjunction with those for the overlapping and adjacent sites as appropriate.

Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

NPWS Documents

Year : 2015

Title : West Connacht Coast SAC (site code: 2998) Conservation objectives supporting document-marine species V1

Author : NPWS

Series : Conservation objectives supporting document

Spatial data sources

Year : 2005

Title : OSi Discovery series vector data

GIS Operations : Low Water Mark (LWM) polyline feature class converted into polygon feature class; clipped to SAC boundary. Expert opinion used as necessary to resolve any issues arising

Used For : 1349 (map 3)

Conservation Objectives for : West Connacht Coast SAC [002998]


1349 Common Bottlenose Dolphin *Tursiops truncatus*


To maintain the favourable conservation condition of Common Bottlenose Dolphin in West Connacht Coast SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Access to suitable habitat	Number of artificial barriers	Species range within the site should not be restricted by artificial barriers to site use. See map 3	See marine supporting document for further details
Disturbance	Level of impact	Human activities should occur at levels that do not adversely affect the bottlenose dolphin population at the site	See marine supporting document for further details



Legend
 West Connacht Coast SAC 002998





Legend

- West Connacht Coast SAC 002998
- 1349 Bottlenose Dolphin - *Tursiops truncatus*
- OSi Discovery Series County Boundary



Conservation objectives for Inishkea Islands SPA [004004]

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Objective: To maintain or restore the favourable conservation condition of the bird species listed as Special Conservation Interests for this SPA:

Bird Code	Common Name	Scientific Name
A018	Shag	<i>Phalacrocorax aristotelis</i>
A045	Barnacle Goose	<i>Branta leucopsis</i>
A137	Ringed Plover	<i>Charadrius hiaticula</i>
A144	Sanderling	<i>Calidris alba</i>
A148	Purple Sandpiper	<i>Calidris maritima</i>
A169	Turnstone	<i>Arenaria interpres</i>
A182	Common Gull	<i>Larus canus</i>
A184	Herring Gull	<i>Larus argentatus</i>



A194	Arctic Tern	<i>Sterna paradisaea</i>
A195	Little Tern	<i>Sterna albifrons</i>
A466	Dunlin	<i>Calidris alpina schinzii</i>

Citation: NPWS (2020) Conservation objectives for Inishkea Islands SPA [004004]. Generic Version 7.0. Department of Culture, Heritage and the Gaeltacht.

National Parks and Wildlife Service

Conservation Objectives Series

Killala Bay/Moy Estuary SPA 004036



*An Roinn
Ealaíon, Oidhreachta agus Gaeltachta*

*Department of
Arts, Heritage and the Gaeltacht*



**National Parks and Wildlife Service,
Department of Arts, Heritage and the Gaeltacht,
7 Ely Place, Dublin 2, Ireland.
Web: www.npws.ie
E-mail: nature.conservation@ahg.gov.ie**

Citation:

**NPWS (2013) Conservation Objectives: Killala Bay/Moy Estuary SPA 004036.
Version 1. National Parks and Wildlife Service, Department of Arts, Heritage
and the Gaeltacht.**

**Series Editor: Rebecca Jeffrey
ISSN 2009-4086**

Introduction

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

* indicates a priority habitat under the Habitats Directive

004036 Killala Bay/Moy Estuary SPA

- A137 Ringed Plover *Charadrius hiaticula*
- A140 Golden Plover *Pluvialis apricaria*
- A141 Grey Plover *Pluvialis squatarola*
- A144 Sanderling *Calidris alba*
- A149 Dunlin *Calidris alpina alpina*
- A157 Bar-tailed Godwit *Limosa lapponica*
- A160 Curlew *Numenius arquata*
- A162 Redshank *Tringa totanus*
- A999 Wetlands

Please note that this SPA overlaps with Killala Bay/Moy Estuary SAC (000458) and Lackan Saltmarsh and Kilcummin Head SAC (000516). See map 2. The conservation objectives for this site should be used in conjunction with those for the overlapping sites as appropriate.

Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

NPWS Documents

Year : 2013
Title : Killala Bay/Moy Estuary SPA (site code 4036) Conservation objectives supporting document V1
Author : NPWS
Series : Conservation objectives supporting document

A137 **Ringed Plover *Charadrius hiaticula***

To maintain the favourable conservation condition of Ringed Plover in Killala Bay/Moy Estuary SPA, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Population trend	Percentage change	Long term population trend stable or increasing	Population trends are presented in part four of the conservation objectives supporting document
Distribution	Number and range of areas used by waterbirds	No significant decrease in the range, timing or intensity of use of areas by ringed plover, other than that occurring from natural patterns of variation	Waterbird distribution from the 2010/2011 waterbird survey programme is discussed in part five of conservation objectives supporting document

A140 Golden Plover *Pluvialis apricaria*

To maintain the favourable conservation condition of Golden Plover in Killala Bay/Moy Estuary SPA, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Population trend	Percentage change	Long term population trend stable or increasing	Population trends are presented in part four of the conservation objectives supporting document
Distribution	Number, range, timing and intensity of use of areas	No significant decrease in the range, timing or intensity of use of areas by golden plover, other than that occurring from natural patterns of variation	Waterbird distribution from the 2010/2011 waterbird survey programme is discussed in part five of the conservation objectives supporting document

A141 Grey Plover *Pluvialis squatarola*

To maintain the favourable conservation condition of Grey Plover in Killala Bay/Moy Estuary SPA, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Population trend	Percentage change	Long term population trend stable or increasing	Population trends are presented in part four of the conservation objectives supporting document
Distribution	Number, range, timing and intensity of use of areas	No significant decrease in the range, timing or intensity of use of areas by grey plover, other than that occurring from natural patterns of variation	Waterbird distribution from the 2010/2011 waterbird survey programme is discussed in part five of the conservation objectives supporting document

A144 Sanderling *Calidris alba*

To maintain the favourable conservation condition of Sanderling in Killala Bay/Moy Estuary SPA, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Population trend	Percentage change	Long term population trend stable or increasing	Waterbird population trends are presented in part four of the conservation objectives supporting document
Distribution	Number, range, timing and intensity of use of areas	No significant decrease in the range, timing or intensity of use of areas by sanderling, other than that occurring from natural patterns of variation	Waterbird distribution from the 2010/2011 waterbird survey programme is discussed in part five of the conservation objectives supporting document

A149 **Dunlin *Calidris alpina alpina***

To maintain the favourable conservation condition of Dunlin in Killala Bay/Moy Estuary SPA, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Population trend	Percentage change	Long term population trend stable or increasing	Population trends are presented in part four of the conservation objectives supporting document
Distribution	Number, range, timing and intensity of use of areas	No significant decrease in the range, timing or intensity of use of areas by dunlin, other than that occurring from natural patterns of variation	Waterbird distribution from the 2010/2011 waterbird survey programme is discussed in part five of the conservation objectives supporting document

A157 Bar-tailed Godwit *Limosa lapponica*

To maintain the favourable conservation condition of Bar-tailed Godwit in Killala Bay/Moy Estuary SPA, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Population trend	Percentage change	Long term population trend stable or increasing	Population trends are presented in part four of the conservation objectives supporting document
Distribution	Number, range, timing and intensity of use of areas	No significant decrease in the range, timing or intensity of use of areas by bar-tailed godwit, other than that occurring from natural patterns of variation	Waterbird distribution from the 2010/2011 waterbird survey programme is discussed in part five of the conservation objectives supporting document

A160 *Curlew Numenius arquata*

To maintain the favourable conservation condition of Curlew in Killala Bay/Moy Estuary SPA, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Population trend	Percentage change	Long term population trend stable or increasing	Population trends are presented in part four of the conservation objectives supporting document
Distribution	Number, range, timing and intensity of use of areas	No significant decrease in the range, timing or intensity of use of areas by curlew, other than that occurring from natural patterns of variation	Waterbird distribution from the 2010/2011 waterbird survey programme is discussed in part five of the conservation objectives supporting document

A162 **Redshank *Tringa totanus***

To maintain the favourable conservation condition of Redshank in Killala Bay/Moy Estuary SPA, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Population trend	Percentage change	Long term population trend stable or increasing	Population trends are presented in part four of the conservation objectives supporting document
Distribution	Number, range, timing and intensity of use of area	No significant decrease in the range, timing or intensity of use of areas by redshank, other than that occurring from natural patterns of variation	Waterbird distribution from the 2010/2011 waterbird survey programme is discussed in part five of the conservation objectives supporting document

A999 Wetlands

To maintain the favourable conservation condition of wetland habitat in Killala Bay/Moy Estuary SPA as a resource for the regularly occurring migratory waterbirds that utilise it. This is defined by the following attribute and target:

Attribute	Measure	Target	Notes
Habitat area	Hectares	The permanent area occupied by the wetland habitat should be stable and not significantly less than the area of 3204 hectares, other than that occurring from natural patterns of variation	The wetland habitat area was estimated as 3204ha using OSi data and relevant orthophotographs. For further information see part three of the conservation objectives supporting document



SPA 004036

SAC 000516

SPA 004036

SAC 000458

Legend

- SPA 004036
- SAC 000458 Killala Bay/Moy Estuary
- SAC 000516 Lackan Saltmarsh And Kilcummin Head
- OSi Discovery Series County Boundaries

National Parks and Wildlife Service

Conservation Objectives Series

Blacksod Bay/Broad Haven SPA 004037



*An Roinn
Ealaíon, Oidhreachta agus Gaeltachta*

*Department of
Arts, Heritage and the Gaeltacht*



**National Parks and Wildlife Service,
Department of Arts, Heritage and the Gaeltacht,**

7 Ely Place, Dublin 2, Ireland.

Web: www.npws.ie

E-mail: nature.conservation@ahg.gov.ie

Citation:

**NPWS (201) Conservation Objectives: Blacksod Bay/Broad Haven SPA 004037.
Version 1. National Parks and Wildlife Service, Department of Arts, Heritage
and the Gaeltacht.**

Series Editor: Rebecca Jeffrey

ISSN 2009-4086

Introduction

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

* indicates a priority habitat under the Habitats Directive

004037 Blacksod Bay/Broad Haven SPA

A003	Great Northern Diver <i>Gavia immer</i>
A046	Brent Goose <i>Branta bernicla hrota</i>
A065	Common Scoter <i>Melanitta nigra</i>
A069	Red-breasted Merganser <i>Mergus serrator</i>
A137	Ringed Plover <i>Charadrius hiaticula</i>
A144	Sanderling <i>Calidris alba</i>
A149	Dunlin <i>Calidris alpina alpina</i>
A157	Bar-tailed Godwit <i>Limosa lapponica</i>
A160	Curlew <i>Numenius arquata</i>
A191	Sandwich Tern <i>Sterna sandvicensis</i>
A466	Dunlin <i>Calidris alpina schinzii</i>
A999	Wetlands

Please note that this SPA overlaps with Mullet/Blacksod Bay Complex SAC (000470), Broadhaven Bay SAC (000472) and Glenamoy Bog Complex SAC (000500). It adjoins Mullet Peninsula SPA (004227). See map 2. The conservation objectives for this site should be used in conjunction with those for overlapping and adjacent sites as appropriate.

Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

NPWS Documents

Year :	2010
Title :	Resurvey of breeding wader populations of machair and associated wet grasslands in north-west Ireland
Author :	Suddaby, D.; Nelson, T.; Veldman, J.
Series :	Irish Wildlife Manual No. 44
<hr/>	
Year :	2014
Title :	Blacksod Bay/Broad Haven SPA (site code: 4037) Conservation objectives supporting document V1
Author :	NPWS
Series :	Conservation objectives supporting document

Other References

Year :	1984
Title :	The 1984 all Ireland tern survey
Author :	Whilde, A.
Series :	Irish Birds 3: 1-32
<hr/>	
Year :	1985
Title :	The birds of the Western Palearctic- Volume IV
Author :	Cramp, S.
Series :	Oxford University Press, Oxford
<hr/>	
Year :	1985
Title :	Breeding waders of sand dune machair in north-west Ireland
Author :	Nairn, R.G.W.; Sheppard, J.R.
Series :	Irish Birds 3: 53-70
<hr/>	
Year :	1995
Title :	Seabird monitoring handbook for Britain and Ireland: a compilation of methods for survey and monitoring of breeding seabirds
Author :	Walsh, P.; Halley, D.J.; Harris, M.P.; del Nevo, A.; Sim, I.M.W.; Tasker, M.L.
Series :	JNCC, Peterborough
<hr/>	
Year :	1997
Title :	The status and distribution of breeding sandwich, roseate, common, arctic and little terns in Ireland in 1995
Author :	Hannon, C.; Berrow, S.D.; Newton, S.F.
Series :	Irish Birds, 6: 1-22
<hr/>	
Year :	1998
Title :	Breeding waders of machair systems in Ireland in 1996
Author :	Madden, B.; Cooney, T.; O'Donoghue, A.; Norriss, D.W.; Merne, O.J.
Series :	Irish Birds 6: 177-191
<hr/>	
Year :	2004
Title :	Seabird Populations of Britain and Ireland
Author :	Mitchell, P.I.; Newton, S.F.; Ratcliffe, N.; Dunn, T.E.
Series :	Poyser, London

Year : 2006
Title : Breeding parameters of selected wader species at machair sites and adjacent areas associated with the Mullet/Blacksod Bay and offshore islands complex, NW Mayo, 2006
Author : Thompson, L.J.; Suddaby, D.; Newton, S.
Series : BirdWatch Ireland Conservation Report No. 06/6

Year : 2014
Title : Seabird Monitoring Programme (SMP) Database
Author : JNCC
Series : <http://jncc.defra.gov.uk/smp/Default.aspx>

Year : 2014
Title : BirdLife International Seabird Ecology and Foraging Range Database
Author : BirdLife International
Series : <http://seabird.wikispaces.com>

Spatial data sources

Year :	2014
Title :	NPWS SPA boundary data
GIS Operations :	SPA boundary polygons divided into two classifications (wetlands, coastal grasslands) based on line identified by expert judgement. Expert opinion used as necessary to resolve any issues arising
Used For :	Wetlands (map 3)

A003 Great Northern Diver *Gavia immer*

To maintain the favourable conservation condition of Great Northern Diver in Blacksod Bay/Broad Haven SPA, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Population trend	Percentage change	Long term population trend stable or increasing	Waterbird population trends are presented in part four of the conservation objectives supporting document
Distribution	Range, timing and intensity of use of areas	No significant decrease in the range, timing or intensity of use of areas by great northern diver, other than that occurring from natural patterns of variation	Waterbird distribution from the 2009/2010 waterbird survey programme is discussed in part five of the conservation objectives supporting document

Conservation Objectives for : Blacksod Bay/Broad Haven SPA [004037]

A046 Brent Goose *Branta bernicla hrota*

To maintain the favourable conservation condition of Light-bellied Brent Goose in Blacksod Bay/Broad Haven SPA, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Population trend	Percentage change	Long term population trend stable or increasing	Waterbird population trends are presented in part four of the conservation objectives supporting document
Distribution	Range, timing and intensity of use of areas	No significant decrease in the range, timing or intensity of use of areas by light-bellied brent goose, other than that occurring from natural patterns of variation	Waterbird distribution from the 2009/2010 waterbird survey programme is discussed in part five of the conservation objectives supporting document

Conservation Objectives for : Blacksod Bay/Broad Haven SPA [004037]

A065 Common Scoter *Melanitta nigra*

To maintain the favourable conservation condition of Common Scoter in Blacksod Bay/Broad Haven SPA, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Population trend	Percentage change	Long term population trend stable or increasing	Waterbird population trends are presented in part four of the conservation objectives supporting document
Distribution	Range, timing and intensity of use of areas	No significant decrease in the range, timing or intensity of use of areas by common scoter, other than that occurring from natural patterns of variation	Waterbird distribution from the 2009/2010 waterbird survey programme is discussed in part five of the conservation objectives supporting document

Conservation Objectives for : Blacksod Bay/Broad Haven SPA [004037]

A069 Red-breasted Merganser *Mergus serrator*

To maintain the favourable conservation condition of Red-breasted Merganser in Blacksod Bay/Broad Haven SPA, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Population trend	Percentage change	Long term population trend stable or increasing	Waterbird population trends are presented in part four of the conservation objectives supporting document
Distribution	Range, timing and intensity of use of areas	No significant decrease in the range, timing or intensity of use of areas by red-breasted merganser, other than that occurring from natural patterns of variation	Waterbird distribution from the 2009/2010 waterbird survey programme is discussed in part five of the conservation objectives supporting document

Conservation Objectives for : Blacksod Bay/Broad Haven SPA [004037]

A137 Ringed Plover *Charadrius hiaticula*

To maintain the favourable conservation condition of Ringed Plover in Blacksod Bay/Broad Haven SPA, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Population trend	Percentage change	Long term population trend stable or increasing	Population trends are presented in part four of the conservation objectives supporting document
Distribution	Range, timing and intensity of use of areas	No significant decrease in the range, timing or intensity of use of areas by ringed plover, other than that occurring from natural patterns of variation	Waterbird distribution from the 2009/2010 waterbird survey programme is discussed in part five of conservation objectives supporting document

Conservation Objectives for : Blacksod Bay/Broad Haven SPA [004037]

A144 Sanderling *Calidris alba*

To maintain the favourable conservation condition of Sanderling in Blacksod Bay/Broad Haven SPA, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Population trend	Percentage change	Long term population trend stable or increasing	Waterbird population trends are presented in part four of the conservation objectives supporting document
Distribution	Range, timing and intensity of use of areas	No significant decrease in the range, timing or intensity of use of areas by sanderling, other than that occurring from natural patterns of variation	Waterbird distribution from the 2009/2010 waterbird survey programme is discussed in part five of the conservation objectives supporting document

A149 **Dunlin *Calidris alpina alpina***

To maintain the favourable conservation condition of Dunlin in Blacksod Bay/Broad Haven SPA, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Population trend	Percentage change	Long term population trend stable or increasing	Population trends are presented in part four of the conservation objectives supporting document
Distribution	Range, timing and intensity of use of areas	No significant decrease in the range, timing or intensity of use of areas by dunlin, other than that occurring from natural patterns of variation	Waterbird distribution from the 2009/2010 waterbird survey programme is discussed in part five of the conservation objectives supporting document

A157 Bar-tailed Godwit *Limosa lapponica*

To maintain the favourable conservation condition of Bar-tailed Godwit in Blacksod Bay/Broad Haven SPA, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Population trend	Percentage change	Long term population trend stable or increasing	Population trends are presented in part four of the conservation objectives supporting document
Distribution	Range, timing and intensity of use of areas	No significant decrease in the range, timing or intensity of use of areas by bar-tailed godwit, other than that occurring from natural patterns of variation	Waterbird distribution from the 2009/2010 waterbird survey programme is discussed in part five of the conservation objectives supporting document

A160 ***Curlew *Numenius arquata****

To maintain the favourable conservation condition of Curlew in Blacksod Bay/Broad Haven SPA, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Population trend	Percentage change	Long term population trend stable or increasing	Population trends are presented in part four of the conservation objectives supporting document
Distribution	Range, timing and intensity of use of areas	No significant decrease in the range, timing or intensity of use of areas by curlew, other than that occurring from natural patterns of variation	Waterbird distribution from the 2009/2010 waterbird survey programme is discussed in part five of the conservation objectives supporting document

Conservation Objectives for : Blacksod Bay/Broad Haven SPA [004037]

A191 Sandwich Tern *Sterna sandvicensis*

To maintain the favourable conservation condition of Sandwich Tern in Blacksod Bay/Broad Haven SPA, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Breeding population abundance: apparently occupied nests (AONs)	Number	No significant decline	Measure based on standard tern survey methods (see Walsh et al., 1995). Hannon et al. (1997) recorded 81 breeding pairs on Inishderry as part of the 1995 All-Ireland Tern Survey. Recent data is lacking for this colony
Productivity rate: fledged young per breeding pair	Mean number	No significant decline	Measure based on standard tern survey methods (see Walsh et al., 1995). The Seabird Monitoring Programme (SMP) online database (JNCC, 2014) provides population data for this species
Distribution: breeding colonies	Number; location; area (Hectares)	No significant decline	Typical sandwich tern breeding sites are located on low-lying offshore islands or islets in bays or brackish lagoons on spits or remote mainland dunes (Cramp, 1985). Wide fluctuations between years in both breeding numbers and colony locations are known to occur for this species (Mitchell et al., 2004). However, a sandwich tern colony has been recorded on Inishderry in the 1990s and on several occasions in the 1980s (see Hannon et al., 1997 and Whilde, 1985)
Prey biomass available	Kilogrammes	No significant decline	Key prey items: Mostly energy-rich fish, some crustaceans and occasionally insects and rag worms. Key habitats: sandwich tern forage in/over shallow marine waters such as bays, inlets and outflows, gullies, shoals, inshore waters, reefs, and sandbanks; also more open waters nearshore and offshore, including open sea. Foraging range: max. 70km, mean max. 42.3km, mean 14.7km (BirdLife International Seabird Database (Birdlife International, 2014))
Barriers to connectivity	Number; location; shape; area (hectares)	No significant increase	Foraging range: Max 70km, mean max 42.3km, mean 14.7km (Birdlife International Seabird Database (Birdlife International, 2014))
Disturbance at the breeding stage	Level of impact	Human activities should occur at levels that do not adversely affect the breeding sandwich tern population	Colonies are typically situated on low-lying offshore islands or islets, in bays or brackish lagoons, on spits or remote mainland dunes (Cramp, 1985). The sandwich tern colony on Inishderry has been recorded in the 1990s and on several occasions in the 1980s (see Hannon et al., 1997 and Whilde, 1984)

Conservation Objectives for : Blacksod Bay/Broad Haven SPA [004037]

A466 Dunlin *Calidris alpina schinzii*



To maintain the favourable conservation condition of Dunlin in Blacksod Bay/Broad Haven SPA, which is defined by the following list of attributes and targets

Attribute	Measure	Target	Notes
Breeding population abundance: apparently occupied territories (AOTs)	Number	Stable or increasing, subject to natural variation	Measure based on standard survey methods (see Suddaby et al. (2010))
Productivity rate: fledged young per breeding pair	Mean number	No significant decline	Measure based on standard survey methods (see Thompson et al. (2007))
Distribution	Number; location	Stable or increasing, subject to natural variation	The distribution of breeding dunlin has contracted since initial surveys were undertaken (Nairn and Shephard, 1985; Madden et al., 1998; Suddaby et al., 2010)
Availability of suitable habitat: area and distribution	Hectares; location	Stable or increasing, subject to natural processes	Ideally, suitable habitat should be at, or close to, existing breeding pairs. Factors that are negatively affecting potentially suitable habitat include fencing, drainage, inappropriate grazing regimes, fertilisation and overgrazing by rabbits (<i>Oryctolagus cuniculus</i>)
Disturbance at the breeding site	Level of impact	Human activities should occur at levels that do not adversely affect the breeding dunlin population	Unsuitable livestock grazing regimes can result in nest trampling and destruction of suitable nesting sites. Agri-environment schemes in Ireland specify less than 1.0 livestock units per hectare during the breeding wader nesting period


A999 Wetlands


To maintain the favourable conservation condition of the wetland habitat in Blacksod Bay/Broad Haven SPA as a resource for the regularly-occurring migratory waterbirds that utilise it. This is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	The permanent area occupied by the wetland habitat should be stable and not significantly less than the area of 8,539 hectares, other than that occurring from natural patterns of variation. See map 3	The wetland habitat area was estimated as 8,539ha using OSi data and relevant orthophotographs. For further information see part three of the conservation objectives supporting document




Legend

 Blacksod Bay / Broad Haven SPA 004037



Legend

- Blacksod Bay / Broad Haven SPA 004037
- Mullet Peninsula SPA 004227
- Mullet/Blacksod Bay Complex SAC 000470
- Broadhaven Bay SAC 000472
- Glenamoy Bog Complex SAC 000500
- OSi Discovery Series County Boundary



Legend

- Blacksod Bay / Broad Haven SPA 004037
- OSi Discovery Series County Boundary

Wetlands and Waterbirds

- Coastal Grasslands
- Wetlands



Conservation objectives for Lough Corrib SPA [004042]

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Objective: To maintain or restore the favourable conservation condition of the bird species listed as Special Conservation Interests for this SPA:

Bird Code	Common Name	Scientific Name
A051	Gadwall	<i>Anas strepera</i>
A056	Shoveler	<i>Anas clypeata</i>
A059	Pochard	<i>Aythya ferina</i>
A061	Tufted Duck	<i>Aythya fuligula</i>
A065	Common Scoter	<i>Melanitta nigra</i>
A082	Hen Harrier	<i>Circus cyaneus</i>
A125	Coot	<i>Fulica atra</i>
A140	Golden Plover	<i>Pluvialis apricaria</i>



A179	Black-headed Gull	<i>Chroicocephalus ridibundus</i>
A182	Common Gull	<i>Larus canus</i>
A193	Common Tern	<i>Sterna hirundo</i>
A194	Arctic Tern	<i>Sterna paradisaea</i>
A395	Greenland White-fronted Goose	<i>Anser albifrons flavirostris</i>

To acknowledge the importance of Ireland's wetlands to wintering waterbirds, "Wetland and Waterbirds" may be included as a Special Conservation Interest for some SPAs that have been designated for wintering waterbirds and that contain a wetland site of significant importance to one or more of the species of Special Conservation Interest. Thus, a second objective is included as follows:

Objective: To maintain or restore the favourable conservation condition of the wetland habitat at Lough Corrib SPA as a resource for the regularly-occurring migratory waterbirds that utilise it.

Citation: NPWS (2020) Conservation objectives for Lough Corrib SPA [004042]. Generic Version 7.0.
Department of Culture, Heritage and the Gaeltacht.



Conservation objectives for Lough Gara SPA [004048]

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Objective: To maintain or restore the favourable conservation condition of the bird species listed as Special Conservation Interests for this SPA:

Bird Code	Common Name	Scientific Name
A038	Whooper Swan	<i>Cygnus cygnus</i>
A395	Greenland White-fronted Goose	<i>Anser albifrons flavirostris</i>



Citation: NPWS (2020) Conservation objectives for Lough Gara SPA [004048]. Generic Version 7.0.
Department of Culture, Heritage and the Gaeltacht.



Conservation objectives for Lough Carra SPA [004051]

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Objective: To maintain or restore the favourable conservation condition of the bird species listed as Special Conservation Interests for this SPA:

Bird Code	Common Name	Scientific Name
A182	Common Gull	<i>Larus canus</i>



Citation: NPWS (2020) Conservation objectives for Lough Carra SPA [004051]. Generic Version 7.0.
Department of Culture, Heritage and the Gaeltacht.



Conservation objectives for Carrowmore Lake SPA [004052]

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Objective: To maintain or restore the favourable conservation condition of the bird species listed as Special Conservation Interests for this SPA:

Bird Code	Common Name	Scientific Name
A191	Sandwich Tern	<i>Sterna sandvicensis</i>



Citation: NPWS (2020) Conservation objectives for Carrowmore Lake SPA [004052]. Generic Version 7.0. Department of Culture, Heritage and the Gaeltacht.



Conservation objectives for Lough Mask SPA [004062]

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Objective: To maintain or restore the favourable conservation condition of the bird species listed as Special Conservation Interests for this SPA:

Bird Code	Common Name	Scientific Name
A061	Tufted Duck	<i>Aythya fuligula</i>
A179	Black-headed Gull	<i>Chroicocephalus ridibundus</i>
A182	Common Gull	<i>Larus canus</i>
A183	Lesser Black-backed Gull	<i>Larus fuscus</i>
A193	Common Tern	<i>Sterna hirundo</i>
A395	Greenland White-fronted Goose	<i>Anser albifrons flavirostris</i>



To acknowledge the importance of Ireland's wetlands to wintering waterbirds, “Wetland and Waterbirds” may be included as a Special Conservation Interest for some SPAs that have been designated for wintering waterbirds and that contain a wetland site of significant importance to one or more of the species of Special Conservation Interest. Thus, a second objective is included as follows:

Objective: To maintain or restore the favourable conservation condition of the wetland habitat at Lough Mask SPA as a resource for the regularly-occurring migratory waterbirds that utilise it.

Citation: NPWS (2020) *Conservation objectives for Lough Mask SPA [004062]. Generic Version 7.0.*
Department of Culture, Heritage and the Gaeltacht.



Conservation objectives for Stags of Broad Haven SPA [004072]

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Objective: To maintain or restore the favourable conservation condition of the bird species listed as Special Conservation Interests for this SPA:

Bird Code	Common Name	Scientific Name
A014	Storm Petrel	<i>Hydrobates pelagicus</i>
A015	Leach's Storm-petrel	<i>Oceanodroma leucorhoa</i>



Citation: NPWS (2020) Conservation objectives for Stags of Broad Haven SPA [004072]. Generic Version 7.0. Department of Culture, Heritage and the Gaeltacht.



Conservation objectives for Illanmaster SPA [004074]

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Objective: To maintain or restore the favourable conservation condition of the bird species listed as Special Conservation Interests for this SPA:

Bird Code	Common Name	Scientific Name
A014	Storm Petrel	<i>Hydrobates pelagicus</i>



Citation: NPWS (2020) Conservation objectives for Illanmaster SPA [004074]. Generic Version 7.0.
Department of Culture, Heritage and the Gaeltacht.



Conservation objectives for Inishglora and Inishkeeragh SPA [004084]

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Objective: To maintain or restore the favourable conservation condition of the bird species listed as Special Conservation Interests for this SPA:

Bird Code	Common Name	Scientific Name
A014	Storm Petrel	<i>Hydrobates pelagicus</i>
A017	Cormorant	<i>Phalacrocorax carbo</i>
A018	Shag	<i>Phalacrocorax aristotelis</i>
A045	Barnacle Goose	<i>Branta leucopsis</i>
A183	Lesser Black-backed Gull	<i>Larus fuscus</i>
A184	Herring Gull	<i>Larus argentatus</i>
A194	Arctic Tern	<i>Sterna paradisaea</i>



Citation: NPWS (2020) Conservation objectives for Inishglora and Inishkeeragh SPA [004084]. Generic Version 7.0. Department of Culture, Heritage and the Gaeltacht.



Conservation objectives for Termoncarragh Lake and Annagh Machair SPA [004093]

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Objective: To maintain or restore the favourable conservation condition of the bird species listed as Special Conservation Interests for this SPA:

Bird Code	Common Name	Scientific Name
A038	Whooper Swan	<i>Cygnus cygnus</i>
A045	Barnacle Goose	<i>Branta leucopsis</i>
A122	Corncrake	<i>Crex crex</i>
A142	Lapwing	<i>Vanellus vanellus</i>
A346	Chough	<i>Pyrrhocorax pyrrhocorax</i>
A395	Greenland White-fronted Goose	<i>Anser albifrons flavirostris</i>
A466	Dunlin	<i>Calidris alpina schinzii</i>

For more information please go to: www.npws.ie/protected-sites/conservation-management-planning



To acknowledge the importance of Ireland's wetlands to wintering waterbirds, “Wetland and Waterbirds” may be included as a Special Conservation Interest for some SPAs that have been designated for wintering waterbirds and that contain a wetland site of significant importance to one or more of the species of Special Conservation Interest. Thus, a second objective is included as follows:

Objective: To maintain or restore the favourable conservation condition of the wetland habitat at Termoncarragh Lake and Annagh Machair SPA as a resource for the regularly-occurring migratory waterbirds that utilise it.

Citation: NPWS (2020) *Conservation objectives for Termoncarragh Lake and Annagh Machair SPA [004093]. Generic Version 7.0. Department of Culture, Heritage and the Gaeltacht.*



Conservation objectives for Owenduff/Nephin Complex SPA [004098]

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Objective: To maintain or restore the favourable conservation condition of the bird species listed as Special Conservation Interests for this SPA:

Bird Code	Common Name	Scientific Name
A098	Merlin	<i>Falco columbarius</i>
A140	Golden Plover	<i>Pluvialis apricaria</i>



Citation: NPWS (2020) Conservation objectives for Owenduff/Nephin Complex SPA [004098]. Generic Version 7.0. Department of Culture, Heritage and the Gaeltacht.



Conservation objectives for Bellanagare Bog SPA [004105]

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Objective: To maintain or restore the favourable conservation condition of the bird species listed as Special Conservation Interests for this SPA:

Bird Code	Common Name	Scientific Name
A395	Greenland White-fronted Goose	<i>Anser albifrons flavirostris</i>



Citation: NPWS (2020) *Conservation objectives for Bellanagare Bog SPA [004105]. Generic Version 7.0. Department of Culture, Heritage and the Gaeltacht.*



Conservation objectives for Duvillaun Islands SPA [004111]

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Objective: To maintain or restore the favourable conservation condition of the bird species listed as Special Conservation Interests for this SPA:

Bird Code	Common Name	Scientific Name
A009	Fulmar	<i>Fulmarus glacialis</i>
A014	Storm Petrel	<i>Hydrobates pelagicus</i>
A045	Barnacle Goose	<i>Branta leucopsis</i>



Citation: NPWS (2020) *Conservation objectives for Duvillaun Islands SPA [004111]. Generic Version 7.0.* Department of Culture, Heritage and the Gaeltacht.



Conservation objectives for Clare Island SPA [004136]

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Objective: To maintain or restore the favourable conservation condition of the bird species listed as Special Conservation Interests for this SPA:

Bird Code	Common Name	Scientific Name
A009	Fulmar	<i>Fulmarus glacialis</i>
A018	Shag	<i>Phalacrocorax aristotelis</i>
A182	Common Gull	<i>Larus canus</i>
A188	Kittiwake	<i>Rissa tridactyla</i>
A199	Guillemot	<i>Uria aalge</i>
A200	Razorbill	<i>Alca torda</i>
A346	Chough	<i>Pyrrhocorax pyrrhocorax</i>



Citation: NPWS (2020) Conservation objectives for Clare Island SPA [004136]. Generic Version 7.0.
Department of Culture, Heritage and the Gaeltacht.



Conservation objectives for High Island, Inishshark and Davillaun SPA [004144]

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Objective: To maintain or restore the favourable conservation condition of the bird species listed as Special Conservation Interests for this SPA:

Bird Code	Common Name	Scientific Name
A009	Fulmar	<i>Fulmarus glacialis</i>
A045	Barnacle Goose	<i>Branta leucopsis</i>
A194	Arctic Tern	<i>Sterna paradisaea</i>



Citation: NPWS (2020) Conservation objectives for High Island, Inishshark and Davillaun SPA [004144]. Generic Version 7.0. Department of Culture, Heritage and the Gaeltacht.



Conservation objectives for Bills Rocks SPA [004177]

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Objective: To maintain or restore the favourable conservation condition of the bird species listed as Special Conservation Interests for this SPA:

Bird Code	Common Name	Scientific Name
A014	Storm Petrel	<i>Hydrobates pelagicus</i>
A204	Puffin	<i>Fratercula arctica</i>



Citation: NPWS (2020) Conservation objectives for Bills Rocks SPA [004177]. Generic Version 7.0.
Department of Culture, Heritage and the Gaeltacht.



Conservation objectives for Connemara Bog Complex SPA [004181]

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Objective: To maintain or restore the favourable conservation condition of the bird species listed as Special Conservation Interests for this SPA:

Bird Code	Common Name	Scientific Name
A017	Cormorant	<i>Phalacrocorax carbo</i>
A098	Merlin	<i>Falco columbarius</i>
A140	Golden Plover	<i>Pluvialis apricaria</i>
A182	Common Gull	<i>Larus canus</i>



Citation: NPWS (2020) Conservation objectives for Connemara Bog Complex SPA [004181]. Generic Version 7.0. Department of Culture, Heritage and the Gaeltacht.